

# The AME2016 atomic mass evaluation <sup>\*</sup>

## (I). Evaluation of input data; and adjustment procedures

W.J. Huang (黄文嘉)<sup>1</sup> G. Audi (欧乔治)<sup>1</sup> Meng Wang (王猛)<sup>2,3,1</sup> F.G. Kondev<sup>4</sup> S. Naimi<sup>5</sup> Xing Xu (徐星)<sup>2</sup>

<sup>1</sup> CSNSM, Univ Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, 91405 Orsay, France

<sup>2</sup> Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, People's Republic of China

<sup>3</sup> Joint Department for Nuclear Physics, Institute of Modern Physics, CAS and Lanzhou University, Lanzhou 730000, China

<sup>4</sup> Argonne National Laboratory, Argonne, IL 60439, USA

<sup>5</sup> RIKEN Nishina Center, Wako, Saitama 351-0198, Japan

**Abstract** This paper is the first of two articles (Part I and Part II) that presents the results of the new atomic mass evaluation, AME2016. It includes complete information on the experimental input data (also including unused and rejected ones), as well as details on the evaluation procedures used to derive the tables of recommended values given in the second part. This article describes the evaluation philosophy and procedures that were implemented in the selection of specific nuclear reaction, decay and mass-spectrometric results. These input values were entered in the least-squares adjustment for determining the best values for the atomic masses and their uncertainties. Details of the calculation and particularities of the AME are then described. All accepted and rejected data, including outweighed ones, are presented in a tabular format and compared with the adjusted values obtained using the least-squares fit analysis. Differences with the previous AME2012 evaluation are discussed and specific information is presented for several cases that may be of interest to AME users. The second AME2016 article gives a table with the recommended values of atomic masses, as well as tables and graphs of derived quantities, along with the list of references used in both the AME2016 and the NUBASE2016 evaluations (the first paper in this issue).

AMDC: <http://amdc.impcas.ac.cn/>

**Keywords:** AME, trends from the mass surface (TMS), atomic-mass tables, least-squares adjustment

**PACS:** 21.10.D-r      **DOI:** 10.1088/1674-1137/41/3/030002

## 1 Introduction

The mass of an atom equals the sum of the masses of its constituents (protons, neutrons and electrons) minus the binding energy, which includes both the atomic and nuclear binding energies. In general, the binding energy of the electrons outside the nucleus is well known. Therefore, the atomic mass reflects the net consequence of all interactions that hold the nucleons together in the nucleus. Since the strong, weak and electromagnetic interactions act among the nucleons, which on the one hand makes the theoretical description of nuclei very complex, on the other hand it gives us natural laboratories to study these fundamental interactions.

As a fundamental property of nuclei, atomic masses are widely used in many domains of science and engineering. A reliable atomic mass table derived from the experimental data, where the atomic masses and the relevant experimental information can be found conveniently, is

in high demand by the research community. To meet the demands, the Atomic Mass Evaluation (AME) was created in 1950's and now serves the research community by providing the most reliable and comprehensive information related to the atomic masses [1].

The last complete evaluation of experimental atomic mass data AME2012 [2, 3] was published in 2012. Since then the experimental knowledge of atomic masses has continuously expanded and a large amount of data relevant to atomic masses has been published in the scientific literature. In this article, general aspects of the development of AME2016 are presented and discussed. In doing this, several local analyses will be given as illustrative examples.

The main AME2016 evaluation table (Table I) is presented in Part I. All accepted and rejected experimental data are given and compared with the adjusted values deduced using a least-squares fit analysis.

As in the previous AME versions, all uncertainties are

Received 10 March 2017

\* This work has been undertaken with the endorsement of the IUPAP Commission on Symbols, Units, Nomenclature, Atomic Masses and Fundamental Constants (SUNAMCO).

1) E-mail: wangm@impcas.ac.cn

©2017 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

one-standard deviation ( $1\sigma$ ).

There is no strict literature cut-off date for the data used in the present AME2016 evaluation: all data, available to the authors until the end of October 2016 were included. Results which were not included for particular reasons, such as the need for a heavy revision of the evaluation at too late a stage, were added in remarks to the relevant data. The final mass-adjustment calculations were performed on February 28, 2017.

The present publication includes updated information presented in the previous AME including the data that were not used in the final adjustment due to specific reasons, e.g. data that have too large uncertainties.

*Remark:* in the following text, several data of general interest will be discussed. Mention of references that can be found in Table I will be avoided. When it is necessary to provide a specific reference, those will be given using the NSR key-numbers [4], listed at the end of Part II, under “References used in the AME2016 and the NUBASE2016 evaluations” (p. 030003-5) (e.g. [2016De15]).

### 1.1 Isomers in the AME and the emergence of NUBASE

In the early AME work, a computer data file (called *Mfile*) that contained the approximate mass values for nuclides in their ground and selected excited isomeric states was maintained. It was used as an approximate input to the adjustment program, which essentially uses the differences between the input and these approximate values in order to improve the precision of the calculations. The other reason for the existence of this file was for isomers, where one has to be careful to identify which state is involved in the reported experimental data, including decay/reaction energies or mass-spectrometric results. Therefore, it was judged necessary to make the *Mfile* as complete as possible. Thus, the NUBASE evaluation was developed to contain values of the main nuclear properties, such as masses, excitation energies of isomers, half-lives, spin and parities, and decay modes and their intensities, for all known nuclides in their ground and excited isomeric states. The NUBASE evaluation was published independently from AME for the first time in 1997 [5], and was greeted with interest from many colleagues working in the areas of nuclear structure physics, nuclear astrophysics and applied nuclear physics. In 2003 and 2012, the NUBASE and AME were published jointly, which is the case again with the present NUBASE2016 evaluation published in the first part of this issue (41-030001).

Isomers may be involved in mass-measurement experiments. This became even more important for new mass-spectrometric methods that were developed to measure masses of exotic nuclides far from the valley of  $\beta$ -stability,

which have, in general, relatively short lifetimes. The resolving power of mass spectrometers has been improved significantly in recent years and many isomers can be clearly separated. But quite often only an isomeric mixture could be measured and an average mass value for a particular isomeric pair can be obtained. Since the mass of the ground state is the primary aim of the present evaluation, it can be derived only in cases where information on the excitation energies and production rates of the isomers is available. When the excitation energy of a particular isomer is not experimentally known, it is estimated from smooth trends in neighboring nuclides (TNN), as explained in NUBASE2016 (p. 030001-4). Two examples are given in Section 6.2.9 (p. 030002-27) and Section 6.8 (p. 030002-32).

### 1.2 Highlights

**The backbone** Nowadays, the highest precision values measured for the atomic masses are obtained by two different experimental techniques: direct mass-spectrometric measurements using Penning traps, and  $\gamma$ -ray energy measurements following neutron capture reactions.

In the present work, results obtained by both methods are combined consistently (with very few exceptions) to improve considerably the precision of the atomic masses for nuclides along the line of stability in a diagram of the atomic number  $Z$  versus neutron number  $N$  [6], thus resulting in a reliable ‘backbone’.

The highest relative mass precision  $\delta m/m$  of  $7 \times 10^{-12}$  has been achieved by a Penning trap spectrometer [2004Ra33]. The masses of some stable alkali-metal nuclides and noble-gas nuclides [2010Mo30] have been determined with relative precision of  $10^{-10}$  or even better, providing reliable reference standards for other mass measurements. While most stable nuclides, and some long-lived ones, could have their mass accuracy improved using Penning traps, the priority has been given by experimentalists to cases where there is a strong motivation from the physics point of view. For example, the  $Q_{\beta\beta}$  values for nuclides relevant to neutrino properties have been determined with very high precision [2011Go23], strengthening at the same time the backbone.

Meanwhile,  $(n,\gamma)$  reactions ([2006De21], [1984Ke15] and [1998Wh01]) determined the binding energies with a relative uncertainty of  $10^{-7}$ , providing mass determination with precisions at the level of  $10^{-10}$ .

**Exotic species** The domain of nuclides with experimentally known masses has extended impressively over the last few years, thanks to the developments of radioactive nuclear beam facilities and novel mass spectrometers. In the past, masses of short-lived nuclides were mainly known from  $Q_{\beta}$  end-point measurements, while

in the present evaluation, mass spectrometry dominates. Classical time-of-flight mass spectrometry stays at the frontier, exploring the light neutron-rich mass region, albeit with larger uncertainties. Penning traps and storage rings keep on playing an important role in mass measurements for short-lived nuclides. Meanwhile, Multi-

Reflection Time-of-Flight spectrometers (MR-TOF) begin to take the stage. It can be concluded that the shape of the atomic mass surface, and hence understanding of nuclear interactions, has been changed significantly over the last 10-20 years.

Table A. Constants used in this work or resulting from the present evaluation.

1 u	=	$M(^{12}\text{C})/12$	=	atomic mass unit				
1 u	=	1 660 539.040	±	0.020	$\times 10^{-33}$ kg	12	ppb	<i>a</i>
1 u	=	931 494.0954	±	0.0057	keV	6.2	ppb	<i>a</i>
1 u	=	931 494.0038	±	0.0004	keV <sub>90</sub>	0.45	ppb	<i>b</i>
1 eV <sub>90</sub>	=	1 000 000.0983	±	0.0061	μeV	6.1	ppb	<i>a</i>
1 MeV	=	1 073 544.1105	±	0.0066	nu	6.2	ppb	<i>a</i>
1 MeV <sub>90</sub>	=	1 073 544.2160	±	0.0004	nu	0.45	ppb	<i>b</i>
$M_e$	=	548 579.909070	±	0.000016	nu	0.03	ppb	<i>a</i>
	=	510 998.9461	±	0.0031	eV	6.2	ppb	<i>a</i>
	=	510 998.89651	±	0.0023	eV <sub>90</sub>	0.45	ppb	<i>b</i>
$M_p$	=	1 007 276 466.93	±	0.09	nu	0.09	ppb	<i>c</i>
$M_\alpha$	=	4 001 506 179.127	±	0.06	nu	0.015	ppb	<i>c</i>
$M_n - M_H$	=	839 883.59	±	0.51	nu	610	ppb	<i>d</i>
	=	782 346.52	±	0.48	eV <sub>90</sub>	610	ppb	<i>d</i>

- a) derived from the work of Mohr and Taylor [7].
- b) for the definition of V<sub>90</sub>, see text.
- c) derived from this work combined with  $M_e$  and total ionization energies for <sup>1</sup>H and <sup>4</sup>He from [7].
- d) this work.

## 2 Units and recalibrations of α- and γ-ray energies

Atomic mass determination for a particular nuclide can be generally performed by establishing an energy relation between the mass we want to deduce and that for a well known nuclide. This energy relation is then expressed in electron-volts (eV). Mass values can also be obtained as an inertial mass from the movement of an ionized atom in an electro-magnetic field. The mass is then derived from a ratio of masses and it is then expressed in ‘unified atomic mass unit’ (u). Those two units are used in the present work.

Since 1960, the mass unit is defined as one twelfth of the mass of one free atom of carbon-12 in its atomic and nuclear ground states,  $1\text{ u} = M(^{12}\text{C})/12$ . Before 1960, two mass units were used: the physics one, defined as <sup>16</sup>O/16, and the chemical one which considered one sixteenth of the average mass of a standard mixture of the three stable oxygen isotopes. This difference was considered as being not at all negligible, when taking into

account the commercial value of all concerned chemical substances. Physicists could not convince the chemists to drop their unit. “The change would mean millions of dollars in the sale of all chemical substances”, said the chemists, which was indeed true! Kohman, Mat- tauch and Wapstra [8] then calculated that, if <sup>12</sup>C/12 was chosen, the change would be ten times smaller for chemists, and in the opposite direction ... This led to an unification; ‘u’ stands therefore, officially, for ‘unified mass unit’. It is worth mentioning that the chemical mass-spectrometry community (e.g. bio-chemistry, polymer chemistry) widely use the dalton unit (symbol Da, named after John Dalton [9]). It allows to express the number of nucleons in a molecule, at least as it is presently used in these domains. It is thus not strictly the same as ‘u’.

The unit for energy is the electron-volt. The choice of the volt for the energy unit is not unambiguous. For example, one may use the *international* volt V, or the volt V<sub>90</sub> as *maintained* in national metrology laboratories and defined by adopting an exact value for the

constant ( $2e/h$ ) in the relation between frequency and voltage from the Josephson effect. Since 1990, by definition  $2e/h = 483597.9$  (exact) GHz/V<sub>90</sub> (see Table B). Already in 1983, an analysis by Cohen and Wapstra [10] showed that all precision measurements of reaction and decay energies were calibrated in such a way that they can be more accurately expressed in *maintained* volts. In fact, the gamma-ray energies determined in wavelength measurements can be expressed in eV<sub>90</sub> without loss in precision, since the conversion coefficient is an exact quantity. Here we take the measurement of the reaction energy for  ${}^1\text{H}(n,\gamma){}^2\text{H}$  as an example. In the experiment, the wavelength of the emitted  $\gamma$  ray is determined by using the Institut Laue-Langevin (ILL) silicon crystal spectrometer. In AME2003, the recom-

mended value was 2224.5660(4) keV<sub>90</sub>, based on the work of Kessler *et al* [1999Ke05]. This result had the highest precision for energy measurement in the input data, with a relative uncertainty of 180 ppb. In the later work from the same group [2006De21], the value was corrected to be 2224.55610(44) keV with new evaluation on the lattice spacing of the crystal. The value of the crystal lattice spacing is used as an adjusted parameter in the CODATA evaluation of Mohr *et al.*, but not expressed explicitly. Using the same value of the wave length in [2006De21], and the length-energy conversion coefficient, we derive 2224.55600(44) keV<sub>90</sub> as an input to our evaluation. During this period, the conversion coefficient with respect to the *international* volt has been changed by  $5.5 \times 10^{-8}$ , which is about one third of the measurement uncertainty.

Table B. Definition of Volt unit, and resulting mass-energy conversion constants.

$2e/h$				u		
1983	483594.21	(1.34)	GHz/V	931501.2	(2.6)	keV
1983	483594	(exact)	GHz/V <sub>86</sub>	931501.6	(0.3)	keV <sub>86</sub>
1986	483597.67	(0.14)	GHz/V	931494.32	(0.28)	keV
1990	483597.9	(exact)	GHz/V <sub>90</sub>	931493.86	(0.07)	keV <sub>90</sub>
1999	483597.9	(exact)	GHz/V <sub>90</sub>	931494.009	(0.007)	keV <sub>90</sub>
2010	483597.9	(exact)	GHz/V <sub>90</sub>	931494.0023	(0.0007)	keV <sub>90</sub>
2012	483597.9	(exact)	GHz/V <sub>90</sub>	931494.0038	(0.0004)	keV <sub>90</sub>

The precision of the conversion factor between mass units and *maintained* volt (V<sub>90</sub>) is higher than that between the former and *international* volt as seen in Table A. Until the end of the last century, the relative precision of  $M - A$  expressed in keV was for several nuclides worse than the same quantity expressed in mass units. Due to the increase of precision of fundamental constants, now the relative precision of  $M - A$  expressed in keV<sub>90</sub> is as good as the same quantity expressed in mass units, whereas the uncertainties expressed in *international* volts are larger than in V<sub>90</sub>. For example, the mass excess of  ${}^4\text{He}$  is  $2\,603\,254.130 \pm 0.063$  nu in mass units,  $2\,424\,915.609 \pm 0.059$  eV<sub>90</sub> in *maintained* volt units and  $2\,424\,915.851 \pm 0.061$  eV in *international* volt units. Therefore, as already adopted in our previous mass evaluations, the V<sub>90</sub> (*maintained* volt) unit is used in the present work.

In the most recent CODATA evaluation by Mohr *et al.* [7], the relation between *maintained* and *international* volts is given as  $V_{90} = [1 + 9.83(0.61) \times 10^{-8}]V$ , which can be expressed as a difference of 98(6) ppb.

In Table A, the relations between *maintained* and *international* volts, and several constants of interest, obtained from the evaluation of Mohr *et al.* [7] are presented. The ratio of mass units to electronvolts for the two Volt units, and the ratio of the two Volts are also given. In addition, values for the masses of the proton, neutron and  $\alpha$  particle, as derived from the present evaluation, are also given, together with the mass difference between the neutron and the light hydrogen atom.

In the earlier mass tables (e.g. AME1993), we used to give values for the binding energies,  $ZM_H + NM_n - M$ . The main reason for this was that the uncertainty of this quantity (in keV<sub>90</sub>) was larger than that of the mass excess,  $M - A$ . However, due to the increased precision of the neutron mass, this is no longer important. Similarly to AME2003 and AME2012, we now give instead the binding energy per nucleon for educational reasons, connected to the Aston Curve and the maximum stability around the ‘iron-peak’ which is of importance in astrophysics. (See also the note in Part II, Section 2, p. 030003-3)

The defining values and the resulting mass-energy conversion factors are given in Table B. Since 2003, the definition has not been modified. Therefore, no recalibration has been necessary in the present AME2016.

Some more historical points are worth mentioning.

In 1986, Taylor and Cohen [11] showed that the empirical ratio between the two types of volts, which had of course been selected to be nearly equal to 1, had changed by as much as 7 ppm. For this reason, in 1990 a new value was chosen [12] to define the *maintained* volt  $V_{90}$ . In their 1998 evaluation, Mohr and Taylor [13] revised the conversion constant to *international* eV. The result was a slightly higher (and 10 times more precise) value for  $V_{90}$ .

Since older high-precision, reaction-energy measurements were essentially expressed in  $\text{keV}_{86}$ , we had to take into account the difference in voltage definition that causes a systematic error of 8 ppm. It was therefore necessary, for the AME2003 tables, to adjust the older precise data to the new  $\text{keV}_{90}$  standard. For  $\alpha$ -particle energies, Rytz [14] has taken this change into account, when updating his earlier evaluation of  $\alpha$ -particle energies. We have used his values in the present input data table (Table I) and indicated this by adding in the reference field the symbol “Z”.

A considerable number of  $(n,\gamma)$  and  $(p,\gamma)$  reactions have precisions not worse than 8 ppm. In 1990, A.H. Wapstra [15] discussed the need for recalibration for several  $\gamma$  rays that are often used as calibration standards. This work has been updated in AME2003 (in a special file dedicated to this study, available on the AMDC website [16]) to evaluate the influence of new calibrators, as well as of the new Mohr and Taylor fundamental constants for  $\gamma$ -ray and particle energies used in  $(n,\gamma)$ ,  $(p,\gamma)$  and  $(p,n)$  reactions. In doing this, the calibration work of Helmer and van der Leun [17], based on the fundamental constant values at that time, was used. For each of the data concerned, the changes were relatively minor. However, it was necessary to make such recalibrations in AME2003, since otherwise they added up to systematic uncertainties that were non-negligible. The calibration for proton energies has also been undertaken in AME2003. As in the case of Rytz’ recalibrations for  $\alpha$ -decay energies, such data are marked by “Z” behind the reference key-number. However, there were cases where it was not possible to do so, for example when this position was used to indicate that a remark was added, the same “Z” symbol was added to the uncertainty value mentioned in the remark.

The list of input values (Table I) for our calculations includes many excitation energies derived from  $\gamma$ -ray measurements that are evaluated and published in Nuclear Data Sheets (NDS) [18]. Only in exceptional cases it made sense to change them to recalibrated results.

### 3 Input data and their representation - connection diagram

As mentioned above, there are two methods that are used in measurements of atomic masses: the mass-spectrometric one (often called a “direct method”), where the inertial mass is determined from the trajectory of the ion in a magnetic field, or from its time-of-flight; and the so-called “indirect method” where the reaction energy, i.e. the difference between several masses, is determined using a specific nuclear reaction or a decay process. In the present work all available experimental data related to atomic masses (both energy and mass-spectrometric data) are considered. The input data are extracted from the available literature, compiled in an appropriate format and then carefully evaluated.

In the AME data treatment, we try our best to enter the true primary experimental information. In this way, the masses can be recalibrated automatically for any future changes, and the original correlation information can be properly preserved.

One example that illustrates our policy of data treatment is the following. In [1986Ma40], the  $Q$  value of the  $^{148}\text{Gd}(p,t)^{146}\text{Gd}$  reaction was measured relative to that of the  $^{65}\text{Cu}(p,t)^{63}\text{Cu}$  reference reaction. In AME2003, the corresponding equation was  $^{148}\text{Gd}(p,t)^{146}\text{Gd} = -7843 \pm 4 \text{keV}$ . However, in the present work, it is presented and used as a differential reaction equation:  $^{148}\text{Gd}(p,t)^{146}\text{Gd} - ^{65}\text{Cu}(p,t)^{63}\text{Cu} = 1500 \pm 4 \text{keV}$ . Strictly speaking, those equations are not exact either. What is measured in the experiment is the energy spectra of the ejected particles. Since there are differences between the masses of the measured nuclides and the reference, the response of the ejected particles to the  $Q$  values are different for the measured nuclides and the reference, depending also on the angle where the spectra are obtained. While the exact equations are quite complex, we believe that the treatment by differential reaction equation represents the original data more reliably and that most of the primary information is preserved.

Nuclear reaction  $A(a,b)B$  and decay  $A(b)B$  energy measurements connect the initial ( $A$ ) and final ( $B$ ) nuclides with one or two reaction or decay particles. With the exception of some reactions between very light nuclides, the precision with which the masses of reaction particles  $a$  and  $b$  are known is much higher than that of the measured reaction and decay energies. Thus, these reactions and decays can each be represented as a link between two nuclides  $A$  and  $B$ . Differential reaction energies  $A(a,b)B - C(a,b)D$  are in principle represented by a combination of four masses.

Direct mass-spectrometric measurements, again with exception of a few cases between very light nuclides, can be separated in a class of connections between two or




Figure 1. (a)–(j). Diagram of connections for input data.

For *primary data* (those checked by other data):

- absolute mass-doublet nuclide (i.e. connected to  $^{12}\text{C}$ ,  $^{35}\text{Cl}$  or  $^{37}\text{Cl}$ );(or nuclide connected by a unique secondary relative mass-doublet to a remote reference nuclide);
- other primary nuclide;
- ◻ ⊙ primary nuclide with relevant isomer;
- // mass-spectrometric connection;
- other primary reaction connection.

Primary connections are drawn with two different thicknesses. Thicker lines represent the highest precision data in the given mass region

- (limits: 1 keV for  $A < 36$ ,
- 2 keV for  $A = 36$  to 165 and
- 3 keV for  $A > 165$ ).

For *secondary data* (cases where masses are known from one type of data and are therefore not checked by a different connection):

- secondary experimental nuclide determined from mass spectrometry;
- secondary experimental nuclide determined by a reaction or a decay;
- nuclide for which mass is estimated from trends from the Mass Surface (TMS);
- connection to a secondary nuclide. Note that an experimental connection may exist between two estimated TMS nuclides when neither of them is connected to the network of primaries.




Figure 1 (b). Diagram of connections for input data — continued.

three nuclides, and a class that essentially determines the absolute mass value (see Section 5). Penning trap measurements almost always give a ratio of masses between two nuclides (inversely proportional to their cyclotron frequencies in the trap). Sometimes these two nuclides can be very far apart. Thus, those measurements are in most cases best represented as a combination of two masses. Other types of direct experimental methods, such as ‘Smith-type’, ‘Schottky’, ‘Isochronous’, ‘Time-of-flight’ and some ‘Multi-reflection Time-of-flight’ mass spectrometers, are calibrated in a more complex way, and are thus published by their authors as absolute mass values. They are then presented in Table I as a difference:  ${}^4\text{El}-u$ .

For completeness, we mention that early mass-spectrometric “triplet” measurements on unstable nuclides (cf. Section 6.2.2, p. 030002-24) can best be represented as linear combinations of masses of three isotopes, with non-integer coefficients [19].

This situation allows us to represent the input data graphically in a diagram of  $(N - Z)$  versus  $(N + Z)$  as shown in Fig. 1. This is straightforward for absolute mass-doublings and for two-nuclide difference cases; but not for spectrometer triplets and differential reaction energies (see Section 3, p. 030002-5). In general, the differential reactions are more important for one of the two reaction energies. Therefore, we present only the more important one in the graphs. For computational reasons, these data are treated as *primaries* (see below) even though the diagrams show only one connection.

In the present work, all input data are evaluated, i.e. calibrations are checked if necessary, and the data are compared with other results and with the trends from the mass surface (TMS, see next Section) in the region. As a consequence, several input data are corrected or even rejected (see below). All input data, including the rejected ones (not presented in Fig. 1), are given in Table I. As can be seen from Fig. 1, the accepted data may




Figure 1 (c). Diagram of connections for input data — continued.

allow determination of the mass of a particular nuclide using several different routes; such a nuclide is called *primary*. Their mass values in the table are then derived by least squares methods. In the other cases, the mass of a nuclide can be derived from only one connection to another nuclide; it is called a *secondary* nuclide. This classification is of importance for our calculation procedure (see Section 5.3, p. 030002-19).

The diagrams in Fig. 1 also show many cases where the relation between two atomic masses is accurately known, but not the actual mass values. Since our policy is to include all available experimental results, we have produced in such cases estimated mass values that are based on TMS in the neighborhood. Also, in this data representation, vacancies occur, which are filled with the estimated values using the same TMS procedure. Estimates of unknown masses are further discussed in the

next section.

Some care should be taken in the interpretation of Fig. 1, since excited isomeric states and data relations involving such isomers are not completely represented on these drawings. This is not considered as a serious defect; readers who want to update such values can conveniently consult Table I, where all relevant information is given.

#### 4 Regularity of the mass surface and the use of TMS

When atomic masses are displayed as a function of  $N$  and  $Z$ , one obtains a *surface* in a 3-dimensional space. However, due to the pairing energy, this surface is divided into four *sheets*. The even-even sheet lies lowest, the odd-odd highest, the other two nearly halfway in-






Figure 1 (d). Diagram of connections for input data — continued.

between, as shown in Fig. 2. The vertical distances from the even-even sheet to the odd-even and even-odd ones are the proton and neutron pairing energies,  $\Delta_{pp}$  and  $\Delta_{nn}$ , respectively, which are nearly equal. The distances of the last two sheets to the odd-odd sheet are equal to  $\Delta_{nn}-\Delta_{np}$  and  $\Delta_{pp}-\Delta_{np}$ , where  $\Delta_{np}$  is the proton-neutron pairing energy due to the interaction between the two odd nucleons, which are generally not in the same shell. These energies are represented in Fig. 2, where a hypothetical zero energy represents a nuclide with no pairing among the last nucleons.

Experimentally, it has been observed that the four sheets run nearly parallel in all directions, which means that the quantities  $\Delta_{nn}$ ,  $\Delta_{pp}$  and  $\Delta_{np}$  vary smoothly and slowly with  $N$  and  $Z$ . In addition, each of the mass sheets also varies smoothly, but rapidly with  $N$  and  $Z$  [20]. The smoothness is also observed for first or-

der derivatives (slopes, e.g. the graphs given in Part II, p. 030003-4) and all second-order derivatives (curvatures of the mass surface). They are only interrupted in places by cusps or bumps associated with important changes in nuclear structure: shell or sub-shell closures, shape transitions (spherical-deformed, prolate-oblate), and the so-called ‘Wigner’ cusp along the  $N = Z$  line.

This observed regularity of the mass sheets in all places, where there is no change in the underlying physics, can be considered as one of the BASIC PROPERTIES of the mass surface. Thus, dependable estimates of unknown, poorly known or questionable masses can be obtained by extrapolation from the well-known mass values on the same sheet. In the evaluation of masses the property of regularity and the possibility to make estimates are used for several purposes:




Figure 1 (e). Diagram of connections for input data — continued.

1. Any coherent deviation from the regularity, in a region  $(N, Z)$  of some extent, could be considered as an indication that some new physics property is being discovered. However, if only one single mass violates the trends from the mass surface (TMS) defined by the neighboring nuclides, then one may seriously question the correctness of the related datum. In such a case, there might be some undetected systematic [21] contribution to the reported experimental results for this mass. We then re-examine with extra care the available experimental information in the literature for possible errors and often consult with the corresponding authors for additional information. Such a process often leads to corrections.
2. There are cases where several experimental data

disagree, but no particular reason can be found for rejecting one, or some of them. In such cases, the measure of agreement with the regularity can be used by the evaluators for selecting which of the conflicting data will be accepted and used in the evaluation, thus following the same policy that was used in our earlier work.

3. There are cases where masses determined from ONLY ONE experiment (or from the same experiments) deviate severely from the smooth surface. Such cases are carefully examined (Section 4.2).
4. Finally, drawings of the mass surface allow to derive estimates for the still unknown masses, either from interpolations or from short extrapolations (see below, Section 4.3).




Figure 1 (f). Diagram of connections for input data — continued.

#### 4.1 Scrutinizing and manipulating the surface of masses

Direct representation of the mass surface is not convenient, since the binding energy varies very rapidly with  $N$  and  $Z$ . Splitting it in four sheets, as mentioned above, complicates even more such a representation. There are two ways that allow to observe the surface of masses with some precision: one of them uses the *derivatives* of this surface, the other is obtained by *subtracting a simple function* of  $N$  and  $Z$  from the masses.

**The derivatives of the mass surface** By *derivative* of the mass surface we mean a specified difference between the masses of two nearby nuclides. These functions are also smooth and have the advantage of displaying much smaller variations. For a derivative defined in

such a way that differences are between nuclides in the same mass sheet, their near parallelism can lead to an (almost) unified surface for the derivative, thus allowing a single display. Therefore, in order to visualize the trends from the mass surface, we found that the derivatives such as  $\alpha$ -decay energies and separation energies of two protons and two neutrons are the best tools to derive such estimates. These three derivatives are plotted against  $N$ , or  $Z$  in Part II, Figs. 1–26, p.030003-74.

However, from the way these derivatives are created, they give information only within one of the four sheets of the mass surface (e-e, e-o, o-e or o-o; e-o standing for even- $N$  and odd- $Z$ ). When examining the mass surface, an increased or decreased spacing of the sheets cannot be observed. Also, when estimating unknown masses, divergences of the four sheets could be unduly created,




Figure 1 (g). Diagram of connections for input data — continued.

which is unacceptable.

Fortunately, other representations are also possible (e.g. separately for odd and even nuclides: one-neutron separation energies versus  $N$ , one-proton separation energy versus  $Z$ ,  $\beta$ -decay energy versus  $A$ , ...). We have prepared a number of such graphs that can be obtained from the AMDC website [22].

The method of ‘derivatives’ suffers from the involvement of two masses for each point to be drawn, which means that if one point deviates from regularity, it could be due to either the mass of the nuclide it represents or that of the related nuclide, rendering the analysis rather complex. Also, reversely, if one mass is moved, then the two related points are changed in opposite directions, causing confusion in the drawing.

**Subtracting a simple function** Since the mass surface is smooth, one can try to define a function of  $N$  and  $Z$  as simple as possible and not too far from the real surface of masses. The difference between the mass surface and this function, while displaying reliably the structure of the former, will vary less rapidly, thus improving its observation.

A first and simple approach is the semi-empirical *liquid drop* formula of Bethe and Weizsäcker [23] with the addition of a pairing term in order to fuse more or less the four sheets of the mass surface. Another possibility, that we prefer [20], is to use the results of one of the modern models. However, we can use only those models that provide masses specifically for the spherical part, forcing the nucleus to be not deformed. The reason is that the models generally describe quite well the shell and sub-




Figure 1 (h). Diagram of connections for input data — continued.

shell closures, and to some extent the pairing energies, but not the locations of deformation. If the theoretical deformations were included and not located at exactly the same position as given by the experimental masses, the mass difference surface would show two dislocations for each shape transition. Interpretation of the resulting surface would then be difficult. In the present work, we make use of such differences with models. The plots that we have prepared can also be retrieved from the AMDC website [22].

**Manipulating the mass surface** In order to make estimates of unknown masses or to test changes on measured ones, an interactive graphical program was developed [20, 24] that allows a simultaneous observation of four graphs, either from the ‘derivatives’ type or from the ‘differences’ type, as a function of any of the variables  $N$ ,  $Z$ ,  $A$ ,  $N - Z$  or  $N - 2Z$ , while drawing iso-lines

(lines connecting nuclides having same value for a parameter) of any of these quantities. The mass of a nuclide can be modified or created in any view and we can determine how much freedom is left in setting a value for this mass. At the same time, interdependence through secondary connections (Fig. 1) are taken into account. In cases where two tendencies may alternate, following the parity of the proton or the neutron numbers, one of the parities may be deselected.

The replaced values for data yielding the ‘irregular masses’ as well as the ‘estimated unknown masses’ (see below) are thus derived by observing the continuous property in several views of the mass surface, with all the consequences due to connections to masses in the same chain. Comparisons with the predictions of 16 nuclear mass-models are presently available in this program.

With this graphical tool, the results of ‘replacement’ analyses are felt to be most robust; and also the estima-




Figure 1 (i). Diagram of connections for input data — continued.




Figure 1 (j). Diagram of connections for input data — continued.




Figure 2. The surface of masses is split into four sheets. This scheme represents the pairing energies responsible for this splitting. The zero energy surface is purely hypothetical with no pairing at all among the outer nucleons.

tion of unknown masses is more reliable.

All mass values dependent on interpolation procedures and indeed all values not derived from experimental data alone have been clearly marked with the sharp ‘#’ symbol in all tables, and elsewhere in the text.

Since publication of AME1983 [25], estimates are also given for the precision of data derived from TMS. These uncertainties are not based on a formalized procedure, but rather on previous experience with such estimates.

In the case of extrapolation, however, the uncertainty of the estimated mass will increase with the distance of extrapolation. These uncertainties are obtained by considering several graphs of TMS with a guess on how much the estimated mass may change without causing the extrapolated surface to look too distorted. This recipe is unavoidably subjective, but has proven to be efficient through the agreement of these estimates with newly measured masses in a large majority of cases [26].

#### 4.2 Irregular mass values

When a single mass deviates significantly from regularity with no similar pattern for nuclides with same  $N$  or with same  $Z$  values, then the correctness of the data determining this mass may be questioned.

Our policy, redefined in AME1995 [27], for those locally *irregular* masses, and only when they are derived from a unique mass relation (i.e., not confirmed by a different experimental method), is to replace them by values derived from trends from the mass surface (TMS). Table C lists 31 such cases in the present evaluation that were removed, to avoid strongly oscillating plots (to be compared to 27 cases in AME2012, 27 in AME2003, 59 in AME1995 and 67 in AME1993). Although these numbers reflects a more strict use of this procedure, the user of our tables should not assume that the remaining 31

items are the same ones carried on from generation to generation. The opposite, however, is true: most of the old ones have been replaced by new data, thus showing that we were correct in our choice. Generally, only one experimental result is reported for such unique mass relation. But sometimes there are two measurements (2 cases) or three (in previous evaluations) that we still treat the same way, since use of the same method and the same type of relation may well lead to the same systematic uncertainty (for example a mis-assignment or ignorance of a final level). Taking into account the connecting chains for secondaries (Figs. 1a–1j) has the consequence that several more ground state masses are affected (and twice as many values in each type of plot of derivatives as given in Part II). It should be stressed that only the most striking cases have been treated in this way, those necessary to avoid, as much as possible, confusions in the graphs in Part II. In particular, the plots of  $\alpha$ -decay energies of light nuclides (Fig. 18 and 19 in Part II, p. 030003–252 and 253) exhibit many overlaps and crossings that obscure the drawings; no attempt was made to locate possible origins of such irregularities.

Replacing these few irregular experimental values by the ones we recommend, in all tables and graphs in AME2016, means also that, as explained already in AME1995, we discontinued an older policy that was introduced in AME1993, where original irregular experimental values were given in all main tables, and ‘recommended’ ones given separately in secondary tables. This policy led to confusion for many users of our tables. Since AME1995, we only give what we consider the “*best recommended values*”, using, when we felt necessary and as explained above, “*values derived from TMS*”. Data which are not used following this policy are given in Table C and they can be easily located in Table I, where they are flagged ‘D’ and always accompanied by a comment



explaining in which direction the value has been changed and by what amount.

Such data, as well as other local irregularities that can be observed in the figures in Part II, could be considered

as incentive for remeasurements, preferably by a different method, in order to remove any doubt and possibly point out true irregularities due to physical properties.

Table C. Experimental data that we recommend replacing by values derived from TMS in AME2016.

Item	Reference <sup>a)</sup>	Experimental value		Recommended value	
<sup>39</sup> Al-u	2007Ju03	21653	676	22170#	430#
<sup>42</sup> Si-u	2007Ju03	16275	623	17680#	540#
<sup>44</sup> P-u	2007Ju03	10070	966	11220#	540#
<sup>47</sup> Cl-u	2007Ju03	-9576	1074	-10500#	430#
<sup>49</sup> Ar-u	2015Me01	-19110	1180	-18450#	430#
<sup>67</sup> Mn-u	2015Me.A	-36600	670	-35920#	320#
<sup>69</sup> Fe-u	2015Me.A	-42240	640	-41900#	430#
<sup>70</sup> Co-u	2011Es06	-50370	320	-50060#	320#
<sup>74</sup> Ni-u	2011Es06	-52830	1060	-52020#	210#
<sup>77</sup> Cu-u	2006Ha62	-51850	540	-52200#	160#
<sup>79</sup> Cu-u	2006Ha62	-46700	540	-44810#	320#
<sup>80</sup> Zr-u	1998Is06	-59600	1600	-58360#	320#
<sup>94</sup> Br-u	2016Kn03	-50242	429	-50890#	320#
<sup>94</sup> Ag <sup>n</sup> (β <sup>+</sup> ) <sup>94</sup> Pd	2004Mu30	17700	500	20180#	300#
<sup>101</sup> Rb(β <sup>-</sup> ) <sup>101</sup> Sr	1992Ba28	11810	110	12480#	200#
<sup>113</sup> Mo-u	2016Kn03	-57317	337	-56350#	320#
<sup>116</sup> Cs(εα) <sup>112</sup> Te	1977Bo28	12300	400	13100#	100#
	1976Jo.A	12400	900	13100#	100#
<sup>118</sup> Ru-u	2016Kn03	-61879	196	-61470#	220#
<sup>118</sup> In <sup>m</sup> (β <sup>-</sup> ) <sup>118</sup> Sn	1964Ka10	4270	100	4530#	50#
<sup>120</sup> In <sup>m</sup> (β <sup>-</sup> ) <sup>120</sup> Sn	1964Ka10	5280	200	5420#	50#
	1978Al18	5340	170	5420#	50#
<sup>124</sup> Pd-u	2016Kn03	-64617	399	-62680#	320#
<sup>126</sup> Ag-u	2016Kn03	-65926	329	-65140#	220#
<sup>129</sup> Nd(εp) <sup>128</sup> Ce	1978Bo.A	5300	300	5930#	200#
<sup>135</sup> Pm <sup>m</sup> (β <sup>+</sup> ) <sup>135</sup> Nd	1995Ve08	6040	150	6390#	50#
<sup>142</sup> Dy(β <sup>+</sup> ) <sup>142</sup> Tb	1991Fi03	7100	200	6440#	200#
<sup>143</sup> I-u	2016Kn03	-53849	495	-54350#	220#
<sup>150</sup> Ba-u	2016Kn03	-55309	371	-53570#	320#
<sup>154</sup> Ce-u	2016Kn03	-56404	619	-56060#	220#
<sup>220</sup> Pa(α) <sup>216</sup> Ac	1987Fa.A	9829	50	9650#	50#
<sup>270</sup> Db(α) <sup>266</sup> Lr	2014Kh04	8019	30	8260#	200#
<sup>281</sup> Rg(α) <sup>277</sup> Mt	2013Og04	9454	40	9900#	400#

a) References are listed in Part II in this issue, Section 6, p.030003-5.



The present authors insist that only the most striking irregularities have been replaced by estimates. Here we give the mass of  $^{84}\text{Nb}$  as an example. The  $Q_\beta$  value was measured in [1996Sh27] with an uncertainty claimed to be 300 keV. However, trends from mass surface strongly suggested that  $^{84}\text{Nb}$  should be 3200 keV less bound. Therefore the experimental value was labeled with “D” in AME2012 and replaced by an estimate. Recently the mass of  $^{84}\text{Nb}$  was measured by mass spectrometry [2016Xi.A], and the value is very close (200 keV) to our estimate, which was given with a precision of 300# keV.

### 4.3 Estimates for unknown masses

Estimates for unknown masses also are made with the use of trends from the mass surface, as explained above, by demanding that all graphs should be as smooth as possible, except where they are expected to be affected by shell closures or nuclear deformation effects. Therefore, we warn the user of our tables that the present extrapolations will be wrong if new regions of deformation or (semi-) magic numbers appear.

In addition to the rather severe constraints imposed by the requirement of simultaneous REGULARITY of all graphs, many further constraints result from knowledge of reaction or decay energies in the regions where these estimates are made. These regions and these constraints are shown in Figs. 1a–1j. Two kinds of constraints are present. In some cases the masses of  $(Z,A)$  and  $(Z,A+4)$  are known but not the mass of  $(Z,A+2)$ . Then, the values of  $S_{2n}(A+2)$  and  $S_{2n}(A+4)$  cannot both be chosen freely from the graphs; their sum is known. In other cases, the mass differences between several nuclides  $(A+4n,Z+2n)$  are known from  $\alpha$ -decays and also those of  $(A-2+4n,Z+2n)$ . Then, the differences between several successive  $S_{2n}(A+4n,Z+2n)$  are known. Similar situations exist for two or three successive  $S_{2p}$  or  $Q_\alpha$  values.

Knowledge of stability or instability against particle emission, or limits on proton or  $\alpha$  emission, yields upper or lower limits on the separation energies.

For proton-rich nuclides with  $N < Z$ , mass estimates can be obtained from the charge symmetry. This feature gives a relation between masses of isobars around the  $N = Z$  line. In several cases, we made a correction by including the Thomas-Ehrman effect [28], which makes proton-unstable nuclides more bound than what follows from the above estimate. For very light nuclides, we can use the estimates of this effect as proposed by Comay *et al.* [29].

Another often good estimate can be obtained from the observation that masses of nuclear states belonging to an isobaric multiplet are represented quite accurately by a quadratic equation of the charge number  $Z$  (or of the third components of the isospin,  $T_z = \frac{1}{2}(N-Z)$ ): the

Isobaric Multiplet Mass Equation (IMME) [30]. The use of this relation is attractive, since it uses experimental information such as the excitation energies of the isobaric analog states (IAS). New mass measurements regularly question the validity of the IMME, followed soon by other work showing that another member of the same multiplet needs to be questioned. For example, [2012Zh34] found, by measuring the mass of  $^{53}\text{Ni}$ , a breakdown of the quadratic form of IMME for the  $A = 53$  quartet, from which a non-zero coefficient  $d$  was derived to be 39(11) keV. In a later experiment, the lowest  $T = 3/2$  state in  $^{53}\text{Co}$  was established via the measurement of the  $\beta$ -delayed  $\gamma$  deexcitation of  $^{53}\text{Ni}$  [2016Su10], which questioned the isobaric-analog state (IAS) assignment in a previous  $\beta$ - $p$  decay experiment [2007Do17]. The validity of the IMME in the quadratic form was thus restored for  $A = 53$ .

Up to AME1983, we indeed used the IMME for deriving mass values for nuclides for which no, or little information was available. This policy was questioned with respect to the correctness in stating as ‘experimental’ a quantity that was derived by combination with a calculation. Since AME1993, it was decided not to present any IMME-derived mass values in our evaluation, but rather use the IMME as a guideline when estimating masses of unknown nuclides. We continue this policy here, and do not replace experimental values by an estimated one from IMME, even if orders of magnitude more precise. Typical examples are  $^{28}\text{S}$  and  $^{40}\text{Ti}$ , for which IMME predicts masses with precisions of 24 keV and 22 keV, respectively, whereas the experimental masses are known for both from double-charge-exchange reactions with 160 keV precision.

The extension of the IMME to higher energy isobaric analog states has been studied by Wapstra [31]. The validity of the method, however, is made uncertain by possible effects spoiling the relation. In the first place, the strength of some IASs at high excitation energies is known to be distributed over several levels with the same spin and parity. Even in cases where this interference effect has not been observed, it remains a possibility, and as such, introduces an uncertainty in the energy level to be attributed to the IAS. In the second place, as argued by Thomas and Ehrman [28], particle-unstable levels must be expected to be shifted somewhat.

It also happens that information on excitation energies of  $T_z = -T + 1$  IASs is available from measurements on proton emission following  $\beta$  decays of their  $T_z = -T$  parents. The authors, in some cases, derived a mass value from their results for the parent nuclide, using a formula, derived by Antony *et al.* [32], from a study of known energy differences between IAS. We observe, however, that one obtains somewhat different mass values by combining Antony differences with the

mass of the mirror nuclide of the mother. Also, earlier considerations did not take into account the difference between proton-pairing and neutron-pairing energies, which A.H. Wapstra noticed have a non-negligible influence on the IMME constants.

Another possibility is to use a relation proposed by Jänecke [33], as done for example by Axelsson *et al.* [34] in the case of  $^{31}\text{Ar}$ . In several cases we have compared the results of different ways for extrapolating, in order to find a best estimate for the desired mass value.

Enough values have been estimated to ensure that every nuclide for which there is any experimental  $Q$ -value is connected to the main group of primary nuclides. In addition, the evaluators want to achieve continuity of the mass surface. Therefore, an estimated value is included for any nuclide if it is between two experimentally studied nuclides on a line defined by either  $Z = \text{constant}$  (isotopes),  $N = \text{constant}$  (isotones),  $N - Z = \text{constant}$  (isodiaspheres), or, in a few cases  $N + Z = \text{constant}$  (isobars). It would have been desirable to also give estimates for all unknown nuclides that are within reach of the present accelerators and mass separator technologies. Unfortunately, such an ensemble is not easy to define. Instead, we estimate mass values for all nuclides for which at least one piece of experimental information is available (e.g. identification or half-life measurement or proof of instability towards proton or neutron emission). Then, the ensemble of experimental and estimated masses has the same contour as in the NUBASE2016 evaluation.

## 5 Calculation Procedures

The atomic mass evaluation is unique when compared to the other evaluations of data [20], in a sense that almost all mass determinations are relative measurements, not absolute ones. Even those called ‘absolute mass doublets’ are relative to  $^{12}\text{C}$ ,  $^{35}\text{Cl}$  or  $^{37}\text{Cl}$ . Each experimental datum sets a relation in mass or in energy among two (in a few cases three or more) nuclides. It can be therefore represented by one link among these two nuclides. The ensemble of these links generates a highly entangled network. Figs. 1a–1j, in Section 3 above, show a schematic representation of such a network.

The masses of a large number of nuclides are multiply determined, entering the entangled area of the canvas, mainly along the backbone. Correlations do not allow determining their masses in a straightforward manner.

To take into account these correlations we use a least-squares method weighed according to the precision with which each piece of data is known. This method allows to determine a set of adjusted masses.

### 5.1 Least-squares method

Each piece of data has a value  $q_i \pm dq_i$  with the accuracy  $dq_i$  (one standard deviation) and makes a relation between two, three or four masses with unknown values  $m_\mu$ . An overdetermined system of  $Q$  data to  $M$  masses ( $Q > M$ ) can be represented by a system of  $Q$  linear equations with  $M$  parameters:

$$\sum_{\mu=1}^M k_i^\mu m_\mu = q_i \pm dq_i, \quad (1)$$

e.g. for a nuclear reaction  $A(a,b)B$  requiring an energy  $q_i$  to occur, the energy balance is written:

$$m_A + m_a - m_b - m_B = q_i \pm dq_i. \quad (2)$$

Thus,  $k_i^A = +1$ ,  $k_i^a = +1$ ,  $k_i^b = -1$  and  $k_i^B = -1$ .

In matrix notation,  $\mathbf{K}$  being the  $(Q, M)$  matrix of coefficients, Eq. 1 is written:  $\mathbf{K}|m\rangle = |q\rangle$ . Elements of matrix  $\mathbf{K}$  are almost all null: e.g. for  $A(a,b)B$ , Eq. 2 yields a line of  $\mathbf{K}$  with only four non-zero elements.

We define the diagonal weight matrix  $\mathbf{W}$  by its elements  $w_i^i = 1/(dq_i dq_i)$ . The solution of the least-squares method leads to a very simple construction:

$${}^t\mathbf{K}\mathbf{W}\mathbf{K}|m\rangle = {}^t\mathbf{K}\mathbf{W}|q\rangle. \quad (3)$$

The NORMAL matrix  $\mathbf{A} = {}^t\mathbf{K}\mathbf{W}\mathbf{K}$  is a square matrix of order  $M$ , positive-definite, symmetric and regular and hence invertible [35]. Thus the vector  $|\bar{m}\rangle$  for the adjusted masses is:

$$|\bar{m}\rangle = \mathbf{A}^{-1} {}^t\mathbf{K}\mathbf{W}|q\rangle \quad \text{or} \quad |\bar{m}\rangle = \mathbf{R}|q\rangle. \quad (4)$$

The rectangular  $(M, Q)$  matrix  $\mathbf{R}$  is called the RESPONSE matrix.

The diagonal elements of  $\mathbf{A}^{-1}$  are the squared errors on the adjusted masses, and the non-diagonal ones  $(a^{-1})_\mu^\nu$  are the coefficients for the correlations between masses  $m_\mu$  and  $m_\nu$ . Values for correlation coefficients for the most precise nuclides are given in Table B of Part II (p.030003-3). Following the advice of B.N. Taylor, we now give on the website of the AMDC [22] the full list of correlation coefficients, allowing any user to perform exact calculation of any combination of masses.

One of the most powerful tools in the least-squares calculation described above is the flow-of-information matrix, discovered in 1984 by one of us [36]. This matrix allows to trace back the contribution of each individual piece of data to each of the parameters (here the atomic masses). The AME uses this method since 1993.

The flow-of-information matrix  $\mathbf{F}$  is defined as follows:  $\mathbf{K}$ , the matrix of coefficients, is a rectangular  $(Q, M)$  matrix. The transpose of the response matrix  ${}^t\mathbf{R}$  is also a  $(Q, M)$  rectangular one. The  $(i, \mu)$  element of  $\mathbf{F}$

is defined as the product of the corresponding elements of  $\mathbf{tR}$  and of  $\mathbf{K}$ . In Ref. [36], it is demonstrated that such an element represents the “*influence*” of datum  $i$  on parameter (mass)  $m_\mu$ . A column of  $\mathbf{F}$  thus represents all the contributions brought by all data to a given mass  $m_\mu$ , and a line of  $\mathbf{F}$  represents all the influences given by a single piece of data. The sum of influences along a line is the “*significance*” of that datum. It has also been proven [36] that the influences and significances have all the expected properties, namely that the sum of all the influences on a given mass (along a column) is unity, that the significance of a datum is always less than unity and that it always decreases when new data are added. The significance defined in this way is exactly the quantity obtained by squaring the ratio of the uncertainty on the adjusted value over that of the input one, which was the recipe used before the discovery of the  $\mathbf{F}$  matrix to calculate the relative importance of data.

A simple interpretation of influences and significances can be obtained in calculating, from the adjusted masses and Eq. 1, the adjusted data:

$$|\bar{q}\rangle = \mathbf{KR}|q\rangle. \quad (5)$$

The  $i^{\text{th}}$  diagonal element of  $\mathbf{KR}$  represents then the contribution of datum  $i$  to the determination of  $\bar{q}_i$  (same datum): this quantity is exactly what is called above the *significance* of datum  $i$ . This  $i^{\text{th}}$  diagonal element of  $\mathbf{KR}$  is the sum of the products of line  $i$  of  $\mathbf{K}$  and column  $i$  of  $\mathbf{R}$ . The individual terms in this sum are precisely the *influences* defined above.

The flow-of-information matrix  $\mathbf{F}$ , provides thus insight on how the information from datum  $i$  flows into each of the masses  $m_\mu$ .

The flow-of-information matrix cannot be given in full in a printed table. It can be observed along lines, displaying for each datum, the nuclides influenced by this datum and the values of these *influences*. It can be observed also along columns to display for each primary mass all contributing data with their *influence* on that mass.

The first display is partly given in the table of input data (Table I) in column ‘Signf.’ for the *significance* of primary data and ‘Main infl.’ for the largest *influence*. Since in the large majority of cases only two nuclides are concerned in each piece of data, the second largest *influence* could easily be deduced. It is therefore not felt necessary to give a table of all *influences* for each primary datum.

The second display is given in Part II, Table II (p.030003-3) for the up to three most important data with their *influence* in the determination of each primary mass.

## 5.2 Consistency of data

The system of equations being largely over-determined ( $Q \gg M$ ) offers the evaluator several interesting possibilities to examine and judge the data. One might for example examine all data for which the adjusted values deviate significantly from the input ones. This helps to locate erroneous pieces of information. One could also examine a group of data in one experiment and check if the uncertainties assigned to them in the experimental paper were not underestimated.

If the precisions  $dq_i$  assigned to the data  $q_i$  were indeed all accurate, the normalized deviations  $v_i$  between adjusted  $\bar{q}_i$  (Eq. 5) and input  $q_i$  data,  $v_i = (\bar{q}_i - q_i)/dq_i$ , would be distributed as a Gaussian function of standard deviation  $\sigma = 1$ , and would make  $\chi^2$ :

$$\chi^2 = \sum_{i=1}^Q \left( \frac{\bar{q}_i - q_i}{dq_i} \right)^2 \quad \text{or} \quad \chi^2 = \sum_{i=1}^Q v_i^2 \quad (6)$$

equal to  $Q - M$ , the number of degrees of freedom, with a standard deviation of  $\sqrt{2(Q - M)}$ .

One can define as above the NORMALIZED CHI,  $\chi_n$  (or ‘consistency factor’ or ‘Birge ratio’):  $\chi_n = \sqrt{\chi^2/(Q - M)}$  for which the expected value is  $1 \pm 1/\sqrt{2(Q - M)}$ .

Another quantity of interest for the evaluator is the PARTIAL CONSISTENCY FACTOR,  $\chi_n^p$ , defined for a (homogeneous) group of  $p$  data as:

$$\chi_n^p = \sqrt{\frac{Q}{Q - M} \frac{1}{p} \sum_{i=1}^p v_i^2}. \quad (7)$$

Of course the definition is such that  $\chi_n^p$  reduces to  $\chi_n$  if the sum is taken over all the input data. One can consider for example the two main classes of data: the reaction and decay energy measurements and the mass-spectrometric data (see Section 5.5). One can also consider groups of data related to a given laboratory and with a given method of measurement and examine the  $\chi_n^p$  of each of them. There are presently 278 groups of data in Table I (among which 185 have at least one measurement used in determining the masses), identified in column ‘Lab’. A high value of  $\chi_n^p$  might be a warning on the validity of the considered group of data within the reported uncertainties. We used such analyses in order to be able to locate questionable groups of data. In bad cases they are treated in such a way that, in the final adjustment, no really serious conflicts occur. Remarks in Table I report where such corrections have been made.

## 5.3 Separating secondary data

In Section 3, while examining the diagrams of connections (Fig. 1), we noticed that, whereas the masses of *secondary* nuclides can be determined uniquely from the

chain of secondary connections going down to a *primary* nuclide, only the latter see the complex entanglement that necessitated the use of the least-squares method.

In terms of equations and parameters, we consider that if, in a collection of equations to be treated with the least-squares method, a parameter occurs in only one equation, removing this equation and this parameter will not affect the result of the fit for all other data. Thus, we can redefine more precisely what was called *secondary* in Section 3: the parameter above is a *secondary* parameter (or mass) and its related equation is a *secondary* equation. After the reduced set has been solved, then the *secondary* equation can be used to determine the final value and uncertainty for that particular *secondary* parameter. The equations and parameters remaining after taking out all secondaries are called *primary*.

Therefore, only the system of *primary* data is overdetermined, and thus will be improved in the adjustment, so that each *primary* nuclide will benefit from all the available information. *Secondary* data will remain unchanged; they do not contribute to  $\chi^2$ .

The diagrams in Fig. 1 show, that many *secondary* data exist. Thus, taking them out simplifies considerably the system. More importantly, if a better value is found for a *secondary* datum, the mass of the *secondary* nuclide can easily be improved (one has only to be careful since the replacement can change other *secondary* masses down the chain, see Fig. 1). The procedure is more complicated for new *primary* data.

We define DEGREES for *secondary* nuclides and *secondary* data. They reflect their distances along the chains connecting them to the network of primaries. The first secondary nuclide connected to a primary one will be a nuclide of degree 2; and the connecting datum will be a datum of degree 2 as well. Degree 1 is for primary nuclides and data. Degrees for secondary nuclides and data range from 2 to 18. It is the heaviest nuclide  $^{295}\text{Ei}$  that has the highest degree number 18. Its mass is determined through a long  $\alpha$  chain, albeit many of them are just estimates. In Table I, the degree of data is indicated in column ‘Dg’. In the table of atomic masses (Part II, Table I, p. 030003-6), each *secondary* nuclide is marked with a label in column ‘Orig.’ indicating from which other nuclide its mass value is determined.

To summarize, separating secondary nuclides and secondary data from primaries allow to significantly reduce the size of the system that will be treated by the least-squares method described above. After treatment of the primary data alone, the adjusted masses for primary nuclides can be easily combined with the secondary data to yield masses of secondary nuclides.

In the next Section we will show methods for reducing further this system, but without loss of any information. Methods that reduce the system of primaries for the benefit of the secondaries not only decrease compu-

tational time (which nowadays is not so important), but allows an easier insight into the relations between data and masses, since no correlation is involved.

*Remark:* the word *primary* used for these nuclides and for the data connecting them does not mean that they are more important than the others, but only that they are subject to the least-squares treatment above. The labels *primary* and *secondary* are not intrinsic properties of data or nuclides. They may change from primary to secondary or vice versa when other information becomes available.

## 5.4 Compacting the set of data

### 5.4.1 Pre-averaging

Two or more measurements of the same physical quantities can be replaced without loss of information by their average value and precision, reducing thus the system of equations to be treated. By extending this procedure, we consider *parallel* data: reaction data occur that give essentially values for the mass-difference between the same two nuclides, except in rare cases where the precision is comparable to that in the masses of the reaction particles. Example:  $^{14}\text{C}(^7\text{Li},^7\text{Be})^{14}\text{B}$  and  $^{14}\text{C}(^{14}\text{C},^{14}\text{N})^{14}\text{B}$ ; or  $^{22}\text{Ne}(t,^3\text{He})^{22}\text{F}$  and  $^{22}\text{Ne}(^7\text{Li},^7\text{Be})^{22}\text{F}$ .

Such data are represented together, in the main least-squares fit calculations, by one of them carrying their average value. If the  $Q$  data to be pre-averaged are strongly conflicting, i.e. if the consistency factor (or Birge ratio)  $\chi_n = \sqrt{\chi^2/(Q-1)}$  resulting in the calculation of the pre-average is greater than 2.5, the (internal) precision  $\sigma_i$  in the average is multiplied by the Birge ratio ( $\sigma_e = \sigma_i \times \chi_n$ ). There are no cases where  $\chi_n > 2.5$ , see Table D (there were 2 cases in AME2012, and 6 in AME2003). The quantity  $\sigma_e$  is often called the ‘external error’. However, this treatment is not used in the rare cases where the precisions of the input values differ too much, since the assigned uncertainties lose any significance. If such a case occurs, considering policies from the Particle Data Group [37] and some statistical-treatment methods reviewed by Rajput and MacMahon [38], we adopt an arithmetic average and the dispersion of values as an uncertainty, which is equivalent to assigning to each of these conflicting data the same uncertainty.

In the present evaluation, we have replaced 2977 data by 1186 averages. As can be seen from Fig. 3, as much as 23% of which have values of  $\chi_n$  (Birge ratio) beyond unity, 1.6% beyond two and none beyond 3, giving an overall very satisfactory distribution for our treatment.

As a matter of fact, in a complex system like the one here, many values of  $\chi_n$  beyond 1 or 2 are expected to exist, and if the uncertainties were multiplied by  $\chi_n$  in all these cases, the  $\chi^2$ -test on the total adjustment would have been invalidated.

Table D. Worst pre-averagings.  $n$  is the number of data in the pre-average.

Item	$n$	$\chi_n$	$\sigma_e$	Item	$n$	$\chi_n$	$\sigma_e$
$^{186}\text{W}(n,\gamma)^{187}\text{W}$	2	2.44	0.11	$^{177}\text{Pt}(\alpha)^{173}\text{Os}$	2	2.06	6.06
$^{144}\text{Ce}(\beta^-)^{144}\text{Pr}$	2	2.44	2.18	$^{244}\text{Cf}(\alpha)^{240}\text{Cm}$	2	2.03	3.97
$^{220}\text{Fr}(\alpha)^{216}\text{At}$	2	2.34	4.66	$^{15}\text{N}(p,n)^{15}\text{O}$	2	2.03	1.28
$^{75}\text{As}(n,\gamma)^{76}\text{As}$	2	2.32	0.17	$^{58}\text{Fe}(t,p)^{60}\text{Fe}$	4	2.03	7.38
$^{110}\text{In}(\beta^+)^{110}\text{Cd}$	3	2.29	28.4	$^{204}\text{Tl}(\beta^-)^{204}\text{Pb}$	2	2.03	0.39
$^{146}\text{Ba}(\beta^-)^{146}\text{La}$	2	2.24	107.4	$^{278}\text{Mt}(\alpha)^{274}\text{Bh}$	3	1.98	43.6
$^{40}\text{Cl}(\beta^-)^{40}\text{Ar}$	2	2.21	76.1	$^{167}\text{Os}(\alpha)^{163}\text{W}$	4	1.98	3.50
$^{219}\text{U}(\alpha)^{215}\text{Th}$	2	2.18	38.5	$^{106}\text{Ag}(\varepsilon)^{106}\text{Pd}$	2	1.98	6.63
$^{153}\text{Gd}(n,\gamma)^{154}\text{Gd}$	2	2.16	0.39	$^{78}\text{Se}(n,\gamma)^{79}\text{Se}$	3	1.96	0.28
$^{36}\text{S}(^{11}\text{B},^{13}\text{N})^{34}\text{Si}$	3	2.14	32.4	$^{46}\text{Ca}(n,\gamma)^{47}\text{Ca}$	2	1.94	0.56
$^{113}\text{Cs}(p)^{112}\text{Xe}$	3	2.10	5.03	$^{181}\text{Pb}(\alpha)^{177}\text{Hg}$	3	1.93	15.2
$^{223}\text{Pa}(\alpha)^{219}\text{Ac}$	2	2.09	10.0	$^{145}\text{Sm}(\varepsilon)^{145}\text{Pm}$	2	1.92	7.92
$^{27}\text{P}^i(2p)^{25}\text{Al}$	2	2.08	74.7	$^{234}\text{Th}(\beta^-)^{234}\text{Pa}^m$	3	1.90	2.10
$^{204}\text{Rn}-^{208}\text{Pb}_{0.981}$	2	2.06	18.2				

This explains the choice we made here of a rather high threshold ( $\chi_n^0 = 2.5$ ), compared e.g. to  $\chi_n^0 = 2$  recommended by Woods and Munster [39] or  $\chi_n^0 = 1$  used in a different context by the Particle Data Group [37], for departing from the rule of ‘internal error’ of the weighted average.




Figure 3. Birge Ratios of all the pre-averaged data.

Besides the computer-automated pre-averaging, we found it convenient, in some  $\beta^+$ -decay cases, to combine results stemming from various capture ratios in an average. These cases are  $^{109}\text{Cd}(\varepsilon)^{109}\text{Ag}$  (average of 3 data),  $^{139}\text{Ce}(\varepsilon)^{139}\text{La}$  (average of 10) and  $^{195}\text{Au}(\varepsilon)^{195}\text{Pt}$  (5 results), and they are detailed in Table I. Four more cases ( $^{147}\text{Tb}$ ,  $^{152}\text{Ho}$ ,  $^{166}\text{Yb}$  and  $^{207}\text{Bi}$ ) occur in our list, but they carry no weight and are labeled with ‘U’ in Table I.

#### 5.4.2 Used policies in treating parallel data

In averaging  $\beta^-$  ( $\alpha^-$ ) decay energies derived from branches observed in the same experiment, to or from different levels in the decay of a given nuclide, the uncertainty we use for further evaluation is not the one resulting from the weighted average adjustment, but instead we use the smallest experimental one. In this way, we avoid decreasing artificially the part of the uncertainty that is not due to statistics. In some cases, however, when it is obvious that the uncertainty is dominated by weak statistics, we do not follow the above rule (e.g.  $^{23}\text{Al}^i(p)^{22}\text{Mg}$  of [1997B104]).

Some quantities have been reported more than once by the same group. If the results are obtained by the same method in different experiments and are published in regular refereed journals, only the most recent one is used in the calculation, unless explicitly mentioned otherwise. There are two reasons for this policy. The first is that one might expect that the authors, who believe their two results are of the same quality, would have averaged them in their latest publication. The second is that if we accept and average the two results, we would have no control on the part of the uncertainty that is not due to statistics. Our policy is different if the newer result is published in a secondary reference (not refereed abstract, preprint, private communication, conference, thesis or annual report). In such cases, the older result is used in the calculations, except when the newer one is an update of the previous value. In the latter case, the original reference in our list mentions the unrefereed paper.

### 5.4.3 Replacement procedure

Large contributions to  $\chi^2$  have been known to be caused by a nuclide  $G$  connected to two other ones  $H$  and  $K$  by reaction links with large uncertainties compared to those deduced from the mass difference between  $H$  and  $K$ , in cases where the two disagreed. Evidently, contributions to  $\chi^2$  of such local discrepancies suggest an unrealistically high value of the overall consistency parameter. This is avoided by a replacement procedure: one of the two links is replaced by an equivalent value for the other. The pre-averaging procedure gives the most reasonable mass value for  $G$  and do not cause undesirably large contributions to  $\chi^2$ .

### 5.4.4 Insignificant data

Another feature to increase the meaning of the final  $\chi^2$  is to not use, in the least-squares procedure, data with weights at least a factor 10 smaller than other data, or combinations of *all* other data giving the same result. Such data were labeled with ‘U’ in the list of input data; comparison with the output values allows to check our judgment. Earlier, data were labeled ‘U’ if their weight was 10 times smaller than that of a *simple* combination of other data. This concept has been extended, since AME1993, to data that weigh 10 times less than the combination of *all* other accepted data. Until the AME2003 evaluation, our policy was not to print data labeled ‘U’ if they already appeared in one of our previous tables, reducing thus the size of the table of data to be printed. This policy has been changed since AME2012, and we try as much as possible to give all relevant data, also including insignificant ones. The reason for this is that conflicts might appear amongst recent results, and access to older ones might shed some light on our judgement, when evaluating the new data.

## 5.5 Used policies – treatment of undependable data

The important interdependence of most data, as illustrated by the connection diagrams (Figs. 1a–1j), allows local and general consistency tests. These can indicate that something may be wrong with the input values. We follow the policy of checking all significant data that differ by more than two (sometimes 1.5) standard deviations from the adjusted values. Fairly often, examination of the experimental results shows that a correction is necessary. Possible reasons could be that a particular decay has been assigned to a wrong final level or that a reported decay energy belongs to an isomer, rather than to a ground state, or even that the mass number assigned to a decay was incorrect. In such cases, the values are corrected and remarks are added below the corresponding  $A$ -group of data in Table I, in order to explain the reasons for the corrections.

It also happens that a careful examination of a particular paper would lead to serious doubts about the validity of the results within the reported precision, but could not permit making a specific correction. Doubts can also be expressed by the authors themselves. The results are given in Table I and compared with the adjusted values. They are labeled ‘F’, and not used in the final adjustment, but always followed by a comment to explain the reason for this label. The reader may observe that in several cases the difference between the experimental and adjusted values is small compared to the experimental uncertainty: this does not disprove the correctness of the label ‘F’ assignment.

It happens quite often that two (or more) pieces of data are discrepant, leading to important contribution to the  $\chi^2$ . A detailed examination of the papers may not allow correction or rejection, indicating that at least the results in one of them could not be trusted within the given uncertainties. Then, based on past experience, we use in the calculations the value that seems to be the most reliable, while the other is labeled ‘B’, if published in a regular refereed journal, or ‘C’ otherwise.

Data with labels ‘F’, ‘B’ or ‘C’ are not used in the calculations. We do not assign such labels if, as a result, no experimental value published in a regular refereed journal could be given for one or more resulting masses. When necessary, the policy defined for ‘irregular masses’ with ‘D’-label assignment may apply (see Section 4.2).

In some cases, detailed analysis of strongly conflicting data could not lead to reasons to assume that one of them is more dependable than the others or could not lead to a rejection of a particular data entry. Also, bad agreement with other data is not the only reason to doubt the correctness of the reported data. As in previous AME, and as explained above (see Section 4), we made use of the property of regularity of the surface of masses in making a choice, as well as in further checks on the other data.

We do not accept experimental results if information on other quantities (e.g. half-lives), derived in the same experiment and for the same nuclide, were in strong contradiction with well established values.

## 5.6 The AME computer program

Our computer program in four phases has to perform the following tasks: **i)** decode and check the data file; **ii)** build up a representation of the connections between masses, allowing thus to separate primary masses and data from secondary ones, to pre-average same and parallel data, and thus to reduce drastically the size of the system of equations to be solved (see Section 5.3 and 5.4), without any loss of information; **iii)** perform the least-squares matrix calculations (see above); and **iv)** deduce

the atomic masses (Part II, Table I), the nuclear reaction and separation energies (Part II, Table III), the adjusted values for the input data (Table I), the *influences* of data on the primary nuclides (Table I), the *influences* received by each primary nuclide (Part II, Table II), and display information on the inversion errors, the correlations coefficients (Part II, Table B), the values of the  $\chi^2$ s, the distribution of the  $v_i$  (see below), among others.

## 5.7 Results of the calculation

In this evaluation, we have 13035 experimental data of which 5663 are labeled ‘U’ (see above), 844 are labeled ‘O’ (old result from same group) and 853 are not accepted and labeled ‘B’, ‘C’, ‘D’ or ‘F’ (respectively 502, 157, 33 and 161 items). In the calculation we have thus 5675 valid input data, compressed to 3884 in the pre-averaging procedure. Separating secondary data, leaves a system of 2023 primary data, representing 1094 primary reactions and decays, and 929 primary mass-spectrometric measurements. To these are added 835 data estimated from TMS trends (see Section 4), p.030002-9), some of which are essential for linking unconnected experimental data to the network of experimentally known masses (see Figs. 1a–1j).

In the atomic mass table (Part II, Table I) there is a total of 3923 masses (including  $^{12}\text{C}$ ) of which 3435 are ground state masses (2497 experimental masses and 938 estimated ones), and 488 are excited isomers (369 experimental and 119 estimated). Among the 2497 experimental ground state masses, 111 nuclides have a precision better than 0.1 keV, 378 better than 1 keV and 1477 better than 10 keV (respectively 87, 315 and 1438 in AME2012). There are 153 nuclides known with uncertainties larger than 100 keV (123 in AME2012). Separating secondary masses in the ensemble of 3923, leaves 1207 primary masses ( $^{12}\text{C}$  not included).

Thus, we have to solve a system of 2023 equations with 1207 parameters. Theoretically, the expectation value for  $\chi^2$  should be  $816 \pm 20$  (and the theoretical  $\chi_n = 1 \pm 0.025$ ).

The total  $\chi^2$  of the adjustment is actually 825 ( $\chi_n = 1.005$ ), thus showing that the ensemble of evaluated data was of excellent quality, and that the adopted criteria of selection and rejection were adequate. In the past this was not always the case and in AME2003 we could observe that on average the uncertainties in the input values were underestimated by 23%. The distribution of the  $v_i$ 's (the individual contributions to  $\chi^2$ , as defined in Eq. 6, and given in Table I) is also acceptable. If we consider all the 11511 data that are used in the adjustment plus the ‘obsolete’ ones (label ‘O’) and the unweighed ones (label ‘U’), the distribution of  $v_i$ 's yields 20% of the cases beyond unity, 3.3% beyond two, and no items beyond 3.

Considering separately the two main classes of data, the partial consistency factors  $\chi_n^p$  are respectively 1.021 and 0.987 for energy measurements and for mass-spectrometric data, showing that both types of selective input data are of excellent quality.

As in our previous works [2, 40, 41], we have estimated the average accuracy for 185 groups of data used in the evaluation that were related to a given laboratory and a specific method of measurement, by calculating their partial consistency factors  $\chi_n^p$  (see Section 5.2). As much as 98 groups have  $\chi_n^p$  larger than unity, and 2 groups larger than 2.

## 6 Discussion of the input data

In most cases, values given by authors in the original publication are accepted, but there are also exceptions. One example is the performed recalibration due to change in the definition of volt, as discussed in Section 2. For somewhat less simple cases, a remark is added in Table I at the end of the concerned A-group. A curious example of combinations of data that cannot be accepted without change follows from the measurements of the Edinburgh-Argonne group [1997Da07]. They reported a series of  $\alpha$ -decay energies, where the ancestors were isomers between which the excitation energy was accurately known from the difference of their proton-decay energies. These authors gave values for the excitation energies between isomeric daughter pairs with considerably smaller uncertainties than those derived from the uncertainties quoted for the measured  $\alpha$ -decay energies. The reason is that the decay energies of two parallel  $\alpha$ -decay chains are correlated; this means that the uncertainties in their differences are relatively small. Unfortunately, the presented data do not allow an exact calculation of both the masses and the isomeric excitation energies. This would have required that, in addition to the two  $E_\alpha$  values of an isomeric pair, the uncertainty of the  $\alpha$ -energy difference should also have been given. Instead, entering all their  $Q_p$  and  $E_1$  (isomeric excitation energies) values in our adjustment would yield outputs with too small uncertainties, while accepting any partial collection makes some uncertainties too large. Therefore, in this case we do enter a selection of the input values, which are slightly changed, but chosen in such a way that our adjusted  $Q_\alpha$  and  $E_1$  values and corresponding uncertainties differ as little as possible from those given by the authors. A further complication could occur if some of the  $Q_\alpha$  values are also measured by other groups. But until now, we found no serious deviations in such cases.

A change in uncertainties, not values, is caused by the fact that, in several cases, we do not necessarily accept the reported  $\alpha$ -energy values as belonging to transitions between ground states. This also causes uncertainties in

the derived proton-decay energies to deviate from those reported by the authors (e.g. in the  $\alpha$ -decay chain of  $^{170}\text{Au}$ ), see also Section 7.9.

## 6.1 Improvements along the backbone

Since AME2012, all new mass-spectrometric data constituting the backbone were obtained from precision measurements of cyclotron frequencies of ions in Penning traps. Like to the classical measurements, where ratios of voltages or resistances were used, we found that the Penning trap results can be converted to a linear combination of masses of electrically neutral atoms (in  $\mu\text{u}$ ), without any loss of accuracy. A special mention is needed for the MIT-FSU group [2005Ra34] which reports their original results as linear equations, including corrections for electron and molecular binding energies. Other groups give their results as ratio of cyclotron frequencies (see also next paragraph), which we convert to linear equations as described in Appendix C, (p. 030002-43) and finally we add corrections for electron and molecular binding energies. In such cases, we added a remark to the equation used in the input data table (Table I), to describe the original data and our treatment. Some authors publish their results directly as masses, but this is not a recommended practice for high-precision mass measurements.

### 6.1.1 Calculation of molecular binding energies for very precise mass measurements

The most precise mass-spectrometric measurements use Penning trap spectrometers, which measure the cyclotron frequency of a reference ion and an ion of interest in a uniform magnetic field  $B$ . Not only the electronic binding energy (single or multiple ionization), but also the molecular binding energy (dissociation energy) can be involved in the measured frequency ratios, as described for example in Ref. [1995Di08]. For most molecules used in the experiments, the binding energy represents typically a correction of a few parts in  $10^{10}$  and its uncertainty only limits the accuracy of the neutral atomic mass to a few parts in  $10^{12}$ , for example in Ref. [2004Ra33]. For measurements with precisions not better than  $10^{-9}$  (100 eV/100 u), the molecule binding energy could be neglected without loss of accuracy. In cases where the precision is better than  $10^{-10}$  (10 eV/100 u), e.g. at MIT [1995Di08] and FSU [2015My03], it is necessary to take into account the molecular binding energy.

## 6.2 Mass spectrometry away from $\beta$ -stability

The reader interested in the history of mass-spectrometric measurements, the resolving powers, resolutions and the discoveries they rendered possible in nuclear physics and cosmology, can refer to the publication by one of us (G.A.) [1].

### 6.2.1 Penning trap spectrometers

Nowadays, seven Penning traps are being operated at the major accelerator facilities around the world: ISOLTRAP-Cern, CPT-Argonne, JYFLTRAP-Jyväskylä, LEBIT-East-Lansing, SHIPTRAP-Darmstadt, TITAN-Vancouver, and TRIGA-TRAP-Mainz. They measure the atomic masses for nuclides farther from the valley of  $\beta$ -stability, using the cyclotron frequencies of charged ions captured in the trap. Such a frequency is always compared to that of a (well) known reference nuclide in order to determine the ratio of two masses, which is converted, without loss of accuracy, to a linear relation between the two masses (see also Section 6.1 above and Appendix C, p. 030002-43). Experimental methods that utilize measurements of cyclotron frequency have an advantage compared to volt or magnetic field measurements in a sense that the observable needed in the former, namely the frequency, is the physical quantity that can be measured with the highest precision. In fact, very high resolving power ( $10^6$ ) and accuracies (up to  $10^{-8}$ ) are routinely achieved for nuclides located quite far from the line of  $\beta$ -stability. Such high resolving power made it possible in 1991 [42], for the first time in the history of mass spectrometry, to resolve nuclear isomers from their ground state ( $^{84}\text{Rb}^m$ ) and to determine their excitation energies. Another beautiful demonstration of complementarity between mass spectrometry and nuclear spectroscopy was given in [2004Va07] for  $^{70}\text{Cu}$ ,  $^{70}\text{Cu}^m$  and  $^{70}\text{Cu}^n$ , where in the same work the masses of the three isomers were determined directly by mass spectrometry, while the excitation energies were measured by  $\beta\gamma$  spectroscopy. Typically, the precision can reach 100 eV or better (60 eV for the difference between  $^6\text{He}$  and  $^7\text{Li}$  at TITAN-Vancouver [2012Br03]). Even the most exotic nuclides, such as  $^{11}\text{Li}$  (8.75 ms) or  $^{74}\text{Rb}$  (64.78 ms), were measured with precisions of 600 eV and 4 keV with the TITAN-Vancouver [2008Sm03] and ISOLTRAP-Cern [2007Ke09] facilities, respectively.

In earlier evaluations, we found it necessary to multiply uncertainties from some groups of mass-spectrometric data [43] with discrete factors ( $F=1.5, 2.5$  or  $4.0$ ) following the partial consistency factors  $\chi_n^p$  we found for these groups (see Section 5.2). Such a treatment is not necessary for most of the Penning trap results, which almost all have  $F=1$ .

### 6.2.2 Double-focussing mass spectrometry

Classical double-focussing mass spectrometry was performed on-line at ISOLDE-CERN to measure masses of nuclides far away from the valley of stability. In these experiments, a relationship between three masses was established. These mass-triplet measurements, in which undetectable systematic effects could build-up in large



deviations when the procedure is iterated [1986Au02], could be recalibrated with the help of the Penning trap measurements. Recalibration was automatically obtained in the evaluation, since each mass-triplet was originally converted to a linear mass relation among the three nuclides, allowing both easy application of least-squares procedures, and automatic recalibration. In the present adjustment of data, most of the 181 original data, performed in the 80's, are now outweighed, except for the most exotic (and thus the most interesting) ones. There are still five of them that contribute to the present adjustment, essentially for some very exotic nuclides:  $^{91}\text{Rb}$  for 12% of the determination of its mass,  $^{95}\text{Rb}$  (49%),  $^{144}\text{Cs}$  (20%). In Table I, the relevant equations are normalized to make the coefficient of the middle isotope unity, so that they read e.g.

$$^{97}\text{Rb} - (0.490 \times ^{99}\text{Rb} + 0.511 \times ^{95}\text{Rb}) = 350 \pm 60 \text{ keV},$$

$$^{145}\text{Cs} - (0.392 \times ^{148}\text{Cs} + 0.608 \times ^{143}\text{Cs}) = -370 \pm 90 \text{ keV},$$

(the  $^{148}\text{Cs}$  symbol represents the mass excess of nuclide  $^{148}\text{Cs}$  in keV). The other two coefficients are three-digit approximations of

$$\frac{A_2}{A_3 - A_1} \times \frac{A_2 - A_1}{A_3} \quad \text{and} \quad \frac{A_2}{A_3 - A_1} \times \frac{A_3 - A_2}{A_1}.$$

We took  $A$  instead of  $M$  in order to arrive at coefficients that do not change if the  $M$ -values change slightly. The difference is, however, unimportant.

### 6.2.3 Radio-frequency mass spectrometry

The Orsay Smith-type mass-spectrometer MISTRAL, which was also connected to ISOLDE, had performed quite precise measurements of very short-lived light nuclides, before the Penning traps could cover all the possibilities that were offered by a transmission mass spectrometer. There are eight of the measurements performed with MISTRAL that are still used in this evaluation for the determination of the masses of  $^{26}\text{Ne}$ ,  $^{26,27,28,29}\text{Na}$  and  $^{29}\text{Mg}$ .

### 6.2.4 Classical time-of-flight

Mass measurements by the time-of-flight mass-spectrometry technique, firstly at SPEG (GANIL) and TOFI (Los Alamos), later at Michigan State University (MSU), also apply to very short-lived nuclides, due to instant measurements, but the precisions are much lower than those obtained with MISTRAL. Masses of almost undecelerated fragment products, coming from thin targets bombarded with heavy ions [44] or high energy protons [45], are determined from a combination of magnetic deflection and time-of-flight measurements. Nuclides in an extended region in  $A/Z$  and  $Z$  are analyzed simultaneously. Each individual ion, even if very short-lived

( $1\mu\text{s}$ ), is identified and has its mass measured. In this way, mass values with precisions of ( $3 \times 10^{-6}$  to  $5 \times 10^{-5}$ ) can be obtained for a large number of neutron-rich nuclides of light elements, up to  $A = 70$ . One difficulty in such experiments is that the obtained value can apply to an isomeric mixture where all isomers with half-lives of the order of, or longer than the time of flight (about  $1\mu\text{s}$ ) may contribute. The limited resolving power, around  $10^4$ , and cross-contaminations can cause significant shifts in masses. The most critical part in these experiments is the calibration, since it is frequently from an empirically determined function, which, in several cases, had to be extrapolated rather far from the calibrating masses. It is possible that, in the future, a few mass-measurements far from stability may provide better calibration points, thus allowing a re-analysis of the concerned data. Such recalibrations require analysis of the raw data and cannot be done by the evaluators. With new data available from other methods, which allow detailed comparisons to be made, we observed strong discrepancies for these groups, and had to increase the associated partial consistency factor to  $F = 1.5$ .

### 6.2.5 Cyclotron time-of-flight

Cyclotrons offer very long time-of-flight basis, yielding high resolving power for ions living longer than  $50\mu\text{s}$ . The accelerator radio-frequency is taken as reference to ensure a precise time determination, but this method implies that the number of turns that an ion has to make inside the cyclotron, should be known exactly. This was achieved successfully at SARA-Grenoble in the mass measurement of  $^{80}\text{Y}$ . Experiments performed at GANIL with the CSS2 cyclotron, could not determine the exact number of turns. In the first experiment around  $^{100}\text{Sn}$  [1996Ch32], a careful simulation was done instead. In the second experiment on  $^{68}\text{Se}$ ,  $^{76}\text{Sr}$ ,  $^{80}\text{Sr}$  and  $^{80}\text{Y}$  [2001La31], a mean value of the number of turns was experimentally determined for the most abundant species only, which mainly involved the calibrants. Penning traps measurements at the CPT-Argonne, JYFLTRAP-Jyväskylä and ISOLTRAP revealed that this method suffered from serious systematic errors. Later, improved measurements at GANIL with the CSS2 cyclotron [2008Go23] were in better agreement with the Penning trap data.

### 6.2.6 Multi-Reflection Time-of-Flight Mass Spectrometer

A new type of instrument, called Multi-Reflection Time-of-Flight mass spectrometer (MR-TOF), has seen the light recently at major nuclear physics facilities. Three MR-TOF's, operated at ISOLDE-CERN, RIBF-RIKEN and at the GSI facility, begin to produce interesting results, which are included in the present evaluation.

Several other MR-TOF's are under construction at other facilities such as TRIUMF, Argonne and SPIRAL2. The MR-TOF mass measurement is based on time-of-flight. Like storage rings, it aims at extending the flight path by reflecting ions back and forth in a static electric field. With this method, a relative mass precision of  $10^{-7}$  is routinely achieved in typically 10 milliseconds. Remarkable results have been achieved recently at ISOLTRAP using the MR-TOF: it was possible to reach the most exotic nuclei in the light mass region,  $^{52,53,54}\text{Ca}$  [2013Wi06] and  $^{52,53}\text{K}$  [2015Ro10], as well as in the heavy mass region,  $^{131}\text{Cd}$  [2015At03]. A precision of 10 keV has been obtained for the less exotic nuclide ( $^{52}\text{Ca}$ ), and a precision of 110 keV for the most exotic one ( $^{53}\text{K}$ ) with half-life of 30 ms.

For mass determination with MR-TOF, the general relation between mass-to-charge ratio ( $m/q$ ) and time-of-flight  $t$  is [2013Wi06]:

$$t = \alpha \sqrt{m/q} + \beta, \quad (8)$$

where  $\alpha$  and  $\beta$  are constants related to the experimental set-up and are the same for the ion of interest and for the reference ions.

At RIKEN and GSI, the  $\beta$  parameter can be determined independently, thus only one reference nuclide is needed for mass calibration. In this way, the ratio of time-of-flight between the ion of interest and the reference ion is used to extract the linear equation, as it is the case for Penning trap measurement (see Appendix C p. 030002-43). Results from RIKEN ([2013It01], [2016Sc.A]) and GSI ([2015Di03]) are presented in this way in the present evaluation.

However, at ISOLDE-CERN, two reference masses are used to determine the mass of interest, and the so-called  $C_{tof}$  method is used. When using this method, we express the linear equation in term of absolute mass.

At first, the two constants ( $\alpha$ ,  $\beta$ ) are extracted from the two reference equations:

$$\begin{aligned} t_1 &= \alpha \sqrt{(m/q)_1} + \beta \\ t_2 &= \alpha \sqrt{(m/q)_2} + \beta, \end{aligned}$$

where  $(m/q)_1$  and  $(m/q)_2$  are the mass-to-charge ratios of reference 1 and reference 2, while  $t_1$  and  $t_2$  are their time-of-flights. We thus obtain:

$$\begin{aligned} \alpha &= \frac{t_1 - t_2}{\sqrt{(m/q)_1} - \sqrt{(m/q)_2}} \\ \beta &= t_1 - \frac{t_1 - t_2}{\sqrt{(m/q)_1} - \sqrt{(m/q)_2}} \sqrt{(m/q)_1}. \end{aligned}$$

To extract the mass of interest,  $\alpha$  and  $\beta$  are replaced in Eq. (8) and the mass can be written:

$$m/q = C_{tof} \Delta_{ref} + \frac{1}{2} \Sigma_{ref}, \quad (9)$$

where  $C_{tof}$ ,  $\Delta_{ref}$  and  $\Sigma_{ref}$  are defined by:

$$\begin{aligned} C_{tof} &= \frac{2t - t_1 - t_2}{2(t_1 - t_2)}, \\ \Delta_{ref} &= \sqrt{(m/q)_1} - \sqrt{(m/q)_2}, \\ \Sigma_{ref} &= \sqrt{(m/q)_1} + \sqrt{(m/q)_2}. \end{aligned}$$

The mass uncertainty is calculated from the uncertainty  $\sigma_c$  of coefficient  $C_{tof}$  and the reference mass uncertainties  $\sigma_1$  and  $\sigma_2$ :

$$\sigma^2 = (m/q) \left\{ \left( C_{tof} - \frac{1}{2} \right)^2 \frac{\sigma_1^2}{(m/q)_1} + \left( C_{tof} + \frac{1}{2} \right)^2 \frac{\sigma_2^2}{(m/q)_2} + 4 \left( \sqrt{(m/q)_2} - \sqrt{(m/q)_1} \right)^2 \sigma_c^2 \right\}.$$

### 6.2.7 Storage-ring time of flight

Similarly, a long flight path can be obtained in a storage ring, which is operated in the mode of isochronous mass spectrometry (IMS). The first set-up of this type was operated at GSI-ESR at Darmstadt. The precision of the measurements could be as good as 90 keV even for nuclides quite far from stability. Recently, [2016Kn03] reanalyzed some data from two earlier experiments and reported some new masses at GSI-ESR. For some of the newly reported nuclides, only two events were recorded. The results show the potential of the IMS method. However the systematic uncertainty is rather large and eight results from that work appear in Table C. The Cooler Storage Ring for experiment (CSRe) at the IMP-LANZHOU is the second spectrometer for IMS mass measurements. Precision better than 10 keV has been achieved. The isomer of  $^{52}\text{Co}$  has been resolved from the ground state, demonstrating the excellent resolving power [2016Xu10].

### 6.2.8 Cooled beam cyclotron frequency

Storage rings could also be used with cooled beams to measure the cyclotron frequency as has been demonstrated since 1997 at the GSI-ESR storage ring, with precisions sometimes as good as 12 keV. Many of the measured nuclides belong to known  $\alpha$ -decay chains. Thus, the available information on masses for proton-rich nuclides is considerably extended.

It must be mentioned that in the first group of mass values as given by GSI authors [2000Ra23], several data could not be accepted without changes. The reason was that in the determination of the mass values they had to combine  $\alpha$ -decay energies between two or more of the occurring nuclides. Evidently, these energies could not be included without corrections in our calculations, where they would be again combined with these  $Q_\alpha$  values. Remarks are added to the data in Table I in order to warn for such cases. Fortunately, this group of data is only of historical interest since they were superseded by more recent high-precision results [2005Li24] using the same

instruments. A wealth of high-quality data were published recently using this technique, see e.g. [2012Ch19] and references therein.

### 6.2.9 Isomeric mixtures

As stated above, many mass-spectrometric results yield an average mass value  $M_{exp}$  for a mixture of isomers. Here, we use a special treatment for the possible mixture of isomers (see Appendix B p.030002-39) and information about these changes are duly explicated in remarks accompanying these data.

The mass  $M_0$  of ground state can be calculated if both the excitation energy  $E_1$  of the upper isomer and the relative production rates of the isomers are known. But often this is not the case. If  $E_1$  is known but not the production ratio, one must assume equal probabilities for all possible relative intensities. In the case of one excited isomer, the estimated mass for  $M_0$  becomes  $M_{exp} - E_1/2$ , and the part of the error due to this uncertainty is  $0.29E_1$  (see Appendix B, Section B.4, p.030002-42). This policy was defined and tested first for the GSI-ESR cooled beam cyclotron frequency data and was discussed with the authors of the measurements. In 15 cases, more than two excited isomers contribute to the measured line.

A further complication arises if  $E_1$  is not known. In such a case, we have to make the best possible estimate for  $E_1$ . As always this estimated value is flagged with ‘#’ This, in addition to questions related to  $\alpha$ -decay chains involving isomers, was a reason for us to consider the matter of isomers with even more attention. Part of the results of our estimates are incorporated in the NUBASE evaluation. In estimating the  $E_1$  values, we first look at experimental data possibly giving lower limits: e.g. if it is known that one of two isomers decays to the other; or if  $\gamma$  rays of known energy occur in such decays. If not, we try to interpolate between  $E_1$  values for neighboring nuclides that can be expected to have the same spin and configuration assignments (for odd  $A$ : isotones if  $Z$  is even, or isotopes if  $Z$  is odd). If such a comparison does not yield useful results, indications from theory were sometimes accepted, including upper limits for transition energies following from the measured half-lives. Values estimated this way were provided with somewhat generous errors, dutifully taken into account in deriving final results.

In several of these measurements, an isomer can only contribute if its lifetime is relatively long (hundreds of milliseconds or longer). However, half-life values given in NUBASE are those for neutral atoms. For bare nuclides, where all electrons are fully stripped from the atom, the lifetimes of such isomers can be considerably longer, since the decay by conversion electrons is switched off. The reported mass measurements [2005Li24] of the 580 ms  $^{151}\text{Er}^m$  isomer at  $E_1=2586.0$  keV excitation energy and

and the 103 ms  $^{117}\text{Te}^m$  isomer at  $E_1=296.1$  keV are two examples.

Considering the isomeric mixtures and combining experimental data from decay spectroscopy and from mass spectrometry can provide valuable information for the atomic masses. This can be demonstrated in the following example. Masses of the nuclides along the  $\alpha$ -decay chain  $^{206}\text{Ac}$ - $^{202}\text{Fr}$ - $^{198}\text{At}$ - $^{194}\text{Bi}$ - $^{190}\text{Tl}$  were deemed unknown in AME2012, while they were considered to be known in AME2003. For these nuclides, two long-lived states exist with high and low spin, respectively, and two  $\alpha$ -decay chains are established in parallel. The excitation energies of the isomeric states are unknown. In AME2003, their masses were determined by storage ring mass spectrometry [2003Li.A], [2005Li24], where the mixture of two states were assumed and the corrections implemented. Later the mass of  $^{190}\text{Tl}$  in its high-spin state was measured with unambiguous assignment of the state from decay spectroscopy. The result was included in AME2012 as private communication [2012Bo.A] (published later as [2014Bo26]). This state was assumed to be the excited isomer, but the excitation energy was unknown. The excitation energy of  $^{190}\text{Tl}$  was estimated from TNN to be 90#(50#) keV in AME2012, and then the masses of these five nuclides in their ground state were deemed unknown. By using resonance ionization laser ion source technique, a specific state can be selected, and it can be identified through decay spectroscopy. In this way, the ground state mass of  $^{198}\text{At}$  was unambiguously measured by ISOLTRAP [2013St25]. The excitation energy of  $^{190}\text{Tl}^m$  was determined to be 89(12) keV for the first time, which agrees with the estimated value. In AME2016, the masses of all five nuclides connected via  $\alpha$  decays are now being experimentally known.

### 6.3 Masses of unbound nuclides

Presently, many nuclides beyond the driplines can be accessed in the light mass region. They can decay via direct proton or neutron emission. The half-lives of these unbound nuclides are too short for them to acquire their outer electrons (which takes around  $10^{-14}$  s), and to form atoms. However, we still convert their masses to “atomic masses” so we can treat them consistently with other nuclides. It is experimentally challenging to study these unbound nuclides far from stability: only very few events can be observed. Frequently, theoretical calculations are required to extract their properties from the experimental data.

On the proton rich side, resonant states could be formed due to the Coulomb barrier. There are different approaches to study these states: transfer reaction with missing mass spectrum, proton scattering, and complete kinematic measurement with invariance mass spec-

trum. For a broad resonant state, the definition of the resonance energy and width is not unique. For example,  $^{15}\text{F}$  was studied in resonant elastic scattering using a thick  $\text{CH}_4$  gas target in inverse kinematics with a  $^{14}\text{O}$  beam [2004Go15]. The proton-decay energy of  $^{15}\text{F}$  was obtained to be  $1.29_{-0.06}^{+0.08}$  MeV from the energy at which the magnitude of the internal wave function is a maximum, or  $1.45_{-0.10}^{+0.16}$  MeV where the elastic scattering cross section is maximum, corresponding to a phase shift of  $\delta = \pi/2$ . Since the latter value is consistent with those obtained in transfer reaction studies, it is adopted in our evaluation.

Some single-proton resonant states can be accessed and studied in two-proton decay experiments. For example, the 1p-decay energies of 1560(130) and 2850(40) keV for the ground and first excited states in  $^{15}\text{F}$  [2004Le12] are well reproduced in the angular-correlation studies of two-proton decays of  $^{16}\text{Ne}$  [2008Mu13].

On the neutron-rich side of the nuclear chart, the mass of unbound nuclides can be determined by means of the missing-mass method using transfer reactions (e.g. [2015Ma54]), or with the invariant-mass method using radioactive-ion beams (e.g. [2012Ko43]). Recently, various such beams and improved detection techniques have been impressively developed, which allows new masses of unbound nuclides to be determined.

In the case of neutron-induced reaction, only the centrifugal barrier plays a role in the formation of a given resonant state. Since no barrier exists at all for a s-wave neutron, the observation of asymmetric peak near the threshold is a general feature of spectra obtained in invariant-mass experiments (e.g. [2010Jo06]). This state is usually referred to as a virtual state, which has no definite lifetime and thus differs significantly from a real resonance state. The virtual state can be characterized by the s-wave neutron-nucleus scattering length; its eigen energy is approximately  $\hbar^2/2\mu a_s^2$ , where  $\mu$  is the reduced mass and  $a_s$  is the scattering length.

#### 6.4 Isobaric Analog states (IAS)

The concept of isospin was introduced by Heisenberg [46] and developed by Wigner [47] to describe the charge independence of nuclear forces. This concept is widely used in particle and nuclear physics. Within the isospin formalism, a nucleus composed of  $Z$  protons and  $N$  neutrons has a fixed isospin projection of  $T_z = (N - Z)/2$ , while all states in the nucleus can have different total isospins  $T \geq |T_z|$ . In other words, states of a given  $T$  can occur in a set of isobaric nuclei with  $T_z = T, T - 1, \dots, -T$ . These states with the same  $T$  and  $J^\pi$  are called Isobaric Analog States (IAS). A set of IASs with fixed  $A$  and  $T$  are believed to have very similar nuclear structure properties and to be energetically degenerate in the framework of isospin symmetry. Their rela-

tive masses can be used to explore the charge symmetry and charge independence of the nuclear interaction via the Isobaric Mass Multiplet Equation (IMME) [48], and with calculations of the Coulomb Displacement Energy (CDE) (see for example [32], and references therein).

As in AME2012, IASs that are determined via external relations were evaluated. In some cases, one IAS can be involved in a local network thus influencing other masses. Such an IAS was included in the present evaluation, although its excitation energy may be determined mainly through an internal relation. An example is  $^{48}\text{Mn}^i$ , which is connected to  $^{47}\text{Cr}$  through proton decay, thus building a loop when its internal transition is included.

#### 6.5 Proton and $\alpha$ decays

In some cases, proton-decay energies can be estimated from proton-decay half-lives. Estimates for the following nuclides can thus be obtained as:

Nuclide	$T_{1/2}$	$S_p$ (keV)	Adopted $S_p$
$^{64}\text{As}$	$40 \pm 30$ ms	$> -100$	$-100\# \pm 200\#$
$^{68}\text{Br}$	$< 1.5$ $\mu\text{s}$	$< -500$	$-500\# \pm 250\#$
$^{73}\text{Rb}$	$< 30$ ns	$< -570$	$-570\# \pm 200\#$
$^{81}\text{Nb}$	$< 44$ ns	$< -600$	$-710\# \pm 500\#$

These limits were used as a guide to obtain estimates for the masses of those nuclides.

These results are important for two main reasons. Firstly, knowledge of proton separation energies just beyond the proton drip line is quite valuable in estimating the mass values for nuclides for which no experimental data are available. Secondly, there are several cases where proton-decay energies from both members of an isomeric pair were measured, so one can determine the excitation energy of a particular isomer. In addition, the lifetime of a proton-emitting nuclide is sensitive to the orbital angular momentum value  $l$  carried by the proton and this can be used in turn to obtain reliable information about the spins and parities of the parent and daughter states. This feature is even more valuable, when  $\alpha$  decays of both members are observed. Combination of long  $\alpha$ -decay chains with proton decays offers a view of extended regions of the chart in the neighborhood of the proton drip-line. These studies showed that several decays that were earlier assigned to ground states actually belong to excited isomers. Also, these measurements are found to yield good values for the excitation energies of the isomers among the descendants. We usually followed the judgment of the authors, including their recommendations about the final levels fed in those  $\alpha$  decays.

Often in  $\alpha$ -decay studies of odd- $N(Z)$  and odd-odd nuclides, the level fed directly by the  $\alpha$  particle is not known. A comprehensive investigation that we per-

formed some time ago suggested that, in most cases, when the decay does not go directly to the ground state, the final level is relatively close to the ground state. In such cases, we adopted the policy of accepting the measured  $E_\alpha$  as feeding the ground state, but assigning a special label to indicate that a close-lying excited level may also be fed. This label, which is not given in Table I, will indicate to our computer program that the uncertainty, after possible pre-averaging of data of the same kind (also given in Table I), is to be increased to 50 keV.

The existence of proton-decay branches, as mentioned above, provided sufficient arguments to omit the mentioned label in several cases. One has also to be careful with the use of this label if mass-spectrometric results with a precision of about 50 keV or better are known for the parent and daughter nuclides. Comparison with theoretical models may also suggest dropping the mentioned above label; or conversely to not accept a reported  $\alpha$ -decay energy.

In some cases, TMS estimates and theoretical predictions of  $\alpha$ -decay energies indicate that the excitation energy  $E_1$  of the final level may be much higher. Then, an estimate for the excited level energy (provided with a generous error) is added as an input value.

In regions where the Nilsson model for deformed nuclides applies, it is expected that the most intense  $\alpha$  transition connects parent and daughter levels that have the same quantum numbers and configurations. In such a case, adding an estimate for  $E_1$  is attractive. Frequently, the energy difference between the excited and ground states can be estimated by comparisons with the energy differences between the corresponding Nilsson levels in nearby nuclides.

For nuclei with  $A > 190$ , as well as for proton-rich nuclei far from the line of stability,  $\alpha$  decay is the main decay mode providing information about atomic masses. Most measurements involve position-sensitive silicon detectors, which require careful energy calibration. Such calibrations usually use the recommended  $E_\alpha$  values evaluated by Rytz [1991Ry01]. The recent development in mass spectrometry allowed independent mass measurements in this region of the nuclear chart using Penning Traps. Such measurements have already been carried out for several No and Lr nuclides [2010Mi.A] and their results were already included in AME2012.

### 6.5.1 Particle energy vs. decay energy

Unfortunately, some authors misuse the meaning of particle and decay energies. Energy values are referred to by some authors as the particle energy  $E$ , while others quote it as decay energy  $Q$ . Actually, the decay energy is the sum of kinetic energies of the emitted particle and the recoiling daughter nuclide. In general, the  $\alpha$  particle car-

ries about 97-98% of its  $Q$  value and the recoiling nuclide accounts for about 2-3%. In the literature one can find too many cases of confusion, especially in proton-decay experiments where  $Q_p$  and  $E_p$  are numerically closer to each other. Sometimes, the confusion could be resolved through a meticulous inspection of the paper and a discussion with the authors. However, ambiguities still remain in many cases.

### 6.5.2 Recalibration of alpha- and proton-decay energies in implantation experiments

In experiments where the  $\alpha$ -emitting nuclei are implanted in a silicon detector, both the  $\alpha$  particle and the recoiling daughter nuclide deposit energies in the detector. Often authors make the simple assumption that only the  $\alpha$ -particle energy is measured in the detector. While in similar cases of proton decays, it is often considered that both the proton and the heavy recoil are detected at the same time. Neither of these statements is correct:  $\alpha$  particles and protons with energies of a few MeV have almost 100% detection efficiency, which is not the case for the heavy recoiling nuclides, where only part of the recoiling energy contributes to the signal.

This effect has been discussed in Ref. [2012Ho12], where approximately 28% of the recoil energy contributes to the signal. In that experiment, the recoil energy of the 11.65 MeV  $\alpha$ -particle line of  $^{212}\text{Po}^m$ , which was used for calibration, is 224 keV, whereas the recoil energy of a superheavy nucleus with mass number 292 is only 162 keV for the same  $\alpha$  energy. Thus the difference of the recoil energies which contribute to the signals is 17 keV, which is larger than the 10 keV energy uncertainty. For this reason, the partial recoil energy of the daughter nuclide has been taken into account in the energy calibration [2012Ho12].

However, not all the experimentalists notice this effect and we need to make our own corrections of the published results. We have developed a procedure [49] to calculate the detection efficiency for heavy nuclides in Si detectors based on Lindhard's integral theory [50], which has been experimentally proven to be reliable [51, 52].

The correction has been done for some of the experimental results included in the present tables. After discussions with the authors of the original publication, the  $\alpha$ -decay energy of  $^{255}\text{Lr}^m$  Ref. [2008Ha31] has been corrected and the difference turns out to be 7(10) keV. The corrected value is used in the current evaluation with a remark given to the relevant data in Table I. Another example is from Ref. [2014De41], where the proton-decay of  $^{69}\text{Br}$  was measured by using  $\beta$ -delayed protons from  $^{20}\text{Mg}$  and  $^{23}\text{Si}$  for the energy calibration. The authors assumed (erroneously) that the recoil energy would be fully recorded at the same time. From our calculations the detection efficiency for the recoil nuclide ( $^{68}\text{Se}$ ) is

about 30% and its neighboring nuclides show similar behavior. Applying the correction, the  $\beta$ -delayed proton-decay energy of  $^{69}\text{Kr}$  changed from 2939 keV to 2916 keV. The difference is 23 keV, comparable with the 22 keV uncertainty. The same correction procedure was also applied to the ground-state proton-decay energy of  $^{69}\text{Br}$  in the same work [2014De41], which changed its proton-decay energy from 641 keV to 631 keV, with uncertainty of 42 keV.

The correction should, in principle, be applied to all implantation  $\alpha$ - and proton-decay experiments of some precision if the recoil effect was not taken into account. This work is not yet complete in the present version of the AME.

Some authors derive a value which they call  $Q_\alpha$  from the measured  $\alpha$ -particle energy by not only correcting for the recoil energy, but also for screening by atomic electrons (see Appendix A p. 030002-38). In our calculations, the latter corrections have been removed.

Finally, some measured  $\alpha$ -particle energies are affected by the coincidence summing between the  $\alpha$  particle that feeds an excited level of the daughter nuclide and the conversion electrons that follow the decay of this level. This is sometimes apparent from the reported  $\alpha$  spectra, since the width of the observed line is larger than that of other ones. In some cases, spurious  $\alpha$  peaks can be observed. When deriving the corresponding  $Q_\alpha$  values, appropriate (small) corrections are made for the escaping X-rays. Those are mentioned in a remark added to such a case.

## 6.6 Decay energies from capture ratios and relative positron feedings

For allowed transitions, the ratio of electron capture in different shells is proportional to the ratio of the squares of the energies of the emitted neutrinos, with a proportionality constant being dependent on  $Z$  [53]. For (non-unique) first forbidden transitions, the ratio is similar, but with a few exceptions. The neutrino energy is determined as the difference of the transition energy  $Q$  and the electron binding energy in the pertinent shell. Especially if the transition energy is not too much larger than the binding energy in, say, the  $K$  shell, it can then be determined rather well from a measurement of the ratio of capture in the  $K$  and  $L$  shells.

The non-linear character of the relation between  $Q$  and the capture ratio introduces two problems. In the first place, a symmetrical error for the ratio is generally transformed in an asymmetrical one for the transition energy. Since our least-squares fit program cannot handle them, we have symmetrized the probability distribution by considering the first and second momenta of the real probability distribution (see NUBASE2016, Appendix A, p. 030001-16). The other problem is related to averaging

of several values that are reported for the same ratio. Since AME1993, our policy is to average the capture ratios, and calculate the decay energy from that average. An example is  $^{139}\text{Ce}(\varepsilon)^{139}\text{La}$  (see p. 030002-211), where 10 results were averaged and the individual values given in the associated remarks. In this procedure we used the best values [53] of the proportionality constant. We also recalculated the older decay energies using the new value for this constant.

The ratio of positron emission and electron capture in the transition to the same final level also depends on the transition energy. It is well known for allowed and not much delayed first forbidden transitions. Thus, the transition energy can be derived from the measured positron intensity to a given level, rather than from the positron spectrum end-point (e.g.  $^{109}\text{Cd}(\varepsilon)^{109}\text{Ag}$ , p. 030002-170). In the case of positron decay, one must remember that it can only occur when the transition energy exceeds  $2m_e c^2 = 1022$  keV. However, in many cases the level fed by positrons is also fed by  $\gamma$ -rays coming from higher levels that are fed by electron capture. Determination of the intensity of this *side* feeding is often difficult. Cases exist where such feeding occurs by a large number of weak  $\gamma$ -rays that can be easily overlooked (the *pandemonium* effect [54]). Then, the reported decay energy may be much lower than the real value. In judging the validity of experimental data, we kept this possibility under consideration.

Total Absorption Spectrometry (TAS) has been applied to overcome the *pandemonium* effect [54]. In some cases the ratio of positron emission and electron capture is measured using TAS, e.g.  $^{103}\text{Sn}$  [2005Ka34]. For this case we adopted the value reported by the authors.

## 6.7 $Q_\beta$ far from $\beta$ -stability

Presently, the mass surface for nuclides far away from the valley of  $\beta$ -stability is observed to be located much higher than was previously believed. This is largely due to the underestimation of the  $Q_\beta$  decay energies, which were measured in the past using the end-point energy method. See the discussion in AME2012, p. 1317.

The deduced higher values of atomic masses for exotic nuclides in the present work will have important consequences for nuclear astrophysics and nuclear energy applications, as discussed in Ref. [55].

To conclude, for nuclei very far from the valley of stability,  $Q_\beta$  results from end-point measurements should be treated with caution. In such cases, data available from Penning traps and/or storage rings facilities should always be given priority.

## 6.8 Superheavy nuclides

The search for superheavy elements (SHE) and elucidation of their properties is one of the prominent areas

of modern nuclear physics research. In the last several years, the nuclear chart was extended impressively in the heaviest mass region up to the element with atomic number  $Z = 118$ . However, the mass surface built with the available data is still rough (see Part II, Fig. 9 and also Fig. 26, p. 030003-243 and 030003-260).

**Names and symbols** At the completion of AME2016, SHE up to  $Z = 118$  were officially named by The Commission on Nomenclature of Inorganic Chemistry of the International Union of Pure and Applied Chemistry (IUPAC) [56]:

113	Nihonium	(Nh),
115	Moscovium	(Mc),
117	Tennesine	(Ts), and
118	Oganesson	(Og).

We were not able to include the new names in AME and NUBASE, but instead we used the provisional symbols Ed, Ef, Eh, and Ei for elements 113, 115, 117, and 118, respectively.

**Experimental methods** Since  $\alpha$  decay is the dominant decay mode in the region of superheavy nuclides, knowledge of masses of SHE is most often obtained from the measured  $\alpha$ -decay energies within a chain that reaches a nuclide with known mass. Position and time-correlated  $\alpha$ -decay and Spontaneous Fission (SF) spectroscopy measurements of SHE continue to provide precious information about their properties. However, it often happens that  $\alpha$  chains end up with a nuclide decaying only by spontaneous fission, offering no link to known masses. For example, the SF decay of  $^{266}\text{Sg}$  does not allow to determine the mass of the doubly magic nuclide  $^{270}\text{Hs}$ . In order to support the discovery of new elements, an indirect method can be applied, where different nuclear reactions are used to produce the daughter nuclide. This was the case in the discovery of the new elements  $^{293}\text{Eh}(\text{Ts})$  and  $^{289}\text{Ef}(\text{Mc})$ , see Ref. [57] and references therein. However, the new analysis of all available data in Ref. [2016Fo16] provided evidence against the proposed cross-reaction link [57] between the  $\alpha$ -decay chains associated with those two nuclides. New studies would be needed to resolve the discrepancies.

A very important development in this mass region was the first direct mass measurements [2010Dw01] of several isotopes of No ( $Z = 102$ ) and Lr ( $Z = 103$ ) by the SHIPTRAP facility at GSI. Those results provided anchor points for atomic masses in this remote region of the nuclear chart. In general, the newly measured masses agree reasonably well with those deduced from known  $Q_\alpha$  values of long  $\alpha$  chains, thus giving confidence not only about the reliability of masses for SHE reported in [2010Dw01], but also the treatment and policies used in

our work. In AME2016 we included new results from such direct mass measurements for the lighter  $^{241,243}\text{Am}$ ,  $^{244}\text{Pu}$  and  $^{249}\text{Cf}$  nuclides [2014Ei01]. However, we have to mention here the disagreement in the  $^{249}\text{Cf} - ^{241}\text{Am}$  mass difference between the Penning trap data and the value deduced using the decay  $Q_\alpha$  values, as discussed in Section 7.10 (p. 030002-36).

**Alpha decay of superheavy nuclides** For even-even nuclides, the strongest (favored) decays connect the parent and daughter ground states. They are directly related to the  $Q_\alpha$  values. As a result, masses determined this way are quite reliable. Unfortunately, some of the nuclides are prone to spontaneous fission decay, thus limiting the number of reliable cases.

For many odd-A nuclides, especially for odd-odd ones, the assignments are frequently complicated. In the region of deformed nuclides,  $\alpha$  decays preferentially connect levels with the same  $J^\pi$  and configurations, and as a consequence the daughter nuclei are often produced in excited states with unknown excitation energies  $E_1$ . Thus, in order to find the corresponding mass difference, we have to estimate these  $E_1$  values. For somewhat lighter nuclides, one may estimate them from known differences in excitation energies for levels with the same Nilsson assignments in neighboring nuclides. But such information is lacking in the SHE region under consideration. Instead, one might consider using values obtained theoretically [58]. We have not done so. However, we have used such theoretical values as a guide, choosing values in such a way that diagrams of  $\alpha$  energies and the mass surface looked smooth. Helpful for this purpose were the experimental  $\alpha$ -decay energies for  $Z = 112, 114$  and  $116$ , especially for the even-even nuclides among them. This is especially true near sub-shell closures, since the favored alpha decay occurs between states that have the same quantum numbers and configurations.

The presence of excited, long-lived isomers can also lead to severe complications. While many dedicated  $\alpha$ - $\gamma$  coincidence studies have been performed for nuclides in the light actinide region, such spectroscopy needs to be extended to the heavier nuclides. In the last several years new results were published in the  $Z = 102 - 104$  region, which resolved some of the ambiguities. However, high quality data are still in demand and such studies would be very beneficial to future mass determination of SHE.

A weak  $\alpha$ -decay branch was observed in the decay of  $^{262}\text{Sg}$  [2010Ac.A], which allowed experimental determination of the mass of  $^{270}_{108}\text{Ds}$ , the heaviest nuclide that has an experimental mass value in AME2016. The new data allowed to establish unambiguously the existence of a significantly deformed sub-shell gap at  $N = 162$  and  $Z = 108$ . This gap appears to be much larger than the

one at  $N = 152$  and  $Z = 100$ .

Our policy in this high- $A$  region, where the  $\alpha$ -decay energies often spread too much, is to adopt the highest  $\alpha$ -decay group as gs-gs transition. The reason is that, even if this group is formed due to  $\alpha$ -electron summing, it is still the closest one to the real gs-gs  $Q$  value.

An interesting case is the determination of the mass of  $^{265}_{106}\text{Sg}$ . This mass was considered as experimentally known in the AME2003 mass table (and was then the heaviest nuclide with known mass), derived from the highest  $\alpha$ -decay group  $E_\alpha = 8940 \pm 30$  keV of [1998Tu01] and adopted as gs-gs transition. With more events [2012Ha05], the status of  $^{265}\text{Sg}$  has been changed and the former  $\alpha$ -decay group assigned to the neighboring isotope  $^{266}\text{Sg}$  was reassigned to the  $^{265}\text{Sg}^m$  state. In the present evaluation we use the strongest group, which may be the unhindered transition, assuming this transition goes to one excited state in the daughter nuclide  $^{261}\text{Rf}$  with unknown energy. This energy is estimated from the trends in the neighboring nuclides (TNN). So, the mass of  $^{265}\text{Sg}$  is now estimated rather than experimental as in AME2003, although the mass value doesn't change much.

With exception of the nuclide  $^{278}_{113}\text{Ed}(\text{Nh})$ , nuclides with atomic number from 113 to 118 are produced by the "hot fusion" method, decaying by  $\alpha$  emission to fissionable nuclides whose masses are unknown experimentally, thus forming a floating island with none of the nuclides having known mass.

## 7 Special cases

Special cases have been discussed in the AME series to highlight the issues raised in the evaluation and to call for more efforts to solve them. Some of the special cases discussed in AME2012 have been solved so we removed them from the current list. Meanwhile some new cases have been added.

### 7.1 $^3\text{H}$ - $^3\text{He}$ atomic mass difference

The  $\beta$ -decay energy of  $^3\text{H}(\beta^-)^3\text{He}$ , which can be deduced from the difference between the  $^3\text{H}$  and  $^3\text{He}$  atomic masses, is very important for neutrino-mass experiments that analyze the shape of the tritium  $\beta$ -decay spectrum near its end-point energy. Because of the significance in the determination of the neutrino mass, the  $Q_\beta$  of  $^3\text{H}$  has been measured by many groups using different methods. About thirty years ago, one of us (G.A.) and colleagues evaluated all of the significant experimental results available at that time on the  $^3\text{H}$ - $^3\text{He}$  mass difference [1985Au07]. In that evaluation, the methods fell into three categories: mass doublet measurements, tritium  $\beta$ -decay measurements in magnetic spectrometers and in implanted detectors. It was found that the data

within each group are consistent with each other, but there were notable discrepancies among the groups.

In the last three decades, Penning trap mass spectrometers have been developed intensively and now dominate mass measurements with the highest precision. Before AME2012, the  $\beta$ -decay measurements always played the most important role in determining the  $Q_\beta$  of  $^3\text{H}$ . In AME2012, the Penning trap results contributed almost as much as the  $\beta$ -decay measurements in this case, thanks to the high-precision results from the SMILETRAP group [2006Na49]. These results were in strong conflict with earlier results from the UW (University of Washington) Penning trap (Seattle group) [1993Va04]. The latter reported that the masses of  $^3\text{He}$  and  $^3\text{H}$  were determined with uncertainties of 1 nu and 1.5 nu, respectively. Thus the  $Q_\beta$  of  $^3\text{H}$  was deduced with a precision of 1.7 eV by this method. The results from [1993Va04] were used in the AME1993 with the originally published values. Later, some serious systematic errors in this measurements were discovered. After discussion with the authors, the result for  $^3\text{H}$  was temporarily discarded. The value for  $^3\text{He}$  was corrected by 3 nu, i.e. 3 sigma away from the original value, and used in AME2003 and AME2012. The results from [2006Na49] supports the correction.

Recently, the  $^3\text{H}$  and  $^3\text{He}$  atomic masses were measured by the Florida State University (FSU) group with precisions of 0.19 nu, as reported in [2015My03]. Because the identical procedures and the same reference ion were used in the mass measurements for these two nuclides, it was concluded that all of the important systematic uncertainties should be cancel. The resulting uncertainty of 0.07 eV is thus much smaller than that for the individual masses.

However, the Seattle group analyzed the results collected in earlier experiments and published the  $^3\text{He}$  mass with an uncertainty of 0.043 nu [2015Za13]. This result still disagrees by 3.3 times the sum of their final uncertainty with the FSU result [2015My03]. In the Seattle experiment, carbon ions were used as reference, whose mass is quite different from the measured nuclide. Thus, their result might be vulnerable to undiscovered systematic errors so is provisionally not used in the present evaluation. In the FSU measurement, the  $\text{HD}^+$  ion was used as the reference, which has similar mass with the measured nuclide. The  $^3\text{H}$ - $^3\text{He}$  mass difference is usually more robust than the absolute mass values, since both nuclides are exposed to the same experimental conditions. This robustness was proven in the work [1993Va04], where although the absolute mass value for  $^3\text{He}$  has been corrected by 3 nu, the mass difference in the original value agrees with the adopted values in series of AME evaluations.

In AME2016, the results from [2015My03] are used



to determine the masses of  ${}^3\text{H}$  and  ${}^3\text{He}$ . Due to the high precision, all of the other results lost their significance. The recommended values for the  $Q_\beta$  decay energy of  ${}^3\text{H}$  from AME1983 to AME2016 are:

AME evaluation	$Q_\beta$ (eV)	Uncertainty (eV)
AME1983	18594	8
AME1993	18591	1
AME2003	18591.3	1.1
AME2012	18590.6	0.8
AME2016	18592.01	0.07

It should be noted that in AME1983, the definition of the *maintained* Volt differed from later evaluations, as explained in Section 2. But its impact on the  $Q$ -value is just 0.15 eV, much smaller than the quoted uncertainty. Because of its importance, the precise determination of the  ${}^3\text{H}$ - ${}^3\text{He}$  mass difference will continue to attract interest. The former Seattle trap is now operational at the Max-Planck Institute in Heidelberg and will be dedicated to such measurements in the future [2016Ho.A].

## 7.2 ${}^9\text{He}$ and ${}^{10}\text{He}$

The knockout reaction on  ${}^{11}\text{Be}$  has been used to produce  ${}^9\text{He}$  [2001Ch31] and its lowest state has been assigned  $l = 0$ . An upper limit of the s-wave scattering length  $a_s = -10$  fm has been obtained, corresponding to an energy for the virtual state below 0.2 MeV. In [2007Go24], the spectrum of  ${}^9\text{He}$  was studied by means of the  ${}^2\text{H}({}^8\text{He}, p){}^9\text{He}$  reaction. The lowest resonant state of  ${}^9\text{He}$  was found at  $2.0 \pm 0.2$  MeV with a width of 2 MeV and has been identified as a  $1/2^-$  state. For the virtual  $1/2^+$  state, a lower limit  $a_s > -20$  fm has been obtained, which is consistent with the result in [2001Ch31]. This assignment has been questioned in [2010Jo06], where  ${}^9\text{He}$  was studied by using knockout reaction from  ${}^{11}\text{Li}$ . The  ${}^8\text{He}+n$  relative-energy spectrum is dominated by a strong peak-like structure at low energy, which may be interpreted within the effective-range approximation as the result of an s-wave interaction with a neutron scattering length  $a_s = -3.17 \pm 0.66$  fm, thus conflicting with [2001Ch31]. It is argued that the s-state might not be the g.s. of  ${}^9\text{He}$ .

This argument is supported by the structure of  ${}^{10}\text{He}$ , which should be similar to the structure of  ${}^9\text{He}$ . If a virtual state in  ${}^9\text{He}$  [2001Ch31] really existed, a narrow near-threshold  $0^+$  state in  ${}^{10}\text{He}$  with a  $[s1/2]^2$  structure would exist in addition to the  $[p1/2]^2$  state [59, 60], in contradiction to the available experimental data on  ${}^{10}\text{He}$ .

Based on these experimental results, we adopt the  $1/2^-$  as the ground state of  ${}^9\text{He}$ . In earlier work [1987Se05], [1988Bo20], and [1991Bo.B], transfer reactions were used, yielding values of  $E_r$  (res-

onance energy) of this state around 1.1 MeV. More recently, [1999Bo26] and [2010Jo06] determined  $E_r \sim 1.3$  MeV. In [2007Go24], the  $1/2^-$  state of  ${}^9\text{He}$  was found at  $E_r = 2.0 \pm 0.2$  MeV with a width  $\sim 2$  MeV in this work, significantly higher than in the other reports. The energy resolution in this experiment was 0.8 MeV (FWHM), which is quite large compared to the energy difference of  $\sim 1.1$  MeV between the  $1/2^-$  and  $3/2^-$  states [1988Bo20], [1999Bo26], [2010Jo06]. Therefore, we suspect this state to be a mixture due to the poor energy resolution in this experiment.

The case related to  ${}^{10}\text{He}$  was discussed in AME2012. At that time four experimental results were known concerning the mass of  ${}^{10}\text{He}$ , and two new results were published since. The six experimental results are:

Reference	$Q_{2n}$ (in keV)	Method of production
1994Os04	$1070 \pm 70$	${}^{10}\text{Be}({}^{14}\text{C}, {}^{14}\text{O}){}^{10}\text{He}$
1994Ko16	$1200 \pm 300$	$\text{C}({}^{11}\text{Li}, {}^{10}\text{He})$
2010Jo06	$1420 \pm 100$	${}^1\text{H}({}^{11}\text{Li}, {}^{10}\text{He})$
2012Si07	$2100 \pm 200$	${}^3\text{H}({}^8\text{He}, p){}^{10}\text{He}$
2012Ko43	$1600 \pm 250$	$\text{C}({}^{14}\text{Be}, {}^{10}\text{He})$
2015Ma54	$1400 \pm 300$	${}^{11}\text{Li}(d, {}^3\text{He}){}^{10}\text{He}$

The mass of  ${}^{10}\text{He}$  from [1994Os04] is significantly lower than the others, with poor statistic compared to the high background. The values obtained in previous invariant-mass measurements [1994Ko16, 2010Jo06] agree with each other, both using  ${}^{11}\text{Li}$  to produce  ${}^{10}\text{He}$ . In [2012Si07] the value is higher than the others, while the authors stated that “the results reported in Refs. [1994Ko16] and [2010Jo06] do not contradict the g.s. energy of  ${}^{10}\text{He}$  obtained in the present work”, based on the calculations of Ref. [60]. They argued that due to the strong initial state effect, the observable g.s. peak position in [1994Ko16] and [2010Jo06] is shifted towards lower energy because of the abnormal size of  ${}^{11}\text{Li}$ , which exhibits one of the most extended known neutron halos. Based on the  ${}^9\text{He}$  spectrum from [2007Go24], the  ${}^{10}\text{He}$  g.s. with structure  $[p1/2]^2$  is predicted to have a 2.0 – 2.3 MeV two-neutron separation energy. However, the result of  ${}^9\text{He}$  from [2007Go24] is not adopted, as discussed earlier. The model has problems in interpreting all of the experimental data, indicating the states may have a more complex structure. In the AME2012 evaluation, we adopted the result from [2010Jo06] provisionally and called for more experiments to clarify this case.

Two experimental results were published after AME2012. In [2012Ko43], a group from MSU has studied  ${}^{10}\text{He}$  using the fragmentation of  ${}^{14}\text{Be}$ . In [2015Ma54], the missing mass spectrum of  ${}^{10}\text{He}$  was measured at RIKEN using the  ${}^{11}\text{Li}(d, {}^3\text{He})$  reaction at 50.4 MeV. Their results support the choice in AME2012. The discrepancy with

the result in [2012Si07] could not be explained simply by the exotic structure of  $^{11}\text{Li}$ , which was the argument used in [2012Si07], since in the later experiments different reaction channels are explored.

The discrepancy among the results for the mass of  $^{10}\text{He}$  has attracted wide attention. To reconcile this case it has been proposed recently that all the experiments were measuring two overlapping  $0^+$  states that were populated with different ratios in different experiments. Thus the lowest state reported up to now was adopted as the ground state in [61]. However, there are too many assumptions in this argument. Four experimental results published up to now are consistent with each other. We use the two most precise ones among them ([2010Jo06] and [2012Ko43]) to determine the mass of  $^{10}\text{He}$ . Meanwhile, we call for more experiments to further clarify this case.

### 7.3 The mass of $^{32}\text{Si}$

In AME2012, we discussed in details the difficulties we met in the determination of the mass of  $^{32}\text{Si}$ . We then decided not to use the  $(n,\gamma)$  data. No new experimental result relevant to this case appeared since then. In the PTB (Physikalisch-Technische Bundesanstalt) experiment [2001Pa52], the nuclide  $^{32}\text{Si}$  is produced by neutron capture on  $^{31}\text{Si}$ , which is radioactive and it is also produced in neutron capture reaction. It seems reasonable to question whether the measurement of the  $\gamma$  rays from  $^{32}\text{Si}$  could involve a substantial background. In the present evaluation, we keep using the Penning trap values, as we did in AME2012.

### 7.4 The $Q$ -value for $^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$

The  $Q$ -value for  $^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$  was adjusted to be 2043(7) and 2044(7) keV in AME2003 and AME2012, respectively. In both evaluations,  $Q_{\beta^+}$  was mainly determined from the two  $\beta$ -decay results in Ref. [1952Sc11] and [1959To25]. A higher  $Q$ -value of 2170(30) keV was reported in the later work [1974An23], but was not used in AME due to its large uncertainty. However, it was found that the  $Q$ -value should be larger than 2059.34 keV because this state was populated in beta-decay experiments. In [1959To25], the authors reported that the highest end-point energy of the  $\beta^+$  spectrum was 1030 keV and no  $\gamma$  ray was observed in coincidence. While [1974An23] reported that the highest energy component of the  $\beta^+$  spectrum was largely connected with the transition to the 89.76 keV level, and the authors suggested that in [1959To25] the  $\beta^+$  spectrum should also be associated with that  $\gamma$  transition. If we accept this explanation, then the deduced  $Q_{\beta^+}$  would be in strong conflict with the result from [1952Sc11], where the  $\beta^+$  spectrum was measured from the excited isomer, whose excitation

energy has been well established from the  $\gamma$ -ray spectroscopy.

Both  $^{99}\text{Rh}$  and  $^{99}\text{Ru}$  are primary nuclides (cf. Section 5.3) in our evaluation. The mass of  $^{99}\text{Ru}$  is mainly determined by  $(n,\gamma)$  reaction to  $^{100}\text{Ru}$ , which in turn is determined from Penning trap measurement. The uncertainty of the adjusted  $^{99}\text{Rh}$  mass is 6.7 keV. The mass of  $^{99}\text{Rh}$  can also be determined through the  $Q_{\beta^+}$  value of  $^{99}\text{Pd}$ , which in turn is measured with a Penning trap with an uncertainty of 5.4 keV. Consequently, even if we don't use any of the experimental results from the beta-spectrum measurements of  $^{99}\text{Rh}$ , we can still determine the  $Q$ -value of  $^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$  to be 2032(21) keV. If we increase the  $Q$ -value of  $^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$ , then strong tension will be built around this region.

Confronting all of the difficulties, in this evaluation we follow the same treatment used in AME2012. Meanwhile, direct mass measurements of  $^{99}\text{Rh}$  with high accuracy are needed, in order to solve this issue.

### 7.5 The mass of $^{100}\text{Sn}$

The determination of the mass of  $^{100}\text{Sn}$  was the subject of a detailed discussion in AME2003 and again in AME2012. This result is particularly interesting due to the doubly magic character of  $^{100}\text{Sn}$  which is, moreover, the heaviest known nuclide with  $N = Z$ . No new results have been reported since then for  $^{100}\text{Sn}$ . We therefore still recommend using the  $Q_{\beta^+}$  value from GSI [2012Hi07], which is also the most precise, for the determination of its mass, and are in demand of more experimental results.

### 7.6 The $^{102}\text{Pd}$ double-electron capture energy

In the AME2012 evaluation, we described the discrepancies we found between data coming from combinations of the very precise  $(n,\gamma)$  reactions with  $\beta^+$ ,  $\beta^-$  and  $\epsilon$  decay energies versus direct Penning trap data [2011Go23]. Having found no reason to distrust any of the measurements involved then, and having on one side one result obtained with a very reliable method, on the other side derived from a combination of several quite trustable measurements, we finally decided at that time to provisionally not use the new Penning trap result and called for more measurements in order to clarify this issue.

Recently, we were aware of the experimental results for  $^{102}\text{Pd}$  and  $^{103}\text{Pd}$  from the ESR mass measurements through a private communication [2014Ya.A]. Their results support the SHIPTRAP datum, although with relatively large uncertainties. We therefore decided to discard two pieces of data:  $^{102}\text{Rh}(\beta^-)^{102}\text{Pd}$  and  $^{103}\text{Pd}(\epsilon)^{103}\text{Rh}$ , and to use the SHIPTRAP result. The local consistency is then restored.




Figure 4. Loops created by three alpha-decay chains interconnected by proton decays and one IT. Each square box represents an individual nuclide. Its mass precision (keV) is given in the lower right corner, its degree in the upper right corner. Along each connection between two nuclides is the type of relation and its precision. ‘m’ stands for the excited isomer of the nuclide below it.

### 7.7 The mass of $^{105}\text{Sb}$

The matter of the determination of the mass of  $^{105}\text{Sb}$  was discussed at length in AME2012. No new data were published since then. Therefore, the  $Q_\alpha$  measurement in [2007Ma35] is again provisionally adopted in the present evaluation, and determines the mass of  $^{105}\text{Sb}$ . We still appeal for direct proton-decay studies of  $^{105}\text{Sb}$  in order to clarify this case.


### 7.8 The $^{163}\text{Ta}(\alpha)^{159}\text{Lu}(\alpha)^{155}\text{Tm}$ decay chain

This  $\alpha$ -decay chain was discussed in the previous AME2003 and AME2012 publications.

To summarize, by combining all available information, and by discarding only one piece of data, we were able to build up a scenario for the double (ground states and excited isomers)  $^{147}\text{Tb}$ - $^{179}\text{Tl}$  decay chain. However, most of the adopted values for excitation energies, and also for the  $^{167}\text{Re}$  ground state are still labeled with the ‘#’ flag, due to the estimated excitation energy of  $^{179}\text{Tl}^m$ . Experimental determination of any of the excitation energy in  $^{159}\text{Lu}$ ,  $^{163}\text{Ta}$ ,  $^{167}\text{Re}$ ,  $^{171}\text{Ir}$ ,  $^{175}\text{Au}$ , or  $^{179}\text{Tl}$  will allow to access all other ones. Future measurements would be beneficial not only in order to firmly establish these excitation energies, but even more importantly, to provide also useful parent-daughter correlations on  $\alpha$  decays that feed the  $^{159}\text{Lu}$  ground state and decays out of the excited isomer.

### 7.9 The $^{170}\text{Au}(\alpha)$ and $^{169}\text{Pt}(\alpha)$ decay chains

It has been previously mentioned that some proton-rich nuclides can decay by both  $\alpha$  and proton emission. In some cases, a loop of interconnected nuclides can be formed. Two long  $\alpha$ -decay chains illustrate this case:



These two chains are connected by  $^{170}\text{Au}(p)^{169}\text{Pt}$  and  $^{166}\text{Ir}(p)^{165}\text{Os}$ , thus forming a loop as shown in Fig. 4.

However, all the masses shown in Fig. 4 are unknown. If the mass of at least one nuclide is measured in the future, then all of the masses along the above two decay chains will be determined. The specific difficulty here, is that if all of the experimental information is used in the evaluation, then a closed loop would be formed, and all nuclides involved would become primaries. The consequence is that two estimated (non-experimental) data would then automatically become primary data (the ones with the ‘#’ flag in Fig. 5).

To avoid this undesirable side-effect, the  $^{170}\text{Au}(\alpha)^{166}\text{Ir}$  value is not used in the mass adjustment, despite its good precision. A local evaluation is carried out in this region, involving all the corresponding nuclides, using least-squares method. The input and adjusted values are listed in Table E, as well as the adjusted values listed in Table I.

Table E. Input data and adjusted values from a Least-Squares Method (LSM) adjustment around  $^{170}\text{Au}$ .

Item	Output AME	Input LSM	Adjusted LSM
$^{166}\text{Ir}(p)^{165}\text{Os}$	1152(8)	1152(8)	1154(6)
$^{166}\text{Ir}^m(\text{IT})^{166}\text{Ir}$	171.5(6.1)	171.5(6.1)	172(6)
$^{169}\text{Pt}(\alpha)^{165}\text{Os}$	6857.6(5.1)	6857.6(5.1)	6856(5)
$^{170}\text{Au}(p)^{169}\text{Pt}$	1471.7(12.0)	1471.7(12.0)	1470(9)
$^{170}\text{Au}^m(p)^{169}\text{Pt}$	1751.4(5.1)	1751.4(5.1)	1751(5)
$^{170}\text{Au}(\alpha)^{166}\text{Ir}$	7177(15)	7170(12)	7172(9)
$^{170}\text{Au}^m(\alpha)^{166}\text{Ir}^m$	7285(12)	7278.5(9.0)	7280(7)

### 7.10 A mass difference between $^{249}\text{Cf}$ and $^{241}\text{Am}$

Recent results from TRIGA-TRAP for  $^{241}\text{Am}$ ,  $^{243}\text{Am}$ ,  $^{244}\text{Pu}$  and  $^{249}\text{Cf}$  [2014Ei01] yield a mass difference between  $^{249}\text{Cf}$  and  $^{241}\text{Am}$ :  $\Delta_{TT} = M(^{249}\text{Cf}) - M(^{241}\text{Am}) = 16781.2(2.2)$  keV. This difference can also be independently obtained from combining the  $Q_\alpha$  values of  $^{249}\text{Cf}$  and  $^{245}\text{Cm}$ , with the  $Q_\beta$  value of  $^{241}\text{Am}$ :  $\Delta_Q = Q_\alpha(^{249}\text{Cf}) + Q_\alpha(^{245}\text{Cm}) + Q_\beta(^{241}\text{Pu}) + 2 \times M(^4\text{He})$ , where  $M(^4\text{He})$  is the mass of  $^4\text{He}$  atom. Using the recommended  $Q_\alpha$  values in AME2012, one obtains  $\Delta_{2012} = 16789.7(1.2)$  keV, which is 8.5(2.5) keV larger (more than  $3\sigma$ ) than the value deduced from the TRIGA-TRAP data [2014Ei01] alone. It should be pointed out that the  $Q_\alpha(^{249}\text{Cf})$  and  $Q_\alpha(^{245}\text{Cm})$  values in AME2012 were heavily weighed by the results of Baranov *et al.* [1971Bb10], [1975Ba65]. Both  $^{249}\text{Cf}$  and  $^{245}\text{Cm}$  are odd-mass nuclides and their  $\alpha$  spectra are complex, involving a number of strong  $\alpha$  lines to excited states in the daughter  $^{245}\text{Cm}$  and  $^{241}\text{Pu}$  nuclides. We note also a number of inconsistencies in the  $\alpha$ -decay energies reported in Refs. [1971Bb10] and [1975Ba65]. For example, in the case of the  $^{245}\text{Cm}$   $\alpha$  decay we obtain  $Q_\alpha(^{245}\text{Cm}) = 5621.2(0.5)$  keV from the ground-state to ground-state  $\alpha$ -decay energy  $E_\alpha = 5529.2(0.5)$  keV [1975Ba65], while from the favored  $\alpha$ -decay energy of  $E_\alpha = 5362.0(1.2)$  keV [1975Ba65] to the excited 175.04 keV level, one gets  $Q_\alpha(^{245}\text{Cm}) = 5626.2(1.2)$  keV, differing by as much as 5.0 keV.

Recently, precise  $\alpha$ -decay energy measurements were carried out at ANL for  $^{249}\text{Cf}$  [2015Ah03] and  $^{245}\text{Cm}$  [2016Ko.A] using mass-separated sources that were calibrated using the absolute measured  $E_\alpha$  values of  $^{244}\text{Cm}$ ,  $^{248}\text{Cm}$  and  $^{250}\text{Cf}$ , as recommended by Rytz [1991Ry01]. As a result, the AME2016 adjusted  $Q_\alpha(^{249}\text{Cf}) = 6293.3(0.5)$  keV and  $Q_\alpha(^{245}\text{Cm}) = 5624.5(0.5)$  keV yield  $\Delta_{2016} = 16788.4(0.7)$  keV, which is still 7.2(2.3) keV larger than the mass-difference from the TRIGA-TRAP

data [2014Ei01]. Nonetheless, all of the relevant data are used in the mass adjustment in AME2016, while strong tensions have been built in this region, especially for  $^{249}\text{Cf}$ . Future precise mass measurements in this region are necessary in order to understand the discrepancy observed between Penning Trap and  $\alpha$ -decay spectroscopy data.

### 7.11 Comparison with the $Q_{\text{EC}}$ evaluation of J.C. Hardy and I.S. Towner

Studies of  $0^+ \rightarrow 0^+$  super-allowed  $\beta$ -decay transitions between isospin analog states can be used to test the validity of the conserved vector current hypothesis that postulates the existence of a universal  $\mathcal{F}t$  value for all such decays. This universal value can be used to determine the  $V_{ud}$  element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix, which can in turn be used to test its unitarity.




Figure 5.  $Q_{\text{EC}}$  difference between the two evaluations. The error bars represent the uncertainties from Ref. [62], while the shaded areas display the uncertainties from AME2016.

The evaluation of super-allowed  $\beta$ -decay properties, such as their transition energies,  $Q_{\text{EC}}$ , and half-lives,  $T_{1/2}$ , has been carried out since a long time by Hardy and Towner and regularly updated. Recently, experimental data for 20 super-allowed  $\beta$ -decay transitions have been evaluated and published [62]. Their evaluation is independent of AME. Although they aim at different goals and use slightly different criteria of evaluation, both are deemed valid and reliable, as can be seen from the comparison of their results for  $Q_{\text{EC}}$  and for half-lives.

In the first paper of this volume, we presented a comparison (cf. 41-030001-8) between the  $T_{1/2}$  values recommended in NUBASE2016 and those given by Hardy and Towner [62]. Below, we report on such a comparison for the  $Q_{\text{EC}}$  values.

The  $Q_{\text{EC}}$  values for all ground-state to ground-state transitions are listed in the atomic mass table (Part II, Table I, p. 030003-6 of this issue). The values for transitions involving excited isomeric states are not included in that table, but they can be easily determined with the information provided in NUBASE2016 and AME2016. When calculating the  $Q_{\text{EC}}$  values from the AME mass tables, the covariances of the mother-daughter mass values must be considered, in order to obtain the correct uncertainties. The importance of the correlation is proven by the examples of the  $Q_{\text{EC}}$  values of  $^{42}\text{Ti}$  and  $^{26}\text{Al}$ , where the uncertainties of their  $Q_{\text{EC}}$  are smaller than the individual masses. Figure 5 shows the differences between the  $Q_{\text{EC}}$  values determined in AME2016 and Ref. [62] for 19 super-allowed  $\beta$ -decay transitions (the one that is missing in the figure is  $^{70}\text{Br}$ , and it will be discussed below). Most of the  $Q_{\text{EC}}$  data are consistent and the absolute differences are in most cases below 0.1 keV. In general, the observed differences can be attributed to two main sources, namely the selection of different input data and the implementation of distinct evaluation policies in AME2016 and in Ref. [62].

As mentioned in Section 5.4.4 (*Insignificant data*, p. 030002-22), data with weights that are a factor of 10 smaller than the other results will not be used in AME and these are labeled with the letter ‘U’ in the AME2016 evaluation. There is no such a threshold defined in the Hardy and Towner’s evaluation [62] and, as a consequence, all experimental data are used, unless rejected for other reasons. For example, the discrepancy for  $^{18}\text{Ne}$  is due to the different masses used in the two evaluations for the daughter  $^{18}\text{F}$  nuclide. There are five experimental results concerning this mass, as reported in [1964Bo13], [1964Ho28], [1967Pr04], [1973Se03] and [1975Ro05], with uncertainties of 0.73, 2.2, 2.8, 3.0 and 0.60 keV, respectively. All of them were used in the Hardy and Towner’s evaluation, while only the two most precise ones were used in AME2016.

The policies for data averaging used in the two evaluations are similar to the one employed by the Particle Data Group [37], where a parameter  $\chi_n$  is used to check for the consistency of data. If  $\chi_n$  is larger than a threshold value of  $\chi_n^0$ , all uncertainties in this data set will be multiplied with a scale factor, assuming that all experimental errors were underestimated by the same factor. In AME,  $\chi_n^0$  is set to be 2.5, as explained in Section 5.4.1 (*Pre-averaging*, p. 030002-20) while a value of  $\chi_n^0 = 1$  has been adopted by Hardy and Towner [62].

The masses of the long-lived isomer  $^{70m}\text{Br}$  and its  $\beta$ -decay daughter  $^{70}\text{Se}$  were measured using Penning-trap spectrometry [2009Sa12]. By adopting the evaluated excitation energy from ENSDF, the  $Q_{\text{EC}}$  value of  $^{70}\text{Br}$  can be determined. However, this value was found to deviate significantly from the systematic behavior of the  $Q_{\text{EC}}$  values in the region and it was rejected in Ref. [62]. Furthermore, Hardy and Towner concluded that “It is likely in this case that the trap actually measured an isomeric state in  $^{70}\text{Br}$  rather its ground state.” In AME2016, the result of Ref. [2009Sa12] was used following the interpretation in the original paper as in AME2012. Additional experimental data are needed to shed more light on this case.

Differences in the selection of the input data also contribute to the observed deviations between the  $Q_{\text{EC}}$  values recommended in the AME2016 and the Hardy and Towner [62] evaluations. For example, some of the experimental results that were used in AME2016 were published after the work of Ref. [62] was completed and therefore they were not included in the latter. One example is the recent result for  $^{14}\text{O}$  from Ref. [2015Va08]. Another one is the case of  $^{30}\text{S}$ , where the level energy of the daughter nuclide that was used in Ref. [62] is outdated, while we implemented an updated value from ENSDF that is listed in the NUBASE2016 table (p. 030001-27 of this issue).

In AME, all experimental information related to atomic masses are collected. Fig. 6 displays connections related to the  $^{42}\text{Ti} \rightarrow ^{42}\text{Sc}$  and  $^{42}\text{Sc} \rightarrow ^{42}\text{Ca}$  super-allowed transitions. In Ref. [62], only results from Penning-trap mass measurements [2009Ku19] were considered and, as a consequence, the excitation energy of  $^{42}\text{Sc}^m$  was determined as  $617.12 \pm 0.39$  keV. In AME2016, we have used the more precise value of  $616.28 \pm 0.06$  keV that was obtained from  $\gamma$ -ray spectroscopy measurements [1989Ki11] and recommended in the latest ENSDF evaluation. Thus, the adoption of different internal transition energies for  $^{42}\text{Sc}^m$  can explain the variance in the  $Q_{\text{EC}}$  values of the  $^{42}\text{Ti} \rightarrow ^{42}\text{Sc}$  and  $^{42}\text{Sc} \rightarrow ^{42}\text{Ca}$  super-allowed  $\beta$  decay transitions in the two evaluations. The same situation occurs in the  $^{26}\text{Si} - ^{26}\text{Al}^m - ^{26}\text{Al} - ^{26}\text{Mg}$  connections.

Another peculiar case involves the  $^{74}\text{Rb}$  nuclide. In AME2016, one input datum is the  $Q_{\text{EC}}$  value of  $^{74}\text{Rb}$

from [2003Pi08]. However, only the intensities of the decay branches were directly measured in this work. The  $Q_{\text{EC}}$  value was deduced from the measured half-life, branching ratios and the average  $\overline{Ft}$  value obtained from the analysis of other super-allowed transitions. Therefore, the recommended  $Q_{\text{EC}}$  value is based on the assumption that  $\mathcal{F}t$  is universal and constant for all super-allowed  $\beta$  decays. This result was used in AME since AME2003 and the original values reported in Ref. [2003Pi08] would be recalculated when new experimental results for the input data were available. We believe, however, that while the deduced  $Q_{\text{EC}}$  value in this way is valuable for mass determination, it should not be considered in Ref. [62], where the primary aim is to recommend a single  $\overline{Ft}$  value.




Figure 6. Schematic diagram of connections involving two super-allowed transitions  $^{42}\text{Ti} \rightarrow ^{42}\text{Sc}$  and  $^{42}\text{Sc} \rightarrow ^{42}\text{Ca}$ . The double-headed arrows represent connections from Penning-Trap mass measurements while the single-headed arrow for the  $\gamma$  transition measurement.

## 8 General information and acknowledgments

The full content of the present issue is accessible on-line at the AMDC website [22]. In addition, several graphs representing the mass surface, beyond the main ones given in Part II, are available on the AMDC website.

Tables of masses (Part II, Table I) and nuclear reaction and separation energies (Part II, Table III) are available in ASCII format to simplify their input to computer programs by the end users. The headers of these files give information on the formats. The first file, named **mass.mas16**, contains the table of masses. The next two files correspond to the table of reaction and separation energies, in two parts of 6 entries each, as in Part II, Table III: **rct1.mas16** for  $S_{2n}$ ,  $S_{2p}$ ,  $Q_{\alpha}$ ,  $Q_{2\beta}$ ,  $Q_{\varepsilon p}$  and  $Q_{\beta n}$  (odd pages in this issue); and **rct2.mas16** for  $S_n$ ,  $S_p$ ,  $Q_{4\beta}$ ,  $Q_{d,\alpha}$ ,  $Q_{p,\alpha}$  and  $Q_{n,\alpha}$  (facing even pages). As explained in Section 4.2, p. 030002-15, since AME1995, we no longer produce special tables.

We wish to thank our many colleagues who provided answers to our questions regarding specific experimental data, as well as those who sent us preprints of their work prior to publication. Continuing interest, discussions, suggestions and encouragements from D. Lunney, A. Lopez-Martens, Yuhu Zhang and Furong Xu are highly appreciated.

This work is supported in part by the National Key Program for S&T Research and Development (Contract No. 2016YFA0400504) and the Major State Basic Research Development Program of China (Contract No. 2013CB834401). The work at ANL was supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Contract No. DE-AC02-06CH11357. W.J. Huang acknowledges the support from the China Scholarship Council, grant No. 201404910496. S. Naimi acknowledges the support of ‘‘RIKEN Pioneering Project Funding’’ from the Riken project. X. Xu acknowledges the support of ‘‘Light of West China Program’’ of the Chinese Academy of Sciences.

## Appendix A The meaning of decay energies

Conventionally, the  $\alpha$ -decay energy,  $Q_{\alpha}$ , is defined as the difference in the atomic masses of the mother and daughter nuclides:

$$Q_{\alpha} = M_{\text{mother}} - M_{\text{daughter}} - M_{4\text{He}}. \quad (10)$$

This value equals the sum of the observed energy of the  $\alpha$  particle and that of the recoiling nuclide (with only a minor correction for the fact that the cortège of atomic electrons in the latter may be in an excited state). Unfortunately, some authors in the literature quote  $Q_{\alpha}$  as a value ‘corrected for screening’, which essentially means that they take for the values  $M$  in the above equation the masses of bare nuclides.

A similar bad habit has been observed for some published proton-decay energies. We very strongly object to this custom; at the very least, the symbol  $Q$  should not be used for the difference in nuclear masses.

### High precision $\alpha$ -decay energies

The most precise  $\alpha$ -decay energies are those measured absolutely using magnetic spectrographs, as summarized in Ref. [1991Ry01] and references therein. All  $\alpha$ -energy standards are determined using this method. The relation between the  $\alpha$ -particle energy  $E_{\alpha}$  and the decay energy,  $Q_{\alpha}$  is:

$$Q_{\alpha} = (M - m_{\alpha}) - \sqrt{(M - m_{\alpha})^2 - 2 \times M \times E_{\alpha}} + 78.6 \text{ eV}. \quad (11)$$

where  $M$  is the mass of the parent nuclide,  $m_{\alpha}$  is the mass of the doubly ionized  $\alpha$  particle and 78.6 eV is the bind-



ing energy energy of two electrons in helium. This formula had been discussed in AME1977 and has been used since then. Unfortunately, the electron binding energy of helium was not taken into account up to now in AME, which caused a significant deviation for the most precise  $\alpha$ -energy values that have uncertainty of tens of eV. This error is corrected in the present mass table.

### Appendix B Mixtures of isomers or of isobars in mass spectrometry

In cases where two or more unresolved lines may combine into a single one in an observed spectrum, while one cannot decide which ones are present and in which proportion, a special procedure has to be used.

The first goal is to determine what is the most probable value  $M_{exp}$  that will be observed in the measurement, and what is the uncertainty  $\sigma$  of this prediction. We assume that all the lines may contribute and that all contributions have equal probabilities. The measured mass reflects the mixing. We call  $M_0$  the mass of the lowest line, and  $M_1, M_2, M_3, \dots$  the masses of the other lines. For a given composition of the mixture, the resulting mass  $m$  is given by

$$m = (1 - \sum_{i=1}^n x_i)M_0 + \sum_{i=1}^n x_i M_i, \quad \text{with } \begin{cases} 0 \leq x_i \leq 1 \\ \sum_{i=1}^n x_i \leq 1 \end{cases} \quad (12)$$

in which the relative unknown contributions  $x_1, x_2, x_3, \dots$  have each a uniform distribution of probability within the allowed range.

If  $P(m)$  is the normalized probability of measuring the value  $m$ , then :

$$\bar{M} = \int P(m) m dm \quad (13)$$

$$\text{and } \sigma^2 = \int P(m) (m - \bar{M})^2 dm. \quad (14)$$

It is thus assumed that the experimentally measured mass will be  $M_{exp} = \bar{M}$ , and that  $\sigma$ , which reflects the uncertainty on the composition of the mixture, will have to be quadratically added to the experimental uncertainties.

The difficult point is to derive the function  $P(m)$ .

#### B.1 Case of 2 spectral lines

In the case of two lines, one simply gets

$$m = (1 - x_1)M_0 + x_1 M_1 \quad \text{with } 0 \leq x_1 \leq 1. \quad (15)$$

The relation between  $m$  and  $x_1$  is biunivocal so that

$$P(m) = \begin{cases} 1/(M_1 - M_0) & \text{if } M_0 \leq m \leq M_1, \\ 0 & \text{elsewhere} \end{cases} \quad (16)$$

*i.e.* a rectangular distribution (see Fig. 7a), and one obtains:

$$\begin{aligned} M_{exp} &= \frac{1}{2}(M_0 + M_1), \\ \sigma &= \frac{\sqrt{3}}{6}(M_1 - M_0) = 0.290 (M_1 - M_0). \end{aligned} \quad (17)$$




Figure 7. Examples of probabilities to measure  $m$  according to an exact calculation in cases of the mixture of two (a) and three (b) spectral lines.




Figure 8. Graphic representation of relations 18 and 19. The length of the segments (full thick lines) inside the triangle are proportional to the probability  $P(m)$ . Three cases are shown corresponding respectively to  $m < M_1$ ,  $m = M_1$ , and to  $m > M_1$ . The maximum of probability is obtained when  $m = M_1$ .

**B.2 Case of 3 spectral lines**

In the case of three spectral lines, we derive from Eq. 12:

$$m = (1 - x_1 - x_2)M_0 + x_1M_1 + x_2M_2, \tag{18}$$

$$\text{with } \begin{cases} 0 \leq x_1 \leq 1 \\ 0 \leq x_2 \leq 1 \\ 0 \leq x_1 + x_2 \leq 1. \end{cases} \tag{19}$$

The relations (18) and (19) may be represented on a  $x_2$  vs  $x_1$  plot (Fig. 8). The conditions (19) define a triangular authorized domain in which the density of probability is uniform. The relation (18) is represented by a straight line. The part of this line contained inside the triangle defines a segment which represents the values of  $x_1$  and  $x_2$  satisfying all relations (19). Since the density of probability is constant along this segment, the probability  $P(m)$  is proportional to its length. After normalization, one gets (Fig. 7b):

$$P(m) = \frac{2k}{M_2 - M_0}, \tag{20}$$

$$\text{with } \begin{cases} k = (m - M_0)/(M_1 - M_0) & \text{if } M_0 \leq m \leq M_1 \\ k = (M_2 - m)/(M_2 - M_1) & \text{if } M_1 \leq m \leq M_2 \end{cases} \tag{21}$$

and finally:

$$M_{exp} = \frac{1}{3}(M_0 + M_1 + M_2) \tag{22}$$

$$\sigma = \frac{\sqrt{2}}{6} \sqrt{M_0^2 + M_1^2 + M_2^2 - M_0M_1 - M_1M_2 - M_2M_0}.$$

**B.3 Case of more than 3 spectral lines**

For more than 3 lines, one may easily infer  $M_{exp} = \sum_{i=0}^n M_i/(n+1)$ , but the determination of  $\sigma$  requires the knowledge of  $P(m)$ . As the exact calculation of  $P(m)$  becomes rather difficult, it is more simple to do simulations.

However, care must be taken that the values of the  $x_i$ 's are explored with an exact equality of chance to occur. For each set of  $x_i$ 's,  $m$  is calculated, and the histogram  $N_j(m_j)$  of its distribution is built (Fig. 9). Calling  $nbin$  the number of bins of the histogram, one gets :

$$P(m_j) = \frac{N_j}{\sum_{j=1}^{nbin} N_j}, \tag{23}$$

and

$$M_{exp} = \sum_{j=1}^{nbin} P(m_j)m_j,$$

$$\sigma^2 = \sum_{j=1}^{nbin} P(m_j)(m_j - M_{exp})^2.$$






Figure 9. Examples of Monte-Carlo simulations of the probabilities to measure  $m$  in cases of two (a), three (b) and four (c) spectral lines.

A first possibility is to explore the  $x_i$ 's step-by-step:  $x_1$  varies from 0 to 1, and for each  $x_1$  value,  $x_2$  varies from 0 to  $(1-x_1)$ , and for each  $x_2$  value,  $x_3$  varies from 0 to  $(1-x_1-x_2)$ , ... using the same step value for all.

A second possibility is to choose  $x_1, x_2, x_3, \dots$  randomly in the range  $[0,1]$  in an independent way, and to keep only the sets of values which satisfy the relation  $\sum_{i=1}^n x_i \leq 1$ . An example of a Fortran program

based on the CERN library is given below for the cases of two, three and four lines. The results are presented in Fig. 9.

Both methods give results in excellent agreement with each other, and as well with the exact calculation in the cases of two lines (see Fig. 7a and 9a) and three lines (see Fig. 7b and 9b).

The Fortran program used to produce the histograms in Fig. 9.

```

program isomers
c-----
c-   October 15, 2003                C.Thibault
c-   Purpose and Methods : MC simulation for isomers (2-4 levels)
c-   Returned value      : mass distribution histograms
c-----
parameter (nwpawc=10000)
    
```

```

common/pawc/hmemor(nwpawc)
parameter (ndim=500000)
dimension xm(3,ndim)
data e0,e1,e31,e41,e42/100.,1100.,400.,200.,400./
call hlimit(nwpawc)
c histograms 2, 3, 4 levels
call hbook1(200,'',120,0.,1200.,0.)
call hbook1(300,'',120,0.,1200.,0.)
call hbook1(400,'',120,0.,1200.,0.)
call hmaxim(200,6500.)
call hmaxim(300,6500.)
call hmaxim(400,2500.)
w=1.
c random numbers [0,1]
ntot=3*ndim
iseq=1
call ranecq(iseed1,iseed2,iseq,' ')
call ranecu(xm,ntot,iseq)
do i=1,ndim
c 2 levels :
t=1-xm(1,i)
e = t*e0 + xm(1,i)*e1
call hfill(200,e,0.,w)
c 3 levels :
if ((xm(1,i)+xm(2,i)).le.1.) then
t=1.-xm(1,i)-xm(2,i)
e= t*e0 + xm(1,i)*e31 + xm(2,i)*e1
call hfill(300,e,0.,w)
end if
c 4 levels
if ((xm(1,i)+xm(2,i)+xm(3,i)).le.1.) then
t=1.-xm(1,i)-xm(2,i)-xm(3,i)
e = t*e0 + xm(1,i)*e41 + xm(2,i)*e42 + xm(3,i)*e1
call hfill(400,e,0.,w)
end if
end do
call hrput(0,'isomers.histo','N')
end

```

#### B.4 Example of application for one, two or three excited isomers

We consider the case of a mixture implying isomeric states. We want to determine the ground state mass  $M_0 \pm \sigma_0$  from the measured mass  $M_{exp} \pm \sigma_{exp}$  and the knowledge of the excitation energies  $E_1 \pm \sigma_1, E_2 \pm \sigma_2, \dots$

With the above notation, we have

$$M_1 = M_0 + E_1,$$

$$M_2 = M_0 + E_2, \dots$$

For a single excited isomer, Eq. 17 can be written:

$$M_0 = M_{exp} - \frac{1}{2}E_1,$$

$$\sigma^2 = \frac{1}{12}E_1^2 \quad \text{or} \quad \sigma = 0.29 E_1,$$

$$\sigma_0^2 = \sigma_{exp}^2 + \left(\frac{1}{2}\sigma_1\right)^2 + \sigma^2.$$

For two excited isomers, Equ. (22) lead to :

$$M_0 = M_{exp} - \frac{1}{3}(E_1 + E_2),$$

$$\sigma^2 = \frac{1}{18}(E_1^2 + E_2^2 - E_1 E_2)$$

$$\text{or} \quad \sigma = 0.236 \sqrt{E_1^2 + E_2^2 - E_1 E_2},$$

$$\sigma_0^2 = \sigma_{exp}^2 + \left(\frac{1}{3}\sigma_1\right)^2 + \left(\frac{1}{3}\sigma_2\right)^2 + \sigma^2.$$

If the levels are regularly spaced, *i.e.*  $E_2 = 2E_1$ ,

$$\sigma = \frac{\sqrt{6}}{12} E_2 = 0.204 E_2,$$

while for a value of  $E_1$  very near 0 or  $E_2$ ,

$$\sigma = \frac{\sqrt{2}}{6} E_2 = 0.236 E_2.$$

For three excited isomers, the example shown in Fig. 9c leads to:

$$\begin{aligned} M_0 &= M_{exp} - \frac{1}{4}(E_1 + E_2 + E_3) = 450, \\ \sigma &= 175, \\ \sigma_0^2 &= \sigma_{exp}^2 + \left(\frac{1}{4}\sigma_1\right)^2 + \left(\frac{1}{4}\sigma_2\right)^2 + \left(\frac{1}{4}\sigma_3\right)^2 + \sigma^2. \end{aligned}$$

## Appendix C Converting frequency ratios to linear equations

In the following, quantities with the subscript  $r$  describe the characteristics of the reference ion in the Penning Trap. Equivalent quantities, with no subscript, describe characteristics of the ion being measured.

In Ref. [2], linear equations deduced from frequency ratios are only valid for atoms. When molecules are involved, an extra term about the molecular binding energy should be included. In this case, Eq. 21 in Ref. [2] should be rewritten as:

$$R = \frac{f_r}{f} = \frac{\mathcal{M} - D - m_e q + B}{\mathcal{M}_r - D_r - m_e q_r + B_r} \frac{q_r}{q}, \quad (24)$$

where  $q$  is the charged state of the given ion,  $D$  is the molecular binding energy (dissociation energy),  $B$  is the electron binding energy,  $m_e$  is the mass of the electron and  $\mathcal{M}$  the total atomic mass. All masses and energies are in atomic mass units ( $u$ ) and so,  $u=1$ . This expression can be written in terms of the mass excess  $M$  and atomic mass number  $A$ :

$$A + M - D - m_e q + B = R \frac{q}{q_r} (A_r + M_r - D_r - m_e q_r + B_r) \quad (25)$$

or, alternatively:

$$\begin{aligned} M - R \frac{q}{q_r} M_r &= m_e q (1 - R) + A_r \left( \frac{q}{q_r} R - \frac{A}{A_r} \right) \\ &+ R \frac{q}{q_r} (B_r - D_r) - (B - D). \end{aligned}$$

The general aim is to establish some quantity  $y$  and its associated precision  $dy$ . We define  $C$  to be a truncated, three-digit decimal approximation of the ratio  $A$  to  $A_r$ , and then we can write:

$$y = M - C M_r \quad (26)$$

and so

$$y = y_1 + y_2 + y_3 + y_4, \quad (27)$$

where

$$y_1 = M_r \left( R \frac{q}{q_r} - C \right), \quad (28)$$

$$y_2 = m_e q (1 - R), \quad (29)$$

$$y_3 = A_r \left( \frac{q}{q_r} R - \frac{A}{A_r} \right), \quad (30)$$

and

$$y_4 = R \frac{q}{q_r} (B_r - D_r) - (B - D). \quad (31)$$

To fix relative orders of magnitude,  $M_r$  is generally smaller than 0.1  $u$ ,  $R - C$  is a few  $10^{-4}$ ,  $(1 - R)$  is usually smaller than unity (and typically 0.2 for a 20% mass change),  $R - \frac{A}{A_r}$  varies from 1 to  $100 \times 10^{-6}$ ,  $(B_r - D_r)$  is generally smaller than 0.1  $\mu u$ , and  $A_r$  is typically 100  $u$  for atomic mass  $A = 100$ . The four terms  $y_1$ ,  $y_2$ ,  $y_3$ , and  $y_4$  take values of the order of 10  $\mu u$ , 100  $\mu u$ , 10 to 10000  $\mu u$ , and 0.1  $\mu u$ , respectively.

The associated precision  $dy$  is written:

$$dy = dy_1 + dy_2 + dy_3 + dy_4, \quad (32)$$

where

$$dy_1 = \frac{q}{q_r} M_r dR + \left( R \frac{q}{q_r} - C \right) dM_r \simeq dR \times 10^5 \mu u, \quad (33)$$

$$dy_2 = m_e q dR \simeq dR \times 10^3 \mu u, \quad (34)$$

$$dy_3 = \frac{q}{q_r} A_r dR \simeq dR \times 10^8 \mu u, \quad (35)$$

and

$$\begin{aligned} dy_4 &= \frac{q}{q_r} (B_r - D_r) dR + R \frac{q}{q_r} (dB_r - dD_r) \\ &- (dB - dD) \simeq dR \times 10^{-1} \mu u. \end{aligned} \quad (36)$$


Consequently, only the 3rd term contributes significantly to the precision of the measurement, and so we write:  $dy = dy_3$

If the two frequencies are measured with a typical precision of  $10^{-7}$  for ions at  $A = 100$ , then the precision on the frequency ratio  $R$  is  $1.4 \times 10^{-7}$  and the precision on the mass is approximately 14  $\mu u$ .

Next we will illustrate how to calculate the molecular binding energy for the most precise Penning Trap experiment.

### C.1 Bond Dissociation Energy

The bond dissociation energy  $D$  is a quantity which signifies the strength of a chemical bond. The bond dissociation energy for a bond A–B, which is broken through reaction:



is defined as the standard enthalpy change at a specified temperature Ref. [63]:

$$D^\circ(AB) = \Delta H f_0^\circ(A) + \Delta H f_0^\circ(B) - \Delta H f_0^\circ(AB), \quad (37)$$

where  $\Delta H f_0^\circ$  is the standard heat of formation and its value is available for a large number of atoms and compounds on NIST Chemistry Webbook [64]. All  $D^\circ$  values refer to the gaseous state at temperature either 0 K or 298 K. When data on the standard heat of formation is absent at 0 K,  $D^\circ(AB)$  will be converted to  $D_{298}^\circ(AB)$  by the approximation:

$$D_{298}^\circ(AB) \approx D^\circ(AB) + 3.72 \text{ kJ/mol}. \quad (38)$$

Unlike diatomic molecules which involve only one bond, polyatomic molecules have several bonds and their  $D$ 's are the sum of all the single bonds. For example, if one wants to know the dissociation energy of  $\text{CH}_4$ , one needs to calculate the dissociation energy of the four bonds  $\text{CH}_3\text{--H}$ ,  $\text{CH}_2\text{--H}$ ,  $\text{CH--H}$ , and  $\text{C--H}$ , respectively:

$$\begin{aligned} D^\circ(\text{CH}_3 - \text{H}) &= \Delta H f_0^\circ(\text{CH}_3) + \Delta H f_0^\circ(\text{H}) - \Delta H f_0^\circ(\text{CH}_4), \\ D^\circ(\text{CH}_2 - \text{H}) &= \Delta H f_0^\circ(\text{CH}_2) + \Delta H f_0^\circ(\text{H}) - \Delta H f_0^\circ(\text{CH}_3), \\ D^\circ(\text{CH} - \text{H}) &= \Delta H f_0^\circ(\text{CH}) + \Delta H f_0^\circ(\text{H}) - \Delta H f_0^\circ(\text{CH}_2), \\ D^\circ(\text{C} - \text{H}) &= \Delta H f_0^\circ(\text{C}) + \Delta H f_0^\circ(\text{H}) - \Delta H f_0^\circ(\text{CH}). \end{aligned} \quad (39)$$

Summing over all four bonds above, we can obtain:

$$\begin{aligned} D^\circ(\text{CH}_4) &= D(\text{CH}_3 - \text{H}) + D(\text{CH}_2 - \text{H}) \\ &\quad + D(\text{CH} - \text{H}) + D(\text{C} - \text{H}) \\ &= \Delta H f_0^\circ(\text{C}) + 4 \times \Delta H f_0^\circ(\text{H}) - \Delta H f_0^\circ(\text{CH}_4). \end{aligned}$$

The dissociation energy can be generalized for a polyatomic molecule which has the form  $\text{A}_n\text{B}_k\text{C}_i$ :

$$\begin{aligned} D^\circ(\text{A}_n\text{B}_k\text{C}_i) &= n\Delta H f_0^\circ(\text{A}) + k\Delta H f_0^\circ(\text{B}) + i\Delta H f_0^\circ(\text{C}) \\ &\quad - \Delta H f_0^\circ(\text{A}_n\text{B}_k\text{C}_i). \end{aligned} \quad (40)$$

The ionization energies for molecules are available on NIST Atomic Spectra Database [65]. We now illustrate this treatment through two examples.

#### C.1.1 $^{13}\text{C}_2\text{H}_2^+$ and $^{14}\text{N}_2^+$ mass doublet

In the work of [2004Ra33], a cyclotron frequency ratio  $R = 0.999421460888(7)$  of relative precision  $7 \times 10^{-12}$  has been obtained for ions  $^{13}\text{C}_2\text{H}_2^+$  and  $^{14}\text{N}_2^+$ . We first calculate  $D^\circ(\text{C}_2\text{H}_2)$  by applying Eq. 40:

$$\begin{aligned} D^\circ(\text{C}_2\text{H}_2) &= 2 \times \Delta H f_0^\circ(\text{C}) + 2 \times \Delta H f_0^\circ(\text{H}) - \Delta H f_0^\circ(\text{C}_2\text{H}_2) \\ &= 2 \times 716.68 + 2 \times 218.998 - 227.40 \\ &= 1641.956 \text{ kJ/mol} \\ &= 17.018 \text{ eV}. \end{aligned}$$

$D^\circ(\text{N}_2) = 944.9 \text{ kJ/mol} = 9.793 \text{ eV}$  is obtained from Ref. [66], since  $\Delta H f_0^\circ(\text{N}_2)$  is not on the list of [64]. Combining the ionization energy  $B(\text{C}_2\text{H}_2) = 11.4 \text{ eV}$  and  $B(\text{N}_2) = 15.6 \text{ eV}$ , we obtain the mass difference:

$$^{13}\text{C} + \text{H} - ^{14}\text{N} = 8105.862995(98) \mu\text{u}.$$

In the original paper [2004Ra33], this equation is given as:

$$^{13}\text{C} + \text{H} - ^{14}\text{N} = 8105.86288(10) \mu\text{u},$$

which differs by  $12(10) \mu\text{u}$  from our calculation. The difference is because we used the updated molecular binding energy from Ref. [64, 66]. In this case, the molecular binding energy is 170 times larger than the uncertainty, and even the updates of the molecular energies have significant impact on the deduced values.

#### C.1.2 Atomic Masses of Tritium and Helium-3

By measuring the cyclotron frequency ratios of  $^3\text{He}^+$  and  $\text{T}^+$  to  $\text{HD}^+$ , using  $\text{HD}^+$  as a mass reference, atomic masses for  $^3\text{He}$  and  $\text{T}$  were obtained Ref. [2015My03]. The essential part here is to calculate the molecule binding energy  $D^\circ(\text{HD})$ . Instead of applying Eq. 37 to calculate the molecule binding energy of  $\text{HD}$ , we used the most recent datum  $D^\circ(\text{HD}) = 36406.78366 \text{ cm}^{-1} = 4.5137 \text{ eV}$  from [67].

Combining the ionization energies of  $B(\text{HD})=15.4445 \text{ eV}$ ,  $B(^3\text{He})=24.5874 \text{ eV}$  and  $B(\text{T}) = 13.5984 \text{ eV}$ , the mass differences and their uncertainties can be derived:

$$^3\text{He} - \text{H} - \text{D} = -5897.487710(144) \mu\text{u}$$

and

$$\text{T} - \text{H} - \text{D} = -5877.528366(144) \mu\text{u}.$$

The correction of the molecule binding energy  $D^\circ(\text{HD}) = 4.5 \text{ eV}$  is 30 times larger than the  $\sim 0.14 \text{ eV}$  uncertainty. We have created a file called "ionization", which includes all the information that is needed to calculate the linear equation.

#### C.2 Program for frequency conversion

Primary data from Penning Trap measurement are typically given in the form of an experimental frequency ratio. An example is given here for a series of nuclides with respect to a various reference nuclides and various charge states. Below is the Fortran frequency conversion program, followed by sample input files and the corresponding output file, all available on the AMDC website [22].

*The frequency conversion program*

```

c                               PTrap17j   G.Audi  WJ.Huang   v 27 jan 2017
c
c Conversion of Frequency Ratios to Linear Equations
c including electron and molecular binding energies
c
  real*8 xzero,mel,mref,smref,mrefk,rap,srap,coef
  real*8 prov,membre,sigmem,m118,sm118,meb,eref,e118,y4
  integer*4 q118,qref,znum, val,nbion,i,qel
  character txref*4,tx118*4,rev*2,ael*4
  character txiref*4,txi118*4
  character*30 filea,fileb,filec
  dimension znum(450),ael(450),qel(450),meb(450)
c
c   mel : electron mass in micro-u
c   mref,smref,m118,sm118 : Masses and uncert. for reference (ref) and mesured (118)
c
  filea='ptkl.equat'           ,           ! output file
  fileb='ptkl.freq'            ,           ! input file
  filec='ionization.data'      ,           ! electron+mol. file
  open(unit=1,file=filea,form='formatted',status='new')
  open(unit=3,file=fileb,form='formatted',status='old')
  open(unit=10,file=filec,form='formatted',status='old')
  mel = 548.57990907d0          ! mass of electron in micro-u
  xzero = 9.314940038d-1       ! conversion factor micro-u to keV
c
  do i=1,450                    ! read electron+mol. file
    read(10,*,iostat=val) znum(i),ael(i),qel(i),meb(i)
    meb(i) = (meb(i)*1.d-3)/xzero ! in micro-u
    if(val.lt.0) then
      nbion = i - 1             ! nbion: number of lines in elec+mol file
      exit
    end if
  end do
  close(10)
c
  12 read(3,1001,err=99) iaref,txref,qref,mref,rev,smref ! read ref.name, mass(micro-u) and charge
1001 format(i4,a4,i4,f19.8,a2,f13.8)
  mrefk = mref * xzero         ! ref. mass in keV
c
  15 read(3,1001,end=90,err=99) ia118,tx118,q118,rap,rev,srap ! read frequency ratio
  if(tx118.eq.'NEW ') go to 12 ! reset reference
  if(rev.eq.' ') then         ! if reversed freq. ratio: rev=-1
    rap = rap / 1.d+6
    srapsrap = srapsrap / 1.d+6
  else
    rap = 1.d+6 / rap
    srapsrap = srapsrap * rap/1.d+6
  endif
  coef = 1000.*ia118/iaref
  coef = anint(coef) / 1000. ! calculate 3-digit coefficient
c
  prov = (ia118*1.d+0)/iaref - rap*q118/qref ! start calculating the equation value
  membre = mref*(rap*q118/qref-coef) + mel*q118*(1-rap)
  * - iaref*1.d+6*prov ! value (in micro-u) for the equation
  sigmem = srapsrap * iaref * 1.d+6 * q118/qref ! its uncertainty
c
c   Correct for electron and molecular binding energies
c
  eref = 0.
  e118 = 0.
  do i=1,nbion                 ! loop over the nbion lines in elec+mol file
    txiref=txiref
    if(txiref(3:3).eq.'m' .or. txiref(3:3).eq.'n' .or. txiref(3:3).eq.'x') then
      txiref(3:4)=txiref(4:4)//' ' ! isomers m n or x : treated as gs
      if(txiref(2:2).eq.'x') txiref(2:3)=txiref(3:3)//' '
    endif
  end do

```

```

txi118=txi118
if(txi118(3:3).eq.'m' .or. txi118(3:3).eq.'n' .or. txi118(3:3).eq.'x') then
  txi118(3:4)=txi118(4:4)/' '
  if(txi118(2:2).eq.'x') txi118(2:3)=txi118(3:3)/' '
endif
if(trim(txiref)==trim(ael(i)).and.qref==qel(i)) eref = meb(i)
if(trim(txii18)==trim(ael(i)).and.q118==qel(i)) e118 = meb(i)
end do

if(eref.ne.0. .and. e118.ne.0.) then
  y4 = eref*rap*q118/qref - e118          ! calculate electronic and molecular correction
else
  y4 = 0.
  write (1,*) "WARNING : el.+mol. binding en. not found"
end if
membre = membre + y4
c
write (1,1020) ial18,txi18,iaref,txref,coef,membre,sigmem
1020 format(5x,i6,a4,'-',i4,a4,'*',f6.3,' =', f15.5,' (' ,f9.5,')')
m118 = membre + coef*mref
sm118 = sqrt(sigmem**2 + (coef*smref)**2)
write (1,1030) ial18,txi18,m118,sm118
1030 format(13x,i4,a4,' =',f15.6,' +/-',f11.6,' micro-u')
m118 = m118 * xzero
sm118 = sm118 * xzero
write (1,1032) m118,sm118
1032 format(13x,8x,' =',f15.6,' +/-',f11.6,' keV',/)
c
go to 15
c
90 write (1,1990)
1990 format(1H0,'Normal End of Freq.Ratios to Equations Conversion')
stop
99 write(1,1999)
1999 format(1H0,'Error in File Reading')
stop
end

```

### A typical frequency ratio input file

6Li	+1	15122.885	..	0.029	1st line : reference nuclide (in micro-u)
4He	+1	665392.8420		0.0077	following lines : frequency ratios
7Li	+1	1166409.2053		0.0131	NEW : new set with new ref. follows
8Li	+1	1333749.8620		0.0180	
NEW			..		column 1 : nuclidic name
7Li	+1	16003.42560		0.00455	column 2 : ionic charge
10Be	+1	700635.628	-1	0.009	column 3 : mass excess for ref. (micro-u)
11Be	+1	636546.859	-1	0.036	or frequency ratio *10 <sup>6</sup>
NEW			..		column 5 : -1 for inverse ratio
39K	+4	-36293.410	..	0.085	column 4 : uncertainty
44K	+4	886306.8169	-1	0.0444	
NEW			..		
85Rb	+9	-88210.26200	..	0.00535	
74Rb	+8	979689.6094		0.0858	
76Rb	+8	1006067.4141		0.0223	
NEW			..		
85Rb	+13	-88210.26200	..	0.00535	
99Sr	+15	1009776.3077		0.0451	
NEW			..		
3HD	+1	21926.81033	..	0.00015	
3He	+1	998048.085153		0.000048	
3H	+1	998054.687288		0.000048	

### The ionization input file

```
1 H +1 13.598434005136
```

1	HD	+1	10.9305
2	He	+1	24.587387936
3	Li	+1	5.391714761
4	Be	+1	9.322699
19	K	+4	142.6857635
37	Rb	+8	505.689068
37	Rb	+9	656.319068
37	Rb	+13	2002.533668
38	Sr	+15	2882.468673

*Corresponding output file*

```

4He - 6Li * 0.667 = -7483.71665 ( 0.04620)
      4He = 2603.247650 +/- 0.050086 micro-u
          = 2424.909576 +/- 0.046655 keV

7Li - 6Li * 1.167 = -1644.99050 ( 0.07860)
      7Li = 16003.416291 +/- 0.085576 micro-u
          = 14907.086316 +/- 0.079714 keV

8Li - 6Li * 1.333 = 2327.42554 ( 0.10800)
      8Li = 22486.231245 +/- 0.114710 micro-u
          = 20945.789572 +/- 0.106852 keV

10Be - 7Li * 1.429 = -9334.15799 ( 0.12834)
      10Be = 13534.737197 +/- 0.128503 micro-u
           = 12607.526542 +/- 0.119700 keV

11Be - 7Li * 1.571 = -3479.82956 ( 0.62193)
      11Be = 21661.552062 +/- 0.621969 micro-u
           = 20177.605859 +/- 0.579360 keV

44K - 39K * 1.128 = 2529.22538 ( 2.20434)
      44K = -38409.741096 +/- 2.206428 micro-u
           = -35778.443518 +/- 2.055275 keV

74Rb - 85Rb * 0.871 = 21096.45221 ( 6.48267)
      74Rb = -55734.685988 +/- 6.482668 micro-u
           = -51916.525801 +/- 6.038567 keV

76Rb - 85Rb * 0.894 = 13930.97003 ( 1.68489)
      76Rb = -64929.004193 +/- 1.684896 micro-u
           = -60480.978079 +/- 1.569470 keV

99Sr - 85Rb * 1.165 = 35661.05986 ( 4.42327)
      99Sr = -67103.895368 +/- 4.423274 micro-u
           = -62506.876167 +/- 4.120253 keV

3He - 3HD * 1.000 = -5897.48771 ( 0.00014)
      3He = 16029.322619 +/- 0.000208 micro-u
           = 14931.217905 +/- 0.000194 keV

3H - 3HD * 1.000 = -5877.52837 ( 0.00014)
      3H = 16049.281964 +/- 0.000208 micro-u
          = 14949.809914 +/- 0.000194 keV

```

ONormal End of Freq.Ratios to Equations Conversion

## References

References in the text such as [2014Ya.A] or [2016Xu10] are listed under "References used in the AME2016 and the NUBASE2016 evaluations", p. 030003-261.

- 1 G. Audi, Int. J. Mass Spectr. **251**: 85 (2006)  
<http://dx.doi.org/10.1016/j.ijms.2006.01.048>
- 2 G. Audi, M. Wang, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu and B. Pfeiffer, Chin. Phys. C, **36**: 1287 (2012)
- 3 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu and B. Pfeiffer Chin. Phys. C, **36**: 1603 (2012)
- 4 B. Pritychenko, E. Běták, M.A. Kellett, B. Singh and J. Totans, Nucl. Instrum. Methods Phys. Res. A, **640**: 213 (2011)  
<http://www.nndc.bnl.gov/nsr/>
- 5 G. Audi, O. Bersillon, J. Blachot and A.H. Wapstra, Nucl. Phys. A, **624**: 1 (1997)  
<http://amdc.impcas.ac.cn/nubase/nubase97.pdf>
- 6 A.H. Wapstra and K. Bos, At. Data Nucl. Data Tables , **20**: 1 (1977)
- 7 P.J. Mohr, D.B. Newell and B.N. Taylor, Rev. Mod. Phys. **88**: 035009 (2016)
- 8 T.P. Kohman, J.H.E. Mattauch and A.H. Wapstra, J. de Chimie Physique **55**: 393 (1958)
- 9 John Dalton, 1766-1844, who first speculated that elements combine in proportions following simple laws, and was the first to create a table of (very approximate) atomic weights.
- 10 E.R. Cohen and A.H. Wapstra, Nucl. Instrum. Methods **211**: 153 (1983)
- 11 E.R. Cohen and B.N. Taylor, CODATA Bull. **63**: (1986) Rev. Mod. Phys. **59**: 1121 (1987)
- 12 T.J. Quinn, Metrologia **26**: 69 (1989)  
B.N. Taylor and T.J. Witt, Metrologia **26**: 47 (1989)
- 13 P.J. Mohr and B.N. Taylor, J. Phys. Chem. Ref. Data **28**: 1713 (1999)
- 14 A. Rytz, At. Data Nucl. Data Tables **47**: 205 (1991)
- 15 A.H. Wapstra, Nucl. Instrum. Methods A, **292**: 671 (1990)
- 16 <http://amdc.impcas.ac.cn/masstables/Ame2003/recalib>
- 17 R.G. Helmer and C. van der Leun, Nucl. Instrum. Methods **422**: 525 (1999)
- 18 Nuclear Data Sheets, <http://www.nndc.bnl.gov/nds/>
- 19 G. Audi, M. Epherre, C. Thibault, A.H. Wapstra and K. Bos, Nucl. Phys. A, **378**: 443 (1982)
- 20 G. Audi, Hyperfine Interactions **132**: 7 (2001) École Internationale Joliot-Curie 2000, Spa, p.103;  
<http://amdc.impcas.ac.cn/masstables/hal.pdf>
- 21 Systematic errors are those due to instrumental drifts or instrumental fluctuations, that are beyond control and are not accounted for in the error budget. They might show up in the calibration process, or when the measurement is repeated under different experimental conditions. The experimentalist adds then quadratically a systematic error to the statistical and the calibration ones, in such a way as to have consistency of his data. If not completely accounted for or not seen in that experiment, they can still be observed by the mass evaluators when considering the mass adjustment as a whole.
- 22 The AME2016 files in the electronic distribution and complementary documents can be retrieved from the Atomic Mass Data Center (AMDC) through the Web: <http://amdc.impcas.ac.cn/> ; and mirror site <https://www-nds.iaea.org/amdc/>
- 23 C.F. von Weizsäcker, Z. Phys. **96**: 431 (1935)  
H.A. Bethe and R.F. Bacher, Rev. Mod. Phys. **8**: 82 (1936)
- 24 C. Borcea and G. Audi, Rev. Roum. Phys. **38**: 455 (1993) *CSNSM Report 92-38*, Orsay 1992:  
<http://amdc.impcas.ac.cn/extrapolations/bernex.pdf>
- 25 A.H. Wapstra, G. Audi and R. Hoekstra, Nucl. Phys. A, **432**: 185 (1985)
- 26 D. Lunney, J.M. Pearson and C. Thibault, Rev. Mod. Phys. **75**: 1021 (2003)  
D. Lunney, Proc. Conf. Advances in Radioactive Isotope Science (ARIS2014), JPS Conf. Proc. **6** 010018 (2015).
- 27 G. Audi and A.H. Wapstra, Nucl. Phys. A, **595**: 409 (1995);  
<http://amdc.impcas.ac.cn/masstables/Ame1995/>
- 28 R.G. Thomas, Phys. Rev. **80**: 136 (1950), **88**: 1109 (1952)  
J.B. Ehrman, Phys. Rev. **81**: 412 (1951)
- 29 E. Comay, I. Kelson and A. Zidon, Phys. Lett. B, **210**: 31 (1988)
- 30 M. MacCormick and G.Audi, Nucl. Phys. A, **925**: 61 (2014)
- 31 A.H. Wapstra, Proc. Conf. Nucl. Far From Stability/AMCO9, Bernkastel-Kues 1992, Inst. Phys. Conf. Series, **132**: 125 (1993)
- 32 M.S. Antony, J. Britz, J.B. Bueb and A. Pape, At. Data Nucl. Data Tables **33**: 447 (1985) M.S. Antony, J. Britz and A. Pape, At. Data Nucl. Data Tables **34**: 279 (1985) A. Pape and M.S. Antony, At. Data Nucl. Data Tables **39**: 201 (1988)  
M.S. Antony, J. Britz and A. Pape, At. Data Nucl. Data Tables **40**: 9 (1988)
- 33 J. Jänecke, in D.H. Wilkinson, 'Isospin in Nuclear Physics', North Holland Publ. Cy. (1969) eq.8.97;  
J. Jänecke, Nucl. Phys. **61**: 326 (1965)
- 34 L. Axelsson, J. Åystö, U.C. Bergmann, M.J.G. Borge, L.M. Fraile, H.O.U. Fynbo, A. Honkanen, P. Hornshøj, A. Jonkinen, B. Jonson, I. Martel, I. Mukha, T. Nilsson, G. Nyman, B. Petersen, K. Riisager, M.H. Smedberg, O. Tengblad and ISOLDE, Nucl. Phys. A, **628**: 345 (1998)
- 35 Y.V. Linnik, Method of Least Squares (Pergamon, New York, 1961); Méthode des Moindres Carrés (Dunod, Paris, 1963).
- 36 G. Audi, W.G. Davies and G.E. Lee-Whiting, Nucl. Instrum. Methods A, **249**: 443 (1986)
- 37 C. Patrignani *et al.* (Particle Data Group), Chin. Phys. C, **40(10)**: 100001 (2016)
- 38 M.U. Rajput and T.D. Mac Mahon, Nucl. Instrum. Methods A, **312**: 289 (1992)
- 39 M.J. Woods and A.S. Munster, NPL Report RS(EXT)95 (1988)
- 40 A.H. Wapstra, G. Audi and C. Thibault, Nucl. Phys. A,



- 729**: 129 (2003)  
<http://amdc.impcas.ac.cn/masstables/Ame2003/Ame2003a.pdf>
- 41 G. Audi, A.H. Wapstra and M. Dedieu, Nucl. Phys. A, **565**: 193 (1993)
- 42 G. Bollen, H.-J. Kluge, M. König, T. Otto, G. Savard, H. Stolzenberg, R.B. Moore, G. Rouleau and G. Audi Phys. Rev. C, **46**: R2140 (1992)
- 43 Each group of mass-spectrometric data is assigned a factor  $F$  according to its partial consistency factor  $\chi_n^p$ , due to the fact that its statistical uncertainties and its internal systematic error alone do not reflect the real experimental situation. From comparison to all other data and more specially to combination of reaction and decay energy measurements, we have assigned factors  $F$  of 1.5, 2.5 or 4.0 to the different labs. Only Penning trap data have almost all been assigned a factor  $F = 1.0$ . Example: the group of data H25 has been assigned  $F = 2.5$ , this means that the total uncertainty assigned to  $^{155}\text{Gd } ^{35}\text{Cl} - ^{153}\text{Eu } ^{37}\text{Cl}$  is  $2.4\mu\text{u} \times 2.5$ . The weight of this piece of data is then very low, compared to  $0.79\mu\text{u}$  derived from all other data. This is why it is labeled “U”.
- 44 A. Gillibert, L. Bianchi, A. Cunsolo, A. Foti, J. Gastebois, Ch. Grégoire, W. Mittig, A. Peghaire, Y. Schutz and C. Stéphan, Phys. Lett. B, **176**: 317 (1986)
- 45 D.J. Vieira, J.M. Wouters, K. Vaziri, R.H. Krauss, Jr., H. Wollnik, G.W. Butler, F.K. Wohn and A.H. Wapstra, Phys. Rev. Lett. **57**: 3253 (1986)
- 46 W. Heisenberg, Z. Phys. **77**: 1 (1932)
- 47 E. P. Wigner, Phys. Rev. **51**: 106 (1937)
- 48 E.P. Wigner, in Proceedings of the Robert A. Welch Foundation Conference on Chemical Research, edited by W.O. Milligan (Houston: Welch Foundation, 1958), **Vol. 1**, p. 88.
- 49 W.J. Huang and G. Audi, ND 2016 International Conference on Nuclear Data for Science and Technology; ArXiv, <http://arxiv.org/abs/1702.01639>
- 50 J. Lindhard, V. Nielsen, M. Schar, and P.V. Thomsen, Mat. Fys. Medd. Vid. Selsk, **33**: 10 (1963)
- 51 A. Ratkowski, Nuclear Instruments and Methods, **130(2)**: 533 (1975)
- 52 S. Hofmann, G. Munzenberg, K. Valli, F. Hessberger, J.R.H. Schneider, P. Armbruster, B. Thuma, and Y. Eyal, GSI-Report, page 241, 1982.
- 53 W. Bambynek, H. Behrens, M.H. Chen, B. Crasemann, M.L. Fitzpatrick, K.W.D. Ledingham, H. Genz, M. Mutterer and R.L. Intemann, Rev. Mod. Phys. **49**: 77 (1977)
- 54 J.C. Hardy, L.C. Carraz, B. Jonson and P.G. Hansen, Phys. Lett. B, **71**: 307 (1977)
- 55 G. Audi, M. Wang, A.H. Wapstra, B. Pfeiffer and F.G. Kondev, J. Korean Phys. Soc. **59**: 1318 (2011)
- 56 L. Öhrström and J. Reedijk, Pure Appl. Chem. **88(12)**: 1225 (2016)
- 57 P.J. Karol, R.C. Barber, B.M. Sherrill, E. Vardaci and T. Yamazaki, Pure Appl. Chem. **88(12)**: 139 (2016)
- 58 S. Cwiok, S. Hofmann and W. Nazarewicz, Nucl. Phys. A, **573**: 356 (1994)  
 S. Cwiok, W. Nazarewicz and P.H. Heenen, Phys. Rev. Lett. **63**: 1108 (1999)
- 59 S. Aoyama, Phys. Rev. Lett. **89**: 052501 (2002)
- 60 L.V. Grigorenko and M.V. Zhukov, Phys. Rev. C, **77**: 034611 (2008)
- 61 H.T. Fortune, Chinese Physics Lett. **33**: 092101 (2016)
- 62 J.C. Hardy and I.S. Towner, Physical Review C **91**: 025501 (2015).
- 63 B.D. Darwent, NSRDS-NBS **31**: 48 (1970)
- 64 P.J. Linstrom and W.G. Mallard, NIST Chemistry WebBook, *NIST Standard Reference Database Number 69*, National Institute of Standards and Technology, Gaithersburg MD, 20899, <http://webbook.nist.gov>, 2016.
- 65 A. Kramida, Yu. Ralchenko, J. Reader, and NIST ASD Team (2015). *NIST Atomic Spectra Database (version 5.3)*, [Online]. Available: <http://physics.nist.gov/asd>. National Institute of Standards and Technology, Gaithersburg, MD.
- 66 Y.R. Luo and J.A. Kerr, Bond dissociation energies. *CRC Handbook of Chemistry and Physics*. **89**, 2012.
- 67 D. Sprecher, J. Liu, C. Jungen, W. Ubachs and F. Merkt, J. Chem. Phys. **133**: 111102 (2010)

**Table I. Input data compared with adjusted values****EXPLANATION OF TABLE**

The ordering is in groups according to highest occurring relevant mass number.

Item	$K^m, Cs^m, Cs^n,$ $In^p, Tl^q:$ higher isomers, see NUBASE.	In nuclear reactions: $\epsilon$ = electron capture,	In mass-doublet equation: $H = {}^1H,$ $N = {}^{14}N,$ $D = {}^2H,$ $O = {}^{16}O,$ $C = {}^{12}C,$ u = absolute mass-doublet.	In mass-triplet equation: $Rb^x, Rb^y:$ different mixtures of isomers or contaminants.
Input value	Mass doublet: value and its standard precision in $\mu u$ . Triplet: value and its standard precision in keV. Reaction: value and its standard precision in keV. The value is the combination of mass excesses $\Delta(M - A)$ given under ‘item’. It is the author’s experimental result and the author’s stated uncertainty, except in a few cases for which comments are given and for some $\alpha$ -reactions: if the $\alpha$ -decay does not clearly feed the ground state, then the precision is increased to 50 keV. If more than one group report such energies, an average is calculated first (mentioned in the Table) and the 50 keV is added to the averaged precision in the adjustment (see Section 6.3).			
Adjusted value	Output of calculation. For secondary data ( $Dg = 2-20$ ) the adjusted value is the same as the input value and is not repeated. Also, the adjusted value is only given once for a group of results for the same reaction or doublet. Values and precisions were rounded off, but not to more than tens of keV. # Value and precision derived not from purely experimental data, but at least partly from trends of the mass surface (TMS). * No mass value has been calculated for one of the masses involved.			
$v_i$	Normalized deviation between input and adjusted value, given as their difference divided by the input precision (see Section 5.2).			
Dg	1 Primary data (see Section 3). 2–13 Secondary data of different degrees. B Well-documented data, or data from regular reviewed journals, which disagree with other well-documented values. C Data from incomplete reports, at variance with other data. o Data included in or superseded by later work of same group. D Data not checked by another method and at large variance with TMS, replaced by an estimated value (see Section 4, p. 030002-15). F Study of paper raises doubts about validity of data within the reported precision. R Item replaced for computational reasons by an equivalent one giving same result. U Data with much less weight than that of a combination of other data. – Data that will be averaged.			
Signf.	<i>Significance</i> ( $\times 100$ ) of primary data only (see Section 5.1); the significance of secondary data is always 100%.			
Main infl.	Largest <i>influence</i> ( $\times 100$ ) and nucleus to which the data contributes the most (see Section 5.1).			
Lab	Identifies the group which measured the corresponding item. Example of Lab key: MA8 Penning Trap data of Mainz-Isolde group. The numbers refer to different experimental conditions.			
F	Multiplying factor for mass spectrometric data (see Section 6.1). The standard precision given in the ‘Input value’ column has been multiplied by this factor before being used in the least-squares adjustment.			

Reference	Reference keys: (in order to reduce the width of the Table, the two digits for the centuries are omitted; at the end of this volume however, the full reference key-number is given: 2003Ba49 and not 03Ba49).
12Na15	Results derived from regular journal. These keys are copied from Nuclear Data Sheets. Where not yet available, the style 12Re.1 has been used.
12Zh.A	Result from abstract, preprint, private communication, conference, thesis or annual report.
Ens169	References to energies of excited states, when of interest, are mentioned in remarks in the Qfile. Their reference-keys refer to the "Evaluated Nuclear Structure Data Files" (ENSDF) (the electronic version of the Nuclear Data Sheets NDS), the reference-keys are indicated Ens169 in which '16' indicates the year (here 2016) and '9' the month (Oct, Nov, Dec indicated a b c) of the released ENSDF file.
Nub169	When the excited energy is derived or estimated in NUBASE2016, it is indicated with 'Nubase' (see previous item).
Averag	Average of data from the following lines.
AHW	(or FGK, GAU, HWJ, MMC, WGM) : comment written by one of the evaluators.
*	A remark on the corresponding item is given below the block of data corresponding to the same (highest) A.
Y	recalibrations of 65Ry01 for charged particle recalibrations, and recalculated triplets for isomeric mixtures.
Z	recalibrations of 91Ry01 for $\alpha$ particles, 90Wa22 for $\gamma$ in (n, $\gamma$ ) and (p, $\gamma$ ) reactions and 91Wa.A for protons and $\gamma$ in (p, $\gamma$ ) reactions (see Section 2).

*Remarks.* For data indicated with a star in the reference column, remarks have been added. They are collected in groups at the end of each block of data in which the highest occurring relevant mass number is the same. They give:

- i) Information explaining how the values in column 'Input value' have been derived for papers not mentioning e.g. the mass differences as derived from measured ratios of voltages or frequencies, or the reaction energies, or values for transitions to excited states in the final nuclei (for which better values of the excitation energies are now known).
- ii) Reasons for changing values (e.g. recalibrations) or precisions as given by the authors or for rejecting them (i.e. for labelling them B, C or F).
- iii) Value suggested by TMS and recommended in this evaluation as the best estimate (see Section 4, p. 030002-9).
- iv) Separate values for capture ratios (see Section 6.4).

Special notation in remarks:

$E_{\beta^-}$ , $Q_{\beta^-}$	$\beta^-$ endpoint energy, $\beta^-$ decay energy
$E_{\beta^+}$ , $Q_{\beta^+}$	$\beta^+$ endpoint energy, $\beta^+$ decay energy
$E_p$ , $Q_p$	proton energy in the laboratory, proton decay energy
$a_s$	scattering length
T	threshold for given reaction
$\varepsilon$	electron capture; $\beta^+ = \varepsilon + e^+$ (see NUBASE2016, p. 030001-20)
$p^+$ , pK, pL	fraction $\beta^+$ , $\varepsilon(K)$ or $\varepsilon(L)$ in transition to mentioned states
L/K, L/M	$\varepsilon(L)/\varepsilon(K)$ , $\varepsilon(L)/\varepsilon(M)$
IBE	internal bremsstrahlung endpoint
M-A, $D_M$	mass excess (in keV), mass difference (in $\mu u$ )
TMS	Trends from Mass Surface
'Z' (after uncertainty)	recalibrated (see above, under 'Reference')

**Table I. Comparison of input data and adjusted values (Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference			
$\pi^+$	140081.18	0.35	140081.2	0.4	0.0	1	100	100 $\pi^+$			06PaDG *			
$\pi^+(2\beta^+)\pi^-$	1021.998	0.001	1021.9980	0.0010	0.0	1	100	100 $\pi^-$			88CoTa			
* $\pi^+$	By convention! This is $M=139570.18(0.35)$ keV + $m(e^-)$										GAu **			
$H_{12}-C$	93902.7	0.4	93900.3869	0.0011	-2.3	U			M17	2.5	66Be10			
	93900.66	0.48			-0.2	U		A2	2.5	70St25				
	93900.32	0.12			0.4	U		B07	1.5	71Sm01				
	93900.391	0.012			-0.3	U		WA1	1.0	95Va38				
	93900.3804	0.0084			0.8	U		MI1	1.0	95Di08				
	93900.3865	0.0017			0.2	1	44	44 $^1H$	WA1	1.0	01Va33			
	93900.3860	0.0042			0.2	U		ST2	1.0	02Be64				
$n(\beta^-)^1H$	782	13	782.3465	0.0005	0.0	U					51Ro50			
$D_6-C$	84610.56	0.12	84610.6687	0.0007	0.4	U			A2	2.5	70St25			
	84610.62	0.09			0.4	U		B07	1.5	71Sm01				
	84611.60	0.34			-1.1	U		J5	2.5	72Ka57 *				
	84611.47	0.40			-0.8	U		J6	2.5	76Ka50				
	84610.644	0.005			4.9	C		WA1	1.0	92Va.A				
	84610.584	0.078			0.4	U		OH1	2.5	93Ma.A				
	84610.662	0.007			1.0	o		WA1	1.0	93Va.C				
	84610.6616	0.0067			1.1	o		WA1	1.0	95Va38				
	84610.6710	0.0054			-0.4	-		MI1	1.0	95Di08				
	84610.6656	0.0036			0.9	-		MI1	1.0	95Di08				
	84610.66897	0.00086			-0.3	-		WA1	1.0	06Va22				
	84610.66834	0.00024			1.4	F		WA1	1.0	15Za13 *				
	ave.	84610.6688			0.0008	-0.2	1	78	78 $^2H$		average			
	$H_2-D$	1547.77			0.28	1548.28637	0.00020	0.7	U			C1	2.5	64Mo.A
1548.22		0.05	0.5	o				M19	2.5	67Jo18				
1548.08		0.08	1.0	o				J2	2.5	69Na21				
1548.286		0.004	0.1	o				B07	1.5	71Sm01				
1548.222		0.063	0.4	o				J5	2.5	72Ka57				
1548.176		0.133	0.3	o				J5	2.5	72Ka57				
1548.298		0.008	-1.0	U				B08	1.5	75Sm02				
1548.301		0.005	-2.0	U				B08	1.5	75Sm02				
1548.190		0.023	1.7	U				J6	2.5	76Ka50				
1548.28		0.05	0.1	U				M25	2.5	78Ha14				
1548.302		0.012	-0.5	U				OH1	2.5	93Go37				
1548.2836		0.0018	1.5	U				MI1	1.0	95Di08				
1548.28649		0.00035	-0.3	1	32			24 $^1H$	ST2	1.0	08So20			
$^1H(n,\gamma)^2H$		2224.564	0.017	2224.5660	0.0004			0.1	U			BNL		80Gr02
		2224.5	0.12					0.5	U		MMn		80Is02	
		2224.561	0.009					0.6	U		Utr		82Va13 Z	
	2224.549	0.009	1.9			U				82Vy10 Z				
	2224.560	0.009	0.7			U				83Ad05 Z				
	2224.5756	0.008	-1.2			U		NBS		86Gr01 *				
	2224.5727	0.0500	-0.1			U		PTc		97Ro26 *				
	2224.5660	0.0004	0.0			o		NBS		99Ke05 *				
	2224.58	0.05	-0.3			U		Bdn		06Fi.A *				
	2224.56600	0.00044	0.0			1	100	100 $1n$	NBS		06De21 *			
* $D_6-C$	For all 72Ka57 doublets, see also reference										72Og03 **			
* $D_6-C$	F : the other result from same paper is not trusted within error bar										GAu **			
* $^1H(n,\gamma)^2H$	Original 2224.5890(0.0022) revised in reference; error increased by evaluator										90Wa22 **			
* $^1H(n,\gamma)^2H$	Original error 0.0005 increased for calibration										GAu **			
* $^1H(n,\gamma)^2H$	More precisely, $H+n-D=2388170.07(0.42)$ nu corrected to 2388169.95(0.42) nu										99Ke05 **			
* $^1H(n,\gamma)^2H$	All errors in reference increased by 20 ppm for calibration										06Fi.A **			
* $^1H(n,\gamma)^2H$	Original 2224.56610(0.00044) recalibrated with 2010 Codata (see text)										WgM129**			

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
${}^3\text{H}_4\text{-C}$	64197.0690	0.0062	64197.1279	0.0009	9.5	B			WA1	1.0	93Va04 *	
	64197.1136	0.0116			1.2	o			ST2	1.0	02Bf02	
	64197.1148	0.0100			1.3	U			ST2	1.0	06Na49	
${}^3\text{He}_4\text{-C}$	64117.2399	0.0039	64117.2906	0.0009	13.0	B			WA1	1.0	93Va04	
	64117.252	0.030			1.3	U			WA1	1.0	93Va04 *	
	64117.294	0.011			-0.3	o			ST2	1.0	01Fr18	
	64117.2868	0.0100			0.4	o			ST2	1.0	06Na49 *	
	64117.28668	0.00017			22.9	B			WA1	1.0	15Za13	
$\text{H}_3\text{-}{}^3\text{He}$	7445.858	0.012	7445.77408	0.00024	-2.8	U			MZ1	2.5	91Ha31 *	
$\text{D}_2\text{-H } {}^3\text{H}$	4329.257	0.003	4329.24200	0.00025	-3.3	B			B08	1.5	75Sm02	
${}^3\text{H-H D}$	-5877.2	0.7	-5877.52837	0.00016	-0.2	U			C1	2.5	64Mo.A	
	-5877.52837	0.00014			0.0	o			FS1	1.0	15My03 *	
${}^3\text{He-H D}$	-5896.84	0.42	-5897.48771	0.00014	-0.6	U			C1	2.5	64Mo.A	
	-5897.512	0.005			3.2	B			B08	1.5	75Sm02	
	-5897.495	0.006			0.8	U			B09	1.5	81Sm02	
	-5897.48771	0.00014			0.0	1	100	$100 {}^3\text{He}$	FS1	1.0	15My03	
${}^3\text{H-}{}^3\text{He}$	19.83	0.18	19.95934	0.00007	0.3	U			C1	2.5	64Mo.A	
	19.951	0.004			0.8	o				2.5	84Ni16 *	
	19.967	0.003			-1.0	o				2.5	84Li24	
	19.967	0.002			-1.5	U				2.5	85Li02	
	19.948	0.003			1.5	U				2.5	85Ta.A	
	19.9570	0.0013			1.8	U			ST2	1.0	06Na49	
	19.95934	0.00007			0.0	1	100	$100 {}^3\text{H}$	FS1	1.0	15My03	
${}^2\text{H}(n,\gamma){}^3\text{H}$	6257.6	0.3	6257.2290	0.0005	-1.2	U					69Pr06	
	6256.96	0.25			1.1	U			ILn		79Br25 Z	
${}^2\text{H}(d,p){}^3\text{H}$	4029	12	4032.66296	0.00024	0.3	U			CIT		49To23 Y	
	4034	6			-0.2	U			MIT		64Sp12	
	4033.7	1.7			-0.6	U			NDm		67Od01	
${}^2\text{H}(d,n){}^3\text{He}$	3260	9	3268.9084	0.0005	1.0	U			CIT		49To23 Y	
	3269	11			0.0	U			Wis		56Do41 Y	
${}^3\text{H}(\beta^-){}^3\text{He}$	18.645	0.016	18.59201	0.00007	-3.3	B					72Be11 *	
	18.619	0.040			-0.7	U					73Pi01 *	
	18.607	0.013			-1.2	o					76Tr07	
	18.614	0.013			-1.7	U					81Lu07 *	
	18.562	0.020			1.5	U					83De47 *	
	18.590	0.008			0.3	U					85Si07 *	
	18.604	0.006			-2.0	o					85Bo34	
	18.603	0.010			-1.1	o					86Fr09 *	
	18.600	0.004			-2.0	U					87Bo07	
	18.598	0.015			-0.4	o					87Bu.A	
	18.603	0.004			-2.7	B					88Ka32	
	18.589	0.003			1.0	o					89St05	
	18.595	0.006			-0.5	o					91Bu12	
	18.592	0.003			0.0	U					91Ka41 *	
	18.591	0.002			0.5	U					91Ro07 *	
	18.595	0.006			-0.5	U					92Bu13 *	
	18.589	0.003			1.0	o					92Ot.A	
	18.593	0.003			-0.3	U					92Ho09 *	
	18.591	0.003			0.3	U					93We03	
18.597	0.014			-0.4	U					95Hi14		
18.5895	0.0025			1.0	U					95St26		
${}^3\text{H}(p,n){}^3\text{He}$	-764.08	0.15	-763.7545	0.0005	2.1	o			Zur		61Ry05	
	-764.39	0.37			1.7	U			NRL		64Bo10	
	-763.82	0.08			0.8	U			Zur		64Sa12	
* ${}^3\text{H}_4\text{-C}$	Item preliminarily disregarded										AHW	**
* ${}^3\text{He}_4\text{-C}$	Original changed after discussion with authors										AHW	**
* ${}^3\text{He}_4\text{-C}$	Use instead the most precise difference between ${}^3\text{H}$ and ${}^3\text{He}$ (see below)										GAu	**
* $\text{H}_3\text{-}{}^3\text{He}$	From ${}^3\text{He}^+/\text{H}_2^+ = 1.496441095(6) + 3\text{eV}$ ionization										AHW	**
* ${}^3\text{H-H D}$	We use already the ${}^3\text{He}$ and ${}^3\text{H-}{}^3\text{He}$ results; only two data are independent										GAu	**
* ${}^3\text{H-}{}^3\text{He}$	Atom mass difference=ion mass difference $18.573 + 0.011$ keV										AHW	**
*	required correction cannot be estimated										85Au07	**
* ${}^3\text{H}(\beta^-){}^3\text{He}$	For corrections to 72Be11 see reference										82Di01	**

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
${}^3\text{H}(\beta^-){}^3\text{He}$	For corrections to 73Pi01 and 81Lu07 see reference										85Au07 **
${}^3\text{H}(\beta^-){}^3\text{He}$	Error for 83De47 increased, see reference										85Au07 **
${}^3\text{H}(\beta^-){}^3\text{He}$	Original value 18580(7) corrected in reference										89Re04 **
${}^3\text{H}(\beta^-){}^3\text{He}$	As calculated from their data in reference										89Re04 **
${}^3\text{H}(\beta^-){}^3\text{He}$	$E_{\beta^-}=18.5721(0.0030)$ , SFS and recoil as in reference										88Ka32 **
${}^3\text{H}(\beta^-){}^3\text{He}$	$E_{\beta^-}=18.5705(0.0020)$ , SFS and recoil as in reference										89St05 **
${}^3\text{H}(\beta^-){}^3\text{He}$	$E_{\beta^-}=18.556(0.006)$ , corrections as in reference										91Bu12 **
${}^3\text{H}(\beta^-){}^3\text{He}$	$E_{\beta^-}=18.5733(0.0002+\text{syst})$ , SFS and recoil as in reference										88Ka32 **
${}^4\text{He}_3\text{-C}$	7809.706	0.009	7809.76239	0.00019	6.3	B			WA1	1.0	92Va.A
	7809.7493	0.0030			4.4	B			WA1	1.0	95Va38
	7809.7704	0.0039			-2.1	U			ST2	1.0	01Fr18
	7809.7620	0.0003			1.3	o			WA1	1.0	01Va.A
	7809.7467	0.0066			1.0	U			MZ2	2.5	01Br27
	7809.76246	0.00019			-0.4	o			WA1	1.0	04Va14
${}^4\text{He-H}_4$	7809.76239	0.00019			0.0	1	100	100 ${}^4\text{He}$	WA1	1.0	06Va22
	-28696.8747	0.0026	-28696.8748	0.0004	-0.1	o			ST2	1.0	06Na49
$\text{D}_2\text{-}{}^4\text{He}$	-28696.8750	0.0026			0.1	U			ST2	1.0	06Na13
	25600.315	0.014	25600.30210	0.00025	-0.6	U			B08	1.5	75Sm02
$\text{H } {}^3\text{H}\text{-}{}^4\text{He}$	25600.331	0.005			-2.3	U			MZ1	2.5	90Ge12 *
	25600.328	0.005			-2.1	U			MZ1	2.5	92Ke06 *
	25600.308	0.005			-0.3	U			BL1	4.0	01He36
${}^4\text{H}(\gamma,n){}^3\text{H}$	21271.075	0.012	21271.0601	0.0003	-0.8	U			B08	1.5	75Sm02
	5200	1700	1600	100	-2.1	U					62Ar05
	2900	500			-2.6	U					69Mi10 *
	8000	3000			-2.1	U					79Me13 *
	2700	600			-1.8	U					81Se11
	2600	200			-5.0	B					85Fr01 *
	3500	500			-3.8	B					86Be35 *
	2600	400			-2.5	U					86Mi14 *
	3000	200			-7.0	B					87Go25 *
	3800	300			-7.3	B					90Am04 *
	3100	300			-5.0	B					91Bl05 *
	2300	300			-2.3	U					95Al31 *
	2670	310			-3.5	B					03Me11
	1600	100				3					09Gu17 *
${}^3\text{He}(\text{d,p}){}^4\text{He}$	18380	10	18353.05346	0.00027	-2.7	U			Mex		64Ma.B
	18382	15			-1.9	U			Mex		64Ma.B
	18350.1	3.9			0.8	U			NDm		67Od01
${}^4\text{Li}(\text{p}){}^3\text{He}$	3300	300	3100	210	-0.7	2					87Br.B
${}^4\text{D}_2\text{-}{}^4\text{He}$	Error has to be confirmed										GAu **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^7\text{Li}(\pi^-,t){}^4\text{H}$ reaction										69Mi10 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^7\text{Li}(\pi^-,t){}^4\text{H}$ reaction										AHW **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^7\text{Li}({}^3\text{He},{}^3\text{He}){}^4\text{H}$ reaction										85Fr01 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^9\text{Be}({}^{11}\text{B},{}^{16}\text{O}){}^4\text{H}$ reaction										86Be35 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^7\text{Li}(n,\alpha){}^4\text{H}$ reaction										86Mi14 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^9\text{Be}(\pi^-,dt){}^4\text{H}$ , same data in reference										91Go19 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^7\text{Li}(\pi^-,t){}^4\text{H}$										90Am04 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^2\text{D}(t,n){}^4\text{H}$										91Bl05 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	From ${}^6\text{Li}({}^6\text{Li},{}^8\text{B})$										95Al31 **
${}^4\text{H}(\gamma,n){}^3\text{H}$	Fit with 3 resonances 1.6(0.1), 3.4(0.1), 6.0(0.1) MeV										09Gu17 **
${}^5\text{H}(\gamma,2n){}^3\text{H}$	1800	800	1800	90	0.0	U					68Yo06
	11000	1500			-6.1	F					81Se.A *
	7400	700			-8.0	F					87Go25 *
	5200	400			-8.5	F					95Al31 *
	1700	300			0.3	U					01Ko52 *
	1800	100			0.0	3					03Go11 *
	1800	200			0.0	3					04St18

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
${}^4\text{He}(n,\gamma){}^5\text{He}$	-890	50	-735	20	3.1	B					66La04 *
	-735	20				3					09Ak03
	-1965	50				3					65Ma32 *
${}^4\text{He}(p,\gamma){}^5\text{Li}$											AHW **
${}^5\text{H}(\gamma,2n){}^3\text{H}$	From ${}^6\text{Li}(\pi^-,p){}^5\text{H}$ . F : private communication by author										
${}^5\text{H}(\gamma,2n){}^3\text{H}$	From ${}^9\text{Be}(\pi^-,pt){}^5\text{H}$ , same data in reference										
${}^5\text{H}(\gamma,2n){}^3\text{H}$	F : probably higher state										
${}^5\text{H}(\gamma,2n){}^3\text{H}$	From ${}^7\text{Li}({}^6\text{Li},{}^8\text{B})$										
${}^5\text{H}(\gamma,2n){}^3\text{H}$	F : probably higher state										
${}^5\text{H}(\gamma,2n){}^3\text{H}$	From $p({}^6\text{He},{}^2\text{He})$										
${}^5\text{H}(\gamma,2n){}^3\text{H}$	From $t(t,p)$										
${}^4\text{He}(n,\gamma){}^5\text{He}$	Average of many reactions leading to ${}^5\text{He}$										
${}^4\text{He}(p,\gamma){}^5\text{Li}$	Average of many reactions leading to ${}^5\text{Li}$										
${}^6\text{Li}_2\text{-C}$	30246.152	0.119	30245.775	0.003	-0.8	F			BL1	4.0	98He.B *
	30245.575	0.034			1.5	U			BL1	4.0	01He36
	30245.7748	0.0031			0.0	1	100	100 ${}^6\text{Li}$	FS1	1.0	10Mo30
${}^6\text{Li-H}_6$	-31827.302	0.040	-31827.3060	0.0016	-0.1	U			ST2	1.0	06Na13
${}^6\text{Li-D}_3$	-27182.500	0.040	-27182.4469	0.0016	0.3	U			BL1	4.0	01He36
${}^4\text{He-}{}^6\text{Li}_{.667}$	-7483.694	0.046	-7483.7117	0.0010	-0.4	U			TT1	1.0	09Br10
${}^6\text{He-}{}^7\text{Li}_{.857}$	5170.947	0.057	5170.95	0.06	0.0	1	100	100 ${}^6\text{He}$	TT1	1.0	12Br03
${}^6\text{H}(\gamma,3n){}^3\text{H}$	2700	400	2710	250	0.0	3					84A108 *
	2600	500			0.2	3					86Be35 *
	2800	500			-0.2	3					92Al.A *
	2850	900			-0.2	3					08Ca22 *
${}^6\text{Li}(n,\alpha){}^3\text{H}$	4794	6	4783.4705	0.0015	-1.8	U			Win		67De15
${}^6\text{Li}(p,\alpha){}^3\text{He}$	4017	12	4019.7160	0.0015	0.2	U			CIT		49To16 Y
	4021	5			-0.3	U			Wis		51Wi26 Y
	4023	2			-1.6	U			Bir		53Co02 Y
	4025	6			-0.9	o			MIT		64Sp12
	4018.2	1.1			1.4	U			MIT		81Ro02
${}^6\text{Li}(d,\alpha){}^4\text{He}$	22396	12	22372.7695	0.0015	-1.9	U			Bir		53Co02 Y
	22376	14			-0.2	U			Ric		53Ph28 Y
	22403	12			-2.5	U			Mex		64Ma.B
${}^6\text{Li}(p,t){}^4\text{Li}$	-18700	300	-18900	210	-0.7	R			Brk		65Ce02
${}^6\text{He}(\beta^-){}^6\text{Li}$	3509.8	3.8	3505.22	0.05	-1.2	U					63Jo04
${}^6\text{Li}(p,n){}^6\text{Be}$	-5074	13	-5071	5	0.3	2			CIT		67Ho01
${}^6\text{Li}({}^3\text{He},t){}^6\text{Be}$	-4306	6	-4307	5	-0.1	2			CIT		66Wh01
${}^6\text{Li}_2\text{-C}$	F : leak during the measurement										
${}^6\text{H}(\gamma,3n){}^3\text{H}$	From ${}^7\text{Li}({}^7\text{Li},{}^8\text{B}){}^6\text{H}$										
${}^6\text{H}(\gamma,3n){}^3\text{H}$	From ${}^9\text{Be}({}^{11}\text{B},{}^{14}\text{O}){}^6\text{H}$										
*	${}^6\text{H}$ not observed in ${}^6\text{Li}(\pi^-, \pi^+)$										
${}^6\text{H}(\gamma,3n){}^3\text{H}$	From ${}^7\text{Li}({}^7\text{Li},{}^8\text{B}){}^6\text{H}$										
${}^6\text{H}(\gamma,3n){}^3\text{H}$	Symmetrized from 2910(+850-950) keV										
${}^7\text{Li-H}_7$	-38771.7889	0.0045	-38771.789	0.004	0.0	1	100	100 ${}^7\text{Li}$	ST2	1.0	06Na13 *
${}^7\text{B-u}$	29712	27				2				1.0	11Ch32 *
${}^7\text{Li-}{}^6\text{Li}_{1.167}$	-1644.991	0.079	-1644.974	0.005	0.2	o			TT1	1.0	09Br10
${}^6\text{Li-}{}^7\text{Li}_{.857}$	1407.954	0.013	1407.942	0.004	-0.9	U			TT1	1.0	09Br.A
${}^3\text{He}(\alpha,\gamma){}^7\text{Be}$	1586.3	0.6	1587.13	0.07	1.4	U					82Kr05
${}^7\text{Li}(p,\alpha){}^4\text{He}$	17364	11	17346.245	0.004	-1.6	U			CIT		51Wh05 Y
	17352	9			-0.6	U			Bir		53Co02 Y
	17345	13			0.1	U			Ric		53Fa18 Y
	17373	6			-4.5	C			Mex		64Ma.B
	17357	14			-0.8	U			MIT		64Sp12

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
${}^7\text{H}(\gamma,2n){}^5\text{H}$	-1100	340	100#	1000#	3.5	F					08Ca22 *
${}^7\text{He}(\gamma,n){}^6\text{He}$	450	20	410	8	-2.0	3			MSU		01Ch31 *
	430	20			-1.0	3					02Me07
	360	50			1.0	U					06Sk03
	400	10			1.0	3					08De29
	388	20			1.1	3					09Ak03
	380	28			1.1	U					16Re14
${}^7\text{Li}(t,\alpha){}^6\text{He}$	9788	30	9839.90	0.05	1.7	U			ChR		54Al35 Y
${}^7\text{Li}(d,{}^3\text{He}){}^6\text{He}-{}^{19}\text{F}({}^{18}\text{O})$	-1981.09	0.42	-1980.36	0.05	1.7	U			MSU		78Ro01 *
${}^6\text{Li}(n,\gamma){}^7\text{Li}$	7250.0	0.5	7251.091	0.005	2.2	U			Utr		68Sp01
	7250.3	0.9			0.9	U					72Op01
	7250.98	0.09			1.2	U			Ptn		85Ko47 *
	7249.94	0.15			7.7	C			Bdn		06Fi.A
${}^6\text{Li}(d,p){}^7\text{Li}$	5028	2	5026.525	0.004	-0.7	U			Bir		53Co02 Y
	5035	5			-1.7	U			Mex		61Ja23
	5024	7			0.4	U			MIT		64Sp12
${}^6\text{Li}(t,d){}^7\text{Li}$	986	7	993.862	0.004	1.1	U			ChR		54Al35
${}^7\text{Li}({}^3\text{He},\alpha){}^6\text{Li}$	13322	10	13326.529	0.004	0.5	U			Mex		64Ma.B
${}^6\text{Li}(n,\gamma){}^7\text{Li}^i$	-3947	50	-4000	30	-1.0	1	39	39 ${}^7\text{Li}^i$			69Pr04 *
${}^6\text{Li}({}^3\text{He},d){}^7\text{Be}$	136	3	113.38	0.07	-7.5	C			Mex		64Ma.B
${}^7\text{Li}(t,{}^3\text{He}){}^7\text{He}$	-11184	30	-11147	8	1.2	U			LAl		69St02
${}^7\text{Be}(\varepsilon){}^7\text{Li}$	866	7	861.89	0.07	-0.6	U					72Pe05
${}^7\text{Li}(p,n){}^7\text{Be}$	-1644.04	0.22	-1644.24	0.07	-0.9	U			Zur		61Ry05 Z
	-1643.68	0.26			-2.2	U			Wis		63Ga09 Y
	-1644.30	0.10			0.6	-			Mar		70Ro07 *
	-1644.18	0.10			-0.6	-			Auc		85Wh03 *
	ave.	-1644.24	0.07		0.0	1	100	100 ${}^7\text{Be}$			average
${}^7\text{Li}(\pi^+,\pi^-){}^7\text{B}$	-11870	100	-11747	25	1.2	U					81Se.A
${}^7\text{Be}^i(\text{IT}){}^7\text{Be}$	11000	50	10980	30	-0.4	3					67Ha08
	10970	40			0.3	3					67Mc14
$*{}^7\text{Li}-\text{H}_7$	$D_M=7016003.4256(45)-7\times 1007825.03207(10)$ using Ame2003										06Na13 **
$*{}^7\text{B}-\text{u}$	Represents ${}^7\text{B} \rightarrow 3p + {}^4\text{He}$ , yielding $M-A=27677(25)$ keV										GAu **
$*{}^7\text{H}(\gamma,2n){}^5\text{H}$	From ${}^7\text{H}(\gamma,4n){}^3\text{H} = 704(323)$ , and ${}^5\text{H}(\gamma,2n){}^3\text{H} = 1800(100)$ keV										08Ca22 **
$*{}^7\text{H}(\gamma,2n){}^5\text{H}$	F : not confirmed in later work of reference with higher statistics										10Ni10 **
$*{}^7\text{Li}(d,{}^3\text{He}){}^6\text{He}-{}^{19}\text{F}({}^{18}\text{O})$	$Q-Q=0.98(0.41)$ to $2^+$ level at $1982.07(0.09)$ keV in ${}^{18}\text{O}$										Ens967 **
$*{}^6\text{Li}(n,\gamma){}^7\text{Li}$	Original 7251.02 recalibrated using ${}^{35}\text{Cl}(n,\gamma)$ of reference										82Kr12 **
$*{}^6\text{Li}(n,\gamma){}^7\text{Li}$	Typo 7250.02 in Ame1986 recalib. 7249.97 in Ame1993, 7249.98 Ame2003										GAu **
$*{}^6\text{Li}(n,\gamma){}^7\text{Li}^i$	IT=11200(50); $Q$ rebuilt with Ame1965										MMC128**
$*{}^7\text{Li}(p,n){}^7\text{Be}$	T=1880.64(0.09,Z); error in $Q$ increased										AHW **
$*{}^7\text{Li}(p,n){}^7\text{Be}$	T=1880.43(0.02,Z); error in $Q$ increased										AHW **
${}^8\text{Li}-\text{u}$	22488.2	4.0	22486.25	0.05	-0.5	U			RI1	1.0	13It01
${}^8\text{C}-\text{u}$	37606	32	37643	20	1.2	1	37	37 ${}^8\text{C}$	1.0	1.0	11Ch32 *
${}^8\text{He}-{}^6\text{Li}_{1.333}$	13776.88	0.72	13775.58	0.10	-1.8	o			TT1	1.0	08Ry03
	13775.50	0.19			0.4	1	25	25 ${}^8\text{He}$	1.0	1.0	08Br.D
${}^8\text{Li}-{}^6\text{Li}_{1.333}$	2327.426	0.034	2327.44	0.05	0.4	o			TT1	1.0	08Sm.A
	2327.42	0.11			0.2	o			TT1	1.0	08Sm03
	2327.42	0.11			0.2	1	21	21 ${}^8\text{Li}$	1.0	1.0	09Br10
${}^8\text{He}-{}^7\text{Li}_{1.143}$	15642.49	0.11	15642.46	0.10	-0.3	1	75	75 ${}^8\text{He}$	1.0	1.0	12Br03
${}^4\text{He}({}^{18}\text{O},{}^{14}\text{O}){}^8\text{He}$	-37967	25	-37975.36	0.09	-0.3	U			MIT		75Ja10
${}^4\text{He}({}^{26}\text{Mg},{}^{22}\text{Mg}){}^8\text{He}$	-44962	30	-44999.4	0.3	-1.2	U			Brk		74Ce05
${}^4\text{He}({}^{64}\text{Ni},{}^{60}\text{Ni}){}^8\text{He}$	-31818	15	-31810.6	0.4	0.5	U			Pri		75Ko18
	-31796	8			-1.8	U			Tex		77Tr07
${}^8\text{Be}(\alpha){}^4\text{He}$	91.88	0.05	91.84	0.04	-0.8	3			Zur		68Be02 *
	91.80	0.05			0.8	3					92Wu09 *
	92.2	0.4			-0.9	U					16Re14



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
${}^6\text{Li}(t,p){}^8\text{Li}$	790	11	801.91	0.05	1.1	U			ChR		54Al35
${}^9\text{Li}({}^3\text{He},p){}^8\text{Be}$	16824	12	16787.46	0.04	-3.0	C			Mex		64Ma.B
${}^6\text{Li}(d,\gamma){}^8\text{Be}^j$	-5216.5	3.0	-5213.4	2.0	1.0	1	43	43 ${}^8\text{Be}^j$			76No07 *
${}^9\text{Li}({}^3\text{He},n){}^8\text{B}$	-1974.8	1.0	-1974.8	1.0	0.0	1	100	100 ${}^8\text{B}$	Nvl		58Du78 Y
${}^7\text{Li}(n,\gamma){}^8\text{Li}$	2032.78	0.15	2032.62	0.05	-1.1	-					74Ju.A *
	2032.77	0.18			-0.8	-			ORn		91Ly01 Z
	2032.57	0.06			0.8	-			Bdn		06Fi.A
${}^7\text{Li}(d,p){}^8\text{Li}$	-192	1	-191.95	0.05	0.1	U			Wis		51Wi26 Y
	-188	7			-0.6	U			MIT		64Sp12
${}^7\text{Li}(n,\gamma){}^8\text{Li}$	ave.	2032.61	0.05	2032.62	0.05	0.1	1	79	79 ${}^8\text{Li}$		average
${}^7\text{Li}({}^3\text{He},d){}^8\text{Be}$	11795	13	11760.93	0.04	-2.6	U			Mex		64Ma.B
${}^8\text{C}-u$	Represents ${}^8\text{C} \rightarrow 4p + {}^4\text{He}$ , yielding $M - A=35030(30)$ keV										
${}^8\text{Be}(\alpha){}^4\text{He}$	For atomic binding energy correction see reference										
${}^6\text{Li}(d,\gamma){}^8\text{Be}^j$	$E_d=6962.8(3.0)$ keV										
${}^7\text{Li}(n,\gamma){}^8\text{Li}$	PrvCom to reference										
${}^9\text{Li}-{}^6\text{Li}_{1,500}$	4105.867	0.092	4105.86	0.20	-0.1	o			TT1	1.0	08Sm.A
	4105.86	0.20				2			TT1	1.0	08Sm03
${}^9\text{Be}-{}^7\text{Li}_{1,286}$	-8397.39	0.10	-8397.35	0.08	0.4	1	67	67 ${}^9\text{Be}$	TT1	1.0	09Ri03
${}^9\text{Be}(p,\alpha){}^6\text{Li}$	2117	7	2125.63	0.08	1.2	U			CIT		49To16 Y
	2130	10			-0.4	U			Chi		51Ca37 Y
	2125	4			0.2	U			Wis		51Wi26 Y
	2126	2			-0.2	U			Bir		53Co02 Y
	2144	6			-3.1	B			MIT		64Sp12
	2125.4	1.8			0.1	U			NDm		67Od01
${}^6\text{Li}(\alpha,p){}^9\text{Be}$	-2125.6	1.2	-2125.63	0.08	0.0	U			NDm		65Br28
${}^6\text{Li}(\alpha,n){}^9\text{B}$	-3974	12	-3976.0	0.9	-0.2	U			Tal		63Me08
${}^7\text{Li}(t,p){}^9\text{Li}$	-2397	20	-2386.96	0.19	0.5	U					64Mi04
	-2385.7	3.0			-0.4	U			MSU		75Ka18
${}^9\text{Be}(d,\alpha){}^7\text{Li}$	7162	10	7152.15	0.08	-1.0	U			CIT		51Wh05 Y
	7153	3			-0.3	U			Bir		53Co02 Y
	7162	4			-2.5	U			Mex		64Ma.B
	7157	8			-0.6	U			MIT		64Sp12
${}^7\text{Li}({}^3\text{He},p){}^9\text{Be}$	11215	15	11200.90	0.08	-0.9	U			Mex		64Ma.B
${}^9\text{Be}(p,{}^3\text{He}){}^7\text{Li}^i$	-22499	50	-22450	30	1.0	o			Brk		65De08
	-22479	40			0.8	1	61	61 ${}^7\text{Li}^i$	Brk		67Mc14
${}^7\text{Be}({}^3\text{He},n){}^9\text{C}$	-6287	5	-6282.1	2.1	1.0	3			CIT		67Ba.A Z
	-6275.2	3.5			-2.0	3			CIT		71Mo01 Z
${}^9\text{He}(\gamma,n){}^8\text{He}$	100	60	1250	50	19.2	B			MSU		01Ch31 *
	1270	100			-0.2	-			Ber		99Bo26
	2000	200			-3.7	B					07Go24
	1330	80			-0.9	-					10Jo06 *
	ave.	1310	60		-0.8	1	56	56 ${}^9\text{He}$			average
${}^9\text{Be}(\gamma,n){}^8\text{Be}$	-1665	1	-1664.54	0.08	0.5	U			Wis		50Mo56 Y
${}^9\text{Be}(p,d){}^8\text{Be}$	557	3	560.03	0.08	1.0	U			CIT		49To16 Y
	558	5			0.4	U			Chi		51Ca37 Y
	557.5	1.			2.5	U			Wis		51Wi26 Y
	560	2			0.0	U			Bir		53Co02 Y
	562	4			-0.5	U			MIT		64Sp12
	559.0	1.1			0.9	U			Zur		66Re02
	559.6	0.6			0.7	U			NDm		67Od01 Z
${}^9\text{Be}(d,t){}^8\text{Be}$	4602	13	4592.69	0.08	-0.7	U			MIT		64Sp12
	4591.7	3.1			0.3	U			NDm		67Od01
${}^9\text{Be}({}^3\text{He},\alpha){}^8\text{Be}$	18931	13	18913.08	0.08	-1.4	U			Mex		64Ma.B
${}^9\text{Be}(\pi^-, \pi^+){}^9\text{He}$	-30472	100	-30610	50	-1.4	-					87Se05

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
${}^9\text{Be}({}^{13}\text{C}, {}^{13}\text{O}){}^9\text{He}$	-50200	600	-49580	50	1.0	o			Ber		88Bo20
	-49470	80			-1.3	o			Ber		91Bo.B
${}^9\text{Be}({}^{14}\text{C}, {}^{14}\text{O}){}^9\text{He}$	-34580	100	-34580	50	0.0	-			Ber		95Bo.B
${}^9\text{Be}(\pi^-, \pi^+){}^9\text{He}$	ave. -30540	70	-30610	50	-0.9	1	44	44 ${}^9\text{He}$			average
${}^9\text{Be}(p,n){}^9\text{B}$	-1850.4	1.0				2			Wis		50Ri59 Z
	-1852	3	-1850.4	0.9	0.5	U			Ric		55Ma84 Z
$*{}^9\text{He}(\gamma,n){}^8\text{He}$	From scattering length $a_s = -10$ fm; questioned in reference										10Jo06 **
$*{}^9\text{He}(\gamma,n){}^8\text{He}$	Scattering length $a_s = -3.17(66)$ fm										10Jo06 **
${}^{10}\text{B} {}^{37}\text{Cl}-\text{C} {}^{35}\text{Cl}$	9987.21	0.56	9986.75	0.07	-0.3	U			H38	2.5	84El05
${}^{10}\text{Be}-{}^7\text{Li}_{1,429}$	-9334.16	0.13	-9334.22	0.09	-0.4	1	44	44 ${}^{10}\text{Be}$	TT1	1.0	09Ri03
${}^{10}\text{B}-\text{u}$	12936.862	0.016	12936.862	0.016	0.0	1	100	100 ${}^{10}\text{B}$	MS1	1.0	16Gu02
${}^{10}\text{C}-{}^{10}\text{B}$	3916.413	0.090	3916.36	0.07	-0.6	1	67	67 ${}^{10}\text{C}$	JY1	1.0	11Er02
${}^7\text{Li}(t,\gamma){}^{10}\text{Be}^i$	-3929.8	21.0				2					73Ab10
${}^{10}\text{B}(n,\alpha){}^7\text{Li}$	2801	4	2789.906	0.016	-2.8	U					67De15
${}^7\text{Li}(\alpha,n){}^{10}\text{B}$	-2787	4	-2789.906	0.016	-0.7	U			Ric		57Bi84 Y
${}^{10}\text{B}(p,\alpha){}^7\text{Be}$	1147	5	1145.67	0.07	-0.3	U			CIT		49Ch35 Y
	1146	6			-0.1	U			CIT		51Br10 Y
	1146	2			-0.2	U			Wis		52Cr30 Y
	1153	4			-1.8	U			MIT		64Sp12
${}^{10}\text{B}({}^3\text{He}, {}^6\text{He}){}^7\text{B}$	-18550	100	-18287	25	2.6	U			Brk		67Mc14
${}^{10}\text{He}(\gamma,2n){}^8\text{He}$	1200	300	1440	90	0.8	U					94Ko16
	1420	100			0.2	2					10Jo06
	2100	200			-3.3	B					12Si07
	1600	250			-0.6	2					12Ko43
	1400	300			0.1	U					15Ma54
${}^{10}\text{Be}(p, {}^3\text{He}){}^8\text{Li}^i$	-26802.3	5.4				2			MSU		75Ro01 *
${}^{10}\text{Be}(p,t){}^8\text{Be}^j$	-27487.0	2.6	-27489.3	2.0	-0.9	1	57	57 ${}^8\text{Be}^j$	MSU		75Ro01 *
${}^{10}\text{B}(d,\alpha){}^8\text{Be}$	17829	10	17819.74	0.04	-0.9	U			Bir		54El10 Y
	17830	6			-1.7	U			Mex		64Ma.B
	17818.6	4.1			0.3	U			NDm		67Od01
${}^{10}\text{B}(p, {}^3\text{He}){}^8\text{Be}$	-535.5	2.5	-533.31	0.04	0.9	U			Wis		52Cr30 Y
${}^{10}\text{Li}(\gamma,n){}^9\text{Li}$	150	150	26	13	-0.8	U					90Am05 *
	25	15			0.1	3					95Zi03 *
	30	24			-0.1	3					08Ak03 *
${}^{10}\text{Li}^m(\gamma,n){}^9\text{Li}$	240	60	220	40	-0.3	3					97Bo10 *
	210	50			0.2	3					97Zi04 *
${}^9\text{Be}({}^9\text{Be}, {}^8\text{B}){}^{10}\text{Li}^n$	-33770	260	-33750	40	0.1	U			Brk		75Wi26 *
${}^9\text{Be}({}^{13}\text{C}, {}^{12}\text{N}){}^{10}\text{Li}^n$	-36370	50	-36390	40	-0.4	2			Ber		93Bo03 *
${}^{10}\text{Be}(d, {}^3\text{He}){}^9\text{Li}$	-14142.8	2.5	-14142.91	0.20	0.0	U			MSU		75Ka18
${}^9\text{Be}(n,\gamma){}^{10}\text{Be}$	6812.33	0.06	6812.28	0.05	-0.8	-			MMn		86Ke14 Z
	6812.10	0.14			1.3	-			Bdn		06Fi.A
${}^9\text{Be}(d,p){}^{10}\text{Be}$	4583	8	4587.72	0.05	0.6	U			Ric		51Ki55 Y
	4595	4			-1.8	U			Mex		64Ma.B
	4590	8			-0.3	U			MIT		64Sp12
${}^9\text{Be}(n,\gamma){}^{10}\text{Be}$	ave. 6812.29	0.06	6812.28	0.05	-0.2	1	88	56 ${}^{10}\text{Be}$			average
${}^9\text{Be}({}^3\text{He}, d){}^{10}\text{B}$	1123	5	1093.34	0.08	-5.9	C			Mex		64Ma.B
${}^{10}\text{B}(d,t){}^9\text{B}$	-2189	10	-2180.0	0.9	0.9	U			MIT		64Sp12
${}^{10}\text{B}({}^3\text{He}, \alpha){}^9\text{B}$	12130	15	12140.4	0.9	0.7	U			Ric		60Sp08
	12171	15			-2.0	U			Mex		64Ma.B
${}^{10}\text{Be}({}^{14}\text{C}, {}^{14}\text{O}){}^{10}\text{He}$	-41190	70	-41580	90	-5.5	B			Ber		94Os04
${}^{10}\text{Be}(\beta^-){}^{10}\text{B}$	560	5	556.88	0.08	-0.6	U					50Hu27
	555	5			0.4	U					52Fe16
${}^{10}\text{C}(\beta^+){}^{10}\text{B}$	3604	16	3648.06	0.07	2.8	U					63Ba52
${}^{10}\text{B}(p,n){}^{10}\text{C}$	-4433.7	1.5	-4430.41	0.07	2.2	U			Har		75Fr.A
	-4430.17	0.34			-0.7	o			Auc		84Ba12 *
	-4430.17	0.09			-2.7	o			Auc		89Ba28 *
	-4430.30	0.12			-0.9	1	33	33 ${}^{10}\text{C}$	Auc		98Ba83 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{10}\text{B}(^3\text{He,t})^{10}\text{C}$	-3667	10	-3666.65	0.07	0.0	U			Brk		68Br23
$^{10}\text{B}(^{14}\text{N},^{14}\text{B})^{10}\text{N}$	-47550	400				2					02Le16
* $^{10}\text{Be}(p,^3\text{He})^8\text{Li}^i$	Original value -26804.1(5.4) recalibrated										
* $^{10}\text{Be}(p,t)^8\text{Be}^j$	Original -27487.6(2.6) recalibrated										
* $^{10}\text{Li}(\gamma,n)^9\text{Li}$	From $^{11}\text{B}(\pi^-,p)^{10}\text{Li}$										
* $^{10}\text{Li}(\gamma,n)^9\text{Li}$	Resonance less than 50 above the one neutron threshold, but could also be final state interaction; then $^{10}\text{Li}$ would be 200 higher										
* $^{10}\text{Li}(\gamma,n)^9\text{Li}$	Deduced from s-state scattering length $a_s = -22.4(4.8)$ fm										
* $^{10}\text{Li}^m(\gamma,n)^9\text{Li}$	From $^{10}\text{Be}(^{12}\text{C},^{12}\text{N})^{10}\text{Li}^m$ ( $1^+$ level)										
* $^{10}\text{Li}^m(\gamma,n)^9\text{Li}$	Theoretical work: $1^+$ level above $1^-$ ground state										
* $^9\text{Be}(^9\text{Be},^8\text{B})^{10}\text{Li}^n$	$Q = -34060(250)$ to $2^+$ level 290(80) above $1^+$ level revised with Breit-Wigner line shape. Probably $2^+$ level										
* $^9\text{Be}(^{13}\text{C},^{12}\text{N})^{10}\text{Li}^n$	Revised with Breit-Wigner line shape (probably $2^+$ level)										
* $^{10}\text{B}(p,n)^{10}\text{C}$	T=4876.90(0.37); withdrawn by author										
* $^{10}\text{B}(p,n)^{10}\text{C}$	T=4876.88(0.10,Z); original T=4876.95(0.10) keV										
* $^{10}\text{B}(p,n)^{10}\text{C}$	Average of two datasets; withdrawn by author										
* $^{10}\text{B}(p,n)^{10}\text{C}$	T=4877.03(0.13); this is the second 89Ba28 dataset, recalibrated by author										
$^{11}\text{B }^{37}\text{Cl}-^{13}\text{C }^{35}\text{Cl}$	2998.15	1.30	3000.22	0.07	0.6	U			H38	2.5	84El05
$^{11}\text{Li}-^6\text{Li}_{1,833}$	16003.5	1.2	16003.3	0.7	-0.1	o			TT1	1.0	08Sm.A
	16003.33	0.66				2			TT1	1.0	08Sm03
$^{11}\text{Be}-^6\text{Li}_{1,833}$	-6059.27	0.28	-6059.17	0.26	0.4	1	83	83 $^{11}\text{Be}$	TT1	1.0	08Br.C
$^{11}\text{Be}-^7\text{Li}_{1,571}$	-3479.83	0.62	-3480.32	0.26	-0.8	1	17	17 $^{11}\text{Be}$	TT1	1.0	09Ri03
$^{11}\text{C}-^{14}\text{N}_{786}$	9016.430	0.064	9016.43	0.06	0.0	1	100	100 $^{11}\text{C}$	MS1	1.0	16Gu02
$^{11}\text{Li}-u$	43780	130	43723.6	0.7	-0.3	U			TO2	1.5	88Wo09
	43805	28			-2.9	U			P40	1.0	03Ba.A
	43715.4	5.0			1.6	o			P40	1.0	04Ba.A
	43714.5	5.1			1.8	U			P40	1.0	09Ga24
$^{11}\text{Be}-u$	21654.0	3.6	21661.08	0.26	2.0	o			P40	1.0	04Ba.A
	21653.5	3.5			2.2	U			P40	1.0	09Ga24
	21658.5	3.8			0.7	U			P40	1.0	09Ga24
$^{11}\text{B}-u$	9305.167	0.013	9305.167	0.013	0.0	1	100	100 $^{11}\text{B}$	MS1	1.0	16Gu02
$^9\text{Li}-^{11}\text{Li}_{491}$ $^6\text{Li}_{600}$	-3949	175	-3494.8	0.4	1.0	U			P12	2.5	75Th08
$^9\text{Li}-^{11}\text{Li}_{409}$ $^7\text{Li}_{643}$	-1250	86	-1288.2	0.3	-0.3	U			P11	1.5	75Th08
	-1223	195			-0.3	U			P13	1.0	75Th08
$^9\text{Li}-^{11}\text{Li}_{273}$ $^8\text{Li}_{750}$	-1928	31	-1873.26	0.25	0.7	U			P12	2.5	75Th08
	-1923	31			1.6	U			P13	1.0	75Th08
$^7\text{Li}(\alpha,\gamma)^{11}\text{B}^i$	-3885.6	20.0	-3896	9	-0.5	1	21	21 $^{11}\text{B}^i$			66Cu02
$^{11}\text{B}(p,\alpha)^8\text{Be}$	8583	15	8590.09	0.04	0.5	U			CIT		51Li26
	8589	4			0.3	U			Bir		53Co02
	8597	6			-1.2	U			Mex		61Ja23
	8575	11			1.4	U			MIT		64Sp12
$^{11}\text{B}(^3\text{He},^6\text{He})^8\text{B}^i$	-27539	8				2			MSU		75Ro01
$^{11}\text{Be}^i(\gamma,d)^9\text{Li}$	3245	20				3					97Te07
$^9\text{Be}(t,p)^{11}\text{Be}$	-1164	15	-1167.87	0.25	-0.3	U			Ald		62Pu01
$^{11}\text{B}^j(2p)^9\text{Li}$	2700	80				3					12Ch40
$^{11}\text{B}(d,\alpha)^9\text{Be}$	8029	4	8030.06	0.08	0.3	U			Bir		54El10
	8035	9			-0.5	U			Mex		64Ma.B
	8024	7			0.9	U			MIT		64Sp12
	8029.7	2.8			0.1	U			NDm		67Od01
$^9\text{Be}(^3\text{He},p)^{11}\text{B}$	10344	13	10322.99	0.08	-1.6	U			Mex		64Ma.B
	10322.1	2.3			0.4	U			NDm		67Od01
$^{11}\text{B}(p,^3\text{He})^9\text{Be}^i$	-24713.3	1.7				2			MSU		74Ka15
$^9\text{Be}(^3\text{He},p)^{11}\text{B}^i$	-2240	12	-2237	9	0.2	-					63Gr.A
	-2240.6	20.			0.2	-			MSU		82Zw02
ave.	-2240	10			0.3	1	79	79 $^{11}\text{B}^i$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{11}\text{B}(\text{p,t})^9\text{B}^i$	-26064.1	2.3				2			MSU		74Ka15 *
$^9\text{Be}(\text{}^3\text{He,n})^{11}\text{C}^i$	-4612	50	-4600	40	0.2	1	50	50 $^{11}\text{C}^i$			71Wa21 *
$^{10}\text{Be}(\text{d,p})^{11}\text{Be}$	-1721	7	-1722.93	0.25	-0.3	U			CIT		70Go11
$^{11}\text{B}(\text{}^7\text{Li},\text{}^8\text{B})^{10}\text{Li}$	-32431	80	-32399	13	0.4	U			MSU		94Yo01 *
$^{11}\text{B}(\text{}^7\text{Li},\text{}^8\text{B})^{10}\text{Li}^n$	-32908	62	-32870	40	0.5	R			MSU		94Yo01
$^{10}\text{Be}(\text{p},\gamma)^{11}\text{B}^i$	-1322	30	-1332	9	-0.3	U					70Go04 *
$^{10}\text{B}(\text{n},\gamma)^{11}\text{B}$	11454.1	0.2	11454.219	0.019	0.6	U			Ptn		86Ko19 Z
	11454.15	0.27			0.3	U			Bdn		06Fi.A
$^{10}\text{B}(\text{d,p})^{11}\text{B}$	9227	5	9229.653	0.019	0.5	U			Bir		54El10 Y
	9234	6			-0.7	U			Mex		64Ma.B
	9244	11			-1.3	U			MIT		64Sp12
	9232.9	2.			-1.6	U			NDm		66Br18
$^{11}\text{B}(\text{}^3\text{He},\alpha)^{10}\text{B}$	9101	20	9123.400	0.019	1.1	U			Man		60Ta12
$^{10}\text{B}(\text{}^3\text{He},\text{d})^{11}\text{C}$	3174	15	3196.71	0.06	1.5	U			Man		60Fo01
	3226	10			-2.9	C			Mex		64Ma.B
$^{11}\text{N}(\text{p})^{10}\text{C}$	1973	180	1320	50	-3.7	B			MSU		74Be20 *
	1300	40			0.4	o			Lis		96Ax01
	1450	400			-0.3	U			MSU		98Az01 *
	1630	50			-6.3	B			Spe		00O101 *
	1350	120			-0.3	2			Lis		00Ma62 *
	1310	50			0.1	2			INS		03Gu06
	1540	20			-11.2	B					06Ca05
$^{11}\text{B}(\pi^-, \pi^+)^{11}\text{Li}$	-33120	50	-33082.5	0.6	0.7	U					91Ko.B
$^{11}\text{B}(\text{}^{14}\text{C}, \text{}^{14}\text{O})^{11}\text{Li}$	-37120	35	-37048.4	0.6	2.0	U			MSU		93Y007
$^{11}\text{B}^i(\text{IT})^{11}\text{B}$	12510	50	12560	9	1.0	U					71Wa21 *
$^{11}\text{C}(\beta^+)^{11}\text{B}$	1982.8	2.6	1981.69	0.06	-0.4	U					75Be28
$^{11}\text{B}(\text{p,n})^{11}\text{C}$	-2759.7	3.	-2764.04	0.06	-1.4	U			Wis		50Ri59 Z
	-2763.2	1.4			-0.6	U			Ric		61Be13 Z
$^{11}\text{B}(\text{}^3\text{He},\text{t})^{11}\text{C}$	-2002.1	1.2	-2000.28	0.06	1.5	U			Str		65Go05 Z
$^{11}\text{B}(\text{}^3\text{He},\text{t})^{11}\text{C}^i$	-14151	50	-14160	40	-0.2	1	50	50 $^{11}\text{C}^i$			71Wa21 *
$^{11}\text{Be}-\text{u}$	Result from the "cooling" experiment										09Ga24 **
$^9\text{Li}-^{11}\text{Li}_{409} \text{}^7\text{Li}_{643}$	Symmetric double-doublet 6-9 8-11 included										GAu **
$^7\text{Li}(\alpha,\gamma)^{11}\text{B}^i$	IT=12550(30); $Q$ rebuilt with Ame1964										GAu **
$^{11}\text{B}(\text{}^3\text{He},\text{}^6\text{He})^8\text{B}^i$	IT=10619(9); rebuilt $Q = -27538.9(8.2)$ keV										GAu **
*	IT=10614(20) keV by a different method										14Br15 **
$^{11}\text{Be}^i(\gamma,\text{d})^9\text{Li}$	$Q(\text{d})=Q(\text{p+n})+2224.5660(0.0004)$ $Q(\text{p+n})=1020(20)$ keV										MMC162**
$^{11}\text{B}(\text{p},\text{}^3\text{He})^9\text{Be}^i$	IT=14392.2(1.8); rebuilt $Q = -24715.2(1.7)$ ; recalibrated +1.87 keV										MMC121**
$^9\text{Be}(\text{}^3\text{He},\text{p})^{11}\text{B}^i$	IT=12565(12); $Q$ rebuilt with Ame1964										MMC121**
$^9\text{Be}(\text{}^3\text{He},\text{p})^{11}\text{B}^i$	IT=12563(20); $Q$ rebuilt with Ame1977										MMC121**
$^{11}\text{B}(\text{p,t})^9\text{B}^i$	IT=14655.4(2.5); rebuilt $Q = -26064.3$ ; recalibrated +0.16 keV										MMC121**
$^9\text{Be}(\text{}^3\text{He},\text{n})^{11}\text{C}^i$	IT=12170(40); $Q$ rebuilt; possibly not pure T=3/2										MMC121**
$^{11}\text{B}(\text{}^7\text{Li},\text{}^8\text{B})^{10}\text{Li}$	Original (>-32471) re-evaluated										GAu **
*	existence of this level not completely certain										94Yo01 **
$^{10}\text{Be}(\text{p},\gamma)^{11}\text{B}^i$	IT=12550(30); $Q$ rebuilt with Ame1964										MMC121**
$^{11}\text{N}(\text{p})^{10}\text{C}$	From $^{14}\text{N}(\text{}^3\text{He},\text{}^6\text{He})^{11}\text{N}$ $Q = -25010(100)$ to 250(150) level										90Aj01 **
$^{11}\text{N}(\text{p})^{10}\text{C}$	From $^9\text{Be}(\text{}^{12}\text{N},\text{}^{10}\text{Be})^{11}\text{N}$										98Az01 **
$^{11}\text{N}(\text{p})^{10}\text{C}$	From $^{10}\text{B}(\text{}^{14}\text{N},\text{}^{13}\text{B})^{11}\text{N}$										00O101 **
$^{11}\text{N}(\text{p})^{10}\text{C}$	From scattering $^{10}\text{C}$ on H; precisely, 1270(+180-50) keV										00Ma62 **
$^{11}\text{B}^i(\text{IT})^{11}\text{B}$	From $^{11}\text{B}(\text{}^3\text{He},\text{}^3\text{He}')$										AHW **
$^{11}\text{B}(\text{}^3\text{He},\text{t})^{11}\text{C}^i$	IT=12150(50); $Q$ rebuilt; possibly not pure T=3/2										MMC121**
$^{12}\text{Be}-\text{u}$	26911.3	14.2	26922.1	2.0	0.8	U			P40	1.0	09Ga24
$^{12}\text{Be}-\text{C}$	26922.4	2.3			-0.1	1	79	79 $^{12}\text{Be}$	TT1	1.0	10Et01
$\text{C}_{14}-^{12}\text{C}_{12}$	1.2	4.9	0.00000	0.00013	-0.2	U			TG1	1.5	09Ke.A
$\text{C}_{14}-^{12}\text{C}_{15}$	2.1	2.7			-0.5	U			TG1	1.5	11Ke03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_{15-^{12}C_{14}}$	-4.0	4.7			0.6	U			TG1	1.5	11Ke03
	-0.5	6.2			0.1	U			TG1	1.5	11Ke03
$C_{15-^{12}C_{16}}$	-1.6	2.1			0.5	U			TG1	1.5	10Ke09
	4.7	4.2			-0.7	U			TG1	1.5	11Ke03
	3.7	4.1			-0.6	U			TG1	1.5	11Ke03
$C_{16-^{12}C_{12}}$	-0.3	6.1			0.0	U			TG1	1.5	09Ke.A
	2.6	5.0			-0.3	U			TG1	1.5	09Ke.A
$C_{16-^{12}C_{15}}$	-1.3	3.9			0.2	U			TG1	1.5	11Ke03
	-0.7	1.3			0.4	U			TG1	1.5	12Sm07
$C_{21-^{12}C_{22}}$	1.6	1.3			-0.8	U			TG1	1.5	14Ei01
$C_{23-^{12}C_{22}}$	1.2	1.7			-0.5	U			TG1	1.5	14Ei01
$C_{23-^{12}C_{21}}$	2.2	1.8			-0.8	U			TG1	1.5	14Ei01
$^7Li(^7Li,2p)^{12}Be$	-9710	100	-9841.5	1.9	-1.3	U			LAI		71Ho26
$^{12}C(\alpha,^8He)^8C$	-64520	200	-64249	18	1.4	U					74Ro17
	-64278	26			1.1	-			Tex		76Tr01
ave.	-64270	23			0.9	1	63	$^{63}C$			average
$^9Be(^7Li,\alpha)^{12}B^i$	-2308.4	50.	-2258	19	1.0	1	14	$^{14}^{12}B^i$	Phi		75Aj03 *
$^{12}C(^3He,^6He)^9C$	-31578	8	-31571.8	2.1	0.8	U			MSU		71Tr03
	-31575.6	3.2			1.2	R			MSU		79Ka.A
$^{10}Be(t,p)^{12}Be$	-4809	15	-4809.4	1.9	0.0	U			Brk		78Al29
	-4808.3	4.2			-0.3	1	21	$^{21}^{12}Be$	Phi		94Fo08
$^{10}B(t,p)^{12}B$	6346	6	6342.1	1.3	-0.7	U			Man		60Ja17
$^{10}B(\alpha,d)^{12}C$	1340.3	0.8	1339.803	0.015	-0.6	U			Wis		56Do41 Z
	1340.6	1.5			-0.5	U			NDm		65Br28
$^{12}C(d,\alpha)^{10}B$	-1340.1	1.2	-1339.803	0.015	0.2	U			NDm		65Br28
$^{10}B(^3He,p)^{12}C$	19694.5	3.6	19692.857	0.015	-0.5	U			NDm		67Od01
	19692.86	0.44			0.0	U			Mun		83Ch08 *
$^{10}B(^3He,p)^{12}C^i$	4585	6	4585	3	-0.1	1	31	$^{31}^{12}C^i$			62Br10
$^{10}B(^3He,n)^{12}N$	1570	25	1572.4	1.0	0.1	U			CIT		64Fi02
	1561	9			1.3	U			CIT		64Ka08
	1568	20			0.2	U			LAI		66Za01
	1574	7			-0.2	U			Har		68Ad03
$^{12}N^i(2p)^{10}B$	2905	29				2					12Ja11 *
$^{12}O(2p)^{10}C$	1770	20	1638	24	-6.6	B					95Kr03
	1638	24				2					12Ja11
$^{12}Li(\gamma,n)^{11}Li$	120	15	210	30	6.0	B					08Ak03 *
	210	30				3					13Ko03
$^{11}B(d,p)^{12}B$	1141	4	1145.1	1.3	1.0	1	11	$^{11}^{12}B$	Mex		61Ja23
	1137	5			1.6	U			MIT		64Sp12
$^{11}B(^3He,d)^{12}C$	10436	17	10463.203	0.012	1.6	U			Man		60Fo01
	10469.7	5.7			-1.1	U			NDm		67Od01
$^{11}B(d,n)^{12}C^i$	-1376.2	4.0	-1376	3	0.0	1	69	$^{69}^{12}C^i$			55Ma76 *
$^{12}N(\beta^+)^{12}C$	17406	15	17338.1	1.0	-4.5	B					63GI04
$^{12}C(p,n)^{12}N$	-18119.9	4.4	-18120.4	1.0	-0.1	U			Yal		69Ov01 Z
$^{12}C(\pi^+,\pi^-)^{12}O$	-31037	48	-30893	24	3.0	B					80Bu15
	-31014	24			5.1	B					92Iv.A
$^9Be(^7Li,\alpha)^{12}B^i$	IT=12770(50) using $Q_{gs}=10461.6(1.6)$ keV										75Aj03 **
	energy and resolution arguments for T=1, not an IAS										08Ch28 **
$^{10}B(^3He,p)^{12}C$	Original $Q=15305.45(0.3)$ revised by authors to $15253.95(31)$ keV										83Vo.A **
	to $2^+$ level at $4438.91(0.31)$ keV										Ens006 **
$^{12}N^i(2p)^{10}B$	$Q_{2p}=1165(29)$ to $^{10}B^i$ at $1740.05(0.04)$										Nub16b **
$^{12}Li(\gamma,n)^{11}Li$	$E_{res}$ derived in reference from scattering length $a_s=-13.7(1.6)$ fm										10Ha04 **
$^{11}B(d,n)^{12}C^i$	$E_{res}=1627(4)$ $Q=-1376(4)$ ; recalibrated $Q=-1376.16(4.00)$ keV										MMC121**

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
C H- <sup>13</sup> C	4500	36	4470.19703	0.00027	-0.6	U			R08	1.5	69De19
	4470.185	0.008			1.0	U			B08	1.5	75Sm02
	4470.10	0.05			0.8	U			M25	2.5	78Ha14
C D- <sup>13</sup> C H	2921.923	0.008	2921.91066	0.00025	-1.0	U			B08	1.5	75Sm02
	2921.87	0.05			0.3	U			M25	2.5	78Ha14
	2921.9086	0.0012			1.7	U			MI1	1.0	95Di08
	2921.9074	0.0015			2.2	U			MI1	1.0	95Di08
<sup>13</sup> C-u	3354.8404	0.0041	3354.83521	0.00023	-1.3	U			WA1	1.0	95Va38
<sup>9</sup> Be( $\alpha,\gamma$ ) <sup>13</sup> C <sup>i</sup>	-4458.4	2.0	-4460.4	1.1	-1.0	2					73Ad02
	-4461.4	1.4			0.7	2					78Hi06
<sup>10</sup> B( $\alpha,p$ ) <sup>13</sup> C	4068	12	4061.546	0.015	-0.5	U			MIT		64Sp12
	4063.4	2.4			-0.8	U			NDm		67Od01
<sup>13</sup> Li( $\gamma,2n$ ) <sup>11</sup> Li	1470	310	110	70	-4.4	B					08Ak03 *
	1470	350			-3.9	B					10Jo07 *
	110	70				3					13Ko03 *
<sup>11</sup> B(t,p) <sup>13</sup> B	-233	4	-233.4	1.0	-0.1	U			Man		60Mu07
	-233.4	1.0				2			Str		83An15
<sup>13</sup> C(d, $\alpha$ ) <sup>11</sup> B	5169	6	5168.108	0.012	-0.1	U			CIT		51Li29 Y
	5166	5			0.4	U			Ric		53Ph28 Y
	5165	10			0.3	U			MIT		64Sp12
	5166.6	2.5			0.6	U			NDm		70Br23
<sup>11</sup> B( <sup>3</sup> He,p) <sup>13</sup> C	13221	10	13184.946	0.012	-3.6	C			Mex		64Ma.B
	13185.4	4.0			-0.1	U			NDm		67Od01
<sup>11</sup> B( <sup>3</sup> He,n) <sup>13</sup> N	10183	11	10182.13	0.27	-0.1	U					71Hs03
<sup>13</sup> Be( $\gamma,n$ ) <sup>12</sup> Be	100	70	510	10	5.9	B					01Th01 *
	60	10			45.0	B					08Ch07 *
	510	10				2					10Ko17
	400	30			3.7	B					14Ra07
<sup>12</sup> C(n, $\gamma$ ) <sup>13</sup> C	4946.47	0.17	4946.3083	0.0005	-1.0	U					67Pr10
	4946.03	0.15			1.9	U			Utr		68Sp01
	4946.51	0.31			-0.7	U			ILn		79Br25 Z
	4946.321	0.024			-0.5	U					80Wa24 *
	4946.337	0.031			-0.9	U			Utr		81Va.B *
	4946.31	0.10			0.0	U			Bdn		06Fi.A
<sup>12</sup> C(d,p) <sup>13</sup> C	2727	6	2721.74226	0.00023	-0.9	o			Ric		51Ki55 Y
	2722	4			-0.1	U			Ric		53Fa18 Y
	2720	2			0.9	U			Bir		54El10 Y
	2725	5			-0.7	U			Mex		61Ja23
	2722	4			-0.1	U			MIT		64Sp12
	2722.3	0.6			-0.9	o			NDm		67Od01
	2721.9	0.8			-0.2	U			NDm		74Jo14
	2721.80	0.50			-0.1	U			Rez		90Pi05 *
	-2722	7	-2721.74226	0.00023	0.0	o			MIT		64Sp12
<sup>13</sup> C(d,t) <sup>12</sup> C	1311	3	1310.92070	0.00029	0.0	U			CIT		51Li29 Y
	1311	6			0.0	U			Mex		64Ma.B
	1311	6			0.0	U			MIT		64Sp12
	1310.9	0.7			0.0	U			NDm		67Od01
<sup>12</sup> C(p, $\gamma$ ) <sup>13</sup> N	1943.24	0.32	1943.49	0.27	0.8	-					77Fr20 Z
	1944.1	0.5			-1.2	-					77He26 Z
<sup>12</sup> C(d,n) <sup>13</sup> N	-280.5	3.	-281.08	0.27	-0.2	U			Ric		49Bo67 Y
<sup>12</sup> C(p, $\gamma$ ) <sup>13</sup> N	ave.	1943.49	1943.49	0.27	0.0	1	100	100 <sup>13</sup> N			average
<sup>12</sup> C(p, $\gamma$ ) <sup>13</sup> N <sup>i</sup>	-13121.62	0.18				2					73Hu07
<sup>13</sup> C( <sup>14</sup> C, <sup>14</sup> O) <sup>13</sup> Be <sup>p</sup>	-37020	50				2			Ber		92Os04
<sup>13</sup> N( $\beta^+$ ) <sup>13</sup> C	2222.3	3.8	2220.47	0.27	-0.5	U					54Ki23
<sup>13</sup> C(p,n) <sup>13</sup> N	-3002.3	1.0	-3002.82	0.27	-0.5	o			Ric		61Be13 Z
	-3004.1	1.5			0.9	U			NRL		64Bo10 Z
	-3002.4	1.0			-0.4	U			Ric		66Bo20

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{13}\text{O}(\beta^+)^{13}\text{N}$	17500	200	17770	10	1.3	U					65Mc09
* $^{13}\text{Li}(\gamma,2n)^{11}\text{Li}$	Corresponds to excited state										
* $^{13}\text{Li}(\gamma,2n)^{11}\text{Li}$	Symmetrized from 120(+60-80)										
* $^{13}\text{Be}(\gamma,n)^{12}\text{Be}$	From scattering length $a_s < -10$ fm; questioned in reference										
* $^{13}\text{Be}(\gamma,n)^{12}\text{Be}$	From scattering length $a_s = -20$ fm; questioned in reference										
*	$a_s = -3.4(0.6)$ fm in 10Ko17; $a_s = -3.2(+.9-1.1)$ fm in 07Si24										
* $^{12}\text{C}(n,\gamma)^{13}\text{C}$	$Q(\gamma)=1261.844(0.006,Z)$ to $3684.477(0.023,Z)$ level										
* $^{12}\text{C}(n,\gamma)^{13}\text{C}$	$Q(\gamma)=1261.844(0.006,Z)$ to $3684.493(0.030,Z)$ level										
* $^{12}\text{C}(d,p)^{13}\text{C}$	Estimated systematic error 0.5 added to statistical error 0.038 keV										
$^{14}\text{Be}-u$	42660	150	42890	140	1.0	2					88Wo09
$\text{C D}_2-^{14}\text{C H}_2$	9311.498	0.006	9311.503	0.004	0.6	1	20	20	$^{14}\text{C}$		B08 1.5 75Sm02
$\text{C H}_2-N$	12576.22	0.10	12576.06002	0.00026	-0.6	U					J2 2.5 69Na21
	12576.086	0.009			-1.9	U					B07 1.5 71Sm01
	12576.0598	0.0008			0.3	U					MI1 1.0 95Di08
$\text{C D}-N$	11027.815	0.018	11027.77365	0.00024	-1.5	o					B07 1.5 71Sm01
	11027.773	0.007			0.1	U					B08 1.5 75Sm02
$\text{C H}_4-N \text{ D}$	14124.17	0.14	14124.3464	0.0004	0.5	U					J6 2.5 76Ka50
$\text{C D}_2-N \text{ H}_2$	9479.68	0.13	9479.4873	0.0003	-0.6	U					J6 2.5 76Ka50
$^{14}\text{N}-u$	3074.014	0.019	3074.00446	0.00021	-0.2	U					OH1 2.5 93Ma.A
	3074.0056	0.0018			-0.6	U					WA1 1.0 95Va38
$^{13}\text{C H}-^{14}\text{N}$	8105.86299	0.00010	8105.86299	0.00010	0.0	1	97	79	$^{13}\text{C}$		MI3 1.0 04Ra33 *
$^{14}\text{C H}_2-N \text{ D}$	1716.269	0.003	1716.270	0.004	0.3	1	80	80	$^{14}\text{C}$		B08 1.5 75Sm02
$^{14}\text{O}-^{14}\text{N}$	5522.702	0.027	5522.702	0.027	0.0	1	100	100	$^{14}\text{O}$		MS1 1.0 15Va08
$^{14}\text{N}(^3\text{He},^9\text{Li})^8\text{C}$	-42214	50	-42225	18	-0.2	R					MSU 76Ro04
$^{11}\text{B}(\alpha,p)^{14}\text{C}$	789	17	783.759	0.013	-0.3	U					MIT 64Sp12
$^{14}\text{B}^i(\gamma,d)^{12}\text{Be}$	2515	20				2					01Ta23 *
$^{14}\text{C}(^{18}\text{O},^{20}\text{Ne})^{12}\text{Be}$	-15770	50	-15798.8	1.9	-0.6	U					ChR 74Ba15
$^{14}\text{C}(d,\alpha)^{12}\text{B}$	361.8	1.4	361.3	1.3	-0.4	1	89	89	$^{12}\text{B}$		Wis 56Do41 Z
$^{14}\text{C}(p,^3\text{He})^{12}\text{B}^i$	-30702.73	19.96	-30711	19	-0.4	1	86	86	$^{12}\text{B}^i$		71Ne.A *
$^{14}\text{C}(p,t)^{12}\text{C}^j$	-32235.9	2.4				2					MSU 78Ro08 *
$^{14}\text{N}(d,\alpha)^{12}\text{C}$	13579	6	13574.22287	0.00024	-0.8	U					Mex 64Ma.B
	13588	6			-2.3	U					MIT 64Sp12
$^{12}\text{C}(^3\text{He},p)^{14}\text{N}$	4779.0	1.4	4778.83059	0.00026	-0.1	U					CIT 62Ba26 Y
	4806	9			-3.0	C					Mex 64Ma.B
	4776.3	1.5			1.7	U					NDm 67Od01
$^{14}\text{N}(p,t)^{12}\text{N}$	-22135.5	1.0	-22135.5	1.0	0.0	1	100	100	$^{12}\text{N}$		MSU 75No.A
$^{12}\text{C}(^3\text{He},n)^{14}\text{O}$	-1146.86	0.72	-1147.880	0.025	-1.4	U					Nvl 61Bu04 *
	-1148.61	0.56			1.3	U					CIT 62Ba26 *
	-1149.01	0.48			2.4	U					Mar 70Ro07 *
$^{14}\text{C}(^7\text{Li},^8\text{B})^{13}\text{Be}$	-39990	500	-38654	10	2.7	U					Dbn 83A120
$^{14}\text{C}(^{11}\text{B},^{12}\text{N})^{13}\text{Be}$	-39600	90	-39310	10	3.2	B					Dbn 98Be28
$^{13}\text{C}(n,\gamma)^{14}\text{C}$	8177	2	8176.433	0.004	-0.3	U					67Th05
	8176.61	0.24			-0.7	U					Bdn 06Fi.A
$^{13}\text{C}(d,p)^{14}\text{C}$	5946	4	5951.867	0.004	1.5	U					CIT 51Li29 Y
	5952	10			0.0	U					Nob 54Ah47 Y
	5951	10			0.1	U					Mex 64Ma.B
	5951	8			0.1	U					MIT 64Sp12
	5951.85	0.54			0.0	U					Rez 90Pi05 *
$^{13}\text{C}(p,\gamma)^{14}\text{N}$	7551.0	0.8	7550.56277	0.00009	-0.5	U					56Ma87 Z
	7551.1	0.5			-1.1	U					63Bo07 *
$^{13}\text{C}(^3\text{He},d)^{14}\text{N}$	2048	14	2057.08833	0.00016	0.6	U					MIT 64Sp12
$^{14}\text{N}(^3\text{He},\alpha)^{13}\text{N}$	10015	10	10024.24	0.27	0.9	U					Ric 59Yo25
$^{14}\text{F}(p)^{13}\text{O}$	1560	40				3					10Go16
$^{14}\text{C}(\pi^-, \pi^+)^{14}\text{Be}$	-38100	170	-37960	130	0.8	R					84Gi09 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{14}\text{C}(^{14}\text{C}, ^{14}\text{O})^{14}\text{Be}^p$	-43440	60				2			Ber		95Bo10
$^{14}\text{C}(^7\text{Li}, ^7\text{Be})^{14}\text{B}$	-21499	30	-21506	21	-0.2	-			ChR		73Ba34
$^{14}\text{C}(^{14}\text{C}, ^{14}\text{N})^{14}\text{B}$	-20494	30	-20487	21	0.2	-			Ors		81Na.A
$^{14}\text{C}(^7\text{Li}, ^7\text{Be})^{14}\text{B}$	ave. -21506	21	-21506	21	0.0	1	100	100 $^{14}\text{B}$			average
$^{14}\text{C}(\beta^-)^{14}\text{N}$	155.2	0.5	156.476	0.004	2.6	U					54Ki23
	155.74	0.08			9.2	F					91Su09 *
	155.95	0.22			2.4	U					95Wi20
	156.27	0.14			1.5	U					00Ku25
$^{14}\text{C}(\text{p}, \text{n})^{14}\text{N}$	-626.15	0.3	-625.870	0.004	0.9	U			Wis		56Sa06
	-625.88	0.09			0.1	U			Zur		73Hi.A
$^{14}\text{N}(\text{p}, \text{n})^{14}\text{O}$	-5930.7	2.8	-5926.710	0.025	1.4	U			Ric		65Ku02
	-5927.6	1.5			0.6	U			Har		73C112
	-5925.6	0.4			-2.8	F			Auc		77Wh01 *
	-5925.41	0.08			-16.3	F			Auc		81Wh03 *
	-5925.41	0.11			-11.8	F			Auc		98Ba83 *
	-5926.68	0.17			-0.2	U			Auc		03To03
$^{14}\text{N}(^3\text{He}, \text{t})^{14}\text{O}$	-5161.3	0.8	-5162.956	0.025	-2.1	F			Mun		77Vo02 *
$^{13}\text{C H} - ^{14}\text{N}$	Original 8105.86288(10) $\mu\text{u}$ recalculated for molecular and ionization										
$^{14}\text{B}^i(\gamma, \text{d})^{12}\text{Be}$	$Q(\text{d})=Q(\text{p+n})+2224.5660(0.0004)$ $Q(\text{p+n})=290(20)$ keV										
$^{14}\text{C}(\text{p}, ^3\text{He})^{12}\text{B}^i$	IT=12710(20); $Q$ rebuilt with Ame1964										
*	energy and resolution arguments for T=1, not an IAS										
$^{14}\text{C}(\text{p}, \text{t})^{12}\text{C}^j$	IT=27595.0(2.4); $Q$ rebuilt										
$^{12}\text{C}(^3\text{He}, \text{n})^{14}\text{O}$	Originals T=1436.2(0.9, Bu) 1437.5(0.7, Ba) 1437.9(0.6, Ro) respectively, recalibrated										
$^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$	Estimated systematic error 0.5 added to statistical error 0.20										
$^{13}\text{C}(\text{p}, \gamma)^{14}\text{N}$	$E_p=1747.06(0.53)$ to $2^+$ level at 9172.25 keV										
$^{14}\text{C}(\pi^-, \pi^+)^{14}\text{Be}$	Original error 160 increased with 60 calibration uncertainty										
$^{14}\text{C}(\beta^-)^{14}\text{N}$	F : find 17 keV neutrino. See also reference										
$^{14}\text{N}(\text{p}, \text{n})^{14}\text{O}$	F : withdrawn by author										
$^{14}\text{N}(\text{p}, \text{n})^{14}\text{O}$	Authors recalibrated 77Wh01 for atomic effects										
$^{14}\text{N}(\text{p}, \text{n})^{14}\text{O}$	F : withdrawn by author										
$^{14}\text{N}(\text{p}, \text{n})^{14}\text{O}$	Original T=6353.08(0.07) recalibrated to T=6352.99(0.12) by author										
$^{14}\text{N}(\text{p}, \text{n})^{14}\text{O}$	F : withdrawn by author										
$^{14}\text{N}(^3\text{He}, \text{t})^{14}\text{O}$	F : rejected in reference of same group										
$\text{C D H} - ^{15}\text{N}$	21817.9119	0.0008	21817.9114	0.0006	-0.6	1	61	61 $^{15}\text{N}$	MI1	1.0	95Di08
$\text{C H}_3 - ^{15}\text{N}$	23366.1979	0.0017	23366.1978	0.0006	-0.1	1	14	13 $^{15}\text{N}$	MI1	1.0	95Di08
$^{15}\text{F} - \text{u}$	17477	86	17785	15	1.4	F				2.5	01Ze.A *
$^{14}\text{N D} - ^{15}\text{N H}$	9242.29	0.11	9241.8514	0.0007	-1.6	U			C5	2.5	71Ke01
	9241.780	0.008			5.9	B			B08	1.5	75Sm02
$^{15}\text{N}(\text{p}, \alpha)^{12}\text{C}$	4966	6	4965.4937	0.0006	-0.1	U			CIT		51Li26 Y
	4962	4			0.9	U			Bir		53Co02 Y
	4954	8			1.4	U			Mex		64Ma.B
	4965	7			0.1	U			MIT		64Sp12
$^{12}\text{C}(\alpha, \text{n})^{15}\text{O}$	-8503	12	-8502.0	0.5	0.1	U			Tal		63Ne05
$^{15}\text{N}(\text{d}, \alpha)^{13}\text{C}$	7675	9	7687.2360	0.0007	1.4	U			Mex		64Ma.B
	7689	6			-0.3	U			MIT		64Sp12
$^{15}\text{Ne}(2\text{p})^{13}\text{O}$	2522	66				3					14Wa09
$^{15}\text{Be}(\gamma, \text{n})^{14}\text{Be}$	1800	100				3					13Sn02
$^{14}\text{C}(\text{d}, \text{p})^{15}\text{C}$	-1006.5	0.8				2			Wis		56Do41 Y
$^{14}\text{C}(\text{p}, \gamma)^{15}\text{N}^i$	-1407.8	3.5				2					59Fe99 *
$^{14}\text{N}(\text{n}, \gamma)^{15}\text{N}$	10833.2	0.6	10833.2951	0.0008	0.2	U					68Gr14
	10833.1	0.7			0.3	U					74Sp04
	10833.5	0.05			-4.1	B			MMn		80Is02
	10833.314	0.012			-1.6	U					97Ju02
	10833.2339	0.0500			1.2	U			PTc		97Ro26 *
	10833.32	0.22			-0.1	U			Bdn		06Fi.A
	10833.3	0.5			0.0	U					06Be33



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{14}\text{N}(\text{d,p})^{15}\text{N}$	8629	11	8608.7292	0.0006	-1.8	U			CIT		52Mi54 Y	
	8614	6			-0.9	U		Mex	64Ma.B			
	8623	3			-4.8	B		MIT	64Sp12			
	8608.83	0.50			-0.2	U		Rez	90Pi05 *			
$^{14}\text{N}(\text{p},\gamma)^{15}\text{O}$	7297.1	0.9	7296.8	0.5	-0.4	1	30	30 $^{15}\text{O}$	CIT		72Ne05	
$^{14}\text{N}(\text{}^3\text{He,d})^{15}\text{O}$	1803	10	1803.3	0.5	0.0	U			Ric		59Yo25 Y	
	1802	15			0.1	U		Man	60Fo01			
$^{15}\text{F}(\text{p})^{14}\text{O}$	1410	150	1270	14	-0.9	o					03Le26	
	1510	110			-2.2	U			03Pe23			
	1490	130			-1.7	U			04Go15 *			
	1560	130			-2.2	U			04Le12			
	1230	50			0.8	U			05Gu25			
	1270	14				3			16De15			
$^{15}\text{C}(\beta^-)^{15}\text{N}$	9810	30	9771.7	0.8	-1.3	U					59Al06	
$^{15}\text{O}(\beta^+)^{15}\text{N}$	2745	5	2754.2	0.5	1.8	U					57Ki22	
$^{15}\text{N}(\text{p,n})^{15}\text{O}$	-3541.7	0.9	-3536.5	0.5	5.8	B					58Jo28 Y	
	-3535.1	1.0			-1.4	-		CIT	72Je02 Z			
	-3537.6	0.8			1.4	-			72Sh08 Z			
	ave.	-3536.6			0.6	0.2	1	70	70 $^{15}\text{O}$	average		
* $^{15}\text{F}-\text{u}$	F : results distrusted (see also $^{18}\text{Na}$ and $^{19}\text{Mg}$ )										GAu	**
* $^{14}\text{C}(\text{p},\gamma)^{15}\text{N}^i$	From a parametrized fit										MMC122**	
* $^{14}\text{N}(\text{n},\gamma)^{15}\text{N}$	Original error 0.0005 increased for calibration										GAu	**
* $^{14}\text{N}(\text{d,p})^{15}\text{N}$	Estimated systematic error 0.5 added to statistical error 0.061 keV										AHW	**
* $^{15}\text{F}(\text{p})^{14}\text{O}$	Symmetrized from 1450(+160-100) keV										04Go15	**
$\text{C H}_2\text{D}-\text{O}$	34837.406	0.033	34837.22300	0.00028	-3.7	B				1.5	71Sm01	
	34837.202	0.020			0.7	U		B08	1.5	75Sm02		
$\text{C D}_2-\text{O}$	33289.129	0.033	33288.93663	0.00029	-3.9	B				1.5	71Sm01	
	33289.061	0.038			-2.2	U		B07	1.5	71Sm01		
$\text{C}_4-\text{O}_3$	33288.940	0.019	15256.1412	0.0005	-0.1	U				1.5	75Sm02	
	15256.131	0.018			0.6	o		WA1	1.0	92Va.A		
	15256.086	0.081			0.3	U		OH1	2.5	93Ma.A		
	15256.121	0.009			2.2	o		WA1	1.0	95Va38		
	15256.1425	0.0008			-1.6	o		WA1	1.0	01Va33		
	15256.1415	0.0005			-0.6	o		WA1	1.0	03Va.A		
	15256.14129	0.00054			-0.2	1	93	93 $^{16}\text{O}$	WA1	1.0	06Va22	
$\text{C H}_4-\text{O}$	36387.55	0.8	36385.5094	0.0004	-1.0	U				2.5	68Ma45	
	36386.01	0.24			-0.8	U		J2	2.5	69Na21		
	36385.644	0.036			-2.5	U		B07	1.5	71Sm01		
	36385.5062	0.0013			2.4	U		MI1	1.0	95Di08		
	36385.5073	0.0019			1.1	U		MI1	1.0	95Di08		
	36385.5060	0.0022			1.5	U		MI1	1.0	95Di08		
$^{16}\text{O}-\text{u}$	-5085.362	0.027	-5085.38040	0.00017	-0.3	U				2.5	93Ma.A	
	-5085.3798	0.0011			-0.2	U			2.5	16Ho.A		
$^{14}\text{C H}_2-\text{O}$	23977.413	0.014	23977.433	0.004	1.0	U			B08	1.5	75Sm02	
$\text{N D}-\text{O}$	22261.160	0.013	22261.16298	0.00024	0.2	U			B08	1.5	75Sm02	
$\text{N}_2-\text{C O}$	11233.57	0.20	11233.3893	0.0004	-0.4	U				2.5	69Na21	
	11233.543	0.025			-4.1	B		B07	1.5	71Sm01		
	11233.43	0.21			-0.1	U		J6	2.5	76Ka50		
	11259	27			-0.4	U		CR1	2.5	89Sh10		
	11233.3909	0.0022			-0.7	U		MI1	1.0	95Di08		
	11233.38932	0.00042		0.0	1	82	81 $^{14}\text{N}$	MI1	1.0	04Th17		
$^{16}\text{O}(\alpha,^8\text{He})^{12}\text{O}$	-66020	120	-65836	24	1.5	U			Brk		78Ke06	
$^{16}\text{O}(\text{p},\alpha)^{13}\text{N}$	-5211	10	-5218.43	0.27	-0.7	U			MIT		64Sp12	
$^{16}\text{O}(\text{}^3\text{He},^6\text{He})^{13}\text{O}$	-30516	14	-30513	10	0.2	2			Brk		70Me11 *	
	-30511	13			-0.2	2		MSU		71Tr03 *		

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{16}\text{Be}(\gamma,2n)^{14}\text{Be}$	1350	100				3					12Sp02
$^{14}\text{C}(^{14}\text{C},^{12}\text{N})^{16}\text{B}$	-48380	60	-48411	25	-0.5	o			Ber		95Bo10
	-48378	60			-0.5	1	17	17 $^{16}\text{B}$	Ber		00Ka21
$^{14}\text{C}(\text{t,p})^{16}\text{C}$	-3015	8	-3013	4	0.2	2			MSU		77Fo09
	-3013	4			-0.1	2			LAI		78Se04
$^{14}\text{C}(^3\text{He,p})^{16}\text{N}$	4983	4	4978.2	2.3	-1.2	R			BNL		66Ga08
$^{14}\text{C}(^3\text{He,p})^{16}\text{N}^i$	-4951	7				2					68He03
$^{14}\text{N}(\text{t,p})^{16}\text{N}$	4853	10	4840.3	2.3	-1.3	U			Ald		66He10
$^{14}\text{C}(^3\text{He,n})^{16}\text{O}^j$	-8100	8	-8104	4	-0.5	1	23	23 $^{16}\text{O}^j$			70Ad01
$^{16}\text{O}(\text{d},\alpha)^{14}\text{N}$	3110.	3.5	3110.38807	0.00024	0.1	U			Wis		52Cr30
	3119	5			-1.7	U			Ric		53Fa18
	3110	6			0.1	U			Mex		64Ma.B
	3113	6			-0.4	U			MIT		64Sp12
$^{14}\text{N}(^3\text{He,p})^{16}\text{O}^i$	2444	6	2447	4	0.5	1	54	54 $^{16}\text{O}^i$			64Br08
$^{14}\text{N}(\text{d},\gamma)^{16}\text{O}^j$	-1986.3	4.4	-1985	4	0.3	1	77	77 $^{16}\text{O}^j$			72Ne10
$^{14}\text{N}(^3\text{He,n})^{16}\text{F}$	-963	40	-957	8	0.2	U			LAI		65Za01
	-970	15			0.9	R			Har		68Ad03
$^{16}\text{Ne}(2\text{p})^{14}\text{O}$	1350	80	1401	20	0.6	U					08Mu13
$^{16}\text{B}(\gamma,\text{n})^{15}\text{B}$	85	15	83	15	-0.1	1	95	83 $^{16}\text{B}$			09Le02
$^{15}\text{N}(\text{d,p})^{16}\text{N}$	286	12	264.3	2.3	-1.8	U			CIT		55Pa50
	269	10			-0.5	U			Pit		57Wa01
	259	6			0.9	2			Mex		64Ma.B
	267	8			-0.3	2			MIT		64Sp12
	270	10			-0.6	U			Pen		66He10
$^{15}\text{N}(\text{p},\gamma)^{16}\text{O}^i$	-665.3	6.6	-669	4	-0.5	1	46	46 $^{16}\text{O}^i$			57Ha99
$^{16}\text{O}(^3\text{He},\alpha)^{15}\text{O}$	4920	10	4913.7	0.5	-0.6	U			Ald		59Hi68
	4907	7			1.0	U			Ric		59Yo25
$^{16}\text{N}(\beta^-)^{16}\text{O}$	10400	20	10420.9	2.3	1.0	U					59A106
$^{16}\text{O}(^3\text{He,t})^{16}\text{F}$	-15430	10	-15436	8	-0.6	2			KVI		80Ja.A
$^{16}\text{O}(\pi^+,\pi^-)^{16}\text{Ne}$	-27766	45	-27702	20	1.4	2					80Bu15
$^{16}\text{O}(^3\text{He},^6\text{He})^{13}\text{O}$	$M - A$ increased by 7 for more recent calibrator $M - A(^9\text{C}) = 28913(2)$										
$^{16}\text{O}(^3\text{He},^6\text{He})^{13}\text{O}$	Recalibrated using their $^{12}\text{C}(^3\text{He},^6\text{He})$ result										
$^{14}\text{C}(^3\text{He,p})^{16}\text{N}^i$	IT=9928(7), $Q$ rebuilt with Ame1965										
$^{14}\text{C}(^3\text{He,n})^{16}\text{O}^j$	IT=22717(8), $Q$ rebuilt with Ame1964										
$^{14}\text{N}(^3\text{He,p})^{16}\text{O}^i$	IT=12798(6), $Q$ rebuilt										
$^{17}\text{O}_2 - ^{28}\text{Si D}_3$	-20968.3557	0.0014	-20968.3560	0.0013	-0.2	1	84	82 $^{17}\text{O}$	FS1	1.0	10Mo29
$^{17}\text{B}-\text{u}$	45970	860	46930	220	0.7	U			GA1	1.5	87Gi05
	46830	180			0.4	2			TO2	1.5	88Wo09
	47127	250			-0.5	2			GA3	1.5	91Or01
$^{17}\text{Ne} - ^{22}\text{Ne}_{.773}$	24373.27	0.38	24373.3	0.4	0.0	1	100	100 $^{17}\text{Ne}$	MA8	1.0	08Ge07
$^{17}\text{Na}-\text{u}$	37760	430				2				2.5	S-u148
$^{17}\text{O}-^{16}\text{O H}$	-3607.8961	0.0016	-3607.8952	0.0007	0.6	1	19	18 $^{17}\text{O}$	FS1	1.0	10Mo29
$^{17}\text{O}(\text{n},\alpha)^{14}\text{C}$	1817.2	3.5	1817.745	0.004	0.2	U					01Wa50
$^{14}\text{C}(\alpha,\text{n})^{17}\text{O}$	-1819.07	2.0	-1817.745	0.004	0.7	U			Wis		56Sa06
$^{17}\text{O}(\text{p},\alpha)^{14}\text{N}$	1200	17	1191.8748	0.0007	-0.5	U			MIT		64Sp12
$^{17}\text{O}(\text{d},\alpha)^{15}\text{N}$	9818	12	9800.6040	0.0009	-1.4	U			Nob		54Pa39
$^{16}\text{O}(\text{n},\gamma)^{17}\text{O}$	4143.24	0.23	4143.0793	0.0008	-0.7	U					77Mc05
	4143.06	0.13			0.1	U			Bdn		06Fi.A
$^{16}\text{O}(\text{d,p})^{17}\text{O}$	1915	8	1918.5133	0.0006	0.4	U			Ric		51K155
	1918	4			0.1	U			MIT		57Br82
	1918	3			0.2	U			Mex		61Ja23
	1920	3			-0.5	U			MIT		64Sp12
	1918.74	0.5			-0.5	U			Rez		90Pi05
$^{16}\text{O}(\text{n},\gamma)^{17}\text{O}^i$	-6935.70	0.17				2					81Hi01
$^{16}\text{O}(\text{p},\gamma)^{17}\text{F}$	600.35	0.28	600.27	0.25	-0.3	-			CIT		75Ro05

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{16}\text{O}(\text{d},\text{n})^{17}\text{F}$	-1626	4	-1624.30	0.25	0.4	U			Ric		51Bo49 Y
	-1624.6	0.5			0.6	-			Nvl		60Bo21 Z
$^{16}\text{O}(\text{p},\gamma)^{17}\text{F}$	ave. 600.27	0.25	600.27	0.25	0.0	1	100	100 $^{17}\text{F}$			average
$^{16}\text{O}(\text{p},\gamma)^{17}\text{F}^i$	-10592.8	1.9				2					76Hi09
$^{16}\text{O}(^3\text{He},2\text{n})^{17}\text{Ne}$	-22420	190	-22448.9	0.4	-0.2	U			BNL		67Es02
$^{17}\text{F}(\beta^+)^{17}\text{O}$	2770	6	2760.47	0.25	-1.6	U					54Wo23
* $^{16}\text{O}(\text{d},\text{p})^{17}\text{O}$	Estimated systematic error 0.5 added to statistical error 0.062 keV										AHW **
* $^{16}\text{O}(\text{n},\gamma)^{17}\text{O}^i$	Original $Q = -6934.41(0.17)$ does not match original $T = 7373.31(0.18)$										MMC129**
$\text{C D}_3 - ^{18}\text{O}$	43145.72216	0.00088	43145.7215	0.0007	-0.7	-			FS1	1.0	09Re15 *
	43145.72116	0.00136			0.3	-			FS1	1.0	09Re15 *
	ave. 43145.7219	0.0007			-0.5	1	87	84 $^{18}\text{O}$			average
$\text{C}_3 - ^{18}\text{O}_2$	1680.7695	0.0038	1680.7743	0.0015	1.3	1	16	16 $^{18}\text{O}$	FS1	1.0	09Re15
$^{18}\text{F}-\text{u}$	943	85	937.3	0.5	0.0	U				2.5	92Ge08
$^{18}\text{Na}-\text{u}$	25969	54	26880	100	6.7	F				2.5	01Ze.A *
	26882	183			0.0	1	30	30 $^{18}\text{Na}$	1.0	1.0	04Ze05 *
$^{18}\text{Ne}-^{22}\text{Ne}_{818}$	12755.68	0.39	12755.7	0.4	0.0	1	100	100 $^{18}\text{Ne}$	MA8	1.0	04Bi20
$^{14}\text{C}(^7\text{Li},^3\text{He})^{18}\text{N}$	-10170	60	-10117	19	0.9	U			Str		80Kr.A
$^{18}\text{O}(^{48}\text{Ca},^{51}\text{V})^{15}\text{B}$	-21760	50	-21762	21	0.0	-			Hei		78Bh02
	-21768	25			0.2	-			Can		83Ho08
	ave. -21766	22			0.2	1	88	88 $^{15}\text{B}$			average
$^{18}\text{O}(\text{p},\alpha)^{15}\text{N}$	3954	9	3979.8007	0.0009	2.9	U			Nob		54Mi60 Y
	3964	10			1.6	U			Mex		64Ma.B
$^{18}\text{O}(\text{d},\alpha)^{16}\text{N}$	4235	7	4244.1	2.3	1.3	R			CIT		55Pa50 Z
	4219	20			1.3	U			Mex		64Ma.B
	4249	15			-0.3	U			Phi		66He10
	4244	4			0.0	R			MIT		67Sp09 Z
$^{16}\text{O}(^3\text{He},\text{p})^{18}\text{F}$	2033	5	2032.1	0.5	-0.2	U			Ric		59Yo25
	2055	5			-4.6	C			Mex		64Ma.B
$^{16}\text{O}(^3\text{He},\text{n})^{18}\text{Ne}$	-3205	13	-3194.7	0.4	0.8	U			Nvl		61Du02 Y
	-3198	6			0.5	U			Ald		61To03 Y
	-3194.0	1.5			-0.5	U					94Ma14
$^{18}\text{B}(\gamma,\text{n})^{17}\text{B}$	5	5				3					10Sp02 *
$^{18}\text{O}(^{48}\text{Ca},^{49}\text{Ti})^{17}\text{C}$	-17465	35	-17476	17	-0.3	2			Hei		77No08
	-17479	20			0.2	2			Can		82Fi10
$^{18}\text{O}(^{207}\text{Pb},^{208}\text{Po})^{17}\text{C}$	-26870	220	-26797	17	0.3	U			ChR		79Ba31
$^{18}\text{O}(\text{t},\alpha)^{17}\text{N}$	3872	15				2			LAL		60Ja13
$^{17}\text{O}(\text{n},\gamma)^{18}\text{O}$	8043.5	1.0	8045.3693	0.0010	1.9	U			Bdn		06Fi.A
$^{17}\text{O}(\text{d},\text{p})^{18}\text{O}$	5820	10	5820.8033	0.0009	0.1	U			Nob		54Ah37 Y
	5820	10			0.1	U			Man		65Mo16
$^{17}\text{O}(\text{p},\gamma)^{18}\text{F}$	5603	3	5607.1	0.5	1.4	U			Str		73Se03
	5606.2	0.6			1.5	1	60	60 $^{18}\text{F}$	CIT		75Ro05 Z
$^{18}\text{Na}(\text{p})^{17}\text{Ne}$	1270	170	1250	90	-0.1	-					04Ze05
	1230	150			0.1	-					12Mu05
	ave. 1250	110			0.0	1	70	70 $^{18}\text{Na}$			average
$^{18}\text{O}(\pi^-, \pi^+)^{18}\text{C}$	-26712	150	-26720	30	-0.1	U					78Se07
$^{18}\text{O}(^{48}\text{Ca},^{48}\text{Ti})^{18}\text{C}$	-21434	30				2			Can		82Fi10
	-21331	300	-21430	30	-0.3	U			Ors		82Na04
$^{18}\text{N}(\beta^-)^{18}\text{O}$	13860	400	13896	19	0.1	U					64Ch19
$^{18}\text{O}(\text{d},2\text{p})^{18}\text{N}$	-15270	100	-15338	19	-0.7	U			Brk		78De.A
$^{18}\text{O}(\text{t},^3\text{He})^{18}\text{N}$	-13917	60	-13877	19	0.7	U			LAL		69St07 *
$^{18}\text{O}(^7\text{Li},^7\text{Be})^{18}\text{N}$	-14761	20	-14758	19	0.2	2			Can		83Pu01
$^{18}\text{O}(^{14}\text{C},^{14}\text{N})^{18}\text{N}$	-13720	50	-13740	19	-0.4	2			Ors		80Na14
$^{18}\text{F}(\beta^+)^{18}\text{O}$	1657	2	1655.9	0.5	-0.5	U					64Ho28
$^{18}\text{O}(\text{p},\text{n})^{18}\text{F}$	-2451	4	-2438.3	0.5	3.2	B			Wis		50Ri59 Y
	-2436.97	0.73			-1.8	1	40	40 $^{18}\text{F}$	Nvl		64Bo13 Z
	-2440.2	2.8			0.7	U					67Pr04

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{18}\text{Ne}(\beta^+)^{18}\text{F}$	4438	9	4444.5	0.6	0.7	U					63Fr10	
*C D <sub>3</sub> - $^{18}\text{O}$	Respectively CD <sub>3</sub> <sup>+</sup> / $^{18}\text{O}^+$ , C <sub>2</sub> D <sub>6</sub> <sup>+</sup> / $^{18}\text{O}^{2+}$ considered independent										GAu	**
* $^{18}\text{Na}-u$	F : results distrusted (see also $^{15}\text{F}$ and $^{19}\text{Mg}$ )										GAu	**
* $^{18}\text{Na}-u$	Other interpretation : $D_M=25969(172) \mu\text{u}$ , $M-A=24190(160) \text{keV}$										04Ze05	**
* $^{18}\text{B}(\gamma, n)^{17}\text{B}$	Decay energy <10 keV, derived from scattering length $a_s < -50 \text{fm}$										10Sp02	**
* $^{18}\text{O}(t, ^3\text{He})^{18}\text{N}$	From $Q=14038(30)$ , reinterpreted as mainly to ( $2^-$ ) level at 114.90 keV										Ens967	**
$^{28}\text{Si H}_3-\text{C } ^{19}\text{F}$	1998.4687	0.0022	1998.4688	0.0010	0.1	1	22	15 $^{19}\text{F}$	FS1	1.0	09Re15	
$^{19}\text{B}-u$	64166	376				2			GA8	1.5	12Ga45	
$^{19}\text{C}-u$	34680	260	34800	110	0.3	o			TO1	1.5	86Vi09	
	35370	450			-0.8	U			GA1	1.5	87Gi05	
	35180	130			-2.0	o			TO2	1.5	88Wo09	
	35506	253			-1.9	U			GA3	1.5	91Or01	
C D <sub>4</sub> -H $^{19}\text{F}$	50178.88	0.05	50178.9173	0.0009	0.5	U			B08	1.5	75Sm02	
$^{19}\text{Mg}-u$	35470	270	34170	50	-1.9	F				2.5	01Ze.A *	
$^{13}\text{C D}_3-^{19}\text{F}$	47257.00669	0.00091	47257.0067	0.0008	0.0	1	87	85 $^{19}\text{F}$	FS1	1.0	09Re15	
$^{19}\text{Ne}-^{22}\text{Ne}_{.864}$	9323.92	0.33	9324.17	0.17	0.8	2			MA8	1.0	04B120	
	9324.26	0.20			-0.5	2			MA8	1.0	08Ge07	
$^{19}\text{F}(p, \alpha)^{16}\text{O}$	8115	10	8113.6122	0.0009	-0.1	U			CIT		50Ch53 Y	
	8115	10			-0.1	U			CIT		57Yo04 Y	
	8122	9			-0.9	U			MIT		64Sp12	
$^{17}\text{O}(t, p)^{19}\text{O}$	3524	7	3519.2	2.6	-0.7	R			Man		65Mo19	
$^{19}\text{F}(d, \alpha)^{17}\text{O}$	10060	12	10032.1254	0.0010	-2.3	U			MIT		64Sp12	
$^{19}\text{Mg}(2p)^{17}\text{Ne}$	750	50				2					07Mu15	
$^{18}\text{C}(n, \gamma)^{19}\text{C}$	530	120	580	90	0.4	3					99Na27 *	
	650	150			-0.5	3					01Ma08 *	
$^{18}\text{O}(^{18}\text{O}, ^{17}\text{F})^{19}\text{N}$	-19374	50	-19374	16	0.0	2			Ors		81Na.A	
	-19334	35			-1.1	2			Can		89Ca25	
$^{18}\text{O}(^{48}\text{Ca}, ^{47}\text{Sc})^{19}\text{N}$	-16540	20	-16527	17	0.6	2			Can		83Ho08	
$^{18}\text{O}(^{208}\text{Pb}, ^{207}\text{Bi})^{19}\text{N}$	-18440	150	-18333	17	0.7	U			ChR		79Ba31	
$^{18}\text{O}(d, p)^{19}\text{O}$	1727	8	1731.1	2.6	0.5	o			Nob		54Mi89 Y	
	1732	8			-0.1	2			CIT		54Th30	
	1731	5			0.0	2			Nob		57Ah19 Y	
	1733	6			-0.3	2			Mex		64Ma.B	
	1727	5			0.8	2			MIT		64Sp12 Z	
	1734	10			-0.3	U			Man		65Mo16	
$^{19}\text{F}(^3\text{He}, \alpha)^{18}\text{F}$	10166	15	10145.7	0.5	-1.4	U			Ald		59Hi67 Y	
$^{19}\text{Na}(p)^{18}\text{Ne}$	160	110	323	11	1.5	U					04Ze05	
	328	22			-0.2	1	23	23 $^{19}\text{Na}$			10Mu12	
$^{19}\text{O}(\beta^-)^{19}\text{F}$	4800	12	4820.3	2.6	1.7	U					59Ai06	
$^{19}\text{Ne}(\beta^+)^{19}\text{F}$	3262	10	3239.49	0.16	-2.3	U					60Wa04	
$^{19}\text{F}(p, n)^{19}\text{Ne}$	-4021.3	4.7	-4021.84	0.16	-0.1	U			Ric		55Ma84	
	-4019.6	1.4			-1.6	U			Ric		61Be13 Z	
	-4021.1	1.0			-0.7	U			Zur		61Ry04 Z	
	-4020.7	0.8			-1.4	U					66Ma60	
	-4019.6	0.7			-3.2	B					69Ov01 Z	
$^{19}\text{F}(^3\text{He}, t)^{19}\text{Ne}^i$	-10759	9				2					98Ut02 *	
* $^{19}\text{Mg}-u$	F : results distrusted (see also $^{15}\text{F}$ and $^{18}\text{Na}$ )										GAu	**
* $^{18}\text{C}(n, \gamma)^{19}\text{C}$	From Coulomb dissociation cross sections and angular distribution										99Na27	**
* $^{18}\text{C}(n, \gamma)^{19}\text{C}$	From momentum distribution following one neutron removal										01Ma08	**
* $^{19}\text{F}(^3\text{He}, t)^{19}\text{Ne}^i$	rebuilt from $E_x=7500(9)$ ; with references from Ame1995; recalibrated +1keV										MMC141**	
$^{20}\text{C}-u$	39940	1210	40260	250	0.2	U			GA1	1.5	87Gi05	
	40360	240			-0.3	2			TO2	1.5	88Wo09	
	40165	491			0.1	2			GA3	1.5	91Or01	
	40420	550			-0.2	2			GA5	1.5	99Sa.A	
	40108	290			0.4	2			GA8	1.5	12Ga45	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{20}\text{N}-u$	23230	280	23370	80	0.3	U			TO1	1.5	86Vi09
	23210	150			0.7	2			GA1	1.5	87Gi05
	23380	130			-0.1	2			TO2	1.5	88Wo09
	23397	69			-0.3	2			GA3	1.5	91Or01
$\text{C D}_4-^{20}\text{Ne}$	63966.9329	0.0026	63966.9363	0.0017	1.3	1	41	40 $^{20}\text{Ne}$	MI1	1.0	95Di08
$^{20}\text{Ne}-u$	-7559.309	0.090	-7559.8238	0.0017	-2.3	U			OH1	2.5	93Ma.A
	-7559.814	0.014			-0.7	U			ST2	1.0	02Bf02
$\text{O D}_2-^{20}\text{Ne}$	30677.497	0.067	30677.9996	0.0017	3.0	B			OH1	2.5	93Go38
$^{20}\text{Mg}-^{23}\text{Na}_{.870}$	27663.8	2.0				2			TT1	1.0	14Ga20
$^{20}\text{Ne}-^{22}\text{Ne}_{.909}$	270.89	0.43	271.111	0.017	0.5	U			MA8	1.0	04B120
	271.16	0.20			-0.2	U			MA8	1.0	08Ge07
$^{20}\text{Ne}(^3\text{He}, ^8\text{Li})^{15}\text{F}$	-29960	200	-29623	14	1.7	U			MSU		78Be26
	-29730	180			0.6	U			Brk		78Ke06
$^{20}\text{Ne}^i(\alpha)^{16}\text{O}$	5548.8	6.3	5542.6	2.0	-1.0	U					73To08
$^{20}\text{Ne}(\alpha, ^8\text{He})^{16}\text{Ne}$	-60150	80	-60213	20	-0.8	U			Brk		78Ke06
	-60197	23			-0.7	R			Tex		83Wo01
$^{20}\text{Ne}(^3\text{He}, ^6\text{He})^{17}\text{Ne}$	-26188	50	-26203.3	0.4	-0.3	U			Brk		70Me11 *
	-26158	32			-1.4	F					98Gu10 *
$^{18}\text{O}(^{48}\text{Ca}, ^{46}\text{Sc})^{20}\text{N}$	-25873	60	-25010	80	14.3	B			Can		89Or03 *
$^{18}\text{O}(t,p)^{20}\text{O}$	3086	15	3081.9	0.9	-0.3	U			LAl		60Ja13
	3076	10			0.6	U			Ald		62Hi06
	3082.4	1.9			-0.3	2			Str		82An12
	3081.7	1.0			0.2	2			Str		85An17
$^{18}\text{O}(^3\text{He}, p)^{20}\text{F}$	6875.2	1.5	6876.895	0.030	1.1	U			NDm		70Ro06
$^{18}\text{O}(^3\text{He}, p)^{20}\text{Fi}$	356.0	3.0				2					76Mi01 *
$^{20}\text{Ne}(d, \alpha)^{18}\text{F}$	2795	9	2795.8	0.5	0.1	U			Nob		54Mi61 Y
	2766	20			1.5	U			Mex		64Ma.B
	2790	10			0.6	U					75Bo59
$^{19}\text{F}(n, \gamma)^{20}\text{F}$	6601.1	0.3	6601.336	0.030	0.8	U			Utr		68Sp01
	6601.29	0.14			0.3	2			ILn		83Hu12 Z
	6601.32	0.05			0.3	2			MMn		87Ke09 Z
	6601.35	0.04			-0.3	2			ORn		96Ra04
	6601.34	0.13			0.0	2			Bdn		06Fi.A
$^{19}\text{F}(d, p)^{20}\text{F}$	4377	7	4376.770	0.030	0.0	U			MIT		64Sp12
	4377.7	0.9			-1.0	U			NDm		70Ro06
$^{19}\text{F}(p, \gamma)^{20}\text{Ne}^j$	-3889.4	2.7				2					67B119
$^{20}\text{Ne}(^3\text{He}, \alpha)^{19}\text{Ne}$	3750	13	3712.32	0.16	-2.9	U			MIT		64Sp12
$^{20}\text{Na}^i(p)^{19}\text{Ne}$	4381	50	4308.0	1.2	-1.5	U			Brk		79Mo02
	4332	16			-1.5	U			MSU		92Go10
	4326	30			-0.6	U			Lis		95Pi03
$^{20}\text{F}(\beta^-)^{20}\text{Ne}$	7053	13	7024.467	0.030	-2.2	U					54Wo23 *
	7050	15			-1.7	U					59Al06 *
	7032	6			-1.3	U					76Ge08 *
	7026.9	1.8			-1.4	U					87Va20 *
	7019.8	1.7			2.7	U					89He11 *
$^{20}\text{Ne}^i(\text{IT})^{20}\text{Ne}$	10274	3	10272.5	2.0	-0.5	2					76In06
	10271.2	2.7			0.5	2					77Fi08
$^{20}\text{Na}(\beta^+)^{20}\text{Ne}$	13906	40	13892.5	1.1	-0.3	U					67Su05
$^{20}\text{Ne}(p, n)^{20}\text{Na}$	-14672.1	7.	-14674.9	1.1	-0.4	U					71Wi07 Z
$^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na}-^{36}\text{Ar}(\text{)}^{36}\text{K}$	-1078.06	1.06	-1078.1	1.1	0.0	1	100	100 $^{20}\text{Na}$	Mun		10Wr01
$^{20}\text{Na}^i(\text{IT})^{20}\text{Na}$	6498.4	0.5				2					15Gi03
$^{20}\text{Ne}(^3\text{He}, ^6\text{He})^{17}\text{Ne}$	Original $M - A = 16479(50)$ but revised calibrator $M(^9\text{C}) = 28910.2(2.1)$										
$^{20}\text{Ne}(^3\text{He}, ^6\text{He})^{17}\text{Ne}$	F : calibrated with $^{24}\text{Mg}(^3\text{He}, ^6\text{He})$ from reference for excitation energies										
*	no details given. No correction possible										
*	Probably to excited levels in $^{20}\text{N}$ and $^{46}\text{Sc}$										
$^{18}\text{O}(^{48}\text{Ca}, ^{46}\text{Sc})^{20}\text{N}$	IT=6519.4(3.0), Q rebuilt with Ame1971										
$^{18}\text{O}(^3\text{He}, p)^{20}\text{Fi}$	E $_{\beta^-}$ = 5419(13) 5416(15) 5398(6) 5392.3(1.8) 5386.1(1.7) respectively,										
$^{20}\text{F}(\beta^-)^{20}\text{Ne}$	to $2^+$ level at 1633.674 keV										
*	Ens992 **										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{21}\text{N}-u$	26580	200	27090	140	1.7	U			TO1	1.5	86Vi09	
	27060	190			0.1	2			GA1	1.5	87Gi05	
	26930	210			0.5	2			TO2	1.5	88Wo09	
	27162	131			-0.4	2			GA3	1.5	91Or01	
$^{21}\text{Na}-^{39}\text{K}_{.538}$	17180.51	0.29	17180.61	0.11	0.4	o			MA8	1.0	04Mu26	
	17180.51	0.29			0.2	U			Ma8	1.5	08Mu05 *	
$\text{H}_3 \text{ }^{18}\text{O}-^{21}\text{Ne}$	28787.76	0.25	28788.02	0.04	1.1	U			CP1	1.0	04Sa53 *	
$^{21}\text{Na}-^{23}\text{Na}_{.913}$	6995.25	0.32	6995.35	0.11	0.3	o			MA8	1.0	04Mu26	
	6995.25	0.32			0.2	U			Ma8	1.5	08Mu05	
$^{21}\text{Mg}-^{23}\text{Na}_{.913}$	21046.41	0.81				2			TT1	1.0	14Ga20	
$^{21}\text{Ne}-^{22}\text{Ne}_{.955}$	2073.82	0.40	2073.91	0.05	0.2	U			MA8	1.0	04B120	
	2074.04	0.26			-0.5	U			MA8	1.0	08Ge07	
$^{21}\text{Na}-^{21}\text{Ne}$	3808.017	0.097	3808.02	0.10	0.0	1	100	100 $^{21}\text{Na}$	MS1	1.0	15Ei01	
$^{21}\text{Na}-^{20}\text{Na}$	-9732	50	-9699.7	1.2	0.3	U			CR1	2.5	89Sh10	
$^{18}\text{O}(^{18}\text{O}, ^{15}\text{O})^{21}\text{O}$	-12574	70	-12483	12	1.3	U			Ors		78Na02	
	-12499	20			0.8	2			Can		89Ca25	
$^{18}\text{O}(^{64}\text{Ni}, ^{61}\text{Ni})^{21}\text{O}$	-11713	15	-11722	12	-0.6	2			Dar		85Wo01	
$^{18}\text{O}(^{208}\text{Pb}, ^{205}\text{Pb})^{21}\text{O}$	-6860	75	-6823	12	0.5	U			ChR		79Ba31	
$^{19}\text{F}(t,p)^{21}\text{F}$	6221.0	1.8				2			Str		84An17	
$^{19}\text{F}(^3\text{He}, p)^{21}\text{Ne}$	11911	15	11886.58	0.04	-1.6	U			Ald		59Hi75 Y	
$^{20}\text{Ne}(n, \gamma)^{21}\text{Ne}$	6760.8	1.5	6761.16	0.04	0.2	U					70Se14	
	6761.16	0.04			0.1	-			MMn		86Pr05 Z	
	6761.19	0.14			-0.2	-			Bdn		06Fi.A	
	4531	9	4536.60	0.04	0.6	U			Nob		55Ah41 Y	
$^{20}\text{Ne}(d,p)^{21}\text{Ne}$	4532	6			0.8	U			Mex		64Ma.B	
	4534	7			0.4	U			MIT		64Sp12	
	ave.	6761.16	0.04	6761.16	0.04	0.0	1	100	100 $^{21}\text{Ne}$		average	
$^{20}\text{Ne}(n, \gamma)^{21}\text{Ne}$	2431.2	0.7	2431.67	0.10	0.7	U					69B103 Z	
$^{20}\text{Ne}(p, \gamma)^{21}\text{Na}^i$	-6547.9	14.3	-6543	4	0.3	U					81Fe05	
$^{21}\text{Na}^i(p)^{20}\text{Ne}$	6543	4				2					73Se08 *	
$^{21}\text{O}(\beta^-)^{21}\text{F}$	8150	175	8110	12	-0.2	U					81Al07	
$^{21}\text{Na}(\beta^+)^{21}\text{Ne}$	3522	30	3547.14	0.09	0.8	U					52Sc15	
	3532	20			0.8	U					60Wa04	
* $^{21}\text{Na}-^{39}\text{K}_{.538}$	CF=1.5 for prelim. results; not trusted within given uncertainties										GAu	**
* $\text{H}_3 \text{ }^{18}\text{O}-^{21}\text{Ne}$	$D_M=28787.78(0.25)$ corrected -0.02 keV for molecular and ionization											04Sa53 **
* $^{21}\text{Na}^i(p)^{20}\text{Ne}$	$Q_p=6548(4), 4904(4)$ to ground state and $2^+$ level at 1633.674 keV											Ens992 **
$^{22}\text{C}-u$	57585	408	57550	250	-0.1	U			GA8	1.5	12Ga45	
	32990	790	34100	220	0.9	U			GA1	1.5	87Gi05	
	34340	250			-0.6	2			TO2	1.5	88Wo09	
	34683	389			-1.0	2			GA3	1.5	91Or01	
	34240	320			-0.3	2			GA5	1.5	99Sa.A	
$^{22}\text{O}-u$	33398	279			1.7	2			GA8	1.5	12Ga45	
	9842	81	9970	60	1.0	R			GA3	1.5	91Or01	
$^{22}\text{Ne}-u$	-8614.885	0.019	-8614.890	0.019	-0.3	1	99	99 $^{22}\text{Ne}$	ST2	1.0	02Bf02	
$^{22}\text{Na}-^{39}\text{K}_{.564}$	14907.33	0.30	14906.96	0.18	-1.2	o			MA8	1.0	04Mu26	
	14907.33	0.30			-0.8	1	17	17 $^{22}\text{Na}$	Ma8	1.5	08Mu05	
$^{22}\text{Mg}-^{39}\text{K}_{.564}$	20040.33	0.35	20040.2	0.3	-0.4	o			MA8	1.0	04Mu26	
	20040.33	0.35			-0.3	1	41	41 $^{22}\text{Mg}$	Ma8	1.5	08Mu05	
$\text{O H}-^{22}\text{Ne}_{.773}$	9398.87	0.19	9398.962	0.015	0.5	U			MA8	1.0	08Ge07	
$^{22}\text{Na}-^{24}\text{Mg}_{.917}$	8153.64	0.31	8154.18	0.18	1.7	o			MA8	1.0	04Mu26	
	8153.64	0.31			1.2	1	16	16 $^{22}\text{Na}$	Ma8	1.5	08Mu05	
$^{22}\text{Na}-^{23}\text{Na}_{.957}$	4228.11	0.29	4228.22	0.18	0.4	o			MA8	1.0	04Mu26	
	4228.11	0.29			0.2	1	18	18 $^{22}\text{Na}$	Ma8	1.5	08Mu05	
$^{22}\text{Na}-^{22}\text{Ne}$	3052.75	0.33	3052.31	0.18	-1.3	1	31	31 $^{22}\text{Na}$	CP1	1.0	04Sa53 *	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{22}\text{Mg}-^{22}\text{Ne}$	8185.79	0.73	8185.5	0.3	-0.3	1	21	21 $^{22}\text{Mg}$	CP1	1.0	04Sa53 *
$^{22}\text{Mg}-^{22}\text{Na}$	5132.99	0.34	5133.2	0.3	0.7	o			MA8	1.0	04Mu26
	5132.99	0.34			0.5	1	46	38 $^{22}\text{Mg}$	Ma8	1.5	08Mu05
$^{22}\text{Ne}-^{20}\text{Ne}$	-1056.415	0.290	-1055.066	0.019	1.9	U			OH1	2.5	93Go38
$^{18}\text{O}(^{18}\text{O},^{14}\text{O})^{22}\text{O}$	-19060	100	-18860	60	2.0	2			Can		76Hi10
$^{18}\text{O}(^{208}\text{Pb},^{204}\text{Pb})^{22}\text{O}$	-6710	180	-6700	60	0.0	2			ChR		79Ba31
$^{22}\text{Mg}^i(\alpha)^{18}\text{Ne}$	5885	40	5902	6	0.4	U			Bor		97B103 *
	5904	8			-0.3	1	60	60 $^{22}\text{Mg}^i$	Bor		06Ac04 *
$^{19}\text{F}(\alpha,p)^{22}\text{Ne}$	1674	11	1673.219	0.018	-0.1	U			MIT		64Sp12
$^{19}\text{F}(\alpha,n)^{22}\text{Na}$	-1958	10	-1952.33	0.17	0.6	U			Duk		60Wi07 Y
$^{22}\text{C}(\gamma,2n)^{20}\text{C}$	-200	120	-35	20	1.4	o					11Ya25 *
	-110	60			1.2	U					12Fo04 *
	-35	20				3					13Mo12 *
$^{20}\text{Ne}(^3\text{He},n)^{22}\text{Mg}$	197	25	217.9	0.3	0.8	U			Har		68Ad03
	209	11			0.8	U			CIT		70Mc06
$^{22}\text{Mg}^i(2p)^{20}\text{Ne}$	6098	13	6108	6	0.8	1	23	23 $^{22}\text{Mg}^i$			06Ac04 *
$^{22}\text{Ne}(t,\alpha)^{21}\text{F}$	4545	10	4547.8	1.8	0.3	U			LAI		61Si03 Y
$^{21}\text{Ne}(n,\gamma)^{22}\text{Ne}$	10364.4	0.3	10364.26	0.04	-0.5	U			MMn		86Pr05 Z
	10363.9	0.5			0.7	U			Bdn		06Fi.A
$^{21}\text{Ne}(d,p)^{22}\text{Ne}$	8152	11	8139.69	0.04	-1.1	U			CIT		52Mi54 Y
$^{21}\text{Ne}(p,\gamma)^{22}\text{Na}$	6739.0	0.7	6738.71	0.18	-0.4	U					70An06 *
$^{22}\text{Mg}^i(p)^{21}\text{Na}$	8547	15	8540	6	-0.5	1	17	17 $^{22}\text{Mg}^i$	Brk		82Ca16 *
$^{22}\text{F}(\beta^-)^{22}\text{Ne}$	11000	150	10818	12	-1.2	U					73Gu05
	10950	120			-1.1	U			ANB		74Da02
$^{22}\text{Ne}(t,^3\text{He})^{22}\text{F}$	-10788	33	-10799	12	-0.3	2					69St07 *
	-10794	18			-0.3	2			Dar		88Cl04 *
$^{22}\text{Ne}(^7\text{Li},^7\text{Be})^{22}\text{F}$	-11691	20	-11680	12	0.6	2			Can		89Or04 *
$^{22}\text{Na}(\beta^+)^{22}\text{Ne}$	2842.2	0.5	2843.21	0.17	2.0	1	12	12 $^{22}\text{Na}$			68Be35 *
	2840.4	1.5			1.9	U					68We02 *
	2841.5	1.0			1.7	U					72Gi17 *
* $^{22}\text{Na}-^{22}\text{Ne}$	$D_M=3052.79(0.33) \mu\text{u}$ , $M-A=-5181.06(0.31) \text{keV}$ ; corrected $-0.04 \text{keV}$ for ion-ion interaction										
*	04Sa53 **										
* $^{22}\text{Mg}-^{22}\text{Ne}$	$D_M=8185.84(0.73) \mu\text{u}$ , $M-A=-399.65(0.68) \text{keV}$ ; corrected $-0.05 \text{keV}$ for ion-ion interaction										
*	04Sa53 **										
* $^{22}\text{Mg}^i(\alpha)^{18}\text{Ne}$	$E_\alpha=3270(40) \text{ to } 2^+$ level at 1887.3 keV										
*	04Sa53 **										
* $^{22}\text{Mg}^i(\alpha)^{18}\text{Ne}$	$Q_\alpha=4017(8) \text{ to } 2^+$ level at 1887.3 keV										
*	Ens967 **										
* $^{22}\text{C}(\gamma,2n)^{20}\text{C}$	From upper limit $S_{2n}<400$										
*	GAu **										
* $^{22}\text{C}(\gamma,2n)^{20}\text{C}$	From upper limit $S_{2n}<220$										
*	GAu **										
* $^{22}\text{C}(\gamma,2n)^{20}\text{C}$	The two items are estimates derived from the experimental result of reference										
*	From upper limit $S_{2n}<70$										
*	GAu **										
* $^{22}\text{Mg}^i(2p)^{20}\text{Ne}$	Original $Q_{2p}=4464(8) \text{ to } 1633.674(0.015) \text{ level}$										
*	06Ac04 **										
* $^{22}\text{Mg}^i(2p)^{20}\text{Ne}$	Estimated systematic 10 keV uncertainty added										
*	16Ma.A **										
* $^{21}\text{Ne}(p,\gamma)^{22}\text{Na}$	$T=701.8(0.5) \text{ to } 1^+$ level at 7408.6(0.5) keV										
*	Ens157 **										
* $^{21}\text{Ne}(p,\gamma)^{22}\text{Na}$	Reanalysis using E(exc) for lower levels of reference										
*	90En08 **										
* $^{22}\text{Mg}^i(p)^{21}\text{Na}$	$E_p=8149(21), 7839(15) \text{ to } 3/2^+$ ground state, $5/2^+$ level at 331.90 keV										
*	Ens04c **										
* $^{22}\text{Ne}(t,^3\text{He})^{22}\text{F}$	Original value $-10834(30)$ re-calculated from $Q$ to $(2^+)$ level at 709.1, $1^+$ at 1627.1 and $1^+$ at 2571.7 keV										
*	GAu **										
* $^{22}\text{Ne}(t,^3\text{He})^{22}\text{F}$	Original value $-10836(12)$ re-calculated										
*	GAu **										
* $^{22}\text{Ne}(^7\text{Li},^7\text{Be})^{22}\text{F}$	$Q=-12400(20) \text{ to } (2^+)$ level at 709.1 keV										
*	Ens157 **										
* $^{22}\text{Na}(\beta^+)^{22}\text{Ne}$	$E_{\beta^+}=545.7(0.5) 543.9(1.5) 545(1)$ respectively, to $2^+$ level at 1274.537 keV										
*	Ens157 **										
$^{23}\text{N}-\text{u}$	37110	2000	39420	450	0.8	o			GA5	1.5	99Sa.A
	39378	923			0.0	U			GA7	1.5	07Ju03
	39421	301				2			GA8	1.5	12Ga45
$^{23}\text{O}-\text{u}$	15700	320	15700	130	0.0	o			TO1	1.5	86Vi09
	15860	320			-0.3	o			GA1	1.5	87Gi05
	15700	150			0.0	2			TO2	1.5	88Wo09
	15621	186			0.3	o			GA3	1.5	91Or01
	15695	107			0.0	2			GA7	1.5	07Ju03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{23}\text{F}-u$		3530	210	3530	40	0.0	U		TO1	1.5	86Vi09
		3553	43			-0.4	-		GT1	1.5	04Ma.A
		3503	48			0.5	-		LZ1	1.0	15Xu14
	ave.	3520	40			0.2	1	86	86 $^{23}\text{F}$		average
$^{23}\text{Na}-u$		-10230.721	0.0037	-10230.7180	0.0019	0.8	-		MI2	1.0	99Br47
		-10230.716	0.0048			-0.4	-		MI2	1.0	99Br47
		-10230.7172	0.0026			-0.3	-		FS1	1.0	10Mo30
	ave.	-10230.7181	0.0019			0.0	1	100	100 $^{23}\text{Na}$		average
$^{23}\text{Ne}-^{22}\text{Ne}_{1.045}$		3469.59	0.37	3469.46	0.11	-0.3	U		MA8	1.0	04Bi20
$^{23}\text{Mg}-^{23}\text{Na}$		4354.80	0.83	4354.66	0.17	-0.2	U		JY1	1.0	09Sa38
		4354.66	0.17				2		TT1	1.0	14Sc09
$^{23}\text{Al}-^{23}\text{Na}$		17475.07	0.37				2		JY1	1.0	09Sa38
$^{23}\text{Na}(p,\alpha)^{20}\text{Ne}$		2377	3	2376.1331	0.0024	-0.3	U		Wis		53Do04 Y
		2373	8			0.4	U		MIT		64Sp12
$^{23}\text{Na}(d,\alpha)^{21}\text{Ne}$		6911	9	6912.73	0.04	0.2	U		Mex		64Ma.B
		6909	10			0.4	U		MIT		64Sp12
$^{22}\text{Ne}(^{18}\text{O},^{17}\text{F})^{23}\text{F}$		-14080	90	-14040	30	0.4	1	14	14 $^{23}\text{F}$		89Or04
$^{22}\text{Ne}(n,\gamma)^{23}\text{Ne}$		5200.2	2.0	5200.65	0.10	0.2	U				70Se14
		5200.65	0.12			0.0	2		MMn		86Pr05 Z
		5200.64	0.20			0.0	2		Bdn		06Fi.A
$^{22}\text{Ne}(d,p)^{23}\text{Ne}$		2967	8	2976.08	0.10	1.1	U		Nob		54Ah20 Y
		2971	9			0.6	o		MIT		60Fr04
		2974	6			0.3	U		Mex		64Ma.B
		2968	7			1.2	U		MIT		64Sp12
		8794.0	1.5	8794.105	0.018	0.1	U				71Pi08 Z
$^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$		8794.26	0.17			-0.9	U				89Ba42 Z
		-10796.3	2.0				2				85Ev01 *
$^{23}\text{Al}(p)^{22}\text{Mg}$		11644	57				2		Bor		97Bi04 *
$^{23}\text{F}(\beta^-)^{23}\text{Ne}$		8510	170	8440	30	-0.4	U				74Go17
$^{23}\text{Ne}(\beta^-)^{23}\text{Na}$		4383	8	4375.80	0.10	-0.9	U				63Ca06
$^{23}\text{Mg}(\beta^+)^{23}\text{Na}$		4121	12	4056.34	0.16	-5.4	B				63Fr10
$^{23}\text{Na}(p,n)^{23}\text{Mg}$		-4832	10	-4838.69	0.16	-0.7	U		Oak		55Ki28 Z
		-4836.5	6.			-0.4	U		Ric		58Bi41 Y
		-4848.0	7.			1.3	U		ChR		58Go77 Y
		-4835.8	2.5			-1.2	U		Har		62Fr09 Z
		-4843.2	5.1			0.9	U		Tkm		63Ok01 Z
$^{*22}\text{Ne}(p,\gamma)^{23}\text{Na}^j$	Original $E_{res}=-10793.6(2.0)$ is a typo										
$^{*23}\text{Al}(p)^{22}\text{Mg}$	$Q_p=11620(100)$ , $10410(70)$ to ground state and $2^+$ level at 1247.02 keV										
*	also $Q_{2p}=6180(100)$ , $5860(100)$ to ground state and 331.90 level in $^{21}\text{Na}$										
$^{24}\text{O}-u$		20080	1070	19860	180	-0.1	o		GA1	1.5	87Gi05
		20000	500			-0.2	U		TO2	1.5	88Wo09
		20659	442			-1.2	o		GA3	1.5	91Or01
		20460	340			-1.2	o		GA5	1.5	99Sa.A
		19861	118				2		GA7	1.5	07Ju03
$^{24}\text{F}-u$		8070	170	8100	100	0.1	U		TO1	1.5	86Vi09
		8450	240			-1.0	U		GA1	1.5	87Gi05
		8135	86			-0.3	2		GA3	1.5	91Or01
		8030	120			0.4	2		TO4	1.5	91Zh24
$^{24}\text{Mg}-\text{H}_{24}$		-202759.080	0.014	-202759.076	0.014	0.3	1	98	98 $^{24}\text{Mg}$		ST2 1.0 03Be02
$^{24}\text{Ne}-^{22}\text{Ne}_{1.091}$		3009.49	0.55				2		MA8	1.0	04Bi20
$^{24}\text{Mg}-^{23}\text{Na}_{1.043}$		-4287.23	0.32	-4287.664	0.014	-0.9	U		Ma8	1.5	08Mu05
$^{24}\text{Al}-^{23}\text{Na}_{1.043}$		10618.18	0.25				2		TT1	1.0	15Ch58
$^{24}\text{Mg}(p,^6\text{He})^{19}\text{Na}$		-37213	70	-37166	11	0.7	U		Brk		69Ce01
$^{24}\text{Mg}(^3\text{He},^8\text{Li})^{19}\text{Na}$		-32876	12	-32878	11	-0.1	1	77	77 $^{19}\text{Na}$		MSU 75Be38



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{24}\text{Mg}(\alpha, ^8\text{He})^{20}\text{Mg}$	-60900	210	-60596.0	1.9	1.4	U					74Ro17	
	-60677	27			3.0	B			Tex		76Tr03	
$^{24}\text{Mg}(^3\text{He}, ^6\text{He})^{21}\text{Mg}$	-27488	40	-27498.3	0.8	-0.3	U			Brk		70Me11	
	-27512	18			0.8	U			MSU		71Tr03	
	5587	10	5587.8	0.5	0.1	U			LAl		61Si03 Z	
$^{24}\text{Mg}(d, \alpha)^{22}\text{Na}$	1955	12	1958.75	0.17	0.3	U			MIT		64Sp12	
$^{24}\text{Mg}(p, t)^{22}\text{Mg}$	-21194	3	-21194.5	0.3	-0.2	U			MSU		74Ha02	
	-21198.3	1.5			2.5	U			MSU		74No07	
	-21193.9	1.0			-0.6	U			Yal		05Pa31	
	6959.50	0.12	6959.365	0.016	-1.1	o			BNn		74Gr37 Z	
$^{23}\text{Na}(n, \gamma)^{24}\text{Na}$	6959.42	0.07			-0.8	2			BNn		80Gr12 *	
	6959.67	0.14			-2.2	U			ILn		83Hu11 Z	
	6959.38	0.08			-0.2	U			Ptn		83Ti02	
	6959.44	0.05			-1.5	2			ORn		04To03	
	6959.59	0.14			-1.6	o			Bdn		06Fi.A	
	6959.352	0.018			0.7	2			Bdn		14Fi01	
	4735	7	4734.799	0.016	0.0	U			CIT		52Mi54 Y	
$^{23}\text{Na}(d, p)^{24}\text{Na}$	4736	5			-0.2	U			Mex		64Ma.B	
	4736	7			-0.2	U			MIT		64Sp12	
	11692.95	0.17	11692.687	0.013	-1.5	U			Wis		67Mo17 Z	
$^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$	11691.2	1.1			1.4	U					72Me09	
	11692.43	0.31			0.8	U					85Uh01 Z	
	-14307.5	1.5	-14306.81	0.16	0.5	U			MSU		74No07	
$^{24}\text{Mg}(p, d)^{23}\text{Mg}$	4051	15	4046.25	0.16	-0.3	U			Man		59Ba13 Y	
$^{24}\text{Mg}(^3\text{He}, \alpha)^{23}\text{Mg}$	-37397	27	-37384.2	0.4	0.5	U					01Ca37	
$^{24}\text{Mg}(^7\text{Li}, ^8\text{He})^{23}\text{Al}$	4086	9	4085	3	-0.1	3			Brk		79Ay01	
	4084.5	3.5			0.1	3			MSU		80Le18	
	4093	20			-0.4	U			Bor		98Cz01	
$^{24}\text{Ne}(\beta^-)^{24}\text{Na}$	2449	50	2466.3	0.5	0.3	U					56Dr11	
$^{24}\text{Na}(\beta^-)^{24}\text{Mg}$	5511.8	2.	5515.669	0.021	1.9	U					61De25 *	
	5515.8	2.			-0.1	U					64Le09 *	
	5516.8	2.			-0.6	U					65Be24 *	
	5511.5	1.0			4.2	B					69Bo48 *	
	5511.8	2.			1.9	U					72Gi17 *	
5512.5	1.2			2.6	U					76Ge06 *		
$^{24}\text{Al}(\beta^+)^{24}\text{Mg}$	13880	50	13884.70	0.23	0.1	U					68Ar03	
$^{24}\text{Mg}(p, n)^{24}\text{Al}$	-14659.0	2.8	-14667.05	0.23	-2.9	B			Yal		69Ov01 Z	
$^{24}\text{Mg}(^3\text{He}, t)^{24}\text{Al}$	-13880	60	-13903.30	0.23	-0.4	U			Brk		66Ma18	
$^{24}\text{Mg}(^3\text{He}, t)^{24}\text{Al}-^{36}\text{Ar}()^{36}\text{K}$	-1071.48	1.05	-1070.2	0.4	1.2	U			Mun		10Wr01	
$^{24}\text{Mg}(\pi^+, \pi^-)^{24}\text{Si}$	-23594	52	-23657	19	-1.2	2					80Bu15	
$^{*23}\text{Na}(n, \gamma)^{24}\text{Na}$	Original value ( $Z$ ) increased by 0.037 for better recoil correction										AHW	**
$^{*24}\text{Na}(\beta^-)^{24}\text{Mg}$	$E_{\beta^-} = 1389(2) 1393(2) 1394(2) 1388.7(1.0) 1389(2) 1389.7(1.2)$ respectively,										GAu	**
*	to $4^+$ level at 4122.889 keV										Ens07a	**
$^{25}\text{F}-u$	12010	220	12170	100	0.5	o			TO1	1.5	86Vi09	
	12010	290			0.4	o			GA1	1.5	87Gi05	
	12210	150			-0.2	2			TO2	1.5	88Wo09	
	12120	151			0.2	o			GA3	1.5	91Or01	
	11990	130			0.9	2			TO4	1.5	91Zh24	
	12249	97			-0.6	2			GA7	1.5	07Ju03	
$^{25}\text{Ne}-u$	-2293	32	-2190	30	3.4	F			P40	1.0	01Lu20 *	
	-2166	41			-0.5	1	58	58 $^{25}\text{Ne}$	LZ1	1.0	15Xu14	
$^{25}\text{Al}-^{25}\text{Mg}$	4591.342	0.048	4591.34	0.05	0.0	1	100	100 $^{25}\text{Al}$	JY1	1.0	16Ca22	
$^{25}\text{Mg}(p, \alpha)^{22}\text{Na}$	-3151	8	-3147.22	0.18	0.5	U			MIT		59Br74 Y	
$^{23}\text{Na}(t, p)^{25}\text{Na}$	7488.8	1.2				2			Str		84An17	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{25}\text{Mg}(d,\alpha)^{23}\text{Na}$	7026	13	7047.88	0.05	1.7	U			MIT		64Sp12
	7048	10			0.0	U					67Ha17
$^{25}\text{O}(\gamma,n)^{24}\text{O}$	776	15	757	8	-1.2	3					08Ho03 *
	740	40			0.4	U					13Ca18 *
	749	10			0.8	3					16Ko11
											69Ha.A
$^{24}\text{Mg}(n,\gamma)^{25}\text{Mg}$	7330.5	9.99	7330.53	0.05	0.0	U					80Is02 Z
	7330.5	0.3			0.1	U			MMn		82Hu02 Z
	7330.78	0.14			-1.8	U			ILn		85Ke.A
	7330.4	0.2			0.7	U			MMn		90Pr02 Z
	7330.64	0.08			-1.4	-			MMn		92Wa06
	7330.69	0.05			-3.2	B			ORn		06Fi.A
	7330.53	0.15			0.0	-			Bdn		61Hi11 Y
$^{24}\text{Mg}(d,p)^{25}\text{Mg}$	5098	12	5105.96	0.05	0.7	U			Har		61Ja23
	5112	12			-0.5	U			Mex		64Sp12
	5102	7			0.6	U			MIT		average
$^{24}\text{Mg}(n,\gamma)^{25}\text{Mg}$	ave.	7330.62	0.07	7330.53	0.05	-1.2	1	45	43 $^{25}\text{Mg}$		71Ev01 Z
$^{24}\text{Mg}(p,\gamma)^{25}\text{Al}$	2271.6	1.1	2271.38	0.07	-0.2	U					72Pi07 Z
	2271.7	0.7			-0.5	U					85Uh01 Z
	2271.4	0.8			0.0	U					73Br27
$^{24}\text{Mg}(^3\text{He},d)^{25}\text{Al}$		-3218.0	4.5	-3222.10	0.07	-0.9	U		NDm		68Te01 *
$^{24}\text{Mg}(p,\gamma)^{25}\text{Al}^i$		-5629.3	5.8	-5629.7	1.8	-0.1	U				73Go11
$^{25}\text{Ne}(\beta^-)^{25}\text{Na}$		7380	300	7322	29	-0.2	U				54Na18
$^{25}\text{Na}(\beta^-)^{25}\text{Mg}$		3650	250	3835.0	1.2	0.7	U				55Ma63
		4000	200			-0.8	U				60Wa04
$^{25}\text{Al}(\beta^+)^{25}\text{Mg}$		4292	30	4276.81	0.04	-0.5	U				69Fr08
$^{25}\text{Mg}(p,n)^{25}\text{Al}$		-5058	6	-5059.15	0.04	-0.2	U		Har		77Ro03
$^{25}\text{Al}^i(\text{IT})^{25}\text{Al}$		7901	2	7901.1	1.8	0.1	1	85	85 $^{25}\text{Al}^i$		06Ga04 **
* $^{25}\text{Ne}-u$	F : rejected by authors: "unreliable double peak"										
* $^{25}\text{O}(\gamma,n)^{24}\text{O}$	Symmetrized from 770(+20-10)										
* $^{25}\text{O}(\gamma,n)^{24}\text{O}$	Symmetrized from 725(+54-29)										
* $^{24}\text{Mg}(p,\gamma)^{25}\text{Al}^i$	IT=7916(6), $Q$ rebuilt with Ame1964, error estimated by evaluator										
$^{26}\text{F}-u$	19800	1000	20020	120	0.1	o			TO1	1.5	86Vi09
	20940	640			-1.0	o			GA1	1.5	87Gi05
	19820	210			0.6	2			TO2	1.5	88Wo09
	19544	300			1.1	o			GA3	1.5	91Or01
	19490	210			1.7	U			TO4	1.5	91Zh24
	20054	86			-0.3	2			GA7	1.5	07Ju03
$^{26}\text{Ne}-u$	448	90	516	20	0.5	2			GA3	1.5	91Or01
	461	33			1.7	o			P40	1.0	01Lu20
	518	20			-0.1	2			P40	1.0	06Ga04
$^{26}\text{Na}-u$	-7367	7	-7365	4	0.2	o			P40	1.0	01Lu17
	-7365	4			-0.1	2			P40	1.0	06Ga04 *
	-7368	11			0.2	2			P40	1.0	06Ga04
$^{26}\text{Mg}-\text{H}_{26}$	-220857.848	0.034	-220857.87	0.03	-0.5	1	89	89 $^{26}\text{Mg}$	ST2	1.0	03Be02
$^{26}\text{Al}-^{23}\text{Na}_{1.130}$	-1547.46	0.24	-1547.43	0.07	0.1	U			MA8	1.0	08Ge08
$^{26}\text{Si}-^{23}\text{Na}_{1.13}$	3895.1	2.1	3894.52	0.12	-0.3	U			MS1	1.0	10Kw02
$^{26}\text{Al}-^{25}\text{Mg}_{1.040}$	1621.46	0.48	1621.42	0.06	-0.1	U			JY1	1.0	06Er08
$^{26}\text{Al}^m-^{25}\text{Mg}_{1.040}$	1867.09	0.53	1866.52	0.06	-1.1	U			JY1	1.0	06Er08
$^{26}\text{Al}-^{26}\text{Mg}$	4299.14	0.17	4298.89	0.07	-1.5	1	16	15 $^{26}\text{Al}$	JY1	1.0	06Er08
$^{26}\text{Al}^m-^{26}\text{Mg}$	4544.09	0.17	4543.99	0.07	-0.6	1	16	16 $^{26}\text{Al}^m$	JY1	1.0	06Er08
$^{26}\text{Al}^m-^{26}\text{Al}$	245.09	0.17	245.096	0.014	0.0	U			JY1	1.0	06Er08
	244.91	0.14			1.3	U			JY1	1.0	09Er02
	245.114	0.049			-0.4	U			JY1	1.0	09Er07
$^{26}\text{Si}-^{26}\text{Al}$	5441.97	0.14	5441.94	0.09	-0.2	2			JY1	1.0	09Er02
	5441.92	0.12			0.2	2			JY1	1.0	09Er02 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{25}\text{Na}-^{26}\text{Na}_{.721} \ ^{22}\text{Na}_{.284}$	-2881	33	-2939.6	2.8	-1.8	U	P13	1.0	75Th08
	-2921	22			-0.8	U	P13	1.0	75Th08
$^{26}\text{Al}(n,\alpha)^{23}\text{Na}$	2966.5	2.5	2966.10	0.07	-0.2	U			01Wa50
$^{23}\text{Na}(\alpha,n)^{26}\text{Al}$	-2968	4	-2966.10	0.07	0.5	U	Duk		60Wi07 Y
$^{26}\text{O}(\gamma,2n)^{24}\text{O}$	90	110	18	5	-0.7	U			12Lu07 *
	18	5				3			16Ko11
$^{24}\text{Mg}(t,p)^{26}\text{Mg}$	9940	12	9941.81	0.03	0.2	U	Har		61Hi11 Y
$^{24}\text{Mg}(^3\text{He},p)^{26}\text{Al}$	5932	15	5918.83	0.07	-0.9	U	Ald		59Hi66 Y
	5922	8			-0.4	U	Phi		72Be51
$^{24}\text{Mg}(^3\text{He},n)^{26}\text{Si}$	85	18	67.35	0.11	-1.0	U	CIT		67Mi02
	75	30			-0.3	U			67Mc03
	95	15			-1.8	U	Har		68Ad03
	65	30			0.1	U	Ber		68Ha09
$^{26}\text{Mg}(^7\text{Li},^8\text{B})^{25}\text{Ne}$	-22050	100	-22194	29	-1.4	-	Brk		73Wi06
$^{26}\text{Mg}(^{13}\text{C},^{14}\text{O})^{25}\text{Ne}$	-19067	50	-19062	29	0.1	-	Can		85Wo04
$^{26}\text{Mg}(^7\text{Li},^8\text{B})^{25}\text{Ne}$	ave. -22170	40	-22194	29	-0.5	1	42	42 $^{25}\text{Ne}$	average
$^{26}\text{Mg}(d,^3\text{He})^{25}\text{Na}$	-8653	10	-8652.2	1.2	0.1	U	MSU		73Be14
$^{26}\text{Mg}(t,\alpha)^{25}\text{Na}$	5664	12	5668.2	1.2	0.3	U	Ald		62Hi01
$^{25}\text{Mg}(n,\gamma)^{26}\text{Mg}$	11092.9	0.3	11093.08	0.04	0.6	U	MMn		80Is02
	11091.84	0.44			2.8	U	ILn		82Hu02 Z
	11093.10	0.06			-0.4	1	55	46 $^{25}\text{Mg}$	MMn 90Pr02 Z
	11093.17	0.06			-1.6	o			ORn 91Ki04 Z
	11093.23	0.05			-3.1	B			ORn 92Wa06 Z
	11093.16	0.22			-0.4	U			Bdn 06Fi.A
$^{25}\text{Mg}(d,p)^{26}\text{Mg}$	8865	12	8868.51	0.04	0.3	U	Ald		61Hi11 Y
	8876	12			-0.6	U	Mex		61Ja23
	8889	12			-1.7	U	MIT		64Sp12
$^{25}\text{Mg}(p,\gamma)^{26}\text{Al}$	6305.0	1.2	6306.34	0.06	1.1	U			74De37
	6304.9	1.1			1.3	U			79E111
	6306.39	0.11			-0.5	-			85Be17 Z
	6306.38	0.08			-0.5	-			Utr 91Ki04 Z
	ave. 6306.38	0.06			-0.7	1	75	64 $^{26}\text{Al}$	average
$^{26}\text{Si}^i(p)^{25}\text{Al}$	7563	15	7553	11	-0.6	2	Brk		83Ca06
	7544	15			0.6	2	Brk		83Ho23 *
$^{26}\text{Mg}(\pi^-, \pi^+)^{26}\text{Ne}$	-17676	72	-17718	18	-0.6	U			80Na12
$^{26}\text{Na}(\beta^-)^{26}\text{Mg}$	9210	200	9354	4	0.7	U			73Al13
$^{26}\text{Mg}(t,^3\text{He})^{26}\text{Na}$	-9292	20	-9335	4	-2.2	U	LAl		74FI01
$^{26}\text{Mg}(^7\text{Li},^7\text{Be})^{26}\text{Na}$	-10182	40	-10216	4	-0.8	U	ChR		72Ba35 *
$^{26}\text{Al}(\beta^+)^{26}\text{Mg}$	3991	8	4004.39	0.06	1.7	U			58Fe16 *
$^{26}\text{Mg}(p,n)^{26}\text{Al}$	-4786.7	10.	-4786.74	0.06	0.0	U	Oak		55Ki28 *
	-4787.04	0.48			0.6	U	Utr		69De27
	-4786.1	1.6			-0.4	U	Har		69Fr08 *
	-4785.66	0.22			-4.9	C	Auc		84Ba.B *
	-4786.57	0.05			-3.4	C	Auc		92Ba.A *
	-4786.25	0.12			-4.1	B	Auc		94Br11 *
$^{26}\text{Mg}(^3\text{He},t)^{26}\text{Al}$	-4023.0	0.6	-4022.98	0.06	0.0	F	Mun		77Vo02 *
$^{26}\text{Mg}(^3\text{He},t)^{26}\text{Al}-^{27}\text{Al}(O)^{27}\text{Si}$	808.2	2.0	807.97	0.12	-0.1	U	ChR		74Ha35
$^{26}\text{Mg}(^3\text{He},t)^{26}\text{Al}-^{14}\text{N}(O)^{14}\text{O}$	1139.43	0.13	1139.97	0.07	4.2	B	ChR		87Ko34 *
$^{26}\text{Al}^m(\text{IT})^{26}\text{Al}$	228.305	0.013	228.306	0.013	0.0	1	99	84 $^{26}\text{Al}^m$	Ens164
$^{26}\text{Si}(\beta^+)^{26}\text{Al}$	5079	13	5069.14	0.08	-0.8	U			63Fr10
* $^{26}\text{Na}-u$	Result from the "Thermo" experiment. Next item from "Rilis"								06Ga04 **
* $^{26}\text{Si}-^{26}\text{Al}$	$D_M=5196.82(0.12) \mu\text{u}$ for $^{26}\text{Al}^m$ at 228.306(0.013); $M-A=-7141.05(0.13) \text{keV}$								Nub16b **
* $^{26}\text{O}(\gamma,2n)^{24}\text{O}$	Symmetrized from 150(+50-150) keV								12Lu07 **
* $^{26}\text{O}(\gamma,2n)^{24}\text{O}$	less than 40 keV								13Ca18 **
* $^{26}\text{Si}^i(p)^{25}\text{Al}$	$E_p=3699(15)$ to 3695.5 level; different from preceding data								Ens098 **
* $^{26}\text{Mg}(^7\text{Li},^7\text{Be})^{26}\text{Na}$	$Q=-10222(30)$ corrected for contribution of unresolved 82.2 level								Ens164 **
* $^{26}\text{Al}(\beta^+)^{26}\text{Mg}$	$E_{\beta^+}=1160(8)$ to $2^+$ level at 1808.74 level								Ens164 **
* $^{26}\text{Mg}(p,n)^{26}\text{Al}$	$T=5191(10,Z)$ to $^{26}\text{Al}^m$ at 228.306 keV								Nub16b **
* $^{26}\text{Mg}(p,n)^{26}\text{Al}$	$T=5209.3(1.6,Z)$ to $^{26}\text{Al}^m$ at 228.306 keV								Nub16b **
* $^{26}\text{Mg}(p,n)^{26}\text{Al}$	$T=5208.86(0.23)$ to $^{26}\text{Al}^m$ at 228.306 keV								Nub16b **
* $^{26}\text{Mg}(p,n)^{26}\text{Al}$	$T=5209.71(0.05)$ to $^{26}\text{Al}^m$ at 228.306 keV								Nub16b **
* $^{26}\text{Mg}(p,n)^{26}\text{Al}$	$T=5209.46(0.12)$ to $^{26}\text{Al}^m$ at 228.306 keV								Nub16b **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* $^{26}\text{Mg}(^3\text{He,t})^{26}\text{Al}$	$Q = -4251.3(0.6, Z)$ to $^{26}\text{Al}^m$ at 228.306 keV										Nub16b **
* $^{26}\text{Mg}(^3\text{He,t})^{26}\text{Al}$	F : rejected in reference of same group										09Fa15 **
* $^{26}\text{Mg}(^3\text{He,t})^{26}\text{Al}-^{14}\text{N}(^{14}\text{O})$	$Q(\text{to } 1057.740(0.023) \text{ level})-^{14}\text{N}(^{14}\text{O})=81.69(0.13)$										82Al19 **
$^{27}\text{F}-\text{u}$	27500	700	27320	420	-0.2	U			TO2	1.5	88Wo09
	26005	770			1.1	U			GA3	1.5	91Or01
	27100	900			0.2	U			TO4	1.5	91Zh24
	26900	580			0.5	o			GA5	1.5	99Sa.A
	26441	204			2.9	B			GA7	1.5	07Ju03
	27322	279				2			GA8	1.5	12Ga45
$^{27}\text{Ne}-\text{u}$	6010	640	7570	100	1.6	F			TO1	1.5	86Vi09 *
	7470	300			0.2	U			GA1	1.5	87Gi05
	7567	172			0.0	o			GA3	1.5	91Or01
	7670	130			-0.5	2			TO4	1.5	91Zh24
	7536	75			0.3	2			GA7	1.5	07Ju03
$^{27}\text{Na}-\text{u}$	-5922	11	-5924	4	-0.1	o			P40	1.0	01Lu17 *
	-5922	4			-0.4	o			P40	1.0	06Ga04 *
$^{27}\text{Al}-^{23}\text{Na}_{1.174}$	-6450.79	0.25	-6450.73	0.05	0.2	U			MA8	1.0	08Ge08
	-6450.754	0.054			0.5	1	89	89 $^{27}\text{Al}$	TT1	1.0	16Kw.A
$^{27}\text{Na}-^{27}\text{Al}$	12538	4				2			P40	1.0	01Lu17
$^{24}\text{Na}-^{27}\text{Na}_{.356} \ ^{22}\text{Na}_{.655}$	-3006	38	-3059.7	1.3	-0.9	U			P10	1.5	75Th08
$^{26}\text{Na}-^{27}\text{Na}_{.770} \ ^{22}\text{Na}_{.236}$	-1437	86	-1389	5	0.6	U			P13	1.0	75Th08
$^{26}\text{Na}-^{27}\text{Na}_{.481} \ ^{25}\text{Na}_{.520}$	676	66	659	4	-0.2	U			P10	1.5	75Th08
	734	86			-0.6	U			P11	1.5	75Th08
$^{23}\text{Na}(\alpha, \gamma)^{27}\text{Al}$	10090.0	1.3	10091.92	0.05	1.5	U			Utr		78Ma23
$^{27}\text{Al}(\text{p}, \alpha)^{24}\text{Mg}$	1601.7	0.7	1600.76	0.05	-1.3	o			Zur		63Ry04
	1598.4	1.0			2.4	U			NDm		65Br28
	1601.3	0.5			-1.1	U			Zur		67St30 Z
	1600.06	0.21			3.3	B			Utr		78Ma23 Z
$^{24}\text{Mg}(\alpha, \text{p})^{27}\text{Al}$	-1598.9	1.0	-1600.76	0.05	-1.9	U			NDm		65Br28
$^{25}\text{Mg}(\text{t}, \text{p})^{27}\text{Mg}$	9055	11	9054.67	0.06	0.0	U			Tal		61Hi11 Y
$^{27}\text{Al}(\text{d}, \alpha)^{25}\text{Mg}$	6699	12	6706.73	0.07	0.6	U			Ald		61Hi11 Y
	6691	11			1.4	U			Tal		62Sh01 Y
	6700	10			0.7	U			MIT		64Sp12
	-23843.4	4.7	-23842.8	1.8	0.1	1	15	15 $^{25}\text{Al}^i$	MSU		73Be14 *
$^{27}\text{P}(^{25}\text{Al})^{25}\text{Al}$	6410	45	6350	30	-1.4	2			Lis		91Bo32
	6270	50			1.5	2					01Ca60 *
$^{26}\text{Mg}(^{18}\text{O}, ^{17}\text{F})^{27}\text{Na}$	-13295	55	-13431	4	-2.5	F			Mun		78Pa12 *
	-13433	60			0.0	U			Can		85Fi08
$^{26}\text{Mg}(\text{n}, \gamma)^{27}\text{Mg}$	6443.35	0.55	6443.39	0.04	0.1	U			ILn		82Hu02
	6443.56	0.25			-0.7	o			MMn		85Ke.A
	6443.26	0.08			1.6	2			MMn		90Pr02 Z
	6443.44	0.05			-1.1	2			ORn		92Wa06 Z
	6443.35	0.13			0.3	2			Bdn		06Fi.A
$^{26}\text{Mg}(\text{d}, \text{p})^{27}\text{Mg}$	4214	12	4218.82	0.04	0.4	U			Ald		61Hi11 Y
	4215	10			0.4	U			Mex		61Ja23
	4211	6			1.3	U			MIT		64Sp12
	8270.8	0.5	8271.29	0.06	1.0	U			Utr		59An33 *
$^{26}\text{Mg}(\text{p}, \gamma)^{27}\text{Al}$	8271.2	0.5			0.2	U					63Va24 Z
	8271.3	0.5			0.0	U			Utr		78Ma24 *
	11541	12	11542.58	0.06	0.1	U			Ald		61Hi11
$^{27}\text{Al}(^3\text{He}, \alpha)^{26}\text{Al}$	7523	15	7519.59	0.08	-0.2	U			Ald		59Hi66
	7519	15			0.0	U			Man		60Ta12
$^{26}\text{Al}(\text{p}, \gamma)^{27}\text{Si}$	7464.9	0.9	7463.32	0.13	-1.8	U					84Bu09 Z
$^{27}\text{Na}(\beta^-)^{27}\text{Mg}$	8930	150	9069	4	0.9	U					73Al13

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{27}\text{Mg}(\beta^-)^{27}\text{Al}$	2600	11	2610.25	0.07	0.9	U					54Da22
$^{27}\text{Si}(\beta^+)^{27}\text{Al}$	4872	20	4812.36	0.10	-3.0	B					60Wa04
$^{27}\text{Al}(\text{p,n})^{27}\text{Si}$	-5573	10	-5594.71	0.10	-2.2	U			Oak		55Ki28 Z
	-5597.5	6.0			0.5	U			Tkm		63Ok01
	-5593.6	4.3			-0.3	U			Ric		65Ku02
	-5585.1	2.3			-4.2	B			Ric		66Bo20 Z
	-5592.0	1.0			-2.7	U			Yal		69Ov01 Z
	-5594.1	3.2			-0.2	U			Har		76Fr13
	-5593.8	0.26			-3.5	F			Auc		77Na24 *
	-5594.27	0.11			-4.0	F			Auc		85Wh03 *
	-5594.72	0.10				2			Auc		94Br37 Z
* $^{27}\text{Ne}-\text{u}$	F : contaminated by $^{27}\text{Na}$										
* $^{27}\text{Na}-\text{u}$	Not independent of $^{27}\text{Na}-^{27}\text{Al}$ from isobaric method, do not use										
* $^{27}\text{Al}(\text{p,t})^{25}\text{Al}^i$	IT=7904(5), rebuilt $Q=-23847.9(4.7)$ ; recalib +4.5keV										
* $^{27}\text{P}^i(2\text{p})^{25}\text{Al}$	And $E_{2p}=5315(60)$ to $3/2^+$ level at 944.9 keV										
* $^{26}\text{Mg}(^{18}\text{O},^{17}\text{F})^{27}\text{Na}$	F : shape of peak raises doubt on centroid determination										
* $^{26}\text{Mg}(\text{p},\gamma)^{27}\text{Al}$	$E_p=338.65(0.12)$ to $8596.8(0.5)$ level										
* $^{26}\text{Mg}(\text{p},\gamma)^{27}\text{Al}$	$E_p=338.21(0.30)$ to $8596.8(0.5)$ level										
* $^{26}\text{Mg}(\text{p},\gamma)^{27}\text{Al}$	$E_p=809.90(0.05,Z)$ to $9050.7(0.5,Z)$ level										
* $^{27}\text{Al}(\text{p,n})^{27}\text{Si}$	F : rejected by same group "measurement contains error"										
											94Br37 **
$^{28}\text{Ne}-\text{u}$	11490	430	12130	140	1.0	o			TO1	1.5	86Vi09
	12270	560			-0.2	o			GA1	1.5	87Gi05
	11958	238			0.5	o			GA3	1.5	91Or01
	12160	140			-0.1	2			TO4	1.5	91Zh24
	12110	118			0.1	2			GA7	1.5	07Ju03
$^{28}\text{Na}-\text{u}$	-1220	190	-1061	11	0.6	o			TO1	1.5	86Vi09
	-1097	96			0.3	U			GA3	1.5	91Or01
	-1062	14			0.1	o			P40	1.0	01Lu17
	-1061	11				2			P40	1.0	06Ga04
$^{28}\text{Si}-\text{u}$	-23073.43	0.30	-23073.4650	0.0005	-0.1	U			ST1	1.0	93Je06
	-23073.4676	0.0020			1.3	U			MII	1.0	95Di08
	-23073.00	0.27			-0.7	U			OH1	2.5	94Go.A
	-23073.466	0.008			0.1	U			ST2	1.0	02Be64 *
$\text{C}_2 \text{D}_2-^{28}\text{Si}$	51277.0224	0.0024	51277.0212	0.0006	-0.5	U			MII	1.0	95Di08
$^{15}\text{N}_2-^{28}\text{Si} \text{H}_2$	7641.2007	0.0024	7641.1984	0.0013	-1.0	1	29	26 $^{15}\text{N}$	MII	1.0	95Di08
$\text{C}_2 \text{H}_4-^{28}\text{Si}$	54373.59360	0.00079	54373.5940	0.0006	0.5	1	54	38 $^{28}\text{Si}$	FS1	1.0	08Re16
$^{13}\text{C}_2 \text{H}_2-^{28}\text{Si}$	45433.19986	0.00071	45433.1999	0.0005	0.1	1	53	34 $^{28}\text{Si}$	FS1	1.0	08Re16
$^{28}\text{Si}_2 \text{ } ^{16}\text{O}-^{35}\text{Cl} \text{ } ^{37}\text{Cl}$	14013.07	0.70	14012.41	0.07	-0.6	U			H46	1.5	93Nx02
$^{25}\text{Na}-^{28}\text{Na}_{.446} \text{ } ^{22}\text{Na}_{.568}$	-5869	75	-5974	5	-0.9	U			P10	1.5	75Th08 *
$^{26}\text{Na}-^{28}\text{Na}_{.619} \text{ } ^{22}\text{Na}_{.394}$	-4229	613	-4207	7	0.0	U			P11	1.5	75Th08
	-4205	128			0.0	U			P12	2.5	75Th08
	-4203	87			-0.1	U			P13	1.0	75Th08
$^{28}\text{Si}(\text{p},^6\text{He})^{23}\text{Al}$	-38569	80	-38544.0	0.3	0.3	U			Brk		69Ce01
$^{28}\text{Si}(^3\text{He},^8\text{Li})^{23}\text{Al}$	-34274	25	-34255.5	0.3	0.7	U			MSU		75Be38
$^{28}\text{Si}(\alpha,^8\text{He})^{24}\text{Si}$	-61433	21	-61423	19	0.5	R			Tex		80Tr04
$^{28}\text{Si}(\text{p},\alpha)^{25}\text{Al}$	-7709.3	2.6	-7712.76	0.06	-1.3	U			NDm		73Br27
$^{28}\text{Si}(^3\text{He},^6\text{He})^{25}\text{Si}$	-27976	50	-27981	10	-0.1	U			Brk		70Me11
	-27981	10				2			MSU		72Be12
$^{26}\text{Mg}(\text{t,p})^{28}\text{Mg}$	6474	12	6465.1	2.0	-0.7	U			Har		61Hi11 Y
$^{26}\text{Mg}(^3\text{He,p})^{28}\text{Al}$	8285	5	8278.35	0.08	-1.3	U			Phi		74Be07
$^{28}\text{Si}(\text{d},\alpha)^{26}\text{Al}$	1429	4	1428.16	0.07	-0.2	U			MIT		64Sp12
$^{28}\text{Si}(\text{p,t})^{26}\text{Si}$	-22009	3	-22012.62	0.11	-1.2	U			MSU		74Ha02
	-22014.1	1.0			1.5	U			Yal		05Pa31
$^{28}\text{F}(\gamma,\text{n})^{27}\text{F}$	220	50				3			MSU		12Ch02

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{27}\text{Al}(n,\gamma)^{28}\text{Al}$	7725.02	0.20	7725.10	0.06	0.4	U			BNn		78St25 Z
	7725.07	0.30			0.1	U			ILn		79Br25 Z
	7725.13	0.3			-0.1	U			MMn		80Is02 Z
	7725.02	0.10			0.8	2					81Su.A Z
	7725.14	0.09			-0.4	2			ILn		82Sc14 Z
	7725.17	0.15			-0.5	2			Bdn		06Fi.A
$^{27}\text{Al}(d,p)^{28}\text{Al}$	5511	5	5500.53	0.06	-2.1	U			Mex		61Ja23
	5503	10			-0.2	U			MIT		64Sp12
$^{27}\text{Al}(p,\gamma)^{28}\text{Si}$	11584.89	0.30	11584.90	0.05	0.0	U			Utr		78Ma23 Z
	11585.09	0.14			-1.3	1	11	11	$^{27}\text{Al}$		28Sir-0
$^{27}\text{Al}(^3\text{He},d)^{28}\text{Si}$	6049	18	6091.43	0.05	2.4	U					60Fo01
$^{27}\text{Al}(p,\gamma)^{28}\text{Si}^r$	-956.15	0.03	-956.139	0.025	0.3	2			Utr		78Ma23 Z
	-956.025	0.020			-5.7	B			Auc		94Br37 Z
	-956.13	0.05			-0.4	2					98Wa.A Z
$^{28}\text{Si}(^3\text{He},\alpha)^{27}\text{Si}$	3407	15	3398.01	0.11	-0.6	U			Ald		59Hi68 Y
$^{28}\text{Si}(^3\text{He},\alpha)^{27}\text{Si}^i$	-3225.5	2.6	-3227.0	2.3	-0.6	1	79	79	$^{27}\text{Si}^i$		86Sc21 *
$^{28}\text{Si}(^7\text{Li},^8\text{He})^{27}\text{P}$	-37513	40	-37473	26	1.0	R					01Ca37
$^{28}\text{P}^i(p)^{27}\text{Si}$	3835	20				3			Lis		89Po10
$^{28}\text{Mg}(\beta^-)^{28}\text{Al}$	1791	10	1831.8	2.0	4.1	B					53Ma23 *
	1831.8	2.0				3					54Ol03 *
$^{28}\text{Al}(\beta^-)^{28}\text{Si}$	4644	10	4642.15	0.08	-0.2	U					52Mo22 *
	4657	14			-1.1	U					54Ol03 *
$^{28}\text{Si}^r(\text{IT})^{28}\text{Si}$	12541.23	0.14	12541.04	0.05	-1.3	R			Utr		90En02 Z
$^{28}\text{P}(\beta^+)^{28}\text{Si}$	14290	40	14345.1	1.2	1.4	U					68Ar03 *
$^{28}\text{Si}(p,n)^{28}\text{P}$	-15118.3	4.1	-15127.4	1.2	-2.2	U			Yal		69Ov01 Z
	-15112.5	5.8			-2.6	U			BNL		71Go18 Z
$^{28}\text{Si}(^3\text{He},t)^{28}\text{P}$	-14380	60	-14363.6	1.2	0.3	U			Brk		66Ma18
$^{28}\text{Si}(^3\text{He},t)^{28}\text{P}-^{36}\text{Ar}(^36\text{K})$	-1530.58	1.10	-1530.6	1.1	0.0	1	100	100	$^{28}\text{P}$		10Wr01
$^{28}\text{Si}(\pi^+, \pi^-)^{28}\text{S}$	-24544	160				2			Mun		82Mo12 *
* $^{28}\text{Si}-u$	Unc. was erroneously 0.0008 in Ame2012, was correct in Ame2003										
* $^{25}\text{Na}-^{28}\text{Na}_{.446}-^{22}\text{Na}_{.568}$	Symmetric double-doublet 22-24 26-28 included										
* $^{28}\text{Si}(^3\text{He},\alpha)^{27}\text{Si}^i$	IT=6626(3), $Q$ rebuilt with Ame1977										
* $^{28}\text{Mg}(\beta^-)^{28}\text{Al}$	$E_{\beta^-}$ =418(10) 459(2) respectively, to $1^+$ level at 1372.917 keV										
* $^{28}\text{Al}(\beta^-)^{28}\text{Si}$	$E_{\beta^-}$ =2865(10) 2878(14) respectively, to $2^+$ level at 1779.030 keV										
* $^{28}\text{P}(\beta^+)^{28}\text{Si}$	$E_{\beta^+}$ =11490(40) to $2^+$ level at 1779.030 keV										
* $^{28}\text{Si}(\pi^+, \pi^-)^{28}\text{S}$	Original -24603(160) recalibrated to $^{16}\text{O}(\pi^+, \pi^-)^{16}\text{Ne}$ $Q=-27704(20)$ keV										
* $^{28}\text{Si}(\pi^+, \pi^-)^{28}\text{S}$	GAu **										
$^{29}\text{F}-u$	43103	376				2			GA8	1.5	12Ga45
$^{29}\text{Ne}-u$	19433	551	19750	160	0.4	o			GA3	1.5	91Or01
	19300	400			0.8	U			TO4	1.5	91Zh24
	19400	410			0.6	o			GA5	1.5	00Sa21
	19753	107				2			GA7	1.5	07Ju03
	2820	230	2877	8	0.2	U			TO1	1.5	86Vi09
$^{29}\text{Na}-u$	2838	143			0.2	U			GA3	1.5	91Or01
	2861	14			1.1	o			P40	1.0	01Lu17
	2866	13			0.9	1	37	37	$^{29}\text{Na}$	1.0	06Ga04
$^{29}\text{Na}-^{39}\text{K}_{.744}$	29885.9	9.9	29879	8	-0.6	1	63	63	$^{29}\text{Na}$	1.0	13Ch49
$^{29}\text{Mg}-u$	-11375	19	-11383	12	-0.4	2			P40	1.0	06Ga04 *
	-11388	16			0.3	2			P40	1.0	06Ga04
$^{29}\text{Al}-\text{O}_{1.812}$	-10328.8	1.0	-10332.1	0.4	-3.3	C			TT1	1.0	12Ch.A
	-10333.5	1.7			0.8	U			TT1	1.0	16Kw.A
$^{29}\text{P}^{40}\text{Ar}-u$	-55816.80	0.50	-55816.5	0.4	0.6	1	59	59	$^{29}\text{P}$	1.0	15Ei01
$^{29}\text{Al}-^{23}\text{Na}_{1.261}$	-6645.90	0.37				2			TT1	1.0	16Ga.1
$^{29}\text{Si}-^{28}\text{Si}^i$	-8256.90198	0.00024	-8256.90198	0.00024	0.0	1	100	100	$^{29}\text{Si}$	1.0	05Ra34
$^{26}\text{Na}-^{29}\text{Na}_{.512}-^{22}\text{Na}_{.506}$	-5763	91	-5611	5	1.1	U			P10	1.5	75Th08
	-6379	293			1.7	U			P11	1.5	75Th08
	-5252	277			-0.5	U			P12	2.5	75Th08
	-5576	66			-0.5	U			P13	1.0	75Th08

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{27}\text{Na}-^{29}\text{Na}_{,466} \ ^{25}\text{Na}_{,540}$	-1708	124	-1713	5	0.0	U			P10	1.5	75Th08
$^{18}\text{O}(^{13}\text{C},2\text{p})^{29}\text{Mg}$	-1456	50	-1633	11	-3.5	B					81Pa17
$^{26}\text{Mg}(^{11}\text{B},^8\text{B})^{29}\text{Mg}$	-19720	50	-19866	11	-2.9	U			Brk		74Sc26
$^{26}\text{Mg}(^{18}\text{O},^{15}\text{O})^{29}\text{Mg}$	-9207	55	-9250	11	-0.8	U			Mun		78Pa12
	-9250	45			0.0	U			Can		85Fi08
$^{26}\text{Mg}(\alpha,\text{p})^{29}\text{Al}$	-2880	40	-2870.8	0.3	0.2	U			Yal		57Gr47 Y
	-2874	10			0.3	U			ANL		68Be13
$^{29}\text{Si}(\text{n},\alpha)^{26}\text{Mg}$	-21	21	-34.135	0.030	-0.6	U			Ham		62An05
$^{27}\text{Al}(\text{t},\text{p})^{29}\text{Al}$	8679.5	1.2	8671.7	0.3	-6.5	B			Str		84An17
$^{29}\text{Si}(\text{d},\alpha)^{27}\text{Al}$	6000	11	6012.59	0.05	1.1	U			MIT		64Sp12
$^{29}\text{Si}(\text{p},\text{t})^{27}\text{Si}^i$	-23802	5	-23796.4	2.3	1.1	1	21	$^{27}\text{Si}^i$	MSU		77Be13
$^{27}\text{Al}(^3\text{He},\text{n})^{29}\text{P}$	6616	30	6615.9	0.4	0.0	U			Oak		72Gr39
$^{28}\text{Si}(\text{n},\gamma)^{29}\text{Si}$	8473.6	0.3	8473.6012	0.0005	0.0	o			MMn		80Is02 Z
	8473.61	0.04			-0.2	U			MMn		90Is02 Z
	8473.55	0.04			1.3	U			ORn		92Ra19 Z
	8473.5509	0.0500			1.0	o			PTc		97Ro26 *
	8473.54	0.17			0.4	U			Bdn		06Fi.A
	8473.551	0.030			1.7	U			PTc		01Pa52 *
	8473.5957	0.0050			1.1	U			NBS		06De21
$^{28}\text{Si}(\text{d},\text{p})^{29}\text{Si}$	6252	10	6249.03522	0.00029	-0.3	U			Mex		64Ma.B
	6252	10			-0.3	U			MIT		64Sp12
	6249.35	0.5			-0.6	U			Rez		90Pi05 *
$^{28}\text{Si}(\text{p},\gamma)^{29}\text{P}$	2747.1	1.7	2749.0	0.4	1.1	-					73Ba35 Z
	2748.8	0.6			0.4	-					74By01 Z
$^{28}\text{Si}(\text{d},\text{n})^{29}\text{P}$	560	30	524.5	0.4	-1.2	U			Ald		60Ma21
$^{28}\text{Si}(^3\text{He},\text{d})^{29}\text{P}$	-2733	12	-2744.5	0.4	-1.0	U			Ald		60Hi03 Y
$^{28}\text{Si}(\text{p},\gamma)^{29}\text{P}$	ave. 2748.6	0.6	2749.0	0.4	0.7	1	40	$^{40}\text{P}$			average
$^{28}\text{Si}(\text{p},\gamma)^{29}\text{P}^i$	-5630	10	-5632.8	2.5	-0.3	U			ANL		66Yo01
	-5631.9	5.0			-0.2	1	24	$^{24}\text{P}^i$			68Te01 *
$^{29}\text{Cl}(\text{p})^{28}\text{S}$	1800	100				3					15Mu13
$^{29}\text{Mg}(\beta^-)^{29}\text{Al}$	7624	400	7605	11	0.0	U					73Go34
$^{29}\text{Al}(\beta^-)^{29}\text{Si}$	3850	100	3687.3	0.3	-1.6	U					54Na14 *
$^{29}\text{P}(\beta^+)^{29}\text{Si}$	4967	20	4942.2	0.4	-1.2	U					55Ro05
$^{29}\text{P}^i(\text{IT})^{29}\text{P}$	8382.1	2.8	8381.8	2.4	-0.1	1	76	$^{76}\text{P}^i$			72Ba26
$^{29}\text{Mg}-\text{u}$	Result from the "Plasma" experiment. Next item from "Rilis"										
$^{28}\text{Si}(\text{n},\gamma)^{29}\text{Si}$	Original error 0.0005 increased for calibration										
$^{28}\text{Si}(\text{n},\gamma)^{29}\text{Si}$	Original error 0.005 increased for calibration										
$^{28}\text{Si}(\text{d},\text{p})^{29}\text{Si}$	Estimated systematic error 0.5 added to statistical error 0.037 keV										
$^{28}\text{Si}(\text{p},\gamma)^{29}\text{P}^i$	IT=8376(6), $Q$ rebuilt with Ame1964, error estimated by evaluator										
$^{29}\text{Al}(\beta^-)^{29}\text{Si}$	$E_{\beta^-}=1550(100)$ to $5/2^+$ level at 2028.16 and $3/2^+$ at 2425.97 keV										
$^{30}\text{Ne}-\text{u}$	23872	884	24990	270	0.8	o			GA3	1.5	91Or01
	25660	850			-0.5	o			GA5	1.5	00Sa21
	24734	301			0.6	-			GA7	1.5	07Ju03
	25024	301			-0.1	-			GA8	1.5	12Ga45
	ave. 24880	320			0.4	1	73	$^{73}\text{Ne}$			average
$^{30}\text{Na}-\text{u}$	7620	540	9098	5	1.8	F			TO1	1.5	86Vi09 *
	9200	370			-0.2	U			GA1	1.5	87Gi05
	9126	218			-0.1	U			GA3	1.5	91Or01
	9330	130			-1.2	U			TO4	1.5	91Zh24
	8976	27			4.5	B			P40	1.0	01Lu17
	8990	25			4.3	B			P40	1.0	06Ga04
$^{30}\text{Na}-\text{O}_{1.876}$	18638.9	5.6	18638	5	-0.1	1	82	$^{82}\text{Na}$	TT1	1.0	13Ch49
$^{30}\text{Na}-^{39}\text{K}_{,769}$	37004	12	37008	5	0.3	1	18	$^{18}\text{Na}$	TT1	1.0	13Ch49
$^{30}\text{Mg}-\text{O}_{1.876}$	3.0	3.7				2			TT1	1.0	13Ch49

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{30}\text{Mg}-u$	-9700	230	-9537	4	0.5	o			TO1	1.5	86Vi09
	-9597	98			0.4	U			GA3	1.5	91Or01
	-9490	110			-0.3	U			TO4	1.5	91Zh24
	-9546	14			0.6	U			P40	1.0	06Ga04
$^{30}\text{Al}-^{39}\text{K}_{769}$	10878.1	3.1				2			TT1	1.0	16Kw.A
$^{30}\text{P}-^{30}\text{Si}$	4543.353	0.066				3			JY1	1.0	16Ca22
$^{30}\text{S}-^{30}\text{P}$	6593.28	0.21				4			JY1	1.0	11So11
$^{26}\text{Na}-^{30}\text{Na}_{433} \quad ^{22}\text{Na}_{591}$	-7454	287	-7468	4	0.0	U			P10	1.5	75Th08
	-8060	641			0.6	U			P11	1.5	75Th08
	-7045	225			-0.8	U			P12	2.5	75Th08
	-7515	117			0.4	U			P13	1.0	75Th08
$^{27}\text{Na}-^{30}\text{Na}_{360} \quad ^{25}\text{Na}_{648}$	-2750	213	-2505	4	0.8	U			P10	1.5	75Th08
$^{26}\text{Mg}(^{18}\text{O}, ^{14}\text{O})^{30}\text{Mg}$	-16234	55	-16121	3	2.0	U			Mun		78Pa12 *
$^{30}\text{Si}(n, \alpha)^{27}\text{Mg}$	-4193	21	-4199.95	0.05	-0.3	U			Ham		62An05
$^{30}\text{Si}(p, \alpha)^{27}\text{Al}$	-2368	10	-2372.04	0.05	-0.4	U			MIT		64Sp12
$^{27}\text{Al}(\alpha, p)^{30}\text{Si}$	2375	8	2372.04	0.05	-0.4	U			Man		59Ba13 Y
$^{30}\text{Si}(d, \alpha)^{28}\text{Al}$	3123	10	3128.49	0.08	0.5	U			MIT		64Sp12
$^{28}\text{Si}(^3\text{He}, n)^{30}\text{S}$	-573	15	-573.64	0.21	0.0	U			CIT		67Mi02
$^{30}\text{Ar}(2p)^{28}\text{S}$	2280	130				3					15Mu13 *
	$^{29}\text{Si}(n, \gamma)^{30}\text{Si}$	10609.6	0.3	10609.199	0.022	-1.3	o		MMn		80Is02 Z
		10609.21	0.04			-0.3	2		MMn		90Is02 Z
		10609.24	0.05			-0.8	2		ORn		92Ra19 Z
		10609.1776	0.0500			0.4	o		PTc		97Ro26 *
		10609.178	0.030			0.7	2		PTc		01Pa52 *
$^{29}\text{Si}(d, p)^{30}\text{Si}$	10609.23	0.21			-0.1	U			Bdn		06Fi.A
	8413	10	8384.633	0.022	-2.8	U			Mex		61Ja23
	8396	13			-0.9	U			MIT		64Sp12
	8384.92	0.53			-0.5	U			Rez		90Pi05 *
$^{29}\text{Si}(p, \gamma)^{30}\text{P}$	5594.5	0.4	5594.75	0.07	0.6	U					85Re02
	5594.5	0.5			0.5	U					96Wa33
$^{30}\text{Na}(\beta^-)^{30}\text{Mg}$	17167	330	17358	6	0.6	U					83De04 *
$^{30}\text{Mg}(\beta^-)^{30}\text{Al}$	6690	240	6981	4	1.2	U					83De04 *
$^{30}\text{Al}(\beta^-)^{30}\text{Si}$	8550	250	8568.1	2.9	0.1	U					61Ro12 *
$^{30}\text{Si}(t, ^3\text{He})^{30}\text{Al}$	-8520	40	-8549.5	2.9	-0.7	U					69Aj03
	-8545	15			-0.3	U					87Pe06
$^{30}\text{P}(\beta^+)^{30}\text{Si}$	4262	40	4232.11	0.06	-0.7	U					56Gr07
	4267	25			-1.4	U					63Fr10
$^{30}\text{Si}(p, n)^{30}\text{P}$	-5012.1	5.	-5014.45	0.06	-0.5	U			Har		75Fr.A Z
$^{30}\text{S}(\beta^+)^{30}\text{P}$	6118	22	6141.60	0.20	1.1	U					63Fr10 *
$^{30}\text{Na}-u$	F : contaminated by $^{30}\text{Mg}$										91Zh24 **
$^{26}\text{Mg}(^{18}\text{O}, ^{14}\text{O})^{30}\text{Mg}$	Tentative, say authors; four counts only										AHW **
$^{30}\text{Ar}(2p)^{28}\text{S}$	Symmetrized from 2250(+150-100)										GAu **
$^{29}\text{Si}(n, \gamma)^{30}\text{Si}$	Original error 0.0005 increased for calibration										GAu **
$^{29}\text{Si}(n, \gamma)^{30}\text{Si}$	Original error 0.005 increased for calibration										GAu **
$^{29}\text{Si}(d, p)^{30}\text{Si}$	Estimated systematic error 0.5 added to statistical error 0.16 keV										AHW **
$^{30}\text{Na}(\beta^-)^{30}\text{Mg}$	Calculated from 3 values used for calibration										GAu **
$^{30}\text{Mg}(\beta^-)^{30}\text{Al}$	Calculated from value used for calibration										GAu **
$^{30}\text{Al}(\beta^-)^{30}\text{Si}$	$E_{\beta^-} = 5050(250)$ to $2^+$ level at 3498.49 keV										Ens109 **
$^{30}\text{S}(\beta^+)^{30}\text{P}$	$E_{\beta^+} = 4422(22)$ to $0^+$ level at 677.01 keV										Ens109 **
$^{31}\text{Ne}-u$	33087	1739	33470	290	0.1	U					GA7 1.5 07Ju03
	33752	333			-0.6	1	33	$^{33} \text{ } ^{31}\text{Ne}$	GA8	1.5	12Ga45
$^{31}\text{Na}-u$	13440	1000	13147	15	-0.2	o			GA1	1.5	87Gi05
	13559	327			-0.8	o			GA3	1.5	91Or01
	13610	210			-1.5	U			TO4	1.5	91Zh24
	13441	118			-1.7	U			GA7	1.5	07Ju03



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{31}\text{Mg}-\text{O}_{1,938}$	6503.7	3.3				2			TT1	1.0	13Ch49
$^{31}\text{Mg}-\text{u}$	-3830	220	-3352	3	1.4	o			TO1	1.5	86Vi09
	-3520	180			0.6	o			GA1	1.5	87Gi05
	-3458	149			0.5	U			GA3	1.5	91Or01
	-3370	120			0.1	U			TO4	1.5	91Zh24
	-3425	18			4.1	B			P40	1.0	06Ga04
$\text{O}_2-^{31}\text{P H}$	8242.20819	0.00086	8242.2083	0.0007	0.2	1	67	61 $^{31}\text{P}$	FS1	1.0	08Re16
$^{31}\text{Na}-^{39}\text{K}_{,795}$	42017	25	42000	15	-0.7	o			TT1	1.0	13Ch49
	42000	15				2			TT1	1.0	16Ga.1
$^{31}\text{Al}-^{39}\text{K}_{,795}$	12803.1	2.4				2			TT1	1.0	16Kw.A
$^{31}\text{P}-^{28}\text{Si H}_3$	-26639.6290	0.0056	-26639.6331	0.0007	-0.7	U			FS1	1.0	06Re19
	-26639.63324	0.00089			0.2	1	64	39 $^{31}\text{P}$	FS1	1.0	08Re16
$^{31}\text{S}-^{31}\text{P}$	5794.98	0.25	5795.01	0.25	0.1	1	97	97 $^{31}\text{S}$	JY1	1.0	10Ka30
$^{31}\text{Cl}-^{31}\text{P}$	18686.1	3.7				2			JY1	1.0	16Ka15
$\text{O}_2-^{31}\text{P}$	16067.228	0.096	16067.2406	0.0007	0.1	U			MS1	1.0	09Kw02 *
$^{26}\text{Na}-^{31}\text{Na}_{,373} \ ^{22}\text{Na}_{,657}$	-7457	286	-8024	6	-0.8	U			P12	2.5	75Th08
$^{18}\text{O}(^{15}\text{N},2\text{p})^{31}\text{Al}$	-170	90	-308.6	2.2	-1.5	U					81Pa11
$^{27}\text{Al}(\alpha,\gamma)^{31}\text{P}$	9667.4	1.3	9668.60	0.05	0.9	U			Utr		78Ma23
$^{31}\text{P}(\text{p},\alpha)^{28}\text{Si}$	1912	5	1916.3084	0.0007	0.9	U			Bar		56Va14 Y
	1919	4			-0.7	U			VUn		64Sm03
	1911	10			0.5	U			MIT		64Sp12
	1915.8	0.2			2.5	U			Zur		67St30
$^{28}\text{Si}(\alpha,\text{n})^{31}\text{S}$	-8135	44	-8096.67	0.23	0.9	U			Tal		63Ne05
$^{31}\text{P}(\text{d},\alpha)^{29}\text{Si}$	8166	11	8165.3436	0.0007	-0.1	U			MIT		64Sp12
$^{31}\text{Cl}^i(2\text{p})^{29}\text{P}$	7700	100	7631	3	-0.7	U					90Bo24 *
	7610	60			0.3	U			Lis		91Bo32
	7643	50			-0.2	U			Lis		92Ba01 *
	7627	15			0.3	o					98Ax02
	7631	3				2					00Fy01 *
$^{30}\text{Ne}(\text{n},\gamma)^{31}\text{Ne}$	190	130	170	130	-0.2	1	95	67 $^{31}\text{Ne}$			14Na10 *
$^{30}\text{Si}(^{18}\text{O},^{17}\text{F})^{31}\text{Al}$	-12200	25	-12216.8	2.2	-0.7	U					88Wo02
	-12237	35			0.6	U			Ber		89Bo.A
$^{30}\text{Si}(\text{n},\gamma)^{31}\text{Si}$	6589.1	0.7	6587.39	0.04	-2.4	U					70Be48
	6587.5	0.8			-0.1	U					70Sp02
	6588.4	0.3			-3.4	B					72Dz13
	6587.32	0.20			0.4	U			MMn		90Is02 Z
	6587.39	0.05			0.1	3			ORn		92Ra19 Z
	6587.3970	0.0500			-0.1	o			PTc		97Ro26 *
	6587.39	0.14			0.0	U			Bdn		06Fi.A
	6587.397	0.057			-0.1	3			PTc		01Pa52
$^{30}\text{Si}(\text{d},\text{p})^{31}\text{Si}$	4368	7	4362.83	0.04	-0.7	U			MIT		64Sp12
	4364.18	0.55			-2.5	U			Rez		90Pi05 *
$^{30}\text{Si}(\text{p},\gamma)^{31}\text{P}$	7297.4	1.2	7296.551	0.022	-0.7	U					68Wo01
$^{31}\text{Cl}^i(\text{p})^{30}\text{S}$	12033	10	12026	3	-0.7	o					98Ax02 *
	12033	14			-0.5	U					00Fy01 *
$^{31}\text{Mg}(\beta^-)^{31}\text{Al}$	10150	700	11829	4	2.4	U					83De04
$^{31}\text{Al}(\beta^-)^{31}\text{Si}$	7940	100	7998.3	2.2	0.6	U					73Go22
$^{31}\text{Si}(\beta^-)^{31}\text{P}$	1471	8	1491.50	0.04	2.6	U					52Mo12
	1486	12			0.5	U					52Wa12
$^{31}\text{S}(\beta^+)^{31}\text{P}$	5412	30	5398.02	0.23	-0.5	U					60Wa04
$^{31}\text{P}(\text{p},\text{n})^{31}\text{S}$	-6212.3	20.	-6180.36	0.23	1.6	o			ChR		58Go77 Y
	-6250	20			3.5	B			ChR		59Br06 Y
* $\text{O}_2-^{31}\text{P}$	For original doublet $^{31}\text{P}-\text{O}_{1,938}$ , $D_M=-16382.522(0.096) \mu\text{u}$										
* $^{31}\text{Cl}^i(2\text{p})^{29}\text{P}$	Large error in $E_{cm}$ due to sequential decay kinematics										
* $^{31}\text{Cl}^i(2\text{p})^{29}\text{P}$	reference also finds 3p emission at 4870										
* $^{31}\text{Cl}^i(2\text{p})^{29}\text{P}$	$Q_{2p}=7620(5), 6245(2), 5679(3), 5223(5) \text{ keV}$										
*	to ground state and levels $3/2^+$ at 1383.55, $5/2^+$ at 1953.91, $3/2^+$ 2422.7 keV										
* $^{30}\text{Ne}(\text{n},\gamma)^{31}\text{Ne}$	Symmetrized from 150(+160-100) keV										
* $^{30}\text{Si}(\text{n},\gamma)^{31}\text{Si}$	Original error 0.0005 increased for calibration										
* $^{30}\text{Si}(\text{d},\text{p})^{31}\text{Si}$	Estimated systematic error 0.5 added to statistical error 0.23 keV										
* $^{31}\text{Cl}^i(\text{p})^{30}\text{S}$	Average of 3 branches										
* $^{31}\text{Cl}^i(\text{p})^{30}\text{S}$	$E_p=11654(28), 9493(20), 8347(15), 8092(14) \text{ keV}$										

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{32}\text{Na}-u$	19720	636	20010	40	0.3	o			GA3	1.5	91Or01
	19900	1100			0.1	U			TO4	1.5	91Zh24
	20980	500			-1.3	o			GA5	1.5	00Sa21
	20193	129			-0.9	U			GA7	1.5	07Ju03
$^{32}\text{Mg}-\text{O}_2$	9280.9	3.5				2			TT1	1.0	13Ch49
$^{32}\text{Mg}-u$	-800	260	-890	4	-0.2	o			TO1	1.5	86Vi09
	-890	270			0.0	U			GA1	1.5	87Gi05
	-924	214			0.1	U			GA3	1.5	91Or01
	-820	130			-0.4	U			TO4	1.5	91Zh24
	-1142	113			2.2	o			P40	1.0	01Lu20
	-966	38			2.0	U			P40	1.0	06Ga04
	-983	22			4.2	B			P40	1.0	06Ga04
$^{32}\text{Al}-\text{O}_2$	-1744	13	-1745	8	-0.1	o			TT1	1.0	12Ch.A
	-1744.9	7.7				2			TT1	1.0	16Kw.A
$^{32}\text{Al}-u$	-12160	220	-11916	8	0.7	U			TO1	1.5	86Vi09
	-11870	200			-0.2	U			GA1	1.5	87Gi05
	-11877	104			-0.2	U			GA3	1.5	91Or01
$^{32}\text{Si O}_2-\text{C}_5\text{H}_4$	-67319.35	0.32				2			MS1	1.0	09Kw02
$\text{O}_2-^{32}\text{S}$	17754.2	1.0	17758.0648	0.0014	1.5	U			J1	2.5	68Ma45
$\text{C}_2\text{H}_8-^{32}\text{S}$	90531.3	1.4	90529.0835	0.0016	-0.6	U			J1	2.5	68Ma45
$^{32}\text{S}-\text{O}_2$	-17758.0663	0.0020	-17758.0648	0.0014	0.8	1	50	48 $^{32}\text{S}$	FS1	1.0	05Sh38
$^{32}\text{S}-\text{C}_2\text{D}_4$	-84335.9367	0.0019	-84335.9380	0.0014	-0.7	1	55	52 $^{32}\text{S}$	FS1	1.0	05Sh38
$^{32}\text{S}-\text{H C F}$	-34156.50	0.57	-34157.0207	0.0016	-0.9	U			MS1	1.0	09Kw02
$^{32}\text{Na}-^{39}\text{K}_{.821}$	49808	40				2			TT1	1.0	16Ga.1
$^{32}\text{Ar}-^{39}\text{K}_{.821}$	27434.8	1.9				2			MA8	1.0	03B117
$\text{C F}_3-^{32}\text{S O}_2\text{ H}$	25483.43	0.34	25484.043	0.003	1.8	U			MS1	1.0	09Kw02
$\text{C F}_3-^{32}\text{S O}_2$	33310.02	0.59	33309.075	0.003	-1.6	U			MS1	1.0	09Kw02
$^{26}\text{Na}-^{32}\text{Na}_{.325}\text{ }^{22}\text{Na}_{.709}$	-8569	354	-9245	13	-0.8	U			P12	2.5	75Th08
$^{32}\text{S}(^3\text{He},^8\text{Li})^{27}\text{P}$	-31277	35	-31308	26	-0.9	2			MSU		77Be13
$^{32}\text{S}(p,\alpha)^{29}\text{P}$	-4171	20	-4198.6	0.4	-1.4	U			Tky		64Ej05
$^{32}\text{S}(^3\text{He},^6\text{He})^{29}\text{S}$	-25520	50				2			MSU		73Be09
$^{30}\text{Si}(t,p)^{32}\text{Si}$	7307	1	7305.56	0.30	-1.4	U			Str		80An.A
$^{32}\text{S}(d,\alpha)^{30}\text{P}$	4892	10	4896.13	0.07	0.4	U			MIT		64Sp12
$^{32}\text{S}(p,t)^{30}\text{S}$	-19614	3	-19617.12	0.21	-1.0	U			MSU		74Ha02
$^{31}\text{Si}(n,\gamma)^{32}\text{Si}$	9203.2180	0.0500	9200.0	0.3	-65.0	B			PTc		97Ro26
	9203.22	0.76			-4.3	B			PTc		01Pa52
$^{31}\text{P}(n,\gamma)^{32}\text{P}$	7935.73	0.16	7935.65	0.04	-0.5	U			MMn		85Ke11
	7935.65	0.04				2			ILn		89Mi16
	7935.60	0.16			0.3	U			Bdn		06Fi.A
$^{31}\text{P}(d,p)^{32}\text{P}$	5712	8	5711.08	0.04	-0.1	U			MIT		64Sp12
$^{31}\text{P}(p,\gamma)^{32}\text{S}$	8864.9	0.9	8863.9632	0.0015	-1.0	U					72Co13
	8862.7	3.			0.4	U					73Ve06
	8865.6	1.0			-1.6	U					73Ve08
	8865.1	0.9			-1.3	U					74Vi02
$^{31}\text{P}(^3\text{He},d)^{32}\text{S}$	3356	13	3370.4888	0.0015	1.1	U			MIT		68Gr17
$^{32}\text{S}(p,d)^{31}\text{S}$	-12817.8	1.5	-12819.76	0.23	-1.3	U			MSU		73Mo23
$^{32}\text{S}(^3\text{He},\alpha)^{31}\text{S}$	5415	15	5533.29	0.23	7.9	B					66Gr26
	5515	15			1.2	U			MIT		67Sp09
	5486	20			2.4	U			Ors		67Ro17
	5538	6			-0.8	U			CIT		70Mo08
	-1583.5	3.1	-1581.1	0.5	0.8	U					85Bj01
$^{32}\text{Cl}(p)^{31}\text{S}$	-1581.9	2.1			0.4	U					93Sc16
	-1581.3	0.6			0.3	1	79	76 $^{32}\text{Cl}$			08Ga.A
	18300	1400	19470	40	0.8	U					83De04
$^{32}\text{Si}(\beta^-)^{32}\text{P}$	213	7	227.2	0.3	2.0	U					64Br09
	221.4	1.2			4.8	B					84Po09

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{32}\text{P}(\beta^-)^{32}\text{S}$	1707.6	0.7	1710.66	0.04	4.4	B					61Ni02	
	1710.1	0.7			0.8	U					68Fi04	
$^{32}\text{Cl}(\beta^+)^{32}\text{S}$	12720	30	12680.9	0.6	-1.3	U					68Ar03 *	
$^{32}\text{S}(\text{p,n})^{32}\text{Cl}$	-13470	14	-13463.2	0.6	0.5	U			Yal		69Ov01 Z	
	-13470	9			0.8	U			BNL		71Go18 Z	
$^{32}\text{S}(^3\text{He,t})^{32}\text{Cl}$	-12699	15	-12699.5	0.6	0.0	U					89Je07	
$^{32}\text{S}(^3\text{He,t})^{32}\text{Cl}-^{36}\text{Ar}(^{36}\text{K})$	133.01	1.10	133.6	0.6	0.6	1	31	24	$^{32}\text{Cl}$	Mun	10Wr01	
$^{32}\text{S}(\pi^+,\pi^-)^{32}\text{Ar}$	-22813	50	-22793.2	1.8	0.4	U					80Bu15	
$^{32}\text{Mg-u}$	Result from the "Plasma" experiment. Next item from "Rilis"										06Ga04 **	
$^{32}\text{Si O}_2-\text{C}_5\text{H}_4$	For original doublet $^{32}\text{Si O}_2\text{H}_3-\text{C}_5\text{H}_7$										GAu **	
$^{32}\text{S}-\text{H C F}$	For original doublet $^{32}\text{S O}_2\text{H}-\text{H}_2\text{C O}_2\text{F}$										GAu **	
$^*\text{C F}_3-^{32}\text{S O}_2\text{H}$	For original doublet $^{32}\text{S O}_2\text{H}-(\text{C F}_3)_{0.942}$ , $D_M=-25761.27(0.34)\mu\text{u}$										GAu **	
$^*\text{C F}_3-^{32}\text{S O}_2$	For original doublet $^{32}\text{S O}_2-(\text{C F}_3)_{0.928}$ , $D_M=-33654.93(0.59)\mu\text{u}$										GAu **	
$^{31}\text{Si}(\text{n},\gamma)^{32}\text{Si}$	Original error 0.0005 increased for calibration										GAu **	
$^{31}\text{P}(\text{p},\gamma)^{32}\text{S}$	$T=3289(3) Q=-3185.3(3.)$ to $^{32}\text{S}^j$ at 12047.96(0.28) keV										Nub16b **	
$^{32}\text{Cl}(\text{p})^{31}\text{S}$	$E_p=3353.5(3.0) Q_p=3462.8(3.1)$ from $^{32}\text{Cl}^i$ 5046.3(0.3) T=2 level										Nub16b **	
$^{32}\text{Cl}(\text{p})^{31}\text{S}$	$E_p=3348.5(2.0) Q_p=3457.6(2.1)$ from $^{32}\text{Cl}^i$ 5046.3(0.3) T=2 level										Nub16b **	
*	corrected to 3464.4(2.1) keV										02Py02 **	
$^{32}\text{Cl}(\text{p})^{31}\text{S}$	$Q_p=3465.0(0.4)$ from $^{32}\text{Cl}^i$ 5046.3(0.3) T=2 level										Nub16b **	
*	this $Q_p$ is quoted in reference as "A.Garcia et al. (in preparation)"										08Bh08 **	
$^{32}\text{Cl}(\beta^+)^{32}\text{S}$	$E_{\beta^+}=9470(30)$ to $2^+$ level at 2230.57 keV										Ens119 **	
$^{33}\text{Na-u}$	27386	1601	25530	480	-0.8	o			GA3	1.5	91Or01	
	26370	1160			-0.5	o			GA5	1.5	00Sa21	
	25142	376			0.7	o			GA7	1.5	07Ju03	
	25529	322				2			GA8	1.5	12Ga45	
$^{33}\text{Mg}-\text{O}_{2.062}$	15813.3	3.1				2			TT1	1.0	13Ch49	
$^{33}\text{Mg-u}$	5460	900	5327	3	-0.1	o			GA1	1.5	87Gi05	
	5203	318			0.3	U			GA3	1.5	91Or01	
	5710	180			-1.4	U			TO4	1.5	91Zh24	
	5311	24			0.7	U			P40	1.0	06Ga04	
$^{33}\text{Al-u}$	-9490	250	-9122	8	1.0	o			TO1	1.5	86Vi09	
	-9250	160			0.5	o			GA1	1.5	87Gi05	
	-9167	142			0.2	U			GA3	1.5	91Or01	
	-9020	120			-0.6	U			TO4	1.5	91Zh24	
	-9125	64			0.0	o			GT1	1.5	04Ma.A	
	-8957	100			-0.7	o			GT2	2.5	08Kn.A	
	-8915	128			-0.6	U			GT2	2.5	08Su19	
	-9209	62			1.4	U			LZ1	1.0	15Xu14	
						2			MS1	1.0	09Kw02 *	
$^{33}\text{Si O}_2-^{13}\text{C C}_4\text{H}_4$	-66848.76	0.75				2			LZ1	1.0	11Tu09	
$^{33}\text{Cl-u}$	-22536.9	7.5	-22548.0	0.4	-1.5	U			TT1	1.0	16Kw.A	
$^{33}\text{Al}-^{39}\text{K}_{.846}$	21582.0	7.5				2			MA8	1.0	03B117	
$^{33}\text{Ar}-^{39}\text{K}_{.846}$	20629.86	0.43				2			MA6	1.0	01He29	
$^{33}\text{Ar}-^{36}\text{Ar}_{.917}$	19689.2	4.5	19686.7	0.4	-0.6	U			MI3	1.0	05Ra34	
$^{33}\text{S}-^{32}\text{S H}$	-8437.29682	0.00030	-8437.2968	0.0003	0.0	1	100	100	$^{33}\text{S}$		ANL	68Be13
$^{30}\text{Si}(\alpha,\text{p})^{33}\text{P}$	-2965	10	-2959.7	1.1	0.5	U					ILL	81Wa31
$^{33}\text{S}(\text{n},\alpha)^{30}\text{Si}$	3497.6	5.	3493.507	0.022	-0.8	U					01Wa50	
	3496.9	5.0			-0.7	U					71Gr04	
$^{31}\text{P}(^3\text{He,p})^{33}\text{S}$	9787	15	9787.5607	0.0015	0.0	U					MMn	80Is02 Z
$^{32}\text{S}(\text{n},\gamma)^{33}\text{S}$	8641.5	0.3	8641.6379	0.0006	0.5	o					ORn	83Ra04 Z
	8641.82	0.10			-1.8	U					MMn	85Ke08 Z
	8641.60	0.03			1.3	U					Bdn	06Fi.A
	8641.81	0.17			-1.0	U					NBS	06De21
	8641.6398	0.0033			-0.6	U					MIT	64Sp12
$^{32}\text{S}(\text{d,p})^{33}\text{S}$	6420	6	6417.0719	0.0003	-0.5	U					59Ku79	
$^{32}\text{S}(\text{p},\gamma)^{33}\text{Cl}$	2276.4	0.9	2276.8	0.4	0.4	-					76Al01	
	2276.8	0.5			-0.1	-						

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{32}\text{S}(\text{d,n})^{33}\text{Cl}$	62	9	52.2	0.4	-1.1	U					72El03
$^{32}\text{S}(^3\text{He,d})^{33}\text{Cl}$	-3218	15	-3216.7	0.4	0.1	U					66Gr26
	-3217	5			0.1	U			CIT		70Mo08
$^{32}\text{S}(\text{p},\gamma)^{33}\text{Cl}$	ave. 2276.7	0.4	2276.8	0.4	0.2	1	80	80 $^{33}\text{Cl}$			average
$^{32}\text{S}(\text{p},\gamma)^{33}\text{Cl}^i$	-3267.0	1.0	-3271.7	0.5	-4.7	B					70Ab15
	-3271.4	2.0			-0.1	U					82Wi.A
	-3271.5	0.8			-0.3	1	37	37 $^{33}\text{Cl}^i$			02Py01
$^{33}\text{Cl}^i(\text{p})^{32}\text{S}$	3266.8	1.0	3271.7	0.5	4.9	B					87Bo21
$^{33}\text{Si}(\beta^-)^{33}\text{P}$	5768	50	5823.0	1.3	1.1	U					73Go33
$^{33}\text{P}(\beta^-)^{33}\text{S}$	249	2	248.5	1.1	-0.2	2					54Ni06
	248.3	1.3			0.2	2					84Po09
$^{33}\text{Cl}(\beta^+)^{33}\text{S}$	5532	50	5582.5	0.4	1.0	U					60Wa04
$^{33}\text{Cl}^i(\text{IT})^{33}\text{Cl}$	5548.5	0.4	5548.4	0.4	-0.1	1	83	63 $^{33}\text{Cl}^i$			06Tr10
$^{*33}\text{Si O}_2 - ^{13}\text{C C}_4 \text{H}_4$	For original doublet $^{33}\text{Si O}_2 \text{H}_3 - ^{13}\text{C C}_4 \text{H}_7$										GAU **
$^{34}\text{Na}-\text{u}$	34010	429				2			GA8	1.5	12Ga45
$^{34}\text{Mg}-\text{O}_{2.126}$	19747	31				2			TT1	1.0	13Ch49
$^{34}\text{Mg}-\text{u}$	8855	476	8940	30	0.1	o			GA3	1.5	91Or01
	9190	350			-0.5	U			TO4	1.5	91Zh24
	9900	350			-1.8	U			GA5	1.5	00Sa21
	9190	97			-1.7	U			GA7	1.5	07Ju03
$^{34}\text{Al}-\text{u}$	-3760	430	-3221	3	0.8	o			TO1	1.5	86Vi09 *
	-3400	250			0.5	o			GA1	1.5	87Gi05 *
	-3262	218			0.1	o			GA3	1.5	91Or01 *
	-2940	120			-1.6	U			TO4	1.5	91Zh24 *
	-3199	97			-0.2	U			GT1	1.5	04Ma.A *
	-3328	86			0.8	U			GA7	1.5	07Ju03 *
$^{34}\text{Al}-^{39}\text{K}_{.872}$	28438.0	7.8	28427	3	-1.4	o			TT1	1.0	16Kw.A
	28427.0	3.3				2			TT1	1.0	16Ga.1
$^{34}\text{Ar}-^{39}\text{K}_{.872}$	11919.02	0.36	11918.04	0.08	-2.7	U			MA8	1.0	02He23
$^{34}\text{Ar}-^{36}\text{Ar}_{.944}$	10907.4	3.8	10907.51	0.09	0.0	U			MA6	1.0	01He29
$^{34}\text{Cl}-^{34}\text{S}$	5895.548	0.058	5895.48	0.04	-1.2	1	49	31 $^{34}\text{Cl}$	JY1	1.0	09Er07
$^{34}\text{Cl}^m-^{34}\text{S}$	6052.575	0.068	6052.60	0.04	0.4	1	41	31 $^{34}\text{Cl}^m$	JY1	1.0	09Er07
$^{34}\text{S}-^{34}\text{Ar}$	-12403.19	0.20	-12403.08	0.08	0.5	1	14	13 $^{34}\text{Ar}$	JY1	1.0	11Er02
$^{34}\text{Cl}^m-^{34}\text{Cl}$	157.05	0.11	157.124	0.029	0.7	U			JY1	1.0	09Er07
	157.30	0.27			-0.7	U			JY1	1.0	11Er02
$^{34}\text{Ar}-^{34}\text{Cl}$	6507.627	0.092	6507.60	0.07	-0.3	1	54	52 $^{34}\text{Ar}$	JY1	1.0	11Er02
$^{34}\text{Cl}^m-^{34}\text{Ar}$	-6350.41	0.11	-6350.48	0.07	-0.6	1	39	35 $^{34}\text{Ar}$	JY1	1.0	11Er02
$\text{C}_4 \text{H}_3 - ^{34}\text{P O}$	54914.59	0.87				2			MS1	1.0	09Kw02 *
$^{30}\text{Si}(^7\text{Li}, ^3\text{He})^{34}\text{P}$	100	40	91.6	0.8	-0.2	U					77Pe17
$^{31}\text{P}(\alpha,\text{p})^{34}\text{S}$	629.9	2.9	627.09	0.04	-1.0	U			Har		73Ry01
$^{31}\text{P}(\alpha,\text{n})^{34}\text{Cl}$	-5632	10	-5646.86	0.05	-1.5	U			Tal		70Um01
	-5641.5	3.7			-1.4	U			Har		73Ry01
$^{34}\text{S}(\text{d},\alpha)^{32}\text{P}$	5096	10	5083.99	0.06	-1.2	U					78Ba30
$^{32}\text{S}(^3\text{He},\text{n})^{34}\text{Ar}$	-759	15	-777.34	0.08	-1.2	U			CIT		67Mi02
$^{34}\text{S}(^{13}\text{C}, ^{14}\text{O})^{33}\text{Si}$	-14243	75	-14300.1	0.7	-0.8	U			Can		86Fi06
$^{33}\text{S}(\text{n},\gamma)^{34}\text{S}$	11417.12	0.10	11417.15	0.04	0.3	-			ORn		83Ra04 Z
	11417.22	0.23			-0.3	-			Bdn		06Fi.A
$^{33}\text{S}(\text{d},\text{p})^{34}\text{S}$	9202	10	9192.58	0.04	-0.9	U			MIT		64Sp12
	9195	6			-0.4	U			Utr		71Va21
$^{33}\text{S}(\text{n},\gamma)^{34}\text{S}$	ave. 11417.14	0.09	11417.15	0.04	0.1	1	24	24 $^{34}\text{S}$			average
$^{33}\text{S}(\text{p},\gamma)^{34}\text{Cl}$	5142.42	0.20	5143.20	0.05	3.9	B			Oak		83Ra04 *
	5142.4	0.3			2.7	U			Utr		83Wa27 Z
	5143.29	0.07			-1.3	1	48	48 $^{34}\text{Cl}$	Auc		94Li20
$^{34}\text{Si}(\beta^-)^{34}\text{P}$	4700	300	4592	14	-0.4	U					77Na05

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{34}\text{P}(\beta^-)^{34}\text{S}$	5383	45	5383.0	0.8	0.0	U			ANB		73Go33
$^{34}\text{S}(\text{t},^3\text{He})^{34}\text{P}$	-5368	20	-5364.4	0.8	0.2	U			LAl		77Aj01
$^{34}\text{S}(^7\text{Li},^7\text{Be})^{34}\text{P}$	-6224	40	-6244.9	0.8	-0.5	U			Can		85Dr06
$^{34}\text{Cl}(\beta^+)^{34}\text{S}$	5522	30	5491.60	0.04	-1.0	U					56Gr07
$^{34}\text{S}(\text{p},\text{n})^{34}\text{Cl}$	-6252	10	-6273.95	0.04	-2.2	U			Tal		70Um01
	-6271.9	1.9			-1.1	U			Har		75Fr.A
	-6274.27	0.56			0.6	U			Auc		77Ba16
	-6273.11	0.25			-3.4	F			Auc		92Ba.A *
$^{34}\text{S}(^3\text{He},\text{t})^{34}\text{Cl}$	-5510.8	0.4	-5510.20	0.04	1.5	F			Mun		77Vo02 *
$^{34}\text{S}(^3\text{He},\text{t})^{34}\text{Cl}-^{27}\text{Al}(^{27}\text{Si})$	-678.7	2.3	-679.24	0.10	-0.2	U			ChR		74Ha35
$^{34}\text{Cl}^m(\text{IT})^{34}\text{Cl}$	146.36	0.03	146.360	0.027	0.0	1	84	65 $^{34}\text{Cl}^m$			Ens126
* $^{34}\text{Al}-\text{u}$	Possible isomeric mixture 26(1) ms, E=550# keV										
* $\text{C}_4\text{H}_3-^{34}\text{P O}$	For original doublet $^{34}\text{P H}_2\text{O}-\text{C}_4\text{H}_5$										
* $^{33}\text{S}(\text{p},\gamma)^{34}\text{Cl}$	$E_p=974.76(0.15,Z)$ to $6088.20(0.10,Z)$ level										
* $^{34}\text{S}(\text{p},\text{n})^{34}\text{Cl}$	F : disturbed by resonance; at least 0.5 keV uncertain										
* $^{34}\text{S}(^3\text{He},\text{t})^{34}\text{Cl}$	F : rejected in reference of same group										
$^{35}\text{Mg}-\text{u}$	18669	1721	16790	290	-0.7	o			GA3	1.5	91Or01
	18830	1070			-1.3	o			GA5	1.5	00Sa21
	16790	193				2			GA7	1.5	07Ju03
$^{35}\text{Al}-\text{u}$	-340	460	-240	8	0.1	o			GA1	1.5	87Gi05
	-296	298			0.1	o			GA3	1.5	91Or01
	80	190			-1.1	U			TO4	1.5	91Zh24
	-236	75			0.0	U			GA7	1.5	07Ju03
$\text{C}_3-^{35}\text{Cl H}$	23320.8	0.3	23322.27	0.04	2.0	U			M17	2.5	66Be10
	23322.239	0.034			0.7	1	56	56 $^{35}\text{Cl}$	B07	1.5	71Sm01
	23321.83	0.63			0.3	U			J5	2.5	72Ka57
	23322.328	0.325			-0.1	U			J5	2.5	72Ka57
$\text{C}_5\text{H}_{10}-^{35}\text{Cl}_2$	140549.37	2.98	140544.93	0.08	-0.6	U			C2	2.5	65De09
	140545.01	0.13			-0.4	1	15	15 $^{35}\text{Cl}$	B07	1.5	71Sm01
$\text{C}_4\text{H}_6\text{O}-^{35}\text{Cl}_2$	104153.75	3.45	104159.42	0.08	0.7	U			C2	2.5	65De09
$\text{C}_2\text{D}_6-^{35}\text{Cl H}$	107934.90	0.54	107932.94	0.04	-1.5	U			J5	2.5	72Ka57
	107933.422	0.538			-0.4	U			J5	2.5	72Ka57
$\text{C}_3\text{H}-\text{D }^{35}\text{Cl}$	24871.92	0.75	24870.56	0.04	-0.7	U			C2	2.5	65De09
$\text{C}_8\text{H}_9-^{35}\text{Cl}_3$	163867.25	0.90	163867.21	0.11	0.0	U			A2	2.5	70St25
$^{35}\text{Ar}-\text{u}$	-24747.3	4.3	-24742.3	0.7	1.2	U			LZ1	1.0	11Tu09
$^{35}\text{Al}-^{39}\text{K}_{897}$	32315.1	7.9				2			TT1	1.0	16Ga.1
$^{35}\text{K}-^{39}\text{K}_{897}$	20560.69	0.55				2			MA8	1.0	07Ya08
$^{35}\text{Cl}(\text{p},\alpha)^{32}\text{S}$	1862	5	1866.06	0.04	0.8	U			Bar		57Va03 Y
	1865	8			0.1	U			MIT		64Sp12
$^{32}\text{S}(\alpha,\text{p})^{35}\text{Cl}$	-1862	17	-1866.06	0.04	-0.2	U			MIT		64Sp12
$^{32}\text{S}(\alpha,\text{n})^{35}\text{Ar}$	-8751	18	-8614.7	0.7	7.6	B			Tal		63Ne05
$^{35}\text{Cl}(\text{d},\alpha)^{33}\text{S}$	8285	10	8283.13	0.04	-0.2	U			MIT		64Sp12
$^{33}\text{S}(^3\text{He},\text{n})^{35}\text{Ar}$	3335	16	3321.3	0.7	-0.9	U					75Da14
$^{35}\text{K}^i(2\text{p})^{33}\text{Cl}$	4311	40				2					85Ay01
$^{34}\text{S}(^{18}\text{O},^{17}\text{F})^{35}\text{P}$	-7796	40	-7808.4	1.9	-0.3	U			Can		88Or01
$^{34}\text{S}(\text{n},\gamma)^{35}\text{S}$	6986.00	0.10	6985.84	0.04	-1.6	-			ORn		83Ra04 Z
	6985.84	0.05			0.0	-			MMn		85Ke08 Z
	6986.09	0.14			-1.8	U			Bdn		06Fi.A
$^{34}\text{S}(\text{d},\text{p})^{35}\text{S}$	4762	10	4761.27	0.04	-0.1	U			MIT		64Sp12
	4757	5			0.9	U			Utr		71Va18
$^{34}\text{S}(\text{n},\gamma)^{35}\text{S}$	ave.	6985.87	6985.84	0.04	-0.8	1	75	46 $^{34}\text{S}$			average
$^{34}\text{S}(\text{p},\gamma)^{35}\text{Cl}$	6367.4	1.6	6370.81	0.04	2.1	U					72Hu10
	6370.7	0.4			0.3	U					76Sp08 Z
	6370.70	0.20			0.6	U			Oak		83Ra04 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{35}\text{Cl}(\gamma,n)^{34}\text{Cl}$	-12660	40	-12644.76	0.05	0.4	U					61Sa11
$^{35}\text{P}(\beta^-)^{35}\text{S}$	3909	75	3988.4	1.9	1.1	U					72Go31
$^{35}\text{S}(\beta^-)^{35}\text{Cl}$	167.4	0.2	167.322	0.026	-0.4	U					57Co62
	166.80	0.15			3.5	B					85A111
	167.288	0.100			0.3	U					85Ap01 *
	166.93	0.2			2.0	o					85Ma59
	167.4	0.1			-0.8	U					85Oh06 *
	166.7	0.2			3.1	B					89Si04 *
	167.56	0.03			-7.9	B					92Ch27 *
	167.35	0.10			-0.3	U					93Ab11 *
	167.23	0.10			0.9	U					93Be21 *
	167.27	0.10			0.5	U					93Mo01 *
	167.334	0.027			-0.5	1	91	71 $^{35}\text{S}$			00Ho13
$^{35}\text{Cl}(n,p)^{35}\text{S}$	612	4	615.025	0.026	0.8	U			BNL		68Se01
$^{35}\text{Ar}(\beta^+)^{35}\text{Cl}$	5980	40	5966.2	0.7	-0.3	U					56Ki29
	5950	50			0.3	U					60Wa04
$^{35}\text{Cl}(p,n)^{35}\text{Ar}$	-6747.2	1.6	-6748.6	0.7	-0.9	2			Har		75Fr.A Z
	-6747.9	1.0			-0.7	2			Auc		77Wh03 Z
	-6750.4	1.2			1.5	2			Mtr		78Az01 *
$^{34}\text{S}(p,\gamma)^{35}\text{Cl}$	$E_p=1264.97(0.13,Z)$ to $7598.91(0.15,Z)$ level										
$^{35}\text{S}(\beta^-)^{35}\text{Cl}$	Original error (0.030) increased to 0.100										
$^{35}\text{Cl}(p,n)^{35}\text{Ar}$	Original T=6942.2(2.2) recalibrated 6945.5(1.2)										
											83Ra04 **
											AHW **
											GAU **
$^{36}\text{Mg}-u$	24930	1610	21880	740	-1.3	o					GA5 1.5 00Sa21
	21879	494				2			GA7 1.5 07Ju03		
$^{36}\text{Al}-u$	6187	421	6390	160	0.3	o					GA3 1.5 91Or01
	6500	400			-0.2	U					TO4 1.5 91Zh24
	6140	310			0.5	o					GA5 1.5 00Sa21
	6388	107				2					GA7 1.5 07Ju03
$^{36}\text{Si}-u$	-13850	640	-13350	80	0.5	U					TO1 1.5 86Vi09
	-13490	320			0.3	o					GA1 1.5 87Gi05
	-13578	191			0.8	o					GA3 1.5 91Or01
	-13110	150			-1.1	2					TO4 1.5 91Zh24
	-13376	75			0.2	2					GT1 1.5 04Ma.A
	-13280	118			-0.4	2					GA7 1.5 07Ju03
	-13484	163			0.8	2					LZ1 1.0 15Xu14
$^{36}\text{Ar}-u$	-32454.895	0.015	-32454.895	0.029	0.0	o					ST2 1.0 02Bf02
	-32454.895	0.029			0.0	1	100	100 $^{36}\text{Ar}$			ST2 1.0 03Fr08
$^{36}\text{K}-^{39}\text{K}_{923}$	14800.99	0.38	14800.9	0.4	-0.2	1	93	93 $^{36}\text{K}$			MA8 1.0 07Ya08
$^{36}\text{Ar}(^3\text{He}, ^8\text{Li})^{31}\text{Cl}$	-29180	50	-29212	3	-0.6	U					MSU 77Be13
$^{36}\text{S}(^{48}\text{Ca}, ^{52}\text{V})^{32}\text{Al}$	-12651	370	-12346	7	0.8	o					Dar 87Ch.A
$^{36}\text{S}(^{48}\text{Ca}, ^{51}\text{V})^{33}\text{Al}$	-14150	140	-14188	7	-0.3	U					Dar 86Wo07
$^{36}\text{S}(^{14}\text{C}, ^{17}\text{O})^{33}\text{Si}$	-6380	20	-6321.1	0.7	2.9	U					Mun 84Ma49
$^{36}\text{S}(^{11}\text{B}, ^{14}\text{N})^{33}\text{Si}$	-4311	30	-4345.5	0.7	-1.2	U					Can 85Fi03
$^{36}\text{Ar}(^3\text{He}, ^6\text{He})^{33}\text{Ar}$	-23512	30	-23508.1	0.4	0.1	U					MSU 74Na07
$^{36}\text{S}(^{11}\text{B}, ^{13}\text{N})^{34}\text{Si}$	-7327	25	-7385	14	-2.3	2					Can 85Fi03
$^{36}\text{S}(^{14}\text{C}, ^{16}\text{O})^{34}\text{Si}$	-2989	20	-2950	14	1.9	2					Mun 84Ma49
$^{36}\text{S}(^{64}\text{Ni}, ^{66}\text{Zn})^{34}\text{Si}$	-8890	41	-8907	14	-0.4	o					Dar 85Wo07 *
	-8903	33			-0.1	2					Dar 86Sm05 *
$^{36}\text{S}(d,\alpha)^{34}\text{P}$	4604.4	5.	4595.4	0.8	-1.8	U					82So.A *
$^{36}\text{Ar}(p,t)^{34}\text{Ar}$	-19513	3	-19514.09	0.08	-0.4	U					MSU 74Ha02
$^{36}\text{Ar}(p,t)^{34}\text{Ar}^i$	-27473	50	-27448	5	0.5	U					69Br21 *
	-27448	5				2					72Pa02 *
$^{36}\text{S}(^{14}\text{C}, ^{15}\text{O})^{35}\text{Si}$	-16184	50	-16110	40	1.5	2					Mun 84Ma49
$^{36}\text{S}(^{13}\text{C}, ^{14}\text{O})^{35}\text{Si}$	-21122	60	-21160	40	-0.6	2					Can 86Fi06
$^{36}\text{S}(^{64}\text{Ni}, ^{65}\text{Zn})^{35}\text{Si}$	-17250	100	-17460	40	-2.1	2					Dar 86Sm05 *
$^{36}\text{S}(d, ^3\text{He})^{35}\text{P}$	-7607	5	-7601.8	1.9	1.0	2					BNL 84Th08
	-7601	2			-0.4	2					Hei 85Kh04

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{36}\text{S}(^{14}\text{C}, ^{15}\text{N})^{35}\text{P}$	-2927	10	-2887.9	1.9	3.9	B			Mun	84Ma49	*	
$^{36}\text{S}(^6\text{Li}, ^7\text{Be})^{35}\text{P}$	-7521	17	-7488.5	1.9	1.9	U			Can	85Dr06		
$^{36}\text{S}(^{64}\text{Ni}, ^{65}\text{Cu})^{35}\text{P}$	-5659	34	-5641.6	2.0	0.5	U			Dar	85Wo.A		
$^{35}\text{Cl}(n, \gamma)^{36}\text{Cl}$	8579.73	0.20	8579.794	0.005	0.3	U			BNn	78St25	Z	
	8579.7	0.3			0.3	o			MMn	80Is02	Z	
	8579.81	0.20			-0.1	U			MMn	81Ke02	Z	
	8579.66	0.10			1.3	U				81Su.A	Z	
	8579.61	0.09			2.0	U			ILn	82Kr12	Z	
	8579.67	0.17			0.7	U			Bdn	06Fi.A		
	8579.7945	0.0048			0.0	1	100	99 $^{36}\text{Cl}$	NBS	06De21		
$^{35}\text{Cl}(d, p)^{36}\text{Cl}$	6360	8	6355.228	0.005	-0.6	U			MIT	64Sp12		
$^{35}\text{Cl}(p, \gamma)^{36}\text{Ar}$	8506.1	0.5	8506.98	0.04	1.8	U				72Ho40	Z	
$^{35}\text{Cl}(p, \gamma)^{36}\text{Ar}^i$	-2346.8	1.5	-2345.2	1.2	1.1	2				76Hu01		
	-2342.5	1.9			-1.4	2				76Ma40		
$^{36}\text{Ar}(d, t)^{35}\text{Ar}$	-9007	10	-8998.3	0.7	0.9	U			Yal	70Wh04		
$^{36}\text{K}^i(p)^{35}\text{Ar}$	2592	21	2623.8	2.3	1.5	U			Brk	81Ay01		
	2623.8	2.3				3				95Ga16		
$^{36}\text{S}(^7\text{Li}, ^7\text{Be})^{36}\text{P}$	-11277	27	-11275	13	0.1	2			Can	85Dr06		
$^{36}\text{S}(^{14}\text{C}, ^{14}\text{N})^{36}\text{P}$	-10256	15	-10257	13	0.0	2			Mun	84Ma49		
$^{36}\text{Cl}(\beta^+)^{36}\text{S}$	1137	18	1142.13	0.19	0.3	U				68Pi03		
$^{36}\text{Cl}(\epsilon)^{36}\text{S}$	1180	15			-2.5	U				64Li10		
	1160	18			-1.0	U				65Be19		
$^{36}\text{S}(p, n)^{36}\text{Cl}$	-1924.64	0.31	-1924.47	0.19	0.5	1	37	36 $^{36}\text{S}$		01Wa50		
$^{36}\text{Cl}(\beta^-)^{36}\text{Ar}$	708.7	0.6	709.53	0.04	1.4	U				67Sp06		
$^{36}\text{Ar}(p, n)^{36}\text{K}$	-13588.3	8.	-13596.8	0.3	-1.1	U			BNL	71Go18	Z	
	-13618	23			0.9	U				71Ja09		
$^{36}\text{Ar}(^3\text{He}, t)^{36}\text{K}$	-12930	40	-12833.1	0.3	2.4	U			Duk	70Dz04		
$^{36}\text{S}(^{64}\text{Ni}, ^{66}\text{Zn})^{34}\text{Si}$	Calibrated with $^{36}\text{S}(^{64}\text{Ni}, ^{62}\text{Ni}) M - A = -26862(12)$ now $-26861(7)$										AHW	**
$^{36}\text{S}(d, \alpha)^{34}\text{P}$	Original error 1.2 judged too small										GAU	**
$^{36}\text{Ar}(p, t)^{34}\text{Ar}^i$	IT=7950(50); $Q$ rebuilt, estimated with 72Pa02 $Q = -19523$ for ground state										MMC12a	**
$^{36}\text{Ar}(p, t)^{34}\text{Ar}^i$	IT=7925(5); $Q$ rebuilt with author's $Q = -19523$ for ground state										MMC128	**
$^{36}\text{S}(^{64}\text{Ni}, ^{65}\text{Zn})^{35}\text{Si}$	$M - A = -14482(59)$ for average of ground state and 54, 114, 207 levels										86Sm05	**
$^{36}\text{S}(^{14}\text{C}, ^{15}\text{N})^{35}\text{P}$	Original report -2693 is a typo										GAU	**
$^{37}\text{Al}-u$	10310	579	10530	190	0.3	o			GA3	1.5	91Or01	
	10900	450			-0.5	o			GA5	1.5	00Sa21	
	10531	129				2			GA7	1.5	07Ju03	
$^{37}\text{Si}-u$	-7550	1410	-7050	120	0.2	o			GA1	1.5	87Gi05	
	-7310	305			0.6	o			GA3	1.5	91Or01	
	-6930	150			-0.6	2			TO4	1.5	91Zh24	
	-7107	97			0.4	2			GA7	1.5	07Ju03	
$^{37}\text{P}-u$	-20740	430	-20390	40	0.5	U			TO1	1.5	86Vi09	
	-19910	190			-1.7	o			GA1	1.5	87Gi05	
	-20442	200			0.2	U			GA3	1.5	91Or01	
$\text{C}_3 \text{H}-^{37}\text{Cl}$	41924.73	1.09	41922.45	0.06	-0.8	U			C2	2.5	65De09	
	41922.2	0.2			0.5	U			M17	2.5	66Be10	
	41922.176	0.305			0.4	U			J5	2.5	72Ka57	
$\text{C}_2 \text{D}_8-^{37}\text{Cl} \text{H}_3$	123436.51	0.12	123436.54	0.06	0.2	U			B07	1.5	71Sm01	
$\text{C}_3 \text{H}_6 \text{O}_2-^{37}\text{Cl}_2$	104974.24	0.08	104974.26	0.11	0.2	1	85	85 $^{37}\text{Cl}$	B07	1.5	71Sm01	
$\text{C}_3 \text{H}_5-\text{D}_2-^{37}\text{Cl}$	45020.96	1.14	45019.02	0.06	-0.7	U			C2	2.5	65De09	
$\text{C}_8 \text{H}_{15}-^{37}\text{Cl}_3$	219665.80	0.90	219667.73	0.17	0.9	U			A2	2.5	70St25	
$\text{C}_3 \text{H}_3-\text{D}-^{37}\text{Cl}$	43473.27	1.33	43470.73	0.06	-0.8	U			C2	2.5	65De09	
$^{37}\text{K}-u$	-26632.5	6.4	-26624.11	0.10	1.3	U			LZ1	1.0	11Tu09	
$\text{H}_3 \text{O}-^{37}\text{Ca}_{.514}$	25638.22	0.35				2			MS1	1.0	07Ri08	
$\text{D}_2-^{35}\text{Cl}-\text{H}_2-^{37}\text{Cl}$	15505.41	0.71	15503.60	0.07	-1.0	U			C2	2.5	65De09	
	15503.80	0.09			-0.9	U			H31	2.5	77So02	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_5 H_{12} - ^{35}Cl - ^{37}Cl$	159145.17	0.12	159145.11	0.07	-0.3	1	13	9 $^{37}Cl$	B07	1.5	71Sm01
$H_2 ^{35}Cl - ^{37}Cl$	18600.0	0.4	18600.17	0.07	0.2	U			M17	2.5	66Be10
$D ^{35}Cl - ^{37}Cl$	17052.95	1.02	17051.89	0.07	-0.4	U			C2	2.5	65De09
	17051.816	0.185			0.2	U			J5	2.5	72Ka57
$^{37}K - ^{39}K_{949}$	7817.98	0.33	7818.43	0.10	1.4	U			MA8	1.0	07Ya08
$^{37}Cl(p, \alpha)^{34}S$	3030	6	3034.20	0.07	0.7	U			Bar		57Va03 Y
	3029	8			0.7	U			MIT		64Sp12
$^{37}Ar(n, \alpha)^{34}S$	4630	7	4630.42	0.21	0.1	U			ILL		78As06
$^{34}S(\alpha, n)^{37}Ar$	-4625	90	-4630.42	0.21	-0.1	U			Tal		63Ne05
$^{37}Cl(d, \alpha)^{35}S$	7791	12	7795.47	0.07	0.4	U			MIT		64Sp12
$^{37}Cl(p, ^3He)^{35}Si$	-19713	10				2					75Gu15
$^{35}Cl(^3He, p)^{37}Ar$	9582	15	9576.38	0.21	-0.4	U			MIT		67Sp09
$^{36}Mg(n, \gamma)^{37}Mg$	240	110				3					14Ko14 *
$^{36}S(^{18}O, ^{17}F)^{37}P$	-14410	40	-14400	40	0.2	2			Can		88Or.A *
$^{36}S(^{48}Ca, ^{47}Sc)^{37}P$	-11490	120	-11560	40	-0.6	2			Dar		88Fi04 *
$^{36}S(n, \gamma)^{37}S$	4303.52	0.12	4303.60	0.06	0.7	2			ORn		84Ra09 Z
	4303.61	0.09			-0.1	2			Bdn		06Fi.A
$^{36}S(d, p)^{37}S$	2079.12	0.13	2079.04	0.06	-0.6	2					84Pi03
$^{36}S(^{14}C, ^{13}C)^{37}S$	-3874	7	-3872.83	0.06	0.2	U			Mun		84Ma49
$^{37}Cl(^{13}C, ^{14}O)^{36}P$	-16433	50	-16393	13	0.8	U			Can		88Or01
$^{36}S(p, \gamma)^{37}Cl$	8386.47	0.23	8386.38	0.19	-0.4	1	65	64 $^{36}S$	Utr		84No05 Z
$^{36}S(p, \gamma)^{37}Cl^i$	-1835.5	0.3				2					84No05
$^{36}Ar(n, \gamma)^{37}Ar$	8791.1	1.0	8787.44	0.21	-3.7	B					68Wi25 Z
	8788.8	1.2			-1.1	U					70Ha56 Z
	8789.9	0.9			-2.7	U			Bdn		06Fi.A
$^{36}Ar(p, \gamma)^{37}K$	1857.3	1.0	1857.63	0.09	0.3	U					64Ar17
	1857.63	0.09				2			Utr		88De03 Z
$^{36}Ar(d, n)^{37}K$	-320	100	-366.94	0.09	-0.5	U			Yal		61Ya01
$^{36}Ar(p, \gamma)^{37}K^i$	-3192.6	0.8				2					88De03
$^{37}S(\beta^-)^{37}Cl$	4750	40	4865.12	0.20	2.9	U					67Wi14
$^{37}Cl(t, ^3He)^{37}S$	-4854	30	-4846.53	0.20	0.2	U			LAI		70Aj01
$^{37}Ar(\epsilon)^{37}Cl$	818	15	813.87	0.20	-0.3	U					53An01
$^{37}Cl(p, n)^{37}Ar$	-1595.5	4.0	-1596.22	0.20	-0.2	U			Wis		50Ri59 Y
	-1595.4	1.0			-0.8	U			MIT		52Sc09 Z
	-1596.9	2.4			0.3	U			Oak		64Jo11
	-1596.8	1.0			0.6	U			Duk		66Pa18 Z
	-1596.22	0.20				2			PTB		98Bo30
	-1596.3	1.0			0.1	U					01Wa50
$^{37}K(\beta^+)^{37}Ar$	6120	70	6147.47	0.23	0.4	U					58Su60
	6170	70			-0.3	U					60Wa04
$^*H_3 O - ^{37}Ca_{514}$	Error in Table II : $M - A = 13135.7(1.4)$ corrected to $-13136.06(0.64)$ keV										
$^*^{36}Mg(n, \gamma)^{37}Mg$	Symmetrized from $220(+120-90)$ keV										
$^*^{36}S(^{18}O, ^{17}F)^{37}P$	And $Q = -13650(40)$ , $M - A = -19750(40)$ if other peak is ground state one										
$^*^{36}S(^{48}Ca, ^{47}Sc)^{37}P$	And $Q = -11569(80)$ , $M - A = -18980(80)$ if other peak due to $^{47}Sc$ 807.89 level										
$^{38}Al-u$	15240	1500	17400	400	1.0	o			GA4	1.5	00Sa21
	17980	920			-0.4	o			GA5	1.5	00Sa21
	17402	268				2			GA7	1.5	07Ju03
$^{38}Si-u$	-4510	180	-4480	110	0.1	o			GA4	1.5	00Sa21
	-4020	290			-1.1	U			TO4	1.5	91Zh24
	-4100	320			-0.8	o			GA5	1.5	00Sa21
	-4477	75				2			GA7	1.5	07Ju03
$^{38}P-u$	-14420	620	-15700	80	-1.4	U			GA1	1.5	87Gi05
	-15910	140			1.0	2			GA4	1.5	00Sa21
	-15530	150			-0.7	2			TO4	1.5	91Zh24
	-16110	310			0.9	U			GA5	1.5	00Sa21
	-15717	75			0.2	o			GT1	1.5	04Ma.A
	-15660	100			-0.1	2			GT2	2.5	08Kn.A
	-15688	97			-0.1	2			LZ1	1.0	15Xu14



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{38}\text{Ca}-\text{H}_6\text{O}_2$	-60460.24	0.30	-60460.21	0.21	0.1	o			MS1	1.0	06Bo11
	-60460.24	0.30			0.1	1	48	48 $^{38}\text{Ca}$	MS1	1.0	07Ri08
$^{38}\text{Ar}-^{39}\text{K}_{.974}$	-1917.88	0.37	-1918.01	0.21	-0.4	1	32	32 $^{38}\text{Ar}$	MA8	1.0	02He23
$^{38}\text{K}-^{39}\text{K}_{.974}$	4430.88	0.44	4431.00	0.21	0.3	1	23	23 $^{38}\text{K}$	MA8	1.0	07Ya08
$^{38}\text{Ca}-^{19}\text{F}-^{39}\text{K}_{1.462}$	27783.80	0.63	27783.51	0.21	-0.5	U			MA8	1.0	07Ge07
$^{38}\text{K}-^{38}\text{Ar}$	6348.974	0.068	6349.01	0.05	0.6	1	50	27 $^{38}\text{K}$	JY1	1.0	09Er07
$^{38}\text{K}^m-^{38}\text{Ar}$	6488.743	0.049	6488.73	0.04	-0.2	1	72	45 $^{38}\text{K}^m$	JY1	1.0	09Er07
$^{38}\text{Ar}-^{38}\text{Ca}$	-13587.18	0.12	-13587.12	0.07	0.5	1	32	17 $^{38}\text{Ar}$	JY1	1.0	11Er02
$^{38}\text{K}^m-^{38}\text{K}$	139.698	0.065	139.72	0.05	0.3	-			JY1	1.0	09Er07
	139.78	0.14			-0.4	-			JY1	1.0	11Er02
ave.	139.71	0.06			0.1	1	60	34 $^{38}\text{K}^m$			average
$^{38}\text{Ca}-^{38}\text{K}$	7238.04	0.10	7238.11	0.07	0.7	1	45	25 $^{38}\text{K}$	JY1	1.0	11Er02
$^{38}\text{K}^m-^{38}\text{Ca}$	-7098.43	0.11	-7098.39	0.07	0.4	1	37	21 $^{38}\text{K}^m$	JY1	1.0	11Er02
$^{24}\text{Mg}(^{16}\text{O},2n)^{38}\text{Ca}$	-12727	30	-12754.71	0.19	-0.9	U					72Zi02 *
$^{35}\text{Cl}(\alpha,p)^{38}\text{Ar}$	837.2	2.4	837.24	0.20	0.0	U			Har		75Sq01
$^{35}\text{Cl}(\alpha,n)^{38}\text{K}$	-5862.1	1.5	-5859.17	0.20	2.0	U			Mun		76Sh24 Z
	-5858.7	2.9			-0.2	U			Har		75Sq01 *
$^{36}\text{S}(t,p)^{38}\text{S}$	3838	30	3858	7	0.7	U					85Da15
$^{36}\text{S}(^{14}\text{C},^{12}\text{C})^{38}\text{S}$	-781	10	-783	7	-0.2	R			Mun		84Ma49
$^{36}\text{Ar}(^3\text{He},n)^{38}\text{Ca}$	-1365	21	-1313.14	0.20	2.5	U			CIT		69Sh04
$^{37}\text{Cl}(n,\gamma)^{38}\text{Cl}$	6107.84	0.30	6107.88	0.08	0.1	U					73Sp06 Z
	6107.95	0.10			-0.7	2			MMn		81Ke02 Z
	6107.73	0.15			1.0	2			Bdn		06Fi.A
$^{37}\text{Cl}(d,p)^{38}\text{Cl}$	3885	8	3883.32	0.08	-0.2	U			MIT		64Sp12
	3883.28	0.50			0.1	U			Rez		90Pi05 *
$^{37}\text{Cl}(p,\gamma)^{38}\text{Ar}$	10243.0	1.0	10242.25	0.20	-0.7	U					68En01 Z
$^{38}\text{S}(\beta^-)^{38}\text{Cl}$	2947	20	2937	7	-0.5	3					71En01
	2936	12			0.1	3					72Vi11
$^{38}\text{Cl}(\beta^-)^{38}\text{Ar}$	4913	5	4916.72	0.22	0.7	U					68Va06
$^{38}\text{K}(\beta^+)^{38}\text{Ar}$	5870	30	5914.07	0.04	1.5	U					56Gr07 *
	5790	50			2.5	U					67Va27 *
$^{38}\text{Ar}(p,n)^{38}\text{K}$	-6695.5	4.	-6696.41	0.04	-0.2	U			Har		75Sq01
	-6695.65	0.70			-1.1	U					78Ja06 Z
$^{38}\text{Ar}(p,n)^{38}\text{K}^m$	-6826.73	0.12	-6826.56	0.04	1.4	U			Auc		98Ha36 Z
$^{24}\text{Mg}(^{16}\text{O},2n)^{38}\text{Ca}$	$E(^{16}\text{O})=24880(30)$ to $2^+$ level at 2213.13(0.10) keV										
$^{35}\text{Cl}(\alpha,n)^{38}\text{K}$	$Q=-5989.1(2.9,Z)$ to $^{38}\text{K}^m$ at 130.15(0.04) keV										
$^{37}\text{Cl}(d,p)^{38}\text{Cl}$	Estimated systematic error 0.5 added to statistical error 0.064 keV										
$^{38}\text{K}(\beta^+)^{38}\text{Ar}$	$E_{\beta^+}=2680(30)$ 2600(50) respectively, to $2^+$ level at 2167.64 keV										
$^{39}\text{Al}-u$	22970	1580	22170#	430#	-0.3	o			GA5	1.5	00Sa21
	21653	676			0.5	D			GA7	1.5	07Ju03 *
$^{39}\text{Si}-u$	1900	540	2490	150	0.7	o			GA4	1.5	00Sa21
	2210	490			0.4	o			GA5	1.5	00Sa21
	2491	97				2			GA7	1.5	07Ju03
$^{39}\text{P}-u$	-13890	140	-13710	120	0.8	2			GA4	1.5	00Sa21
	-13580	160			-0.6	2			TO4	1.5	91Zh24
	-13870	280			0.4	2			GA5	1.5	00Sa21
	-13602	140			-0.5	2			GT1	1.5	04Ma.A
$^{39}\text{K}-^{23}\text{Na}_{1.696}$	-18942.88	0.58	-18942.216	0.006	0.8	U			Ma8	1.5	08Mu05
$^{39}\text{Ca}-u$	-29278.8	6.4	-29289.2	0.6	-1.6	U			LZ1	1.0	11Tu09
$^{39}\text{K}-^{36}\text{Ar}_{1.083}$	-1144.65	0.44	-1144.86	0.03	-0.5	U			MA8	1.0	02He23
	-1144.83	0.40			-0.1	U			MA8	1.0	03B117
$^{39}\text{K}-^{37}\text{K}_{1.054}$	-8231.29	0.53	-8231.70	0.11	-0.5	U			Ma8	1.5	08Mu05
$^{39}\text{Ca}-^{19}\text{F}-^{39}\text{K}_{1.487}$	23082.43	0.64	23082.4	0.6	0.0	1	100	100 $^{39}\text{Ca}$	MA8	1.0	08Ge08
$^{39}\text{K}-^{40}\text{Ar}$	1323.3631	0.0043	1323.363	0.004	-0.1	1	100	100 $^{39}\text{K}$	FS1	1.0	10Mo30

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{39}\text{K}(\text{p},\alpha)^{36}\text{Ar}$	1287	7	1288.405	0.027	0.2	U			MIT		64Sp12
$^{37}\text{Cl}(\text{t},\text{p})^{39}\text{Cl}$	5701.9	2.5	5699.5	1.7	-1.0	2			Str		84An03
$^{39}\text{K}(\text{p},^3\text{He})^{37}\text{Ar}^i$	-15493.4	6.				2			MSU		73Be23 *
$^{39}\text{K}(\text{p},\text{t})^{37}\text{K}^i$	-21713.1	3.	-21718.1	0.8	-1.7	U			MSU		73Be23 *
$^{39}\text{Sc}^i(2\text{p})^{37}\text{K}$	4969	120	5170	40	1.7	U			Lis		90De43 *
	4877	40			7.3	B			Brk		92Mo15 *
	5146	40			0.6	o			Bor		01Gi01 *
	5170	40				3			Bor		07Do17 *
$^{38}\text{Ar}(\text{p},\gamma)^{39}\text{K}$	6380.9	1.1	6381.34	0.19	0.4	U					70Ma31 Z
	6382.2	0.8			-1.1	U					84Ha27 Z
$^{39}\text{K}(\text{p},\text{d})^{38}\text{K}$	-10851	2	-10853.19	0.20	-1.1	U			MSU		74Wi17
$^{39}\text{K}(^3\text{He},\alpha)^{38}\text{K}$	7498	15	7499.87	0.20	0.1	U			Roc		66Bl04
	7483	10			1.7	U			Roc		72Fe06
$^{39}\text{Cl}(\beta^-)^{39}\text{Ar}$	3440	20	3442	5	0.1	U					56Pe38
$^{39}\text{Ar}(\beta^-)^{39}\text{K}$	565	5				2					50Br66
$^{39}\text{Ca}(\beta^+)^{39}\text{K}$	6512	25	6524.5	0.6	0.5	U					58Ki40
$^{39}\text{K}(\text{p},\text{n})^{39}\text{Ca}$	-7302.5	6.	-7306.8	0.6	-0.7	U			Tal		70Ke08
	-7314.9	1.8			4.5	B					78Ra15 Z
* $^{39}\text{Al}-\text{u}$	Trends from Mass Surface TMS suggest $^{39}\text{Al}$ 480 less bound										GAu **
* $^{39}\text{K}(\text{p},^3\text{He})^{37}\text{Ar}^i$	$M-A=-25954(6)$ ; rebuilt $Q=-15493.8(6.)$ with Ame1971; recalibration +0.35										MMC123**
* $^{39}\text{K}(\text{p},\text{t})^{37}\text{K}^i$	$M-A=-19753(3)$ ; $Q$ rebuilt with Ame1971										MMC123**
* $^{39}\text{Sc}^i(2\text{p})^{37}\text{K}$	$E_{2p}=3600(120)$ to $1/2^+$ level at 1370.85 keV										Ens123 **
* $^{39}\text{Sc}^i(2\text{p})^{37}\text{K}$	Other possibility $^{39}\text{Sc}^i(\alpha)^{35}\text{K}=3600(120)$ keV										90De43 **
* $^{39}\text{Sc}^i(2\text{p})^{37}\text{K}$	$E_{2p}=4750(40)$ p+p at 90 degrees; deduced $Q=E_{2p}[1 + \text{Mp}/\text{M}(^{37}\text{K})]$										MMC123**
* $^{39}\text{Sc}^i(2\text{p})^{37}\text{K}$	$E_{2p}=4880(40)$ + recoil 266 keV; data reanalysed and included in 07Do17										MMC135**
* $^{39}\text{Sc}^i(2\text{p})^{37}\text{K}$	IAS identification not sure										MMC135**
$^{40}\text{Si}-\text{u}$	5290	1010	5830	370	0.4	o			GA4	1.5	00Sa21
	6180	740			-0.3	o			GA5	1.5	00Sa21
	5829	247				2			GA7	1.5	07Ju03
$^{40}\text{P}-\text{u}$	-8800	200	-8710	160	0.3	o			GA4	1.5	00Sa21
	-8950	210			0.8	2			TO4	1.5	91Zh24
	-8200	320			-1.1	o			GA5	1.5	00Sa21
	-8621	129			-0.5	2			GA7	1.5	07Ju03
$^{40}\text{S}-\text{u}$	-24440	190	-24517	4	-0.3	o			GA4	1.5	00Sa21
	-24530	250			0.0	U			TO4	1.5	91Zh24
	-24910	340			0.8	o			GA5	1.5	00Sa21
	-24627	129			0.6	U			GA7	1.5	07Ju03
$\text{C}_3 \text{H}_4-^{40}\text{Ar}$	68917.0053	0.0035	68917.0052	0.0024	0.0	1	46	46 $^{40}\text{Ar}$	MI1	1.0	95Di08
$\text{C}_2 \text{D}_8-^{40}\text{Ar}$	150431.1045	0.0040	150431.1011	0.0024	-0.8	1	36	33 $^{40}\text{Ar}$	MI1	1.0	95Di08
$^{20}\text{Ne}_2-^{40}\text{Ar}$	22497.2245	0.0042	22497.228	0.003	1.0	-			MI1	1.0	95Di08
	22497.2280	0.0060			0.1	-			MI1	1.0	95Di08
ave.	22497.226	0.003			0.9	1	74	60 $^{20}\text{Ne}$			average
$^{40}\text{Ar}-\text{u}$	-37616.878	0.040	-37616.8762	0.0024	0.0	U			ST2	1.0	02Bf02
$^{40}\text{Ca}-\text{H}_{40}$	-350410.425	0.022	-350410.424	0.022	0.0	1	99	99 $^{40}\text{Ca}$	ST2	1.0	06Na18
$^{40}\text{S O}-^{41}\text{K}_{1,366}$	22541	16	22544	4	0.2	U			MS1	1.0	09Ri12
$^{40}\text{S}-^{41}\text{K}_{976}$	12752.0	9.4	12741	4	-1.2	1	21	21 $^{40}\text{S}$	MS1	1.0	09Ri12
$^{40}\text{S}-^{40}\text{Ar}$	13096.6	4.8	13099	4	0.6	1	79	79 $^{40}\text{S}$	MS1	1.0	09Ri12
$^{40}\text{Ca}-^{40}\text{Ar}$	208.2	0.5	207.742	0.022	-0.4	U			J3	2.5	68Fu11
$^{40}\text{Ca}(^3\text{He},^8\text{Li})^{35}\text{K}$	-29693	20	-29688.1	0.5	0.2	U			MSU		76Be08
$^{40}\text{Ca}(\alpha,^8\text{He})^{36}\text{Ca}$	-57580	40				2			Tex		77Tr03
$^{40}\text{Ar}(\text{n},\alpha)^{37}\text{S}$	-2500	50	-2497.08	0.20	0.1	U					55Be78
	-2490	50			-0.1	U			Ric		64Da11
$^{40}\text{K}(\text{n},\alpha)^{37}\text{Cl}$	3866	7	3872.45	0.08	0.9	U			BNL		68Sc01
$^{40}\text{Ca}(\text{p},\alpha)^{37}\text{K}$	-5179	9	-5182.13	0.10	-0.3	U			CIT		66Mc13

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{40}\text{Ca}(^3\text{He}, ^6\text{He})^{37}\text{Ca}$	-24270	50	-24371.2	0.6	-2.0	U	Brk		68Bu02
	-24368	25			-0.1	U	MSU		73Be23 *
$^{40}\text{Ar}(p, ^3\text{He})^{38}\text{Cl}^i$	-21092	24				2	Brk		70Ha10 *
$^{40}\text{Ar}(p,t)^{38}\text{Ar}^j$	-26765	31				2			70Ha10 *
$^{40}\text{Ca}(d,\alpha)^{38}\text{K}$	4655	10	4665.18	0.20	1.0	U	MIT		64Sp12
$^{40}\text{Ca}(p,t)^{38}\text{Ca}$	-20459	25	-20448.72	0.20	0.4	U			66Ha32
	-20428	11			-1.9	U	MSU		72Pa02
	-20452	5			0.7	U	MSU		74Se05
$^{40}\text{Ar}(^{18}\text{O}, ^{19}\text{Ne})^{39}\text{S}$	-14504	200	-14410	50	0.5	U	Can		84Ho.B
$^{40}\text{Ar}(^{13}\text{C}, ^{14}\text{O})^{39}\text{S}$	-16760	50				2	Can		89Dr03
$^{40}\text{Ar}(t,\alpha)^{39}\text{Cl}$	7256	40	7285.2	1.7	0.7	U	LAI		61Ja07
$^{40}\text{Ar}(d, ^3\text{He})^{39}\text{Cl}-^{36}\text{Ar}(^3\text{He})^{35}\text{Cl}$	-4024.13	2.42	-4021.7	1.7	1.0	R	Hei		93Ma50
$^{40}\text{Ar}(^3\text{He}, \alpha)^{39}\text{Ar}^i$	1604	19	1627	7	1.2	2			67Gr01 *
	1631.3	8.0			-0.5	2			72Wi07 *
$^{39}\text{K}(n,\gamma)^{40}\text{K}$	7799.50	0.08	7799.62	0.06	1.5	-	ILn		84Vo01 Z
	7799.56	0.16			0.4	-	Bdn		06Fi.A
$^{39}\text{K}(d,p)^{40}\text{K}$	5579	10	5575.05	0.06	-0.4	U	MIT		64Sp12
$^{39}\text{K}(n,\gamma)^{40}\text{K}$	ave.	7799.51	7799.62	0.06	1.5	1	61	61 $^{40}\text{K}$	average
$^{39}\text{K}(p,\gamma)^{40}\text{Ca}$	8329.5	0.9	8328.165	0.021	-1.5	U			68Do12
	8329.6	0.9			-1.6	U			68Li12 *
	8328.24	0.09			-0.8	U	Utr		90Ki07 Z
$^{39}\text{K}(^3\text{He}, d)^{40}\text{Ca}$	2845	8	2834.690	0.021	-1.3	U	Oak		67Se10
$^{40}\text{Ca}(^3\text{He}, \alpha)^{39}\text{Ca}$	4950	20	4942.6	0.6	-0.4	U	Ald		66Hi06
	4919	15			1.6	U	MIT		71Ra35
$^{40}\text{Ca}(^7\text{Li}, ^8\text{He})^{39}\text{Sc}$	-37400	40	-37376	24	0.6	2	MSU		88Mo18
$^{40}\text{Ca}(^{14}\text{N}, ^{15}\text{C})^{39}\text{Sc}$	-27670	30	-27683	24	-0.4	2	Can		88Wo07
$^{40}\text{Sc}^i(p)^{39}\text{Ca}$	3840	120	3830	6	-0.1	o	Lis		90De43
	3820	30			0.3	U			90Zh.A *
	3827.7	10.			0.2	2	GSI		97Li25 *
	3830.8	7.			-0.1	2	Lis		98Bh12
	3841	20			-0.6	U	Bor		07Do17
$^{40}\text{Cl}(\beta^-)^{40}\text{Ar}$	7320	80	7480	30	2.0	2			89Mi03
$^{40}\text{Ar}(^7\text{Li}, ^7\text{Be})^{40}\text{Cl}$	-8375	35	-8340	30	0.9	2			84Fi02
$^{40}\text{K}(\epsilon)^{40}\text{Ar}$	1504	7	1504.40	0.06	0.1	U			67Mc10 *
	1497	8			0.9	U			68Az01 *
$^{40}\text{K}(n,p)^{40}\text{Ar}$	2270	5	2286.75	0.06	3.3	B	BNL		68Sc01
	2286.7	1.0			0.0	U	ILL		81We12
$^{40}\text{Ar}(p,n)^{40}\text{K}$	-2286.3	1.0	-2286.75	0.06	-0.4	U	Duk		66Pa18 Z
	-2286.3	1.0			-0.4	U			01Wa50
$^{40}\text{K}(\beta^-)^{40}\text{Ca}$	1325	15	1310.89	0.06	-0.9	U			52Fe16
	1350	20			-2.0	U			59Ke26
$^{40}\text{Sc}(\beta^+)^{40}\text{Ca}$	14330	40	14323.0	2.8	-0.2	U			68Ar03 *
$^{40}\text{Ca}(p,n)^{40}\text{Sc}$	-15105.4	2.9				2	Yal		69Ov01 Z
$^{40}\text{Ca}(^3\text{He}, t)^{40}\text{Sc}$	-14490	60	-14341.6	2.8	2.5	U	Bld		65Ri06
$^{40}\text{Ca}(\pi^+, \pi^-)^{40}\text{Ti}$	-24974	160				2			82Mo12 *
$^{40}\text{Ca}(^3\text{He}, ^6\text{He})^{37}\text{Ca}$	Average of 2 values with small calibration correction								
$^{40}\text{Ar}(p, ^3\text{He})^{38}\text{Cl}^i$	IT=8216(25); rebuilt $Q=-21093.65(23.68)$ ; recalibrated for $^{10}\text{B}+1.5$ keV								
$^{40}\text{Ar}(p,t)^{38}\text{Ar}^j$	IT=18784(30); $Q$ rebuilt with $^{10}\text{C}=15702.5(1.8)$ from reference								
$^{40}\text{Ar}(^3\text{He}, \alpha)^{39}\text{Ar}^i$	IT=9089(20); $Q$ rebuilt with Ame1961								
$^{40}\text{Ar}(^3\text{He}, \alpha)^{39}\text{Ar}^i$	IT=9075(10); $Q$ rebuilt with Ame1964								
$^{39}\text{K}(p,\gamma)^{40}\text{Ca}$	$E_{res}=1345.4(0.5)$ to $2^-$ level at $9641.1(0.8)$ keV								
$^{40}\text{Sc}^i(p)^{39}\text{Ca}$	Uncertainty not given, estimated from graph: stat(9keV), calib(11)								
$^{40}\text{Sc}^i(p)^{39}\text{Ca}$	$E_p=3731(10)$ ; also $E_p=1330(20)$ $Q_p=1364.4(20)$ keV to 2468.5 level								
$^{40}\text{Sc}^i(p)^{39}\text{Ca}$	IT=4370(10) in original paper								
$^{40}\text{K}(\epsilon)^{40}\text{Ar}$	LMK=0.34(0.08) 0.47(0.16) respectively, to $2^+$ level at 1460.851, recalculated $Q$								
$^{40}\text{Sc}(\beta^+)^{40}\text{Ca}$	$E_{\beta^+}=9580(40)$ to $3^-$ level at 3736.69 keV, and other $E_{\beta^+}$								
$^{40}\text{Ca}(\pi^+, \pi^-)^{40}\text{Ti}$	Recalibrated to $^{16}\text{O}(\pi^+, \pi^-) Q=-27704(20)$ keV								

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{41}\text{Si}-u$	14560	1980	13010	600	-0.5	o			GA5	1.5	00Sa21
	13011	397				2			GA7	1.5	07Ju03
$^{41}\text{P}-u$	-5930	300	-5350	130	1.3	o			GA4	1.5	00Sa21
	-5200	500			-0.2	U			TO4	1.5	91Zh24
	-5290	420			-0.1	o			GA5	1.5	00Sa21
	-5346	86				2			GA7	1.5	07Ju03
$^{41}\text{S}-u$	-20500	150	-20407	4	0.4	U			GA4	1.5	00Sa21
	-19970	230			-1.3	U			TO4	1.5	91Zh24
	-20430	330			0.0	U			GA5	1.5	00Sa21
	-20494	75			0.8	U			GT1	1.5	04Ma.A
$^{41}\text{S}-\text{C}_2\text{H O}$	-23146.2	4.4				2			MS1	1.0	09Ri12
$^{41}\text{Cl}-u$	-29620	190	-29320	70	1.1	2			TO3	1.5	90Tu01
	-29500	270			0.5	2			TO4	1.5	91Zh24
$^{41}\text{Sc}-u$	-30741	12	-30748.90	0.09	-0.7	U			LZ1	1.0	11Tu09
$^{41}\text{Ti}-u$	-16200	390	-16850	30	-1.1	U			GT1	1.5	04St05
	-16852	30				2			LZ1	1.0	12Zh34
$^{41}\text{K}-^{39}\text{K}_{1.051}$	-30.05	0.32	-30.260	0.006	-0.7	U			MA8	1.0	02He23
	-29.5	2.4			-0.3	U			MA8	1.0	09Na.A
$^{41}\text{K}-^{40}\text{Ar H}$	-8382.9005	0.0061	-8382.898	0.003	0.4	-			FS1	1.0	10Mo30
$^{41}\text{K}-^{40}\text{Ar}$	-557.8652	0.0039	-557.866	0.003	-0.2	-			FS1	1.0	10Mo30
$^{41}\text{K}-^{40}\text{Ar H}$	ave. -8382.898	0.003	-8382.898	0.003	0.1	1	100	100 $^{41}\text{K}$			average
$^{41}\text{K}(\text{p},\alpha)^{38}\text{Ar}$	4002	20	4019.33	0.20	0.9	U			ChR		60CI02
	4018	10			0.1	U			MIT		64Sp12
$^{41}\text{K}(\text{d},\alpha)^{39}\text{Ar}$	8397	15	8393	5	-0.2	U			MIT		67Sp09
$^{39}\text{K}(\text{}^3\text{He},\text{p})^{41}\text{Ca}$	8920	20	8972.95	0.14	2.6	U			MIT		67Sp09
$^{40}\text{Ar}(\text{}^{18}\text{O}, \text{}^{17}\text{F})^{41}\text{Cl}$	-10530	83	-10470	70	0.8	R			Can		84Ho.B
$^{40}\text{Ar}(\text{n},\gamma)^{41}\text{Ar}$	6098.4	0.7	6098.9	0.3	0.8	2					70Ha56
	6099.1	0.4			-0.4	2			Bdn		06Fi.A
$^{40}\text{Ar}(\text{d},\text{p})^{41}\text{Ar}$	3878	6	3874.4	0.3	-0.6	U			MIT		64Sp12
$^{40}\text{Ar}(\text{p},\gamma)^{41}\text{K}$	7807.8	0.3	7808.619	0.003	2.7	U					89Sm06
$^{40}\text{Ar}(\text{}^3\text{He},\text{d})^{41}\text{K}^i$	-6034	15				2					75Me10
$^{40}\text{K}(\text{n},\gamma)^{41}\text{K}$	10095.19	0.10	10095.37	0.06	1.8	-			ILn		84Kr05
	10095.25	0.20			0.6	-			Bdn		06Fi.A
$^{40}\text{Ca}(\text{n},\gamma)^{41}\text{Ca}$	ave. 10095.20	0.09			1.9	1	39	39 $^{40}\text{K}$			average
	8363.0	0.5	8362.82	0.14	-0.4	-					69Ar.A
	8362.5	0.5			0.6	-					70Cr04
	8362.72	0.3			0.3	-			MMn		80Is02
	8362.86	0.17			-0.2	-			Bdn		06Fi.A
$^{40}\text{Ca}(\text{d},\text{p})^{41}\text{Ca}$	6134	4	6138.26	0.14	1.1	U			MIT		68Be36
$^{40}\text{Ca}(\text{n},\gamma)^{41}\text{Ca}$	ave. 8362.81	0.14	8362.82	0.14	0.1	1	100	100 $^{41}\text{Ca}$			average
	1085.7	1.4	1085.00	0.08	-0.5	U					73Al11
$^{40}\text{Ca}(\text{p},\gamma)^{41}\text{Sc}$	1085.09	0.09			-1.0	1	79	79 $^{41}\text{Sc}$	Utr		87Zi02
$^{40}\text{Ca}(\text{d},\text{n})^{41}\text{Sc}$	-1145	15	-1139.57	0.08	0.4	U					61Ma08
$^{40}\text{Ca}(\text{p},\gamma)^{41}\text{Sc}^r$	-1796.4	1.5	-1797.33	0.09	-0.6	U					77Ko10
$^{40}\text{Ca}(\text{p},\gamma)^{41}\text{Sc}^i$	-4851.4	4.9	-4854	3	-0.5	2					76Fo01
	4855.6	5.	4854	3	-0.4	2			Jyp		97Ho12
$^{41}\text{Sc}^i(\text{p})^{40}\text{Ca}$	4855.6	8.			-0.2	2			Lis		98Bh12
	4857	16			-0.2	U			Bor		07Do17
$^{41}\text{Cl}(\beta^-)^{41}\text{Ar}$	5670	150	5760	70	0.6	R					74Gu10
$^{41}\text{Ar}(\beta^-)^{41}\text{K}$	2492.0	1.1	2492.0	0.3	0.0	U					64Pa03
$^{41}\text{K}(\text{p},\text{n})^{41}\text{Ca}$	-1209.6	1.5	-1204.00	0.14	3.7	B			Oak		64Jo11
	-1203.8	0.5			-0.4	U			Can		70Kn03
$^{41}\text{Sc}(\beta^+)^{41}\text{Ca}$	6630	100	6495.48	0.16	-1.3	U					62Cr04
$^{41}\text{Sc}^r(\text{IT})^{41}\text{Sc}$	2882.6	0.3	2882.32	0.05	-0.9	U					77Ko10
	2882.39	0.10			-0.7	-			Utr		87Zi02
	2882.26	0.06			1.1	-			Utr		89Ki11
	ave. 2882.29	0.05			0.6	1	93	72 $^{41}\text{Sc}^r$			average
$^{41}\text{S}-\text{C}_2\text{H O}$	For original doublet $^{41}\text{S C H}-\text{C}_3\text{H}_2\text{O}$										
$^{41}\text{Ti}-u$	Same result in reference										
$^{40}\text{Ar}(\text{}^3\text{He},\text{d})^{41}\text{K}^i$	IT=8349(15); $Q$ rebuilt with Ame1971										
$^{40}\text{Ca}(\text{p},\gamma)^{41}\text{Sc}$	$E_p=647.25(0.05,Z)$ to $1716.43(0.08,Z)$ level										
$^{41}\text{Ar}(\beta^-)^{41}\text{K}$	$E_{\beta^-}=1198.3(1.1)$ to $7/2^-$ level at 1293.609 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{42}\text{Si}-u$	20860	3990	17680#	540#	-0.5	o			GA5	1.5	99Sa.A
	16275	623			1.5	D			GA7	1.5	07Ju03 *
$^{42}\text{P}-u$	260	740	1080	340	0.7	o			GA4	1.5	00Sa21
	1550	630			-0.5	o			GA5	1.5	00Sa21
	1084	225				2			GA7	1.5	07Ju03
$^{42}\text{S}-u$	-18940	150	-18935	3	0.0	U			GA4	1.5	00Sa21
	-18510	350			-0.8	U			TO4	1.5	91Zh24
	-19390	350			0.9	U			GA5	1.5	00Sa21
	-18934.9	3.0				2			MS1	1.0	09Ri12 *
$^{42}\text{Cl}-u$	-27000	190	-26660	60	1.2	U			TO3	1.5	90Tu01
	-26870	190			0.7	U			TO4	1.5	91Zh24
	-26658	64				2			LZ1	1.0	15Xu14
$^{42}\text{Ti}-u$	-26973	25	-26950.98	0.30	0.9	U			LZ1	1.0	16Zh.A
$^{42}\text{Ar}-^{36}\text{Ar}_{1.167}$	920.6	6.2				2			MA6	1.0	01He29
$^{42}\text{S}-^{41}\text{K}_{1.024}$	20151.8	9.5	20156	3	0.4	U			MS1	1.0	09Ri12
$^{42}\text{Sc}-^{42}\text{Ca}$	6898.74	0.22	6898.69	0.10	-0.2	1	22	19 $^{42}\text{Sc}$	JY1	1.0	06Er08
$^{42}\text{Sc}^m-^{42}\text{Ca}$	7560.68	0.23	7560.34	0.11	-1.5	1	25	22 $^{42}\text{Sc}^m$	JY1	1.0	06Er08
$^{42}\text{Ti}-^{42}\text{Ca}$	14431.69	0.71	14431.19	0.26	-0.7	1	13	13 $^{42}\text{Ti}$	JY1	1.0	09Ku19
$^{42}\text{Sc}^m-^{42}\text{Sc}$	661.97	0.24	661.65	0.06	-1.3	U			JY1	1.0	06Er08
	662.50	0.42			-2.0	U			JY1	1.0	09Ku19
$^{42}\text{Ti}-^{42}\text{Sc}$	7532.92	0.34	7532.50	0.24	-1.2	1	50	49 $^{42}\text{Ti}$	JY1	1.0	09Ku19
$^{42}\text{Ti}-^{42}\text{Sc}^m$	6870.19	0.38	6870.85	0.24	1.7	1	40	38 $^{42}\text{Ti}$	JY1	1.0	09Ku19
$^{28}\text{Si}(^{16}\text{O},2n)^{42}\text{Ti}$	-17250	13	-17267.76	0.28	-1.4	U					72Zi02
$^{42}\text{Ca}(p,\alpha)^{39}\text{K}$	118	7	124.00	0.15	0.9	U			MIT		64Sp12
$^{39}\text{K}(\alpha,n)^{42}\text{Sc}$	-7160	60	-7332.44	0.17	-2.9	U			Yal		61Sm05
	-7455	30			4.1	B			Tal		65Ne02
$^{40}\text{Ar}(t,p)^{42}\text{Ar}$	7043	40	7044	6	0.0	U			LAI		61Ja07
$^{40}\text{Ca}(^3\text{He},p)^{42}\text{Sc}$	4966	20	4917.02	0.17	-2.4	U			MIT		64Sp12
	4905	5			2.4	U			ANL		74Ha55
$^{40}\text{Ca}(^3\text{He},n)^{42}\text{Ti}$	-2865	6	-2881.81	0.28	-2.8	U			CIT		67Mi02
$^{41}\text{K}(n,\gamma)^{42}\text{K}$	7533.78	0.15	7533.80	0.11	0.1	2			ILn		85Kr06 Z
	7533.82	0.15			-0.1	2			Bdn		06Fi.A
$^{41}\text{K}(d,p)^{42}\text{K}$	5314	12	5309.23	0.11	-0.4	U			MIT		64Sp12
$^{41}\text{K}(p,\gamma)^{42}\text{Ca}$	10275.5	3.4	10276.67	0.15	0.3	U					71Vi14
$^{41}\text{Ca}(n,\gamma)^{42}\text{Ca}$	11480.63	0.06	11480.67	0.06	0.7	1	91	90 $^{42}\text{Ca}$	ORn		89Ki11 Z
$^{42}\text{Ca}(^3\text{He},\alpha)^{41}\text{Ca}$	9102	15	9096.95	0.06	-0.3	U			MIT		71Ra35
$^{41}\text{Ca}(p,\gamma)^{42}\text{Sc}^r-^{40}\text{Ca}(^{41}\text{Sc}^r$	-6.67	0.05	-6.70	0.05	-0.6	1	94	66 $^{42}\text{Sc}^r$	Utr		89Ki11 *
$^{42}\text{Cl}(\beta^-)^{42}\text{Ar}$	9760	220	9590	60	-0.8	U					89Mi03
$^{42}\text{K}(\beta^-)^{42}\text{Ca}$	3519.	3.5	3525.22	0.18	1.8	U					68Va06
	3524	6			0.2	U					75Ra09
$^{42}\text{Sc}(\beta^+)^{42}\text{Ca}$	6342	100	6426.09	0.10	0.8	U					61Ja22
	6486	100			-0.6	U					63Ro10 *
$^{42}\text{Ca}(p,n)^{42}\text{Sc}$	-7213.7	2.3	-7208.44	0.10	2.3	U			Har		75Fr.A
$^{42}\text{Ca}(^3\text{He},t)^{42}\text{Sc}$	-6442.3	0.4	-6444.68	0.10	-6.0	F			Mun		77Vo02 *
$^{42}\text{Ca}(^3\text{He},t)^{42}\text{Sc}-^{27}\text{Al}(^{27}\text{Si}$	-1611.7	2.6	-1613.73	0.14	-0.8	U			ChR		74Ha35
$^{42}\text{Ca}(^3\text{He},t)^{42}\text{Sc}-^{26}\text{Mg}(^{26}\text{Al}$	-2417.8	3.5	-2421.70	0.11	-1.1	U			ChR		74Ha35
	-2421.83	0.23			0.6	1	23	14 $^{42}\text{Sc}$	ChR		87Ko34 *
$^{42}\text{Sc}^m(\text{IT})^{42}\text{Sc}$	616.28	0.06	616.32	0.06	0.7	1	93	76 $^{42}\text{Sc}^m$			Ens167
$^{42}\text{Sc}^r(\text{IT})^{42}\text{Sc}$	6076.33	0.08	6076.26	0.07	-0.9	1	84	50 $^{42}\text{Sc}$	Utr		89Ki11 Z
* $^{42}\text{Si}-u$	Trends from Mass Surface TMS suggest $^{42}\text{Si}$ 1310 less bound										
* $^{42}\text{S}-u$	For original doublet $^{42}\text{S}-(\text{C}_2\text{H O})_{1.024}$ , $D_M=-21740.2(3.1)\mu\text{u}$										
* $^{41}\text{Ca}(p,\gamma)^{42}\text{Sc}^r-^{40}\text{Ca}(^{41}\text{Sc}^r$	Calculated from resonance energy difference = 5.73(0.05) keV										
* $^{42}\text{Sc}(\beta^+)^{42}\text{Ca}$	$E_{\beta^+}=2870(100)$ from $^{42}\text{Sc}^m$ at 616.28 to $6^+$ level at 3189.26 keV										
* $^{42}\text{Ca}(^3\text{He},t)^{42}\text{Sc}$	F : rejected in reference of same group										
* $^{42}\text{Ca}(^3\text{He},t)^{42}\text{Sc}-^{26}\text{Mg}(^{26}\text{Al}$	$Q=-2193.52(0.23)$ to $^{26}\text{Al}^m$ at 228.306 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{43}\text{P}-u$	4220	1620	5020	600	0.3	U			GA4	1.5	00Sa21
	6190	1040			-0.7	o			GA5	1.5	00Sa21
	5024	397				2			GA7	1.5	07Ju03
$^{43}\text{S}-u$	-12810	250	-13092	5	-0.8	o			GA4	1.5	00Sa21
	-13400	900			0.2	U			TO4	1.5	91Zh24
	-12900	460			-0.3	o			GA5	1.5	00Sa21
	-12958	107			-0.8	U			GA7	1.5	07Ju03
	-13087	22			-0.2	2			MS1	1.0	09Ri12 *
	-13092.7	5.5			0.1	2			MS1	1.0	09Ri12 *
	-13022	236			-0.2	U			GA8	1.5	12Ga45
$^{43}\text{Cl}-u$	-26090	300	-25940	70	0.3	o			GA4	1.5	00Sa21
	-25740	200			-0.7	o			TO3	1.5	90Tu01
	-25970	350			0.1	U			TO4	1.5	91Zh24
	-26010	330			0.1	o			GA5	1.5	00Sa21
	-25905	86			-0.2	o			GT1	1.5	04Ma.A
	-25894	140			-0.2	2			GA7	1.5	07Ju03
	-26361	100			1.7	C			GT2	2.5	08Kn.A
	-25941	70			0.1	2			LZ1	1.0	15Xu14
$^{43}\text{V}-u$	-19234	46				2		LZ1	1.0	13Ya03	
$^{43}\text{Ar}-^{36}\text{Ar}_{1.194}$	4387.2	5.7				2		MA6	1.0	01He29	
$^{43}\text{K}-^{39}\text{K}_{1.103}$	766.45	0.44				2		MA8	1.0	07Ya08	
$^{43}\text{Ca}(\alpha, \alpha)^{40}\text{K}$	-14	8	-9.28	0.23	0.6	U			MIT		64Sp12
$^{40}\text{Ca}(\alpha, \alpha)^{43}\text{Sc}$	-3470	30	-3522.3	1.9	-1.7	U					61Ma03
$^{40}\text{Ca}(\alpha, n)^{43}\text{Ti}$	-11169.9	10.	-11172	7	-0.2	2			Tal		67Al08
$^{41}\text{K}(\beta^-, \text{He}, p)^{43}\text{Ca}^i$	2452	30	2497	14	1.5	1	23	23 $^{43}\text{Ca}^i$	MIT		68Do02
$^{43}\text{V}^i(2p)^{41}\text{Sc}$	4320	50	4359	15	0.8	U			Lis		92Bo37
	4503	22			-6.5	B			Bor		01Gi01 *
	4348	16			0.7	1	89	89 $^{43}\text{V}^i$	Bor		07Do17
$^{42}\text{Ca}(n, \gamma)^{43}\text{Ca}$	7933.1	0.5	7932.89	0.17	-0.4	-					69Ar.A Z
	7933.1	0.5			-0.4	-			Ptn		69Gr08 Z
	7933.1	0.4			-0.5	-					71Bi.A
	7932.73	0.23			0.7	-			Bdn		06Fi.A
$^{42}\text{Ca}(d, p)^{43}\text{Ca}$	5716	10	5708.33	0.17	-0.8	U			MIT		64Sp12
	5707	12			0.1	U			MIT		66Do02
$^{43}\text{Ca}(d, t)^{42}\text{Ca}$	-1672	10	-1675.67	0.17	-0.4	U			Ald		64Bj02
$^{42}\text{Ca}(n, \gamma)^{43}\text{Ca}$	ave.	7932.89	0.17	7932.89	0.17	0.0	1	99	99 $^{43}\text{Ca}$		average
	$^{42}\text{Ca}(p, \gamma)^{43}\text{Sc}$	4935	5	4929.8	1.9	-1.0	2				65Br31
$^{42}\text{Ca}(\beta^-, \text{He}, d)^{43}\text{Sc}^i$	4929	2			0.4	2					69Wa19
	-4808	8	-4795	3	1.6	1	17	17 $^{43}\text{Sc}^i$			66Sc17 *
$^{43}\text{V}^i(p)^{42}\text{Ti}$	8200	45	8111	15	-2.0	1	11	11 $^{43}\text{V}^i$	Bor		01Gi01 *
$^{43}\text{K}(\beta^-)^{43}\text{Ca}$	1817	20	1833.4	0.5	0.8	U					54Li24 *
	1815	10			1.8	U					59Be72 *
$^{43}\text{Sc}(\beta^+)^{43}\text{Ca}$	2200	20	2220.7	1.9	1.0	U					52Ha44
	2220	10			0.1	U					54Li42
	-3005	10	-3003.1	1.9	0.2	U			Har		60Mc12 Y
$^{43}\text{Ca}(p, n)^{43}\text{Sc}$	-2998	10			-0.5	U					67Mc07
	-6467	8	-6471	3	-0.5	-					71Al19 *
$^{43}\text{Ca}(\beta^-, \text{He}, t)^{43}\text{Sc}^i$	-6469	4			-0.5	-					71Be29 *
	ave.	-6469	4		-0.7	1	83	83 $^{43}\text{Sc}^i$			average
	$^{43}\text{S}-u$	For original doublet $^{43}\text{S}-(\text{C}_3 \text{H}_5 \text{O})_{0.754}$ , $D_M=-38753(22) \mu\text{u}$									
$^{43}\text{S}-u$	For original doublet $^{43}\text{S C H}-(\text{C}_3 \text{H}_5 \text{O})_{0.982}$ , $D_M=-38694.8(5.5) \mu\text{u}$										
$^{43}\text{V}^i(2p)^{41}\text{Sc}$	$E_{2p}=4292(22) + \text{recoil}=211$ ; data reanalysed and included in 07Do17										
$^{42}\text{Ca}(\beta^-, \text{He}, d)^{43}\text{Sc}^i$	IT=4238(8); $Q$ rebuilt with Ame1961										
$^{43}\text{V}^i(p)^{42}\text{Ti}$	$Q_p=4590(45)$ followed by $\gamma$ 's 1938+1554 keV + recoil=118 keV										
$^{43}\text{K}(\beta^-)^{43}\text{Ca}$	$E_{\beta^-}=827(20) 825(10)$ respectively, to $3/2^+$ level at 990.257 keV										
$^{43}\text{Ca}(\beta^-, \text{He}, t)^{43}\text{Sc}^i$	IT=4226(8); $Q$ rebuilt with Ame1965										
$^{43}\text{Ca}(\beta^-, \text{He}, t)^{43}\text{Sc}^i$	CDE=7238(4) $Q=-6474(4)$ ; recalibration +6 keV for $^{42}\text{Ca}(p, n)^{42}\text{Sc}$ from Ame1961										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{44}\text{P}-u$	10070	966	11220#	540#	0.8	D			GA7	1.5	07Ju03 *
$^{44}\text{S}-u$	-10510	580	-9881	6	0.7	o			GA4	1.5	00Sa21
	-8960	620			-1.0	o			GA5	1.5	00Sa21
	-9769	150			-0.5	U			GA7	1.5	07Ju03
	-10027	301			0.3	U			GA8	1.5	12Ga45
$^{44}\text{S}-\text{C}_2\text{H}_4\text{O}$	-36095.9	5.6				2			MS1	1.0	09Ri12 *
$^{44}\text{Cl}-u$	-21700	130	-21880	150	-0.9	2			GA4	1.5	00Sa21
	-21500	500			-0.5	U			TO3	1.5	90Tu01
	-21450	270			-1.1	U			TO4	1.5	91Zh24
	-22150	370			0.5	2			GA5	1.5	00Sa21
	-22115	161			1.0	2			GT1	1.5	04Ma.A
$^{44}\text{Ca}-u$	-44515.3	10.0	-44518.5	0.3	-0.3	U			MR1	1.0	15Wi.A
$^{44}\text{Sc}-u$	-40480	410	-40597.1	1.9	-0.2	U			TO6	1.5	98Ba.A *
$^{44}\text{V}-u$	-25890	130				2			GT1	1.5	04St05 *
$^{44}\text{Ar}-^{39}\text{K}_{1.128}$	5862.9	1.7				2			MA8	1.0	03B117
$^{44}\text{K}-^{39}\text{K}_{1.128}$	2526.07	0.45				2			MA8	1.0	07Ya08
	2529.2	2.2	2526.1	0.5	-1.4	o			TT1	1.0	10La.A
	2529.1	1.7			-1.8	U			TT1	1.0	12La05
$^{40}\text{Ca}(\alpha,\gamma)^{44}\text{Ti}$	5127.1	0.7				2					82Di05
$^{44}\text{Ca}(\text{p},\alpha)^{41}\text{K}$	-1058	10	-1045.1	0.3	1.3	U			MIT		64Sp12
$^{41}\text{K}(\alpha,\text{n})^{44}\text{Sc}$	-3420	60	-3390.0	1.8	0.5	U			Yal		61Sm05
$^{44}\text{Ca}(\text{d},\alpha)^{42}\text{K}$	4273	20	4264.2	0.3	-0.4	U					77Pa24
$^{42}\text{Ca}(\text{t},\text{p})^{44}\text{Ca}$	10593	15	10582.27	0.29	-0.7	U			Ald		67Bj06
$^{42}\text{Ca}(\text{}^3\text{He},\text{p})^{44}\text{Sc}$	6920	20	6911.0	1.7	-0.5	U			Hei		70Sc22
$^{43}\text{Ca}(\text{n},\gamma)^{44}\text{Ca}$	11130.6	0.5	11131.17	0.23	1.1	-					69Ar.A Z
	11130.1	0.7			1.5	-					72Wh02 Z
	11131.54	0.29			-1.3	-			Bdn		06Fi.A
$^{43}\text{Ca}(\text{d},\text{p})^{44}\text{Ca}$	8922	14	8906.60	0.23	-1.1	U			MIT		64Sp12
	8920	10			-1.3	U			Kop		67Bj02
$^{44}\text{Ca}(\text{}^3\text{He},\alpha)^{43}\text{Ca}$	9452	15	9446.45	0.23	-0.4	U			MIT		71Ra35
$^{43}\text{Ca}(\text{n},\gamma)^{44}\text{Ca}$	ave. 11131.17	0.24	11131.17	0.23	0.0	1	99	98 $^{44}\text{Ca}$			average
$^{44}\text{Ca}(\text{p},\text{d})^{43}\text{Ca}^i$	-16880	30	-16901	14	-0.7	-					72Ma23 *
$^{44}\text{Ca}(\text{d},\text{t})^{43}\text{Ca}^i$	-12858.7	19.7	-12869	14	-0.5	-					76Do05 *
$^{44}\text{Ca}(\text{p},\text{d})^{43}\text{Ca}^i$	ave. -16888	16	-16901	14	-0.8	1	77	77 $^{43}\text{Ca}^i$			average
$^{43}\text{Ca}(\text{p},\gamma)^{44}\text{Sc}$	6694	2	6696.1	1.7	1.1	2					71Po.A
$^{43}\text{Ca}(\text{}^3\text{He},\text{d})^{44}\text{Sc}^i$	-1583	5	-1575.1	2.5	1.6	1	24	24 $^{44}\text{Sc}^i$			68Sc15
$^{44}\text{V}^i(\text{p})^{43}\text{Ti}$	950	50	908	11	-0.8	U			Lis		92Bo37
	908	11				3			Bor		07Do17
	917	53			-0.2	U					14Po05
$^{44}\text{K}(\beta^-)^{44}\text{Ca}$	5580	80	5687.2	0.5	1.3	U					70Le05
$^{44}\text{Ca}(\text{t},\text{}^3\text{He})^{44}\text{K}$	-5660	40	-5668.6	0.5	-0.2	U			LAl		70Aj01
$^{44}\text{Sc}(\beta^+)^{44}\text{Ca}$	3642	5	3652.7	1.8	2.1	R					50Br52 *
	3650	5			0.5	R					55B123 *
$^{44}\text{Ca}(\text{p},\text{n})^{44}\text{Sc}$	-4410	15	-4435.0	1.8	-1.7	U			Har		60Mc12 Y
	-4447	10			1.2	U					67Mc07
$^{44}\text{Ca}(\text{}^3\text{He},\text{t})^{44}\text{Sc}^i$	-6444	4	-6449.0	2.5	-1.3	-					71Be29 *
	-6449	4			0.0	-					72Ma50 *
	ave. -6446.5	2.8			-0.9	1	76	76 $^{44}\text{Sc}^i$			average
$^{*44}\text{P}-u$	Trends from Mass Surface TMS suggest $^{44}\text{P}$ 1070 less bound										
$^{*44}\text{S}-\text{C}_2\text{H}_4\text{O}$	For original doublet $^{44}\text{S C H}-\text{C}_3\text{H}_5\text{O}$										
$^{*44}\text{Sc}-u$	$M-A=-37570(370)$ keV for mixture gs+p at 271.240 keV										
$^{*44}\text{V}-u$	$M-A=-23980(80)$ keV for mixture gs+m at 270#100 keV										
$^{*44}\text{V}-u$	Authors have unduely increased the lower error to 380 keV										
$^{*44}\text{Ca}(\text{p},\text{d})^{43}\text{Ca}^i$	IT=7970(30); $Q$ rebuilt with Ame1965										
$^{*44}\text{Ca}(\text{d},\text{t})^{43}\text{Ca}^i$	IT=7980(20); $Q$ rebuilt with Ame1971										
$^{*43}\text{Ca}(\text{}^3\text{He},\text{d})^{44}\text{Sc}^i$	IT=2796(5); $Q$ rebuilt with Ame1965										
$^{*44}\text{Sc}(\beta^+)^{44}\text{Ca}$	$E_{\beta^+}=1463(5)$ 1471(5) respectively, to $2^+$ level at 1157.019 keV										
$^{*44}\text{Ca}(\text{}^3\text{He},\text{t})^{44}\text{Sc}^i$	CDE=7214(4) $Q=-6450(4)$ ; recalibration +6 keV for $^{42}\text{Ca}(\text{p},\text{n})^{42}\text{Sc}$ from Ame1961										
$^{*44}\text{Ca}(\text{}^3\text{He},\text{t})^{44}\text{Sc}^i$	IT=2781(5); $Q$ rebuilt with Ame1971										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{45}\text{S}-u$	-3610	2460	-4280	1110	-0.2	o			GA4	1.5	00Sa21
	-3330	2880			-0.2	o			GA5	1.5	00Sa21
	-4283	741				2			GA7	1.5	07Ju03
$^{45}\text{Cl}-u$	-19690	140	-19610	150	0.4	o			GA4	1.5	00Sa21
	-20300	700			0.7	U			TO3	1.5	90Tu01
	-19850	460			0.4	o			GA5	1.5	00Sa21
	-19710	107			0.7	2			GA7	1.5	07Ju03
	-19098	236			-1.4	2			GA8	1.5	12Ga45
$^{45}\text{V}-u$	-34225.7	9.7	-34231.0	0.9	-0.6	U			LZ1	1.0	11Tu09
$^{45}\text{Cr}-u$	-20390	540	-20950	40	-0.7	U			GT1	1.5	04St05 *
	-20950	38				2			LZ1	1.0	12Zh34 *
$^{45}\text{Ar}-^{39}\text{K}_{1.154}$	9922.45	0.55				2			MA8	1.0	03B117
$^{45}\text{K}-^{39}\text{K}_{1.154}$	2574.21	0.56				2			MA8	1.0	07Ya08
$^{45}\text{V}-^{45}\text{Ti}$	7647.74	0.23	7647.74	0.23	0.0	1	100	100 $^{45}\text{V}$	JY1	1.0	14Ka22
$^{45}\text{Sc}(p,\alpha)^{42}\text{Ca}$	2343	8	2339.4	0.7	-0.4	U			MIT		64Sp12
$^{45}\text{Sc}(d,\alpha)^{43}\text{Ca}$	8028	12	8047.7	0.7	1.6	U			MIT		64Sp12
	8059	12			-0.9	U			Kop		67Ha.A
$^{43}\text{Ca}(^3\text{He},p)^{45}\text{Sc}$	10310	20	10305.3	0.7	-0.2	U			Hei		70Sc22
$^{45}\text{Fe}(2p)^{43}\text{Cr}$	1140	40	1154	16	0.3	o					02Gi09
	1100	100			0.5	U					02Pf02
	1154	16				3					05Do20
$^{44}\text{Ca}(n,\gamma)^{45}\text{Ca}$	7414.8	1.0	7414.82	0.17	0.0	U					69Ar.A Z
	7414.83	0.3			0.0	-			MMn		80Is02 Z
	7414.79	0.21			0.1	-			Bdn		06Fi.A
$^{44}\text{Ca}(d,p)^{45}\text{Ca}$	5184	4	5190.25	0.17	1.6	U			MIT		68Be36
$^{44}\text{Ca}(n,\gamma)^{45}\text{Ca}$	ave.	7414.80	0.17	7414.82	0.17	0.1	1	99	97 $^{45}\text{Ca}$		average
$^{44}\text{Ca}(p,\gamma)^{45}\text{Sc}$	6887.8	1.2	6892.2	0.7	3.7	B					74Sc02 Z
$^{45}\text{Sc}(^3\text{He},\alpha)^{44}\text{Sc}$	9249	15	9250.4	1.9	0.1	U			MIT		71Ra09
$^{45}\text{Sc}(d,t)^{44}\text{Sc}^i$	-7846	10	-7847.7	2.6	-0.2	U					71Oh01 *
$^{45}\text{V}^i(p)^{44}\text{Ti}$	3190	50	3170	9	-0.4	U					74Ja10 *
	3170	9				3			Bor		07Do17 *
$^{45}\text{K}(\beta^-)^{45}\text{Ca}$	4180	200	4196.5	0.6	0.1	U					64Mo18
$^{45}\text{Ca}(\beta^-)^{45}\text{Sc}$	258	2	259.7	0.7	0.9	1	14	11 $^{45}\text{Sc}$			65Fr12
$^{45}\text{Ti}(\beta^+)^{45}\text{Sc}$	2066	5	2062.1	0.5	-0.8	U					66Po04
$^{45}\text{Sc}(p,n)^{45}\text{Ti}$	-2844.2	4.	-2844.4	0.5	-0.1	U			Ric		55Br16 Y
	-2843.6	4.0			-0.2	U			Can		70Kn03
	-2844.4	0.5			0.0	1	100	100 $^{45}\text{Ti}$	PTB		85Sc16 Z
$^{45}\text{Sc}(^3\text{He},t)^{45}\text{Ti}^i$	-6801	4	-6800	3	0.3	1	61	60 $^{45}\text{Ti}^i$			71Be29 *
* $^{45}\text{Cr}-u$	$M - A = -18940(500)$ keV for mixture gs+m at 107(1) keV										
* $^{45}\text{Cr}-u$	Same result in reference										
* $^{45}\text{Sc}(d,t)^{44}\text{Sc}^i$	IT=2784(10) combined with $Q = -5062$ ; $Q$ rebuilt										
* $^{45}\text{V}^i(p)^{44}\text{Ti}$	$Q_p = 2060(50)$ 2087(9) respectively, to $2^+$ level at 1083.06 keV										
* $^{45}\text{Sc}(^3\text{He},t)^{45}\text{Ti}^i$	CDE=7571(4) $Q = -6807(4)$ ; recalibration +6 keV for $^{42}\text{Ca}(p,n)^{42}\text{Sc}$ from Ame1961										
$^{46}\text{Cl}-u$	-16000	860	-14880	220	0.9	o			GA4	1.5	00Sa21
	-14940	1730			0.0	o			GA5	1.5	00Sa21
	-14826	172			-0.2	2			GA7	1.5	07Ju03
	-15040	301			0.4	2			GA8	1.5	12Ga45
$^{46}\text{Ar}-u$	-32013	107	-31962.6	1.2	0.3	U			GT1	1.5	04Ma.A
$^{46}\text{Sc}-u$	-44650	230	-44832.5	0.7	-0.5	U			TO6	1.5	98Ba.A *
$\text{C}_2 \text{H}_8 \text{N}-^{46}\text{Ti}$	113071	7	113047.41	0.18	-0.8	U			R09	4.0	72De11
$\text{C}^{13} \text{C} \text{H}_5 \text{O}-^{46}\text{Ti}$	84799	13	84767.76	0.18	-0.6	U			R09	4.0	72De11
$\text{C} \text{H}_4 \text{N} \text{O}-^{46}\text{Ti}$	76672	8	76661.90	0.18	-0.3	U			R09	4.0	72De11
$\text{C}_5 \text{H}_2-^{46}\text{Ti} \text{O}$	68145	15	68108.59	0.18	-0.6	U			R09	4.0	72De11
$\text{C} \text{H}_2 \text{O}_2-^{46}\text{Ti}$	52881	14	52852.45	0.18	-0.5	U			R09	4.0	72De11



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{13}\text{C H O}_2-^{46}\text{Ti}$	48423	9	48382.25	0.18	-1.1	U	R09	4.0	72De11
$^{46}\text{Ti}-^{22}\text{Ne}_{2,091}$	-29358.74	0.48	-29359.41	0.18	-1.4	1	14	13	$^{46}\text{Ti}$ CP1 1.0 05Sa44
$^{46}\text{V}-^{22}\text{Ne}_{2,091}$	-21787.10	0.58	-21788.29	0.22	-2.1	1	14	14	$^{46}\text{V}$ CP1 1.0 05Sa44
$^{46}\text{Cr}-\text{u}$	-31638	15	-31639	12	-0.1	1	67	67	$^{46}\text{Cr}$ LZ1 1.0 16Zh.A
$^{46}\text{Ar}-^{39}\text{K}_{1,179}$	10827.5	1.2				2			MA8 1.0 15Mo.A
$^{46}\text{K}-^{39}\text{K}_{1,179}$	4771.64	0.78				2			MA8 1.0 07Ya08
$^{46}\text{V}-^{46}\text{Ti}$	7571.67	0.41	7571.12	0.10	-1.4	U			CP1 1.0 05Sa44
	7571.41	0.33			-0.9	o			JY1 1.0 06Er08
	7571.10	0.11			0.1	1	83	100	$^{46}\text{V}$ JY1 1.0 11Er02
$^{32}\text{S}(^{16}\text{O},2\text{n})^{46}\text{Cr}$	-17421.6	20.	-17424	11	-0.1	1	33	33	$^{46}\text{Cr}$ 72Zi02
$^{46}\text{Ti}(\text{p},\alpha)^{43}\text{Sc}$	-3065	14	-3075.6	1.9	-0.8	U			MIT 64Sp12 *
	-3083	10			0.7	U			Tal 65Pl01
$^{46}\text{Ti}(^3\text{He},^6\text{He})^{43}\text{Ti}$	-17470	12	-17468	7	0.2	R			MSU 77Mu03 *
$^{44}\text{Ca}(\text{t},\text{p})^{46}\text{Ca}$	9339	20	9331.5	2.3	-0.4	U			Kop 67Bj06
$^{44}\text{Ca}(^3\text{He},\text{p})^{46}\text{Sc}$	7940	20	7934.8	0.7	-0.3	U			Hei 70Sc22
$^{46}\text{Ti}(\text{d},\alpha)^{44}\text{Sc}$	4400	12	4399.0	1.8	-0.1	U			Kop 67Ha.A
$^{46}\text{Ti}(\text{p},\text{t})^{44}\text{Ti}$	-14235	10	-14240.1	0.7	-0.5	U			Oak 72Ra05
$^{46}\text{Ca}(\text{t},\alpha)^{45}\text{K}$	5998	10	6001.2	2.3	0.3	U			Ald 68Sa09
$^{46}\text{Ca}(\text{d},\text{t})^{45}\text{Ca}$	-4144	10	-4141.3	2.3	0.3	U			Ald 67Bj05
$^{46}\text{Ca}(^3\text{He},\alpha)^{45}\text{Ca}$	10194	10	10179.1	2.3	-1.5	U			MIT 71Ra35
$^{45}\text{Sc}(\text{n},\gamma)^{46}\text{Sc}$	8760.61	0.3	8760.64	0.10	0.1	2			BNn 80Li07 Z
	8760.58	0.14			0.4	2			Utr 82Ti02 Z
	8760.75	0.18			-0.6	2			Bdn 06Fi.A
$^{45}\text{Sc}(\text{d},\text{p})^{46}\text{Sc}$	6541	8	6536.07	0.10	-0.6	U			MIT 64Sp12
	6543	8			-0.9	U			Kop 67Ha.A
$^{45}\text{Sc}(\text{p},\gamma)^{46}\text{Ti}$	10344.7	0.7	10344.9	0.7	0.2	1	89	88	$^{45}\text{Sc}$ 71Gu.A
$^{46}\text{Ti}(\text{p},\text{d})^{45}\text{Ti}^i$	-15682	5	-15684	3	-0.4	1	40	40	$^{45}\text{Ti}^i$ 78Ko27
$^{46}\text{Cr}^i(\text{p})^{45}\text{V}$	4350	50	4269	13	-1.6	U			Lis 92Bo37
	4269	13				2			Bor 07Do17 *
$^{46}\text{Mn}^i(\text{p})^{45}\text{Cr}$	3520	100	4840	30	13.2	B			Lis 92Bo37
	4840	33				3			Bor 07Do17 *
$^{46}\text{K}(\beta^-)^{46}\text{Ca}$	7650	300	7725.4	2.4	0.3	U			
$^{46}\text{Ca}(^3\text{He},\text{t})^{46}\text{Sc}^i$	-6407	4	-6410	3	-0.8	1	72	63	$^{46}\text{Sc}^i$ 66Pa20
$^{46}\text{Sc}(\beta^-)^{46}\text{Ti}$	2367	3	2366.6	0.7	-0.1	U			
	2364	6			0.4	U			
$^{46}\text{Ti}(\text{p},\text{n})^{46}\text{V}$	-7844	9	-7834.80	0.09	1.0	U			Tal 63Ja12
	-7835.8	1.8			0.6	U			Har 76Sq01 Z
$^{46}\text{Ti}(^3\text{He},\text{t})^{46}\text{V}$	-7069.0	0.6	-7071.04	0.09	-3.4	F			Mun 77Vo02 *
$^{46}\text{Ti}(^3\text{He},\text{t})^{46}\text{V}-^{27}\text{Al}(^27\text{Si})$	-2230.8	2.7	-2240.09	0.13	-3.4	B			ChR 74Ha35
$^{46}\text{Ti}(^3\text{He},\text{t})^{46}\text{V}-^{47}\text{Ti}(^47\text{V})$	-4121.62	0.19	-4121.70	0.14	-0.4	1	56	39	$^{47}\text{V}$ Mun 09Fa15 *
$^{46}\text{Ti}(^3\text{He},\text{t})^{46}\text{V}-^{48}\text{Ti}(^48\text{V}^i)$	-18.57	0.20	-18.58	0.20	0.0	1	100	100	$^{48}\text{V}^i$ Mun 09Fa15
$^{46}\text{Ti}(^3\text{He},\text{t})^{46}\text{V}-^{50}\text{Ti}(^50\text{V}^i)$	-31.21	0.25	-31.21	0.25	0.0	1	100	100	$^{50}\text{V}^i$ Mun 09Fa.A
$^{46}\text{Sc}-\text{u}$	$M-A=-41520(210)$ keV for mixture gs+n at 142.528 keV								Nub16c **
$^{46}\text{Ti}(\text{p},\alpha)^{43}\text{Sc}$	$Q=-3217$ probably to $^{43}\text{Sc}^m$ at 151.79 keV								Nub16b **
$^{46}\text{Ti}(^3\text{He},^6\text{He})^{43}\text{Ti}$	Averaged with reference ; $Q$ reduced by 3 for recalibration $^{27}\text{Al}(^3\text{He},^6\text{He})$								75Mu09 **
$^{46}\text{Cr}^i(\text{p})^{45}\text{V}$	$Q_p=4254(15)$ 3494(25) 3003(13) to ground state, $(5/2^+)$ level at 797.2, $(7/2^+)$ at 1272.2 keV								MMC128**
*									Ens082 **
$^{46}\text{Mn}^i(\text{p})^{45}\text{Cr}$	$Q_p=4239(33)$ to $(5/2^+)$ state at 493.6 + 107(1)								11Ho02 **
$^{46}\text{Ca}(^3\text{He},\text{t})^{46}\text{Sc}^i$	CDE=7177(4) $Q=-6413(4)$ ; recalibration +6 keV for $^{42}\text{Ca}(\text{p},\text{n})^{42}\text{Sc}$ from Ame1961								MMC123**
$^{46}\text{Sc}(\beta^-)^{46}\text{Ti}$	$E_{\beta^-}=357(3)$ to $4^+$ level at 2009.846 keV								Ens00b **
$^{46}\text{Sc}(\beta^-)^{46}\text{Ti}$	$E_{\beta^-}=1475(6)$ to $2^+$ level at 889.286 keV								Ens00b **
$^{46}\text{Ti}(^3\text{He},\text{t})^{46}\text{V}$	F : rejected in reference of same group								09Fa15 **
$^{46}\text{Ti}(^3\text{He},\text{t})^{46}\text{V}-^{47}\text{Ti}(^47\text{V})$	$Q-Q=28.73(0.16)$ keV to $^{47}\text{V}^i$ IAS at 4150.35(0.11) keV								Ens075 **
$^{47}\text{Cl}-\text{u}$	-9576	1074	-10500#	430#	-0.6	D			GA7 1.5 07Ju03 *
$^{47}\text{Ar}-\text{u}$	-25400	600	-27231.9	1.2	-2.0	U			TO3 1.5 90Tu01
	-26570	1360			-0.3	U			GA5 1.5 00Sa21
	-26903	204			-1.1	U			GA8 1.5 12Ga45

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{47}\text{Sc}-u$	-47630	230	-47597.3	2.1	0.1	U			TO6	1.5	98Ba.A *
$\text{C }^{35}\text{Cl}-^{47}\text{Ti}$	17085.94	0.82	17094.94	0.13	4.4	B			H32	2.5	79Ko10
$\text{C }^{13}\text{C H}_8 \text{N}-^{47}\text{Ti}$	117329	14	117271.34	0.12	-1.0	U			R09	4.0	72De11
$\text{C}_2 \text{H}_7 \text{O}-^{47}\text{Ti}$	98012	7	97932.09	0.12	-2.9	B			R09	4.0	72De11
$\text{C}_5 \text{H}_3 -^{47}\text{Ti O}$	76869	10	76802.72	0.12	-1.7	U			R09	4.0	72De11
$\text{C H}_3 \text{O}_2 -^{47}\text{Ti}$	61608	10	61546.58	0.12	-1.5	U			R09	4.0	72De11
$^{47}\text{Cr}-u$	-37103.8	8.6	-37104	6	-0.1	1	57	$^{57} ^{47}\text{Cr}$	LZ1	1.0	11Tu09
$^{47}\text{Mn}-u$	-24226	34				2			LZ1	1.0	13Ya03
$^{47}\text{Ar}-^{39}\text{K}_{1,205}$	16501.8	1.2				2			MA8	1.0	15Mo.A
$^{47}\text{K}-^{39}\text{K}_{1,205}$	5398.5	2.7	5395.3	1.5	-1.2	o			TT1	1.0	10La.A
	5395.3	1.5				2			TT1	1.0	12La05
$^{46}\text{Ti }^{13}\text{C}-^{47}\text{Ti C}$	4218.03	0.94	4223.94	0.14	2.5	U			H32	2.5	79Ko10
$^{47}\text{Ti}-^{46}\text{Ti}$	-929	41	-869.10	0.14	0.4	U			R09	4.0	72De11
$^{47}\text{Ti}(d,\alpha)^{45}\text{Sc}$	6830	12	6845.3	0.7	1.3	U			Kop		67Ha.A
$^{46}\text{Ar}(d,p)^{47}\text{Ar}$	1327	80	1440.2	1.6	1.4	U					06Ga28
$^{46}\text{Ca}(n,\gamma)^{47}\text{Ca}$	7277.4	0.6	7276.37	0.27	-1.7	-					70Cr04 Z
	7276.1	0.3			0.9	-			Bdn		06Fi.A
$^{46}\text{Ca}(d,p)^{47}\text{Ca}$	5055	8	5051.81	0.27	-0.4	U			Kop		67Ha.A
	5044	4			2.0	U			MIT		68Be36
$^{46}\text{Ca}(n,\gamma)^{47}\text{Ca}$	ave.	7276.36	0.27	7276.37	0.27	0.1	1	100	$^{90} ^{46}\text{Ca}$		average
$^{46}\text{Ti}(n,\gamma)^{47}\text{Ti}$	8875.1	3.0	8880.88	0.13	1.9	U					69Te01 Z
	8880.5	0.3			1.3	1	20	$^{18} ^{46}\text{Ti}$	Bdn		06Fi.A
$^{46}\text{Ti}(d,p)^{47}\text{Ti}$	6658	6	6656.32	0.13	-0.3	U			MIT		67Ba32 *
	6659	8			-0.3	U			Kop		67Ba32 *
	6654.3	1.7			1.2	U			NDm		76Jo01
$^{46}\text{Ti}(d,p)^{47}\text{Ti}-^{48}\text{Ti}(^{49}\text{Ti})$	738.15	0.25	738.49	0.14	1.3	1	30	$^{25} ^{46}\text{Ti}$	Mun		09Fa15
$^{46}\text{Ti}(p,\gamma)^{47}\text{V}$	5167.80	0.07	5167.79	0.07	-0.2	1	94	$^{61} ^{47}\text{V}$	Utr		86De13 *
$^{46}\text{Ti}(^3\text{He},d)^{47}\text{V}$	-317	15	-325.69	0.07	-0.6	U			MIT		67Do03
$^{47}\text{Mn}^i(p)^{46}\text{Cr}$	6867	20	6992	13	6.2	B			Bor		01Gi01 *
	6992	13				2			Bor		07Do17 *
$^{47}\text{K}(\beta^-)^{47}\text{Ca}$	6700	300	6632.4	2.6	-0.2	U					64Ku02 *
$^{47}\text{Ca}(\beta^-)^{47}\text{Sc}$	1984.6	5.	1992.2	1.2	1.5	U					67Hs03 *
	1992.3	5.			0.0	U					68Fi04 *
	1991.9	1.2			0.2	1	97	$^{91} ^{47}\text{Ca}$			87Ju04
$^{47}\text{Sc}(\beta^-)^{47}\text{Ti}$	600	2	600.8	1.9	0.4	1	93	$^{93} ^{47}\text{Sc}$			56Gr12
$^{47}\text{V}(\beta^+)^{47}\text{Ti}$	2912	10	2930.75	0.14	1.9	U					54Da31
$^{47}\text{Ti}(p,n)^{47}\text{V}$	-3706	13	-3713.09	0.14	-0.5	U			Har		60Mc12 Y
$^{*47}\text{Cl}-u$	Trends from Mass Surface TMS suggest $^{47}\text{Cl}$ 860 more bound										
$^{*47}\text{Sc}-u$	$M - A = -44320(210)$ keV for mixture gs+m at 766.83 keV and										
*	assuming ratio $R=0.07(3)$ , from half-life=272 ns and TOF=1 $\mu\text{s}$										
$^{*46}\text{Ti}(d,p)^{47}\text{Ti}$	All 67Ba32 results decreased 0.2% for recalibration										
$^{*46}\text{Ti}(p,\gamma)^{47}\text{V}$	$E_p=985.94(0.05,Z)$ to $1/2^+$ level at 6132.60(0.09) keV										
$^{*47}\text{Mn}^i(p)^{46}\text{Cr}$	$Q_p=5975(25)$ 4880(20) to $2^+$ level at 892.16 and ( $4^+$ ) at 1987.1 keV;										
*	data reanalysed and included in 07Do17										
$^{*47}\text{Mn}^i(p)^{46}\text{Cr}$	$Q_p=6104(24)$ 5000(15) to $2^+$ level at 892.16 and ( $4^+$ ) at 1987.1 keV										
*	also tentatively $Q_p=3973(20)$ to ( $3^-$ ) at 3196.5 keV, not used										
$^{*47}\text{K}(\beta^-)^{47}\text{Ca}$	$E_{\beta^-}=4100(300)$ to 2578.33 $3/2^+$ and 2599.53 $1/2^+$ levels										
$^{*47}\text{Ca}(\beta^-)^{47}\text{Sc}$	Original values increased by 4(4) for shape factor										
											87Ju04 **
$^{48}\text{Ar}-u$	-23920	330				2			MT1	1.0	15Me01
$^{48}\text{K}-^{39}\text{K}_{1,231}$	10017.7	2.5	10018.5	0.8	0.3	o			TT1	1.0	10La.A
	10018.50	0.83				2			TT1	1.0	12La05
$^{48}\text{Ca}-^{39}\text{K}_{1,231}$	-2799.93	0.22	-2799.78	0.10	0.7	1	22	$^{22} ^{48}\text{Ca}$	MS1	1.0	12Re17
$^{48}\text{Ca}-\text{N }^{18}\text{O O}$	-44625.15	0.29	-44625.33	0.10	-0.6	1	13	$^{13} ^{48}\text{Ca}$	TT1	1.0	14Kw04
$^{13}\text{C }^{35}\text{Cl}-^{48}\text{Ti}$	24261.73	0.75	24266.60	0.12	2.6	U			H32	2.5	79Ko10

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_5 H_4 -^{48}Ti O$	88492	24	88444.58	0.12	-0.5	U			R09	4.0	72De11
	88494	27			-0.5	U			R09	4.0	72De11
$C_4 H_2 N -^{48}Ti O$	75935	17	75868.52	0.12	-1.0	U			R09	4.0	72De11
$C_4 -^{48}Ti$	52109	19	52059.07	0.12	-0.7	U			R09	4.0	72De11
$^{48}Ti O -^{85}Rb_{.753}$	9277.7	1.2	9277.88	0.12	0.2	U			MA8	1.0	12Na15
$^{48}Ti - N^{18}O O$	-49207.30	0.23	-49207.30	0.12	0.0	1	26	26 $^{48}Ti$	TT1	1.0	14Kw04
$^{48}Mn - u$	-31480	120	-31451	7	0.2	U			GT1	1.5	04St05
	-31454	10			0.3	1	55	55 $^{48}Mn$	LZ1	1.0	16Zh.A
$^{48}Ca - ^{40}Ca_{1.200}$	-2586.23	0.23	-2586.13	0.11	0.4	1	21	20 $^{48}Ca$	MS1	1.0	12Re17
$^{48}Ca - ^{41}K_{1.171}$	-2774.65	0.22	-2774.47	0.10	0.8	1	22	22 $^{48}Ca$	MS1	1.0	12Re17
$^{48}Ti O - ^{55}Mn_{1.164}$	14972.6	1.2	14973.3	0.4	0.6	1	11	10 $^{55}Mn$	MA8	1.0	12Na15
$^{46}Ti - ^{37}Cl - ^{48}Ti - ^{35}Cl$	1726.8	1.1	1735.81	0.16	2.0	U			H18	4.0	64Ba03
	1730.29	0.87			2.5	U			H32	2.5	79Ko10
$^{48}Ti - ^{48}Ca$	-4582.018	0.086	-4581.97	0.08	0.5	1	88	65 $^{48}Ti$	MS1	1.0	13Bu12
	-4581.86	0.22			-0.5	U			TT1	1.0	14Kw04
$^{48}Ti - ^{47}Ti$	-3791	48	-3816.82	0.04	-0.1	U			R09	4.0	72De11
$^{48}Ca(^3He, ^{11}C)^{40}S$	-17416	35	-17105	4	8.9	F			Pri		79Ko.B *
$^{48}Ca(^3He, ^8B)^{43}Cl$	-29070	60	-28060	60	16.9	F			MSU		76Ka24 *
$^{48}Ca(\alpha, ^9Be)^{43}Ar$	-21165.5	70.2	-21138	5	0.4	U			Brk		74Je01
$^{48}Ca(^3He, ^7Be)^{44}Ar$	-12362	20	-12389.2	1.6	-1.4	U			MSU		76Cr03 *
$^{48}Ca(\alpha, ^7Be)^{45}Ar$	-27840.4	60.2	-27797.9	0.5	0.7	U			Brk		74Je01
$^{48}Ti(p, \alpha)^{45}Sc$	-2560	5	-2556.8	0.7	0.6	U			ANL		64Yn03
	-2545	15			-0.8	U			Tal		65Pl01
$^{48}Ca(^6Li, ^8B)^{46}Ar$	-23324.8	70.2	-23286.4	1.5	0.5	U			Brk		74Je01
$^{48}Ca(^{14}C, ^{16}O)^{46}Ar$	-6739	50	-6694.8	1.1	0.9	U			Mun		80Ma40
$^{48}Ca(d, \alpha)^{46}K$	1915	15	1900.1	0.7	-1.0	U			ANL		65Ma07
$^{46}Ca(t, p)^{48}Ca$	8752	20	8746.1	2.2	-0.3	U			Ald		67Bj06
$^{48}Ti(d, \alpha)^{46}Sc$	3967	12	3979.3	0.7	1.0	U			Kop		67Ha.A
$^{48}Ti(p, ^3He)^{46}Sc^i$	-19394	6	-19387	4	1.2	1	37	37 $^{46}Sc^i$			78Ko27
$^{48}Ti(p, t)^{46}Ti^i$	-21192	7				2					78Ko27
$^{48}Ti(p, t)^{46}Ti^j$	-26177	6				2					78Ko27
$^{46}Ti(^3He, n)^{48}Cr$	5550	18	5554	7	0.2	R			CIT		67Mi02
$^{48}Ni(2p)^{46}Fe$	1400	100	1310	40	-0.9	3					05Gi15 *
	1280	120			0.2	o					11Po09
	1280	60			0.4	o					12Po03 *
	1290	40			0.4	3					14Po05
$^{48}Ca(^{14}C, ^{15}O)^{47}Ar$	-18142	100	-18694.0	1.2	-5.5	B			MSU		85Be50
$^{48}Ca(d, ^3He)^{47}K$	-10304	12	-10308.1	1.4	-0.3	U			ANL		66Ne01
$^{48}Ca(t, \alpha)^{47}K$	4006	15	4012.2	1.4	0.4	U			LAI		66Wi11
	4001	10			1.1	U			Ald		68Sa09
$^{48}Ca(d, t)^{47}Ca$	-3699	10	-3694.3	2.2	0.5	U			ANL		66Er02
$^{48}Ca(^3He, \alpha)^{47}Ca$	10630	12	10626.1	2.2	-0.3	U			ANL		66Er02
	10642	10			-1.6	U			MIT		71Ra35
$^{47}Ti(n, \gamma)^{48}Ti$	11626.39	0.3	11626.66	0.04	0.9	U			MMn		80Is02 Z
	11626.65	0.04			0.3	1	99	91 $^{47}Ti$	Ptn		84Ru06 Z
	11626.66	0.23			0.0	U			Bdn		06Fi.A
$^{47}Ti(d, p)^{48}Ti$	9401	8	9402.10	0.04	0.1	U			Kop		67Ba32
	9403	6			-0.2	U			MIT		67Ba32
$^{47}Ti(^3He, d)^{48}V$	1337	15	1335.8	1.0	-0.1	U			MIT		68Do06
$^{47}Ti(^3He, d)^{48}V^i$	-1706	20	-1683.03	0.22	1.1	U					68Do06
$^{48}Mn^i(p)^{47}Cr$	979.6	32.	1014	6	1.1	o			Bor		95B105
	979.6	33.			1.0	o			Bor		96Fa09 *
	1013	12			0.1	-			Bor		07Do17 *
	1018	10			-0.4	-					16Or03
$^{48}K(\beta^-)^{48}Ca$	ave.	1016	8		-0.3	1	70	45 $^{48}Mn^i$			average
	12000	500	11940.2	0.8	-0.1	U					75Mu08

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{48}\text{Ca}(^7\text{Li}, ^7\text{Be})^{48}\text{K}$	-12959	27	-12802.0	0.8	5.8	B			Can		78We14
$^{48}\text{Ca}(^{14}\text{C}, ^{14}\text{N})^{48}\text{K}$	-11910	50	-11783.7	0.8	2.5	U			Mun		80Ma40
$^{48}\text{Ca}(\text{p}, \text{n})^{48}\text{Sc}$	-534	15	-503	5	2.1	U					67Mc07 Z
	-506	7			0.4	1	50	50 $^{48}\text{Sc}$			68Mc10
$^{48}\text{Sc}(\beta^-)^{48}\text{Ti}$	3986	7	3989	5	0.4	1	50	50 $^{48}\text{Sc}$			57Va08 *
$^{48}\text{V}(\beta^+)^{48}\text{Ti}$	4008	5	4015.0	1.0	1.4	U					53Ma64 *
	4013.6	3.			0.5	1	10	10 $^{48}\text{V}$			67Ko01 *
	4014	7			0.1	U					74Me15 *
$^{48}\text{Ti}(\text{p}, \text{n})^{48}\text{V}$	-4803	10	-4797.4	1.0	0.6	U			Tal		62Ne08 Y
$^{48}\text{Ti}(^3\text{He}, \text{t})^{48}\text{V}^i$	-7048	4	-7052.46	0.22	-1.1	U					71Be29 *
$^{48}\text{V}^i(\text{IT})^{48}\text{V}$	3018.7	1.0	3018.9	0.9	0.2	1	90	90 $^{48}\text{V}$			Ens067
$^{48}\text{Mn}^i(\text{IT})^{48}\text{Mn}$	3036.7	0.9	3036.7	0.9	0.0	1	100	55 $^{48}\text{Mn}^i$			07Do17 *
$^{*48}\text{Ca}(^3\text{He}, ^{11}\text{C})^{40}\text{S}$	F : possible $^{40}\text{Ca}$ contamination; mismatch in cross-sections										
$^{*48}\text{Ca}(^3\text{He}, ^8\text{B})^{43}\text{Cl}$	F : poor spectrum. Authors say: possibly not to ground state										
$^{*48}\text{Ca}(^3\text{He}, ^7\text{Be})^{44}\text{Ar}$	$M - A = -32270(20) Q = -12791(20)$ for $^7\text{Be}$ 429 keV level										
$^{*48}\text{Ni}(2\text{p})^{46}\text{Fe}$	From only 1 event, Si detector										
$^{*48}\text{Ni}(2\text{p})^{46}\text{Fe}$	From 4 events, gaseous detector										
$^{*48}\text{Mn}^i(\text{p})^{47}\text{Cr}$	Unexpectedly low intensity 3.6(1.1)%										
$^{*48}\text{Mn}^i(\text{p})^{47}\text{Cr}$	Measured intensity 1.8(0.3)%										
$^{*48}\text{Sc}(\beta^-)^{48}\text{Ti}$	$E_{\beta^-} = 654(7)$ to $6^+$ level at 3333.196 keV										
$^{*48}\text{V}(\beta^+)^{48}\text{Ti}$	$E_{\beta^+} = 692(5) 698(3) 698(7)$ respectively, to $4^+$ level at 2295.654 keV										
$^{*48}\text{Ti}(^3\text{He}, \text{t})^{48}\text{V}^i$	CDE=7818(4) $Q = -7054(4)$ ; recalibration +6 keV for $^{42}\text{Ca}(\text{p}, \text{n})^{42}\text{Sc}$ from Ame1961										
$^{*48}\text{Mn}^i(\text{IT})^{48}\text{Mn}$	$\gamma$ cascade 2633.5(0.5)+89.9(0.6)+313.3(0.4)										
$^{49}\text{Ar}-\text{u}$	-19110	1180	-18450#	430#	0.6	D			MT1	1.0	15Me01 *
$^{49}\text{K}-\text{u}$	-31981	225	-31789.2	0.9	0.6	U			GT1	1.5	04Ma.A
$^{49}\text{K}-^{39}\text{K}_{1.256}$	13794.9	2.8	13795.4	0.9	0.2	o			TT1	1.0	10La.A
	13795.41	0.86				2			TT1	1.0	12La05
$^{49}\text{Ca}-^{39}\text{K}_{1.256}$	1247.1	2.9	1247.53	0.22	0.1	o			TT1	1.0	10La.A
	1247.1	1.2			0.4	U			TT1	1.0	12La05
$\text{C H}_2 ^{35}\text{Cl}-^{49}\text{Ti}$	36637	13	36638.13	0.13	0.0	U			R09	4.0	72De11
$\text{C}_4 \text{H}-^{49}\text{Ti}$	59967	10	59960.40	0.12	-0.2	U			R09	4.0	72De11
$\text{C}_5 \text{H}_5-^{49}\text{Ti O}$	96348	19	96345.91	0.12	0.0	U			R09	4.0	72De11
$\text{C H}_5 ^{32}\text{S}-^{49}\text{Ti}$	63365	14	63331.71	0.12	-0.6	U			R09	4.0	72De11
$^{49}\text{Mn}-\text{u}$	-40410	12	-40387.4	2.4	1.9	U			LZ1	1.0	11Tu09
$^{49}\text{Fe}-\text{u}$	-26571	26				2			LZ1	1.0	12Zh34 *
$^{47}\text{Ti } ^{37}\text{Cl}-^{49}\text{Ti } ^{35}\text{Cl}$	946.4	1.1	943.02	0.09	-0.8	U			H18	4.0	64Ba03
	944.46	0.35			-1.7	U			H32	2.5	79Ko10
$^{48}\text{Ti } ^{13}\text{C}-^{49}\text{Ti C}$	3432.64	0.80	3431.14	0.03	-0.7	U			H32	2.5	79Ko10
$^{48}\text{Ti H}-^{49}\text{Ti}$	7876	7	7901.34	0.03	0.9	U			R09	4.0	72De11
	7874	27			0.3	U			R09	4.0	72De11
$^{49}\text{Mn}-^{49}\text{Cr}$	8279.63	0.25	8279.63	0.25	0.0	1	100	100 $^{49}\text{Mn}$	JY1	1.0	14Ka22
$^{49}\text{Ti}-^{48}\text{Ti}$	-43	36	-76.31	0.03	-0.2	U			R09	4.0	72De11
$^{49}\text{Ti}(\text{d}, \alpha)^{47}\text{Sc}$	6476	12	6483.6	1.9	0.6	U			Kop		67Ha.A
$^{48}\text{Ca}(\text{n}, \gamma)^{49}\text{Ca}$	5146.6	0.7	5146.45	0.18	-0.2	2					69Ar.A Z
	5146.38	0.30			0.2	2					70Cr04 Z
	5146.48	0.23			-0.1	2			Bdn		06Fi.A
$^{48}\text{Ca}(\text{d}, \text{p})^{49}\text{Ca}$	2917	7	2921.89	0.18	0.7	U			ANL		66Er02
	2917	4			1.2	U			MIT		68Be36
$^{48}\text{Ca}(\text{p}, \gamma)^{49}\text{Sc}$	9628.7	3.6	9625.6	2.7	-0.9	-					68Vi01 Z
$^{48}\text{Ca}(\text{d}, \text{n})^{49}\text{Sc}$	7404	7	7401.0	2.7	-0.4	-					68Gr09
$^{48}\text{Ca}(^3\text{He}, \text{d})^{49}\text{Sc}$	4150	12	4132.1	2.7	-1.5	U			ANL		66Er02
$^{48}\text{Ca}(\text{p}, \gamma)^{49}\text{Sc}$	ave.	9629	9625.6	2.7	-1.0	1	71	71 $^{49}\text{Sc}$			average
$^{48}\text{Ti}(\text{n}, \gamma)^{49}\text{Ti}$	8142.22	0.3	8142.395	0.030	0.6	U			MMn		80Is02 Z
	8142.39	0.03			0.2	1	99	100 $^{49}\text{Ti}$	Ptn		83Ru08 Z
	8142.35	0.16			0.3	U			Bdn		06Fi.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{48}\text{Ti}(\text{d,p})^{49}\text{Ti}$	5907	8	5917.829	0.030	1.4	U			Kop		67Ba32
	5918	6			0.0	U			MIT		67Ba32
	5918.6	1.7			-0.5	U			NDm		76Jo01
$^{48}\text{Ti}(\text{p},\gamma)^{49}\text{V}$	6756.8	1.5	6758.2	0.8	0.9	R					72Ki06
$^{49}\text{Mn}^i(\text{p})^{48}\text{Cr}$	2712.2	50.	2729	16	0.3	U					70Ce02 *
	2730	29			0.0	o			Bor		96Fa09 *
	2729	16				3			Bor		07Do17 *
$^{49}\text{K}(\beta^-)^{49}\text{Ca}$	10970	70	11688.3	0.8	10.3	B					86Mi08
$^{49}\text{Ca}(\beta^-)^{49}\text{Sc}$	5200	100	5261.5	2.7	0.6	U					56Ma27
	4970	50			5.8	B					56Ok02
$^{49}\text{Sc}(\beta^-)^{49}\text{Ti}$	2010	5	2002.5	2.7	-1.5	1	29	29 $^{49}\text{Sc}$			61Re06
	1983	7			2.8	B					69Fi02
$^{49}\text{V}(\epsilon)^{49}\text{Ti}$	626	10	601.9	0.8	-2.4	U					56Ha59
$^{49}\text{Ti}(\text{p,n})^{49}\text{V}$	-1383	9	-1384.2	0.8	-0.1	U			Har		60Mc12 Z
	-1383.6	1.0			-0.6	2			Oak		64Jo11 Z
$^{49}\text{Ti}(\text{}^3\text{He,t})^{49}\text{V}^i$	-7052	4				2					71Be29 *
$^{49}\text{Cr}(\beta^+)^{49}\text{V}$	2590	20	2628.9	2.4	1.9	U					53Cr18 *
* $^{49}\text{Ar}-\text{u}$	Trends from Mass Surface TMS suggest $^{49}\text{Ar}$ 610 less bound										
* $^{49}\text{Fe}-\text{u}$	Same result in reference										
* $^{49}\text{Mn}^i(\text{p})^{48}\text{Cr}$	$Q_p=1960(50)$ 1978(29) 1977(16) respectively, to $2^+$ level at 752.19(0.11) keV										
* $^{49}\text{Ti}(\text{}^3\text{He,t})^{49}\text{V}^i$	CDE=7822(4) $Q=-7058(4)$ ; recalibration +6 keV for $^{42}\text{Ca}(\text{p,n})^{42}\text{Sc}$ from Ame1961										
* $^{49}\text{Cr}(\beta^+)^{49}\text{V}$	$E_{\beta^+}=1540(10)$ 1390(20) to $(7/2^-)$ ground state + $(5/2^-)$ 90.6392 and $3/2^-$ at 152.9282										
$^{50}\text{K}-\text{u}$	-26100	800	-27620	8	-1.3	U			TO3	1.5	90Tu01
$^{50}\text{K}-^{39}\text{K}_{1,282}$	18899	11	18908	8	0.8	o			TT1	1.0	10La.A
	18908.3	8.3				2			TT1	1.0	12La05
$^{50}\text{Ca}-^{39}\text{K}_{1,282}$	4027.0	4.0	4027.5	1.7	0.1	o			TT1	1.0	10La.A
	4027.5	1.7				2			TT1	1.0	12La05
$^{50}\text{Sc}-\text{u}$	-47940	250	-47824	16	0.3	U			TO6	1.5	98Ba.A *
$\text{C H}_3 \text{}^{35}\text{Cl}-^{50}\text{Ti}$	47550	23	47541.95	0.13	-0.1	U			R09	4.0	72De11
$\text{C}_4 \text{ H}_2-^{50}\text{Ti}$	70860	8	70864.23	0.13	0.1	U			R09	4.0	72De11
$\text{C}_5 \text{ H}_6-^{50}\text{Ti O}$	107253	18	107249.73	0.13	0.0	U			R09	4.0	72De11
$\text{C}_3 \text{}^{13}\text{C H}-^{50}\text{Ti}$	66401	21	66394.03	0.13	-0.1	U			R09	4.0	72De11
$\text{C}_3 \text{ N}-^{50}\text{Ti}$	58279	43	58288.17	0.13	0.1	U			R09	4.0	72De11
$\text{C}_4 \text{ H}_2-^{50}\text{V}$	68485	14	68494.2	0.4	0.2	U			R09	4.0	72De11
$\text{C}_3 \text{ N}-^{50}\text{V}$	55903	23	55918.2	0.4	0.2	U			R09	4.0	72De11
$\text{C H}_3 \text{}^{35}\text{Cl}-^{50}\text{V}$	45158	17	45171.9	0.4	0.2	U			R09	4.0	72De11
$\text{C}_4 \text{ H}_2-^{50}\text{Cr}$	69608	8	69608.6	0.5	0.0	U			R09	4.0	72De11
$\text{C}_3 \text{ N}-^{50}\text{Cr}$	57051	7	57032.6	0.5	-0.7	U			R09	4.0	72De11
$\text{C H}_3 \text{}^{35}\text{Cl}-^{50}\text{Cr}$	46290	14	46286.3	0.5	-0.1	U			R09	4.0	72De11
$^{50}\text{Fe}-\text{u}$	-37012	9				2			LZ1	1.0	16Zh.A
$^{49}\text{Ti} \text{}^{13}\text{C}-^{50}\text{Ti C}$	6440.47	0.88	6433.62	0.04	-3.1	B			H32	2.5	79Ko10
$^{50}\text{Mn}-^{50}\text{Cr}$	8195.91	0.10	8195.95	0.07	0.4	1	52	$^{50}\text{Mn}$	JY1	1.0	08Er04
$^{50}\text{Mn}^m-^{50}\text{Cr}$	8437.852	0.065	8437.83	0.06	-0.3	1	81	$^{50}\text{Mn}^m$	JY1	1.0	08Er04
$^{50}\text{Mn}^m-^{50}\text{Mn}$	241.840	0.100	241.88	0.07	0.4	1	55	$^{50}\text{Mn}$	JY1	1.0	08Er04
$^{50}\text{Ti}-^{49}\text{Ti}$	-3075	38	-3078.79	0.04	0.0	U			R09	4.0	72De11
$^{50}\text{Cr}(\text{p},\text{}^6\text{He})^{45}\text{V}$	-28686	17	-28679.2	1.0	0.4	U			MSU		75Mu09 *
$^{50}\text{Ti}(\text{p},\alpha)^{47}\text{Sc}$	-2231	15	-2231.0	1.9	0.0	U			Tal		65Pi01
$^{50}\text{V}(\text{p},\alpha)^{47}\text{Ti}$	572	23	577.4	0.4	0.2	U			MIT		67Sp09
$^{50}\text{Cr}(\text{p},\alpha)^{47}\text{V}$	-3387	10	-3391.4	0.5	-0.4	U			Ald		66Br06
$^{50}\text{Cr}(\text{}^3\text{He},\text{}^6\text{He})^{47}\text{Cr}$	-18365	14	-18360	6	0.3	1	19	$^{47}\text{Cr}$	MSU		77Mu03 *
$^{48}\text{Ca}(\text{t,p})^{50}\text{Ca}$	3012	15	3025.4	1.6	0.9	U			Ald		66Hi01
	3020	10			0.5	U			LAI		66Wi11
$^{48}\text{Ca}(\text{}^3\text{He,p})^{50}\text{Sc}$	7965	15				2			ANL		69Oh01
$^{50}\text{V}(\text{d},\alpha)^{48}\text{Ti}$	9982	15	9979.5	0.4	-0.2	U			MIT		66Do06
	9988	20			-0.4	U			Kop		67Ha.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{50}\text{Cr}(d,\alpha)^{48}\text{V}$	4928	12	4926.4	1.1	-0.1	U			Kop		67Ha.A
	4923	15			0.2	U			MIT		68Do03
$^{50}\text{Cr}(d,\alpha)^{48}\text{V}^i$	1880	11	1907.6	0.5	2.5	U			MIT		68Do03 *
$^{50}\text{Cr}(p,t)^{48}\text{Cr}$	-15100	8	-15101	7	-0.1	2			Oak		71Do18
	-15100	30			0.0	U			Bld		72Sh27
$^{50}\text{Cr}(p,t)^{48}\text{Cr}^j$	-23861	15				2			MSU		75Mo26 *
$^{50}\text{Co}^i(2p)^{48}\text{Mn}$	1972	13				2			Bor		07Do17
$^{50}\text{Ti}(t,\alpha)^{49}\text{Sc}$	7644	25	7654.5	2.7	0.4	U			LAl		66Wi11
$^{49}\text{Ti}(n,\gamma)^{50}\text{Ti}$	10939.6	0.3	10939.19	0.04	-1.4	U			MMn		80Is02 Z
	10939.19	0.04			0.0	1	100	100 $^{50}\text{Ti}$	Ptn		84Ru06 Z
	10939.20	0.22			0.0	U			Bdn		06Fi.A
$^{49}\text{Ti}(d,p)^{50}\text{Ti}$	8723	8	8714.62	0.04	-1.0	U			Kop		67Ba32
	8721	6			-1.1	U			MIT		67Ba32
$^{50}\text{Cr}(p,d)^{49}\text{Cr}$	-10790	30	-10775.8	2.2	0.5	U			Pri		67Wh03
$^{50}\text{Cr}(d,t)^{49}\text{Cr}$	-6743.1	2.2	-6743.1	2.2	0.0	1	100	100 $^{49}\text{Cr}$	NDm		76Jo01
$^{50}\text{Fe}^i(p)^{49}\text{Mn}$	4389	41	4332	10	-1.4	o			Bor		96Fa09 *
	4332	10				2			Bor		07Do17 *
$^{50}\text{K}(\beta^-)^{50}\text{Ca}$	14050	300	13861	8	-0.6	U					86Mi08
$^{50}\text{Sc}(\beta^-)^{50}\text{Ti}$	6500	200	6884	15	1.9	U					63Ch03
	6260	100			6.2	B					69Wa24
$^{50}\text{V}(n,p)^{50}\text{Ti}$	2979	15	2990.0	0.4	0.7	U			ILL		81Wa31
	2984	10			0.6	U			ILL		94Wa17
$^{50}\text{Ti}(p,n)^{50}\text{V}$	-2991	10	-2990.0	0.4	0.1	U			Har		60Mc12 Y
$^{50}\text{Ti}(^3\text{He},t)^{50}\text{V}^i$	-7032	4	-7039.83	0.27	-2.0	U					71Be29 *
$^{50}\text{Cr}(p,n)^{50}\text{Mn}$	-8416.1	1.9	-8416.82	0.07	-0.4	U			Har		75Fr.A
$^{50}\text{Cr}(^3\text{He},t)^{50}\text{Mn}$	-7650.5	0.4	-7653.07	0.07	-6.4	F			Mun		77Vo02 *
$^{50}\text{Cr}(^3\text{He},t)^{50}\text{Mn}-^{27}\text{Al}(^27\text{Si})$	-2820.0	2.8	-2822.12	0.12	-0.8	U			ChR		74Ha35
$^{50}\text{Cr}(^3\text{He},t)^{50}\text{Mn}-^{42}\text{Ca}(^42\text{Sc})$	-1207.6	2.3	-1208.38	0.12	-0.3	U			ChR		74Ha35
$^{50}\text{Cr}(^3\text{He},t)^{50}\text{Mn}-^{54}\text{Fe}(^54\text{Co})$	610.09	0.17	610.07	0.10	-0.1	1	35	23 $^{54}\text{Co}$	ChR		87Ko34 *
$^{50}\text{Sc}-u$	$M-A=-44530(220)$ keV for mixture gs+m at 256.895 keV										
$^{50}\text{Cr}(p,^6\text{He})^{45}\text{V}$	Original $Q$ increase by 1 for recalibration										
$^{50}\text{Cr}(^3\text{He},^6\text{He})^{47}\text{Cr}$	Original $Q$ reduced by 3, see $^{46}\text{Ti}(^3\text{He},^6\text{He})$										
$^{50}\text{Cr}(d,\alpha)^{48}\text{V}^i$	IT=3043(9); rebuilt from their $Q_{gs}=4923(15)$ keV										
$^{50}\text{Cr}(p,t)^{48}\text{Cr}^j$	Strongest of two fragments given as IT=8760(15); $Q$ rebuilt with Ame1971										
$^{50}\text{Fe}^i(p)^{49}\text{Mn}$	$E_p=2790(41)$ to $11/2^{(-)}$ level at 1541.3125 keV										
$^{50}\text{Fe}^j(p)^{49}\text{Mn}$	$Q_p=2770(12)$ 41.1%, 1874(16) 1.0% to $11/2^{(-)}$ level at 1541.3125, and										
*	$13/2^{(-)}$ at 2481.3 keV										
$^{50}\text{Ti}(^3\text{He},t)^{50}\text{V}^i$	CDE=7802(4) $Q=-7038(4)$ ; recalibration +6 keV for $^{42}\text{Ca}(p,n)^{42}\text{Sc}$ from Ame1961										
$^{50}\text{Cr}(^3\text{He},t)^{50}\text{Mn}$	F : rejected in reference of same group										
$^{50}\text{Cr}(^3\text{He},t)^{50}\text{Mn}-^{54}\text{Fe}(^54\text{Co})$	$Q-Q=40.90(0.16)$ to $650.99(0.06)$ level in $^{50}\text{Mn}$										
$^{51}\text{Ca}-^{39}\text{K}_{1.308}$	8467.58	0.56				2			MA8	1.0	13Wi06
$^{51}\text{Ca}-u$	-38800	350	-39004.3	0.6	-0.4	U			TO3	1.5	90Tu01
	-38900	400			-0.2	U			TO5	1.5	94Se12
	-39249	183			0.9	U			GT1	1.5	04Ma.A
$^{51}\text{V}-^{39}\text{K}_{1.308}$	-8571.23	0.73	-8571.2	0.4	0.0	-			TT1	1.0	14Ma21
	-8570.68	0.99			-0.5	-			TT1	1.0	14Ma21
	ave.	-8571.0	0.6			-0.3	1	54	54 $^{51}\text{V}$		average
$\text{C}_4 \text{H}_3-^{51}\text{V}$	79526	9	79518.2	0.4	-0.2	U			R09	4.0	72De11
$\text{C}_5 \text{H}_7-^{51}\text{V O}$	115921	13	115903.7	0.4	-0.3	U			R09	4.0	72De11
$\text{C}_4 \text{H}_5 \text{N}-^{51}\text{V O}$	103334	13	103327.7	0.4	-0.1	U			R09	4.0	72De11
$\text{C}_3 \text{H N}-^{51}\text{V}$	66943	7	66942.2	0.4	0.0	U			R09	4.0	72De11
$^{51}\text{Cr}-^{39}\text{K}_{1.308}$	-7763.66	0.78	-7763.4	0.4	0.3	-			TT1	1.0	14Ma21
	-7764.0	1.2			0.5	-			TT1	1.0	14Ma21
	ave.	-7763.8	0.7			0.5	1	43	43 $^{51}\text{Cr}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{51}\text{Fe}-u$	-43148	12	-43159	10	-0.9	1	64	$^{64}\text{Fe}$	LZ1	1.0	11Tu09
$^{51}\text{Co}-u$	-29353	52				2			LZ1	1.0	14Sh14
$^{51}\text{Ca}-^{58}\text{Ni}_{.879}$	17823	24	17830.2	0.7	0.3	U			TT1	1.0	12Ga29
$^{47}\text{Ti } ^{37}\text{Cl}_2-^{51}\text{V } ^{35}\text{Cl}_2$	1906.	1.8	1900.7	0.5	-0.7	U			H18	4.0	64Ba03
$^{49}\text{Ti } ^{37}\text{Cl}-^{51}\text{V } ^{35}\text{Cl}$	956.7	0.7	957.6	0.5	0.3	U			H18	4.0	64Ba03
$^{51}\text{K}-^{51}\text{V}$	31871	14				2			TT1	1.0	12Ga29
$^{51}\text{Cr}-^{51}\text{V}$	807.46	0.71	807.78	0.23	0.5	-			TT1	1.0	14Ma21
	806.3	1.2			1.2	-			TT1	1.0	14Ma21
	ave.	807.2			1.0	1	14	$^{51}\text{V}$			average
$^{48}\text{Ca}(^{14}\text{C}, ^{11}\text{C})^{51}\text{Ca}$	-15900	150	-15521.8	0.5	2.5	U			Mun		80Ma40 *
	-16886	100			13.6	B			MSU		85Be50
$^{48}\text{Ca}(^{18}\text{O}, ^{15}\text{O})^{51}\text{Ca}$	-12040	120	-11530.7	0.7	4.2	B			Hei		85Br03 *
	-13900	40			59.2	B			Can		88Ca21
$^{48}\text{Ca}(\alpha, p)^{51}\text{Sc}$	-5860	20				2			ANL		66Er02
$^{51}\text{V}(p, \alpha)^{48}\text{Ti}$	1162	10	1152.9	0.4	-0.9	U			MIT		64Sp12
$^{51}\text{V}(d, \alpha)^{49}\text{Ti}$	7066	12	7070.8	0.4	0.4	U			Kop		67Ha.A
$^{50}\text{Ti}(n, \gamma)^{51}\text{Ti}$	6372.3	1.2	6372.5	0.5	0.2	2					71Ar39 Z
	6372.6	0.6			-0.2	2			Bdn		06Fi.A
$^{50}\text{Ti}(d, p)^{51}\text{Ti}$	4143	6	4147.9	0.5	0.8	U			MIT		67Ba32
	4148	8			0.0	U			Kop		67Ba32
	4147.7	1.2			0.2	2			NDm		76Jo01
$^{50}\text{Ti}(p, \gamma)^{51}\text{V}$	8063.3	2.0	8061.2	0.4	-1.1	U					70K105 Z
	8063.6	2.0			-1.2	U					70Ma36 Z
$^{50}\text{Ti}(^3\text{He}, d)^{51}\text{V}$	2555	15	2567.7	0.4	0.8	U			MIT		67Ob04
$^{50}\text{V}(n, \gamma)^{51}\text{V}$	11051.18	0.10	11051.15	0.08	-0.3	2			MMn		78Ro03 Z
	11051.05	0.17			0.6	2			ILn		91Mi08 Z
	11051.14	0.22			0.0	2			Bdn		06Fi.A
$^{51}\text{V}(\gamma, n)^{50}\text{V}$	-11040	60	-11051.15	0.08	-0.2	U			Phi		60Ge01
$^{50}\text{V}(d, p)^{51}\text{V}$	8840	15	8826.58	0.08	-0.9	U			MIT		67De02
	8828	20			-0.1	U			Kop		67Ha.A
$^{51}\text{V}(p, d)^{50}\text{V}$	-8815	20	-8826.58	0.08	-0.6	U			Oak		65Ba29
$^{50}\text{V}(^3\text{He}, d)^{51}\text{Cr}$	4031	12	4022.88	0.23	-0.7	U			MIT		69Do01
$^{50}\text{Cr}(n, \gamma)^{51}\text{Cr}$	9261.71	0.30	9260.64	0.20	-3.6	B			MMn		80Is02 Z
	9260.63	0.20			0.1	1	98	$^{87}\text{Cr}$	Bdn		06Fi.A
$^{50}\text{Cr}(d, p)^{51}\text{Cr}$	7049	8	7036.07	0.20	-1.6	U			Kop		67Ha.A
	7041	6			-0.8	U			MIT		68Ro09
$^{50}\text{Cr}(p, \gamma)^{51}\text{Mn}$	5270.8	0.3	5270.78	0.29	-0.1	1	95	$^{81}\text{Mn}$			72Fo25 Z
$^{50}\text{Cr}(^3\text{He}, d)^{51}\text{Mn}$	-206	15	-222.70	0.29	-1.1	U			MIT		67Sp09
$^{50}\text{Cr}(p, \gamma)^{51}\text{Mn}^i$	819	2	820.2	1.5	0.6	2					72Fo25
$^{50}\text{Cr}(^3\text{He}, d)^{51}\text{Mn}^i$	-4652	20	-4673.3	1.5	-1.1	U			MIT		67Ra14
	-4671.7	2.3			-0.7	2					79Pa14 *
$^{51}\text{Co}^i(p)^{50}\text{Fe}$	6513	16				3			Bor		07Do17 *
$^{51}\text{Ti}(\beta^-)^{51}\text{V}$	2440	30	2471.0	0.6	1.0	U					55Bu01
	2450	30			0.7	U					55Ma01
$^{51}\text{Cr}(e)^{51}\text{V}$	756	5	752.45	0.21	-0.7	U					55Bi29
$^{51}\text{V}(p, n)^{51}\text{Cr}$	-1533.5	2.0	-1534.79	0.21	-0.6	U			Nvl		59Go68 Z
	-1533.3	1.8			-0.8	U			Oak		64Jo11 Z
	-1533.7	1.5			-0.7	U			Can		70Kn03 Z
	-1534.93	0.24			0.6	1	78	$^{39}\text{V}$	PTB		89Sc24 Z
$^{51}\text{V}(^3\text{He}, t)^{51}\text{Cr}^i$	-7384	5				2					71Be29
$^{51}\text{Mn}(\beta^+)^{51}\text{Cr}$	3232	20	3207.5	0.3	-1.2	U					66G102
$^{48}\text{Ca}(^{14}\text{C}, ^{11}\text{C})^{51}\text{Ca}$	May be a $^{40}\text{Ca}$ contamination. There is a -16900(150) peak										85Be50 **
$^{48}\text{Ca}(^{18}\text{O}, ^{15}\text{O})^{51}\text{Ca}$	Proposed 970(90) level reinterpreted as ground state in reference										85Be50 **
$^{48}\text{Ca}(^{18}\text{O}, ^{15}\text{O})^{51}\text{Ca}$	Weak $M - A = -36120(120)$ level disregarded										AHW **
$^{50}\text{Cr}(^3\text{He}, d)^{51}\text{Mn}^i$	IT=4449(3); $Q$ rebuilt with Ame1977										MMC124**
$^{51}\text{Co}^i(p)^{50}\text{Fe}$	$Q_p=4662(16)$ to $(4^+)$ level at 1851.5 keV										Ens10c **
$^{51}\text{Co}^i(p)^{50}\text{Fe}$	$Q_p=6153$ was a misprint in Ame2012										XuX16b **
$^{51}\text{V}(^3\text{He}, t)^{51}\text{Cr}^i$	CDE=8145(5) $Q = -7881(5)$ ; recalibration -3 keV for $^{50}\text{Cr}(p, n)^{50}\text{Mn}$ from Ame1961										MMC123**

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{52}\text{K}-u$	-18398	36				2			MR1	1.0	15Ro10	
$^{52}\text{Ca}-^{39}\text{K}_{1.333}$	11592.90	0.72				2			MA8	1.0	13Wi06	
$^{52}\text{Ca}-u$	-34900	500	-36786.4	0.7	-2.5	U			TO3	1.5	90Tu01	
	-36792	11			0.5	U			MR1	1.0	13Wi06	
$^{52}\text{Sc}-u$	-43500	230	-43420	90	0.2	2			TO3	1.5	90Tu01	
	-43350	250			-0.2	2			TO5	1.5	94Se12	
	-43110	240			-0.9	2			TO6	1.5	98Ba.A	
	-43438	97			0.2	2			LZ1	1.0	15Xu14	
	-43260	560			-0.3	U			MT1	1.0	15Me08	
$^{52}\text{Cr}-^{39}\text{K}_{1.333}$	-11116.33	0.48	-11115.8	0.4	1.2	1	58	58 $^{52}\text{Cr}$	MA8	1.0	17Mo.A	
$\text{C}_4 \text{H}_4-^{52}\text{Cr}$	90826	9	90795.1	0.4	-0.9	U			R09	4.0	72De11	
$\text{C}_3 \text{}^{13}\text{C} \text{H}_3-^{52}\text{Cr}$	86373	18	86324.9	0.4	-0.7	U			R09	4.0	72De11	
$\text{C}_3 \text{H}_2 \text{N}-^{52}\text{Cr}$	78253	6	78219.1	0.4	-1.4	U			R09	4.0	72De11	
$^{52}\text{Co}-u$	-36888	9				2			LZ1	1.0	16Xu10	
$^{52}\text{Co}^m-u$	-36473	11				2			LZ1	1.0	16Xu10	
$^{52}\text{Ca}-^{58}\text{Ni}_{.897}$	21220	110	21212.1	0.8	-0.1	U			TT1	1.0	12Ga29	
$^{52}\text{Ca}-^{52}\text{Cr}$	22740	82	22708.7	0.8	-0.4	U			TT1	1.0	12Ga29	
$^{52}\text{Cr}-^{50}\text{Cr}$	-5566	41	-5536.5	0.6	0.2	U			R09	4.0	72De11	
$^{52}\text{Cr}(p,\alpha)^{49}\text{V}$	-2596	10	-2593.3	0.9	0.3	U			Ald		66Br06	
$^{50}\text{Ti}(t,p)^{52}\text{Ti}$	5698	10	5699	7	0.1	2			LAI		66Wi11	
	5700	10			-0.1	2			LAI		71Ca19	
$^{50}\text{Ti}(\text{}^3\text{He},p)^{52}\text{V}$	7653	15	7654.4	0.4	0.1	U			Phi		75Ca07	
$^{52}\text{Cr}(d,\alpha)^{50}\text{V}$	4517	12	4515.6	0.5	-0.1	U			Kop		67Ha.A	
$^{52}\text{Cr}(p,\text{}^3\text{He})^{50}\text{V}^i$	-18645	6	-18651.1	0.4	-1.0	U					78Ko27	
$^{52}\text{Cr}(p,t)^{50}\text{Cr}^i$	-21244	7				2					78Ko27	
$^{52}\text{Cr}(p,t)^{50}\text{Cr}^j$	-26041	6				2					78Ko27	
$^{51}\text{V}(n,\gamma)^{52}\text{V}$	7311.2	0.5	7311.24	0.13	0.1	2					84De15	
	7311.18	0.26			0.2	2			ILn		91Mi08	
	7311.27	0.15			-0.2	2			Bdn		06Fi.A	
$^{51}\text{V}(d,p)^{52}\text{V}$	5098	9	5086.68	0.13	-1.3	U			MIT		64Sp12	
	5086	8			0.1	U			Kop		67Ha.A	
$^{51}\text{V}(p,\gamma)^{52}\text{Cr}$	10500.7	2.8	10504.4	0.5	1.3	U					74Ro44	
$^{52}\text{Co}'(p)^{51}\text{Fe}$	1367	60	1487	14	2.0	o			Bor		94Fa06	
	1349	10			13.8	B			Bor		07Do17	
	1352	10			13.5	B					16Or03	
$^{52}\text{Ca}(\beta^-)^{52}\text{Sc}$	5700	200	6180	80	2.4	B					85Hu03	
$^{52}\text{Sc}(\beta^-)^{52}\text{Ti}$	8020	250	9030	80	4.0	B					85Hu03	
$^{52}\text{Ti}(\beta^-)^{52}\text{V}$	1940	200	1974	7	0.2	U					67Mo11	
$^{52}\text{V}(\beta^-)^{52}\text{Cr}$	3904	30	3975.5	0.5	2.4	U					65Ko09	
	3854	30			4.0	B					67Va27	
$^{52}\text{Mn}(\beta^+)^{52}\text{Cr}$	4710.9	4.	4712.0	1.9	0.3	R					58Ko57	
	4707.9	6.			0.7	R					60Ka20	
$^{52}\text{Cr}(p,n)^{52}\text{Mn}$	-5479	10	-5494.3	1.9	-1.5	U			Ric		66Ri09	
$^{52}\text{Cr}(\text{}^3\text{He},t)^{52}\text{Mn}^i$	-7653	5				2					71Be29	
$^{52}\text{Fe}(\beta^+)^{52}\text{Mn}$	2372	10	2377	5	0.5	-					56Ar33	
	2229	130			1.1	U					79Ge02	
	2510	100			-1.3	U					95Ir01	
ave.	2375	6			0.4	1	65	61 $^{52}\text{Fe}$			average	
$^{52}\text{Co}'(IT)^{52}\text{Co}^m$	2548	2				3					16Or03	
$^{52}\text{Ti}(\beta^-)^{52}\text{V}$	$E_{\beta^-}=1800(200)$ to $1^+$ level at 141.610 keV										Ens159	**
$^{52}\text{V}(\beta^-)^{52}\text{Cr}$	$E_{\beta^-}=2470(30)$ 2420(30) respectively, to $2^+$ level at 1434.091 keV										Ens159	**
$^{52}\text{Mn}(\beta^+)^{52}\text{Cr}$	$E_{\beta^+}=575(4)$ and $572(6)$ respectively, to $6^+$ level at 3113.858 keV										Ens159	**
$^{52}\text{Cr}(\text{}^3\text{He},t)^{52}\text{Mn}^i$	CDE=8414(5) $Q=-7650(5)$ ; recalibration -3 keV for $^{50}\text{Cr}(p,n)^{50}\text{Mn}$ from Ame1961										MMC123**	
$^{52}\text{Fe}(\beta^+)^{52}\text{Mn}$	$E_{\beta^+}=804(10)$ to $1^+$ level at 546.438 keV										Ens159	**
$^{52}\text{Fe}(\beta^+)^{52}\text{Mn}$	$E_{\beta^+}=5350(130)$ from $^{52}\text{Fe}^m$ $12^+$ at 6958.0 to $11^+$ level at 3837.2 keV										Ens159	**



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{53}\text{K}-u$	-13200	120				2			MR1	1.0	15Ro10
$^{53}\text{Ca}-u$	-31549	47				2			MR1	1.0	13Wi06
$^{53}\text{Sc}-u$	-41440	260	-41770	100	-0.8	o			TO3	1.5	90Tu01
	-41830	280			0.1	U			TO5	1.5	94Se12
	-41100	400			-1.1	U			TO6	1.5	98Ba.A
	-41694	118			-0.4	2			GT1	1.5	04Ma.A
	-40910	290			-3.0	B			MT1	1.0	11Es06
	-41804	123			0.3	2			LZ1	1.0	15Xu14
	-40980	610			-1.3	U			MT1	1.0	15Me08
$\text{C}_4 \text{H}_5 - ^{53}\text{Cr}$	98529	8	98478.2	0.4	-1.6	U			R09	4.0	72De11
$\text{C}_3 \text{H}_3 \text{N} - ^{53}\text{Cr}$	85958	10	85902.1	0.4	-1.4	U			R09	4.0	72De11
$\text{C}_2 \text{ }^{13}\text{C} \text{H}_2 \text{N} - ^{53}\text{Cr}$	81507	27	81431.9	0.4	-0.7	U			R09	4.0	72De11
$\text{C}_3 \text{H} \text{O} - ^{53}\text{Cr}$	62152	14	62092.7	0.4	-1.1	U			R09	4.0	72De11
$^{53}\text{Co}-u$	-45783	18	-45796.8	1.8	-0.8	U			LZ1	1.0	11Tu09
$^{53}\text{Ni}-u$	-31810	27				2			LZ1	1.0	12Zh34 *
$^{53}\text{Co}-^{53}\text{Fe}$	8897.67	0.49	8897.6	0.5	-0.1	1	94	$93 \text{ }^{53}\text{Co}$	JY1	1.0	10Ka26
$^{53}\text{Co}^m - ^{53}\text{Fe}$	12305.2	1.3	12305.3	1.0	0.1	1	59	$58 \text{ }^{53}\text{Co}^m$	JY1	1.0	10Ka26
$^{53}\text{Co}^m - ^{53}\text{Co}$	3407.9	1.5	3407.6	1.0	-0.2	1	46	$39 \text{ }^{53}\text{Co}^m$	JY1	1.0	10Ka26
$^{53}\text{Cr}-^{52}\text{Cr}$	115	46	141.97	0.15	0.1	U			R09	4.0	72De11
$^{51}\text{V}(t,p)^{53}\text{V}$	7325	25	7308	3	-0.7	U			Ald		67Hi02
$^{53}\text{Cr}(d,\alpha)^{51}\text{V}$	7635	12	7627.6	0.5	-0.6	U			Kop		67Ha.A
$^{52}\text{Cr}(n,\gamma)^{53}\text{Cr}$	7939.52	0.3	7939.07	0.14	-1.5	-			MMn		80Is02 Z
	7939.01	0.2			0.3	-			BNn		80Ko01 Z
	7939.10	0.28			-0.1	-			Bdn		06Fi.A
$^{52}\text{Cr}(d,p)^{53}\text{Cr}$	5725	6	5714.51	0.14	-1.7	U			MIT		64Sp12
	5719	8			-0.6	U			Kop		67Ha.A
$^{52}\text{Cr}(n,\gamma)^{53}\text{Cr}$	ave.	7939.15	7939.07	0.14	-0.5	1	95	$62 \text{ }^{53}\text{Cr}$			average
$^{52}\text{Cr}(p,\gamma)^{53}\text{Mn}$	6559.1	1.1	6559.8	0.3	0.7	U					70Ma25 Z
	6559.72	0.36			0.3	1	86	$77 \text{ }^{53}\text{Mn}$			79Sw01 Z
$^{52}\text{Cr}(^3\text{He},d)^{53}\text{Mn}$	1070	15	1066.4	0.3	-0.2	U			MIT		67Ob04
$^{53}\text{Co}^m(p)^{52}\text{Fe}$	1559.7	40.	1556	5	-0.1	o					70Ja22
	1600.5	30.			-1.5	U					70Ce04
	1590	30			-1.1	U					72Ce01
	1590	30			-1.1	U					76Vi02
	1552.3	8.0			0.5	1	42	$39 \text{ }^{52}\text{Fe}$			15Sh16 *
$^{53}\text{Co}^i(p)^{52}\text{Fe}$	2789.5	50.	2707	6	-1.7	U					76Vi02 *
	2778.5	18.			-4.0	B			Bor		07Do17 *
$^{53}\text{Ti}(\beta^-)^{53}\text{V}$	5020	100				3			ANB		77Pa01
$^{53}\text{V}(\beta^-)^{53}\text{Cr}$	3536	50	3436	3	-2.0	U					56Sc.A *
$^{53}\text{Cr}(p,n)^{53}\text{Mn}$	-1379	8	-1379.2	0.4	0.0	U			MIT		52Lo06 Y
	-1381.1	1.6			1.2	U			Oak		64Jo11 Z
$^{53}\text{Cr}(^3\text{He},t)^{53}\text{Mn}^i$	-7589	4				2					71Be29 *
$^{53}\text{Fe}(\beta^+)^{53}\text{Mn}$	3860	100	3742.6	1.7	-1.2	U					59Ju40
	3820	100			-0.8	U					75BI01
$^{53}\text{Co}^i(\text{IT})^{53}\text{Co}$	4325	2				2					16Su10
$^{53}\text{Ni}-u$	Same result in reference										
$^{53}\text{Co}^m(p)^{52}\text{Fe}$	Original $Q=1558(8)$ corrected for recoil										
$^{53}\text{Co}^i(p)^{52}\text{Fe}$	$Q_p=1940(50)$ $1929(18)$ respectively, to $2^+$ level at 849.45 keV										
$^{53}\text{V}(\beta^-)^{53}\text{Cr}$	$E_{\beta^-}=2530(50)$ to $5/2^-$ level at 1006.27 keV										
$^{53}\text{Cr}(^3\text{He},t)^{53}\text{Mn}^i$	CDE=8350(4) $Q=-7586(4)$ ; recalibration -3 keV for $^{50}\text{Cr}(p,n)^{50}\text{Mn}$ from Ame1961										
$^{54}\text{Ca}-u$	-27011	52				2			MR1	1.0	13Wi06
$^{54}\text{Sc}-u$	-36060	500	-36380	290	-0.4	o			TO3	1.5	90Tu01 *
	-37060	500			0.9	o			TO5	1.5	94Se12 *
	-36960	400			1.0	U			TO6	1.5	98Ba.A *
	-37059	225			2.0	U			GT1	1.5	04Ma.A *
	-36070	390			-0.8	2			MT1	1.0	11Es06 *
	-37202	585			1.4	2			LZ1	1.0	15Xu14 *
	-36230	680			-0.2	2			MT1	1.0	15Me08
$^{54}\text{Ti}-u$	-48820	230	-48980	90	-0.5	2			TO3	1.5	90Tu01
	-49130	250			0.4	2			TO5	1.5	94Se12

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{54}\text{Ti}-u$	-48820	280	-48980	90	-0.4	2			TO6	1.5	98Ba.A
	-48998	118			0.2	2			LZ1	1.0	15Xu14
$\text{C}_4 \text{H}_6-^{54}\text{Cr}$	108018	17	108072.2	0.4	0.8	U			R09	4.0	72De11
$\text{C}_3 \text{}^{13}\text{C} \text{H}_5-^{54}\text{Cr}$	103569	15	103602.0	0.4	0.5	U			R09	4.0	72De11
$\text{C}_3 \text{H}_4 \text{N}-^{54}\text{Cr}$	95445	13	95496.1	0.4	1.0	U			R09	4.0	72De11
$\text{C}_2 \text{}^{13}\text{C} \text{H}_3 \text{N}-^{54}\text{Cr}$	90960	24	91025.9	0.4	0.7	U			R09	4.0	72De11
$\text{C}_2 \text{N O}-^{54}\text{Cr}$	59057	26	59110.6	0.4	0.5	U			R09	4.0	72De11
$^{13}\text{C} \text{}^{37}\text{Cl}_3-^{54}\text{Fe} \text{}^{35}\text{Cl}_2$	23744.46	1.26	23748.9	0.4	1.4	U			H39	2.5	84Ha20
$\text{C}_4 \text{H}_6-^{54}\text{Fe}$	107368	11	107341.9	0.4	-0.6	U			R09	4.0	72De11
$\text{C}_3 \text{H}_4 \text{N}-^{54}\text{Fe}$	94791	8	94765.8	0.4	-0.8	U			R09	4.0	72De11
$\text{C}_2 \text{N O}-^{54}\text{Fe}$	58411	8	58380.3	0.4	-1.0	U			R09	4.0	72De11
$\text{C}_3 \text{}^{13}\text{C} \text{H}_5-^{54}\text{Fe}$	102908	48	102871.7	0.4	-0.2	U			R09	4.0	72De11
$^{54}\text{Ni}-u$	-42167	5				2			LZ1	1.0	16Zh.A
$^{54}\text{Co}-^{54}\text{Fe}$	8850.94	0.14	8850.89	0.10	-0.4	1	47	47 $^{54}\text{Co}$	JY1	1.0	08Er04
$^{54}\text{Co}^m-^{54}\text{Fe}$	9062.960	0.092	9062.99	0.08	0.3	1	81	81 $^{54}\text{Co}^m$	JY1	1.0	08Er04
$^{54}\text{Co}^m-^{54}\text{Co}$	212.18	0.15	212.10	0.10	-0.5	1	49	30 $^{54}\text{Co}$	JY1	1.0	08Er04
$^{54}\text{Cr}-^{53}\text{Cr}$	-1662	48	-1768.95	0.13	-0.6	U			R09	4.0	72De11
$^{54}\text{Fe}(p, ^6\text{He})^{49}\text{Mn}$	-28943	24	-28937.0	2.3	0.3	U			MSU		75Mu09 *
$^{54}\text{Fe}(\alpha, ^8\text{He})^{50}\text{Fe}$	-50950	60	-50963	8	-0.2	U			Tex		77Tr05
$^{54}\text{Cr}(p, \alpha)^{51}\text{V}$	130	30	133.1	0.5	0.1	U			Kop		64Ve02
$^{54}\text{Fe}(p, \alpha)^{51}\text{Mn}$	-3145	9	-3146.6	0.6	-0.2	U			Ald		66Br05
	-3146.9	1.1			0.3	1	28	19 $^{51}\text{Mn}$	NDm		74Jo14
$^{54}\text{Fe}(p, \alpha)^{51}\text{Mn}^i$	-7606.6	5.0	-7597.1	1.6	1.9	U					79Ta22 *
$^{54}\text{Fe}(^3\text{He}, ^6\text{He})^{51}\text{Fe}$	-18694	15	-18713	9	-1.3	1	36	36 $^{51}\text{Fe}$	MSU		77Mu03 *
$^{54}\text{Cr}(d, \alpha)^{52}\text{V}$	5225	12	5219.8	0.5	-0.4	U			Kop		67Ha.A
$^{52}\text{Cr}(t, p)^{54}\text{Cr}$	9171	10	9176.36	0.18	0.5	U			LAL		71Ca19
$^{52}\text{Cr}(^3\text{He}, p)^{54}\text{Mn}$	7785	15	7780.6	1.0	-0.3	U			MIT		69Ly06
	7788	9			-0.8	U			Phi		72Be07
$^{52}\text{Cr}(^3\text{He}, p)^{54}\text{Mn}^i$	1633.6	3.9	1634.5	2.8	0.2	1	51	51 $^{54}\text{Mn}^i$			72Be07 *
$^{52}\text{Cr}(^3\text{He}, n)^{54}\text{Fe}^j$	-7173	20				2					75Bo14
$^{54}\text{Fe}(d, \alpha)^{52}\text{Mn}$	5169	12	5163.6	1.8	-0.5	U			Kop		67Ha.A
	5159	15			0.3	U			MIT		67Sp09
	5163.3	2.2			0.1	-			NDm		76Jo01
ave.	5163.8	1.8			-0.1	1	97	97 $^{52}\text{Mn}$			average
$^{54}\text{Fe}(p, t)^{52}\text{Fe}$	-15584	8	-15585	5	-0.1	R					78Ko27 *
$^{54}\text{Fe}(p, t)^{52}\text{Fe}^j$	-24139	7	-24140	6	-0.1	2					78Ko27
	-24141.3	11.0			0.1	2					78De18 *
$^{54}\text{Zn}(2p)^{52}\text{Ni}$	1480	20	1480	20	0.0	o					05Gi15
	1480	20				3					05B115
	1280	210			1.0	U					11As08
$^{54}\text{Cr}(d, ^3\text{He})^{53}\text{V}$	-6879.2	3.1				2			NDm		79Br.B
$^{53}\text{Cr}(n, \gamma)^{54}\text{Cr}$	9719.30	0.16	9719.08	0.12	-1.4	-					68Wh03 Z
	9718.3	0.4			2.0	-					72Lo26 Z
	9718.91	0.27			0.6	-			MMn		80Is02 Z
	9718.0	0.2			5.4	B					87Mh.A
	9719.7	0.5			-1.2	-			SAn		89Ho15 Z
	9720.00	0.20			-4.6	C			Bdn		06Fi.A
$^{53}\text{Cr}(d, p)^{54}\text{Cr}$	7480	12	7494.52	0.12	1.2	U			MIT		64Sp12
	7514	10			-1.9	U			Kop		67Ha.A
ave.	9719.14	0.13	9719.08	0.12	-0.5	1	96	58 $^{54}\text{Cr}$			average
$^{53}\text{Cr}(p, \gamma)^{54}\text{Mn}$	7559.6	1.0				2					75We10 Z
$^{53}\text{Cr}(^3\text{He}, d)^{54}\text{Mn}$	2080	12	2066.1	1.0	-1.2	U			MIT		69Ly06
$^{54}\text{Fe}(d, t)^{53}\text{Fe}$	-7121.5	2.1	-7121.1	1.6	0.2	-			NDm		74Jo14
$^{54}\text{Fe}(^3\text{He}, \alpha)^{53}\text{Fe}$	7197	20	7199.3	1.6	0.1	U			MIT		68Tr01
	7199.6	2.6			-0.1	-			NDm		74Jo14
ave.	-7121.2	1.6	-7121.1	1.6	0.1	1	98	98 $^{53}\text{Fe}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{54}\text{Ti}(\beta^-)^{54}\text{V}$	4280	160	4270	80	-0.1	R					96Do23
$^{54}\text{V}(\beta^-)^{54}\text{Cr}$	7000	100	7042	15	0.4	U					70Wa14
$^{54}\text{Cr}(t, ^3\text{He})^{54}\text{V}$	-7023	15				2			LAI		77F103
$^{54}\text{Mn}(\epsilon)^{54}\text{Cr}$	1359	8	1377.1	1.0	2.3	U					72Ko47 *
	1379	8			-0.2	U					00Hi08 *
$^{54}\text{Cr}(\text{p},\text{n})^{54}\text{Mn}$	-2160	5	-2159.5	1.0	0.1	U			MIT		52Lo06 Z
$^{54}\text{Cr}(^3\text{He}, t)^{54}\text{Mn}^i$	-7541	4	-7541.9	2.8	-0.2	1	49	49 $^{54}\text{Mn}^i$			71Be29 *
$^{54}\text{Co}(\beta^+)^{54}\text{Fe}$	8023	110	8244.55	0.09	2.0	U					59Su.A *
	8459	41			-5.2	C					60Mi.A *
$^{54}\text{Fe}(\text{p},\text{n})^{54}\text{Co}$	-9031.1	2.5	-9026.89	0.09	1.7	U			Yal		69Ov01 *
	-9023.7	1.8			-1.8	U			Har		74Ho21 Z
$^{54}\text{Fe}(^3\text{He}, t)^{54}\text{Co}$	-8261.2	1.0	-8263.14	0.09	-1.9	F			Mun		77Vo02 *
$^{54}\text{Fe}(^3\text{He}, t)^{54}\text{Co}-^{27}\text{Al}(^{27}\text{Si})$	-3432.5	3.0	-3432.19	0.13	0.1	U			ChR		74Ha35
$^{54}\text{Fe}(^3\text{He}, t)^{54}\text{Co}-^{42}\text{Ca}(^{42}\text{Sc})$	-1817.24	0.18	-1818.45	0.13	-6.7	B			ChR		87Ko34
* $^{54}\text{Sc}-\text{u}$	Original -36000(500) $\mu\text{u}$ or $M-A=-33500(470)$ keV										GAu **
* $^{54}\text{Sc}-\text{u}$	Original -37000(500) $\mu\text{u}$ or $M-A=-34470(470)$ keV										GAu **
* $^{54}\text{Sc}-\text{u}$	$M-A=-34370(370)$ keV for mixture gs+m at 110.5 keV										Nub16b **
* $^{54}\text{Sc}-\text{u}$	$M-A=-33540(360)$ keV for mixture gs+m at 110.5 keV										Nub16b **
* $^{54}\text{Sc}-\text{u}$	No isomeric correction needed										HWJ151 **
* $^{54}\text{Fe}(\text{p}, ^6\text{He})^{49}\text{Mn}$	$Q$ increased 1 for recalibration										AHW **
* $^{54}\text{Fe}(\text{p}, \alpha)^{51}\text{Mn}^i$	IT=4459(5); $Q$ rebuilt with Ame1977										MMC124**
* $^{54}\text{Fe}(^3\text{He}, ^6\text{He})^{51}\text{Fe}$	Averaged with reference See $^{46}\text{Ti}(^3\text{He}, ^6\text{He})$										75Mu09 **
* $^{52}\text{Cr}(^3\text{He}, \text{p})^{54}\text{Mn}^i$	IT=6151(5); $Q$ rebuilt with Ame1971										MMC124**
* $^{54}\text{Fe}(\text{p}, t)^{52}\text{Fe}$	$Q=-21239(8)$ to $^{52}\text{Fe}^i$ at 5654.5										Nub16c **
* $^{54}\text{Fe}(\text{p}, t)^{52}\text{Fe}^j$	IT=8561(5); $Q$ rebuilt with Ame1977										MMC124**
* $^{54}\text{Mn}(\epsilon)^{54}\text{Cr}$	IBE=518(8) to $2^+$ level at 834.855 keV, B(K)=5.99										Ens148 **
* $^{54}\text{Mn}(\epsilon)^{54}\text{Cr}$	IBE=544(8) to $2^+$ level at 834.855 keV										Ens148 **
* $^{54}\text{Cr}(^3\text{He}, t)^{54}\text{Mn}^i$	CDE=8302(4) $Q=-7538(4)$ ; recalibration -3 keV for $^{50}\text{Cr}(\text{p}, \text{n})^{50}\text{Mn}$ from Ame1961										MMC123**
* $^{54}\text{Co}(\beta^+)^{54}\text{Fe}$	$E_{\beta^+}=4250(110)$ from $^{54}\text{Co}^m$ at 197.57 to 2949.2 $6^+$ level										Nub16b **
* $^{54}\text{Fe}(\text{p}, \text{n})^{54}\text{Co}$	Uncorrected for resonance. Orig T=9204.1(1.8) corrected in reference										76Fr13 **
* $^{54}\text{Fe}(^3\text{He}, t)^{54}\text{Co}$	Original value -8260.2(0.6) recalibrated										AHW **
* $^{54}\text{Fe}(^3\text{He}, t)^{54}\text{Co}$	F : rejected in reference of same group										09Fa15 **
$^{55}\text{Sc}-\text{u}$	-30600	1100	-32380	490	-1.1	2			TO3	1.5	90Tu01
	-32100	600			-0.3	2			TO6	1.5	98Ba.A
	-32460	640			0.1	o			MT1	1.0	11Es06
	-32760	620			0.6	2			MT1	1.0	15Me08
$^{55}\text{Ti}-\text{u}$	-44650	280	-44730	170	-0.2	-			TO3	1.5	90Tu01
	-44880	260			0.4	-			TO5	1.5	94Se12
	-44360	350			-0.7	-			TO6	1.5	98Ba.A
ave.	-44680	250			-0.2	1	48	48 $^{55}\text{Ti}$			average
$^{55}\text{Cr}-^{85}\text{Rb}_{.647}$	-2093.4	1.3	-2090.7	0.4	2.1	U			MA8	1.0	17Mo.A
$\text{C}_4 \text{H}_7-^{55}\text{Mn}$	116757	8	116732.1	0.3	-0.8	U			R09	4.0	72De11
$\text{C}_3 \text{ }^{13}\text{C} \text{H}_6-^{55}\text{Mn}$	112281	25	112261.9	0.3	-0.2	U			R09	4.0	72De11
$\text{C}_3 \text{H}_5 \text{N}-^{55}\text{Mn}$	104202	10	104156.0	0.3	-1.2	U			R09	4.0	72De11
$\text{C}_2 \text{H}_3 \text{N}_2-^{55}\text{Mn}$	91618	28	91579.9	0.3	-0.3	U			R09	4.0	72De11
$\text{C}_3 \text{H}_3 \text{O}-^{55}\text{Mn}$	80372	10	80346.5	0.3	-0.6	U			R09	4.0	72De11
$^{55}\text{Mn}-^{85}\text{Rb}_{.647}$	-4884.41	0.84	-4884.8	0.3	-0.4	1	15	15 $^{55}\text{Mn}$	MA8	1.0	12Na15
	-4887.0	1.3			1.7	U			MA8	1.0	17Mo.A
$^{55}\text{Ni}-\text{u}$	-48678	18	-48670.0	0.8	0.4	U			LZ1	1.0	11Tu09
$^{55}\text{Cu}-\text{u}$	-33962	167				2			LZ1	1.0	13Ya03
$^{55}\text{Ni}-^{55}\text{Co}$	9333.43	0.62				2			JY1	1.0	10Ka26
$^{55}\text{Mn}(\text{p}, \alpha)^{52}\text{Cr}$	2570	8	2570.9	0.3	0.1	U			MIT		64Sp12
	2600	10			-2.9	U			ANL		67Ka11
$^{55}\text{Mn}(\text{d}, \alpha)^{53}\text{Cr}$	8283	8	8285.4	0.3	0.3	U			MIT		64Sp12
	8277	15			0.6	U			Kop		67Ha.A
$^{54}\text{Cr}(\text{n}, \gamma)^{55}\text{Cr}$	6246.2	0.4	6246.26	0.19	0.2	-					72Wh05 Z
	6246.28	0.21			-0.1	-			Bdn		06Fi.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{54}\text{Cr}(\text{d,p})^{55}\text{Cr}$	4027	8	4021.70	0.19	-0.7	U			MIT		64Sp12
	4035	8			-1.7	U			Kop		67Ha.A
	4022.1	1.2			-0.3	U			NDm		74Jo14
$^{54}\text{Cr}(\text{n},\gamma)^{55}\text{Cr}$	ave.	6246.26	0.19	6246.26	0.19	0.0	1	100	100 $^{55}\text{Cr}$		average
$^{54}\text{Cr}(\text{p},\gamma)^{55}\text{Mn}$		8067.2	0.4	8066.6	0.3	-1.5	1	63	42 $^{54}\text{Cr}$		78We12
$^{54}\text{Cr}(\text{}^3\text{He,d})^{55}\text{Mn}$		2568	18	2573.1	0.3	0.3	U		MIT		69Ra02
$^{55}\text{Mn}(\gamma,\text{n})^{54}\text{Mn}$		-10192	20	-10226.1	1.1	-1.7	U		Phi		60Ge01
$^{54}\text{Fe}(\text{n},\gamma)^{55}\text{Fe}$		9297.91	0.3	9298.12	0.19	0.7	-		MMn		80Is02 Z
		9298.53	0.27			-1.5	-		Bdn		06Fi.A
$^{54}\text{Fe}(\text{d,p})^{55}\text{Fe}$		7084	8	7073.55	0.19	-1.3	U		MIT		64Sp12
		7083	10			-0.9	U		Kop		67Ha.A
		7072.3	1.7			0.7	U		NDm		74Jo14
$^{54}\text{Fe}(\text{n},\gamma)^{55}\text{Fe}$	ave.	9298.25	0.20	9298.12	0.19	-0.7	1	90	71 $^{54}\text{Fe}$		average
$^{54}\text{Fe}(\text{p},\gamma)^{55}\text{Co}$		5064.0	0.7	5064.35	0.30	0.5	-				77Er02 Z
		5063.9	0.4			1.1	-				80Ha36 Z
$^{54}\text{Fe}(\text{}^3\text{He,d})^{55}\text{Co}$		-428	15	-429.12	0.30	-0.1	U		MIT		67Ob04
		-426.9	2.2			-1.0	U		NDm		74Jo14
$^{54}\text{Fe}(\text{p},\gamma)^{55}\text{Co}$	ave.	5063.9	0.3	5064.35	0.30	1.2	1	74	55 $^{55}\text{Co}$		average
$^{55}\text{Ti}(\beta^-)^{55}\text{V}$		7440	200	7480	160	0.2	1	62	52 $^{55}\text{Ti}$		96Do23
$^{55}\text{V}(\beta^-)^{55}\text{Cr}$		5956	100	5970	100	0.1	1	90	90 $^{55}\text{V}$	ANB	77Na17
$^{55}\text{Cr}(\beta^-)^{55}\text{Mn}$		2500	40	2602.7	0.4	2.6	U				63Me06
		2494	25			4.3	B				65Ko09
$^{55}\text{Fe}(\epsilon)^{55}\text{Mn}$		224.5	4.	231.11	0.18	1.7	U				65Be19
		224.5	3.			2.2	U				69Ka13
		231.4	0.4			-0.7	-				89Zl.A
		230.7	1.9			0.2	U				90Is06
		231.0	1.0			0.1	U				93Wi05 *
		231.37	0.30			-0.9	-				95Da14 *
		231.0	0.3			0.4	-				95Sy01 *
		232.36	0.64			-1.9	U				01Ke14
$^{55}\text{Mn}(\text{p,n})^{55}\text{Fe}$		-1015.7	2.	-1013.46	0.18	1.1	U		Nvl		59Go68 Z
		-1014.6	0.8			1.4	U		Oak		64Jo11 Z
$^{55}\text{Fe}(\epsilon)^{55}\text{Mn}$	ave.	231.23	0.19	231.11	0.18	-0.6	1	91	82 $^{55}\text{Fe}$		average
$^{55}\text{Mn}(\text{}^3\text{He,t})^{55}\text{Fe}^i$		-7883	6			2					71Be29 *
$^{55}\text{Co}(\beta^+)^{55}\text{Fe}$		3466	2	3451.4	0.3	-7.3	B				66Fi06 *
$^{55}\text{Fe}(\epsilon)^{55}\text{Mn}$		Error estimated by evaluator									AHW **
$^{55}\text{Fe}(\epsilon)^{55}\text{Mn}$		Original error 0.10 increased by evaluator									GAu **
$^{55}\text{Fe}(\epsilon)^{55}\text{Mn}$		Original statistical error 0.10 increased by evaluator									GAu **
$^{55}\text{Mn}(\text{}^3\text{He,t})^{55}\text{Fe}^i$		CDE=8654(6) $Q = -7890(6)$ ; recalibration +7 keV for $^{54}\text{Fe}(\text{p,n})^{54}\text{Co}$ from Ame1961									MMC123**
$^{55}\text{Co}(\beta^+)^{55}\text{Fe}$		$E_{\beta^+} = 1513(2)$ to $5/2^-$ level at 931.29 keV									Ens097 **
$^{56}\text{Sc}-u$		-26680	630				2		MT1	1.0	15Me08 *
$^{56}\text{Ti}-u$		-41300	350	-42210	130	-1.7	-		TO3	1.5	90Tu01
		-42010	300			-0.4	-		TO5	1.5	94Se12
		-41770	270			-1.1	-		TO6	1.5	98Ba.A
		-42319	129			0.6	-		GT1	1.5	04Ma.A
		-42700	290			1.7	-		LZ1	1.0	15Xu14
	ave.	-42240	140			0.2	1	90	90 $^{56}\text{Ti}$		average
$^{56}\text{V}-u$		-49470	250	-49550	190	-0.2	-		TO3	1.5	90Tu01
		-49640	260			0.2	-		TO5	1.5	94Se12
		-49310	250			-0.6	-		TO6	1.5	98Ba.A
	ave.	-49470	220			-0.4	1	75	75 $^{56}\text{V}$		average
$^{56}\text{Cr}-^{85}\text{Rb}_{.659}$		-1216.3	2.0	-1220.3	0.6	-2.0	U		MA8	1.0	05Gu27
$^{56}\text{Mn}-^{85}\text{Rb}_{.659}$		-2965.1	1.5	-2966.5	0.4	-0.9	U		MA8	1.0	05Gu37
$^{56}\text{Mn}-^{39}\text{K}_{1.436}$		-8979.0	2.7	-8979.6	0.4	-0.2	U		MA8	1.0	09Na.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_4 H_8 -^{56}Fe$	127754	10	127664.6	0.3	-2.2	U			R09	4.0	72De11
$C_3 ^{13}C H_7 -^{56}Fe$	123300	47	123194.4	0.3	-0.6	U			R09	4.0	72De11
$C_3 H_6 N -^{56}Fe$	115171	13	115088.6	0.3	-1.6	U			R09	4.0	72De11
$C_3 H_4 O -^{56}Fe$	91381	15	91279.1	0.3	-1.7	U			R09	4.0	72De11
$C_2 H_2 N O -^{56}Fe$	78790	24	78703.1	0.3	-0.9	U			R09	4.0	72De11
$C_2 O_2 -^{56}Fe$	54990	9	54893.6	0.3	-2.7	B			R09	4.0	72De11
$^{56}Fe - ^{85}Rb_{.659}$	-6933.79	0.62	-6933.8	0.3	-0.1	1	27	27 $^{56}Fe$	MA8	1.0	17Mo.A
$^{56}Cu - u$	-41485	16				2			LZ1	1.0	16Zh.A
$^{56}Fe - ^{58}Ni_{.966}$	-2604.70	0.47	-2604.54	0.26	0.3	1	31	26 $^{58}Ni$	JY1	1.0	10Ka26
$^{56}Co - ^{58}Ni_{.966}$	2297.85	0.55	2298.0	0.4	0.3	1	58	51 $^{56}Co$	JY1	1.0	10Ka26
$^{56}Ni - ^{55}Co_{1.018}$	1176.23	0.48	1175.4	0.4	-1.7	1	60	33 $^{55}Co$	JY1	1.0	10Ka26
$^{56}Cr - ^{56}Fe$	5713.49	0.56				2			MA8	1.0	17Mo.A
$^{56}Ni - ^{56}Fe$	7192.00	0.52	7192.3	0.3	0.5	1	42	40 $^{56}Ni$	JY1	1.0	10Ka26
$^{56}Ni - ^{56}Co$	2289.61	0.49	2289.7	0.4	0.2	1	67	49 $^{56}Co$	JY1	1.0	10Ka26
$^{56}Fe - ^{54}Fe$	-4755	47	-4672.69	0.30	0.4	U			R09	4.0	72De11
$^{56}Fe(p, \alpha)^{53}Mn$	-1060	9	-1052.9	0.4	0.8	U			MIT		64Sp12
	-1056	9			0.3	U			Ald		66Br05
	-1052.3	0.8			-0.8	1	29	23 $^{53}Mn$	NDm		74Jo14
$^{54}Cr(t, p)^{56}Cr$	5995	30	6011.1	0.6	0.5	U			Ald		68Ch20
	6024	10			-1.3	U			LAL		71Ca19
$^{56}Fe(d, \alpha)^{54}Mn$	5662	12	5661.4	1.1	-0.1	U			Kop		67Ha.A
	5673	30			-0.4	U					67Hj01
$^{54}Fe(^3He, p)^{56}Co$	7410	10	7428.1	0.5	1.8	U			CIT		67Mi02
	7408	15			1.3	U			MIT		68Be10
$^{54}Fe(^3He, n)^{56}Ni$	4513	14	4512.9	0.4	0.0	U			CIT		67Mi02
$^{55}Mn(n, \gamma)^{56}Mn$	7270.53	0.3	7270.44	0.13	-0.3	2			MMn		80Is02 Z
	7270.42	0.15			0.1	2			Bdn		06Fi.A
$^{55}Mn(d, p)^{56}Mn$	5052	5	5045.88	0.13	-1.2	U			MIT		64Sp12
	5053	8			-0.9	U			Kop		67Ha.A
$^{55}Mn(p, \gamma)^{56}Fe$	10189	7	10183.64	0.16	-0.8	U					69Fr22
	10193.7	4.5			-2.2	U					70Sa19 *
	10195.7	3.6			-3.4	B					74Pe15 *
	10183.80	0.17			-0.9	1	86	44 $^{55}Mn$	Utr		92Gu03 Z
$^{56}Fe(d, t)^{55}Fe$	-4938.3	1.3	-4939.87	0.23	-1.2	U			NDm		74Jo14
$^{56}Ni(p)^{55}Co$	-7148.5	30.	-7166.6	0.3	-0.6	U					08Jo04 *
$^{56}Cu^i(p)^{55}Ni$	2929	31	2948	10	0.6	U			Bor		07Do17 *
	2948	10				3					14Or04 *
$^{56}Ti(\beta^-)^{56}V$	7030	330	6830	190	-0.6	1	35	25 $^{56}V$			96Do23
$^{56}Cr(\beta^-)^{56}Mn$	1610	150	1626.5	0.6	0.1	U					60Dr03
$^{56}Mn(\beta^-)^{56}Fe$	3685	5	3695.54	0.21	2.1	U					62Ho14 *
$^{56}Co(\beta^+)^{56}Fe$	4566.0	2.0	4566.7	0.4	0.3	U					65Pe18 *
$^{56}Fe(p, n)^{56}Co$	-5351	10	-5349.0	0.4	0.2	U			Tal		62Ne08 Y
$^{56}Fe(^3He, t)^{56}Co^i$	-8178	9				2					71Be29 *
$^{56}Sc - u$	$M - A = -24850(590)$ keV for mixture gs+m at 0#(100#) keV										15Me08 **
$^{55}Mn(p, \gamma)^{56}Fe$	$E_p = 1537(2)$ to $11703(4)$ level										70Sa19 **
$^{55}Mn(p, \gamma)^{56}Fe$	$E_p = 1537(2)$ to $11705(3)$ level										74Pe15 **
$^{56}Ni(p)^{55}Co$	$E_p = 2540(30)$ from 9735 level										08Jo04 **
$^{56}Cu^i(p)^{55}Ni$	Strongest fragment (strength 2.7); weaker fragment 85(70) keV lower (strength 1.3)										14Or04 **
$^{56}Mn(\beta^-)^{56}Fe$	$E_{\beta^-} = 2838(5)$ to $2^+$ level at 846.7778 keV										Ens115 **
$^{56}Co(\beta^+)^{56}Fe$	$E_{\beta^+} = 1459(3)$ to $4^+$ level at 2085.1045 keV										Ens115 **
$^{56}Fe(^3He, t)^{56}Co^i$	Strongest fragment given as CDE=8950(6); $Q = -8186(6)$ rebuilt with Ame1965										71Be29 **
*	recalibration +7 keV for $^{54}Fe(p, n)^{54}Co$ from Ame1961										MMC123**
$^{57}Sc - u$	-22540	1400				2			MT1	1.0	15Me08
$^{57}Ti - u$	-35700	1000	-36410	280	-0.5	2			TO3	1.5	90Tu01
	-36200	400			-0.3	2			TO6	1.5	98Ba.A
	-37102	408			1.1	2			GT1	1.5	04Ma.A
	-36280	370			-0.4	2			MT1	1.0	11Es06

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{57}\text{V}-u$	-47300	400	-47680	90	-0.6	U			TO3	1.5	90Tu01	
	-47640	270			-0.1	2			TO5	1.5	94Se12	
	-47320	250			-1.0	2			TO6	1.5	98Ba.A	
	-47703	91			0.3	2			LZ1	1.0	15Xu14	
$^{57}\text{Cr}-u$	-56240	250	-56387.6	1.1	-0.4	U			TO3	1.5	90Tu01	
	-56300	260			-0.2	U			TO5	1.5	94Se12	
	-56170	270			-0.5	U			TO6	1.5	98Ba.A	
$^{57}\text{Cr}-^{85}\text{Rb}_{.671}$	2802.1	2.0	2801.5	1.1	-0.3	2			MA8	1.0	05Gu27	
	2801.2	1.4			0.2	2			MA8	1.0	17Mo.A	
$^{57}\text{Mn}-^{85}\text{Rb}_{.671}$	-2525.1	2.3	-2524.9	1.6	0.1	1	49	49 $^{57}\text{Mn}$	MA8	1.0	05Gu37	
$^{57}\text{Mn}-^{39}\text{K}_{1.462}$	-8650.7	2.8	-8652.9	1.6	-0.8	1	33	33 $^{57}\text{Mn}$	MA8	1.0	12Na15	
$\text{C}_7 \text{H}_8-^{57}\text{Fe} \text{ } ^{35}\text{Cl}$	158378.5	3.5	158355.4	0.3	-2.6	U			M18	2.5	68Hu05	
$\text{C}_4 \text{H}_9-^{57}\text{Fe}$	135085	11	135033.2	0.3	-1.2	U			R09	4.0	72De11	
$\text{C}_3 \text{H}_7 \text{N}-^{57}\text{Fe}$	122500	10	122457.1	0.3	-1.1	U			R09	4.0	72De11	
$\text{C}_3 \text{H}_5 \text{O}-^{57}\text{Fe}$	98684	8	98647.6	0.3	-1.1	U			R09	4.0	72De11	
$\text{C}_2 \text{H}_3 \text{N O}-^{57}\text{Fe}$	86104	17	86071.6	0.3	-0.5	U			R09	4.0	72De11	
$^{57}\text{Ni}-^{85}\text{Rb}_{.671}$	-1019.8	2.7	-1019.4	0.6	0.2	U			MA8	1.0	07Gu09	
$^{57}\text{Cu}-u$	-50772	43	-50788.2	0.6	-0.4	U			LZ1	1.0	11Tu09	
	$^{56}\text{Fe} \text{ } ^{13}\text{C}-^{57}\text{Fe} \text{ } \text{C}$	2897.67	0.47	2898.32	0.04	0.6	U		H30	2.5	77Ba10	
	2897.68	0.40			0.6	U		H30	2.5	77Ba10		
$^{56}\text{Fe} \text{H}-^{57}\text{Fe}$	7325	7	7368.52	0.04	1.6	U			R09	4.0	72De11	
$^{57}\text{Fe}-^{56}\text{Fe}_{1.018}$	1627.95	0.46	1627.68	0.04	-0.6	U			JY1	1.0	10Ka26	
$^{57}\text{Fe}-^{58}\text{Ni}_{.983}$	-1048.75	0.46	-1048.84	0.27	-0.2	1	34	28 $^{58}\text{Ni}$	JY1	1.0	10Ka26	
$^{57}\text{Ni}-^{58}\text{Ni}_{.983}$	3350.77	0.72	3350.6	0.5	-0.3	1	55	50 $^{57}\text{Ni}$	JY1	1.0	10Ka26	
$^{57}\text{Cu}-^{56}\text{Ni}_{1.018}$	8126.29	0.55	8125.6	0.4	-1.2	1	63	48 $^{57}\text{Cu}$	JY1	1.0	10Ka26	
$^{57}\text{Cu}-^{57}\text{Fe}$	13817.80	0.86	13819.7	0.5	2.2	1	29	28 $^{57}\text{Cu}$	JY1	1.0	10Ka26	
$^{57}\text{Cu}-^{57}\text{Ni}$	9420.42	0.55	9420.3	0.5	-0.2	1	74	50 $^{57}\text{Ni}$	JY1	1.0	10Ka26	
$^{56}\text{Fe} \text{ } ^{37}\text{Cl}-^{57}\text{Fe} \text{ } ^{35}\text{Cl}$	-3413.7	4.3	-3406.63	0.08	0.7	U			M18	2.5	68Hu05	
$^{57}\text{Fe}-^{56}\text{Fe}$	456.6	1.4	456.52	0.04	0.0	U			M18	2.5	68Hu05	
	453.2	2.1			0.6	U			M18	2.5	68Hu05	
	491	39			-0.2	U			R09	4.0	72De11	
	$^{54}\text{Cr}(\alpha,p)^{57}\text{Mn}$	-4308	8	-4312.6	1.5	-0.6	U		NDm		76Ma03	
		-4302	8			-1.3	U		Can		78An10	
	$^{57}\text{Fe}(\text{p},\alpha)^{54}\text{Mn}$	237	9	239.8	1.1	0.3	U		MIT		64Sp12	
	$^{54}\text{Fe}(\alpha,p)^{57}\text{Co}$	-1770.3	1.8	-1773.0	0.5	-1.5	U		NDm		74Jo14	
	$^{55}\text{Mn}(\text{t},\text{p})^{57}\text{Mn}$	7438.2	3.6	7434.7	1.5	-1.0	1	18	17 $^{57}\text{Mn}$	NDm		77Ma12
	$^{57}\text{Fe}(\text{d},\alpha)^{55}\text{Mn}$	8246	15	8241.38	0.16	-0.3	U		Kop		67Ha.A	
	$^{56}\text{Fe}(\text{n},\gamma)^{57}\text{Fe}$	7645.9	0.5	7646.07	0.04	0.3	U			Utr		68Sp01
7646.10		0.17			-0.2	o			BNn		76Al16	
7645.96		0.20			0.6	U			BNn		78St25	
7646.13		0.21			-0.3	U			MMn		80Is02	
7645.93		0.15			1.0	-			Ptn		80Ve05	
7646.0956		0.0500			-0.4	-			PTc		97Ro26	
7646.08		0.09			-0.1	-					02Bo11	
7646.10		0.15			-0.2	-			Bdn		06Fi.A	
$^{56}\text{Fe}(\text{d},\text{p})^{57}\text{Fe}$		5425	8	5421.51	0.04	-0.4	U			MIT		64Sp12
		5425	8			-0.4	U			Kop		67Ha.A
	5419.8	1.3			1.3	U			NDm		74Jo14	
$^{56}\text{Fe}(\text{n},\gamma)^{57}\text{Fe}$	ave.	7646.08	0.04	7646.07	0.04	-0.2	1	99	83 $^{57}\text{Fe}$		average	
$^{56}\text{Fe}(\text{p},\gamma)^{57}\text{Co}$		6027.7	1.0	6027.5	0.4	-0.2	-				70Ob02	
		6029.3	1.5			-1.2	-				71Le21	
$^{56}\text{Fe}(\text{ } ^3\text{He},\text{d})^{57}\text{Co}$	538	20	534.0	0.4	-0.2	U			LAI		65B113	
$^{56}\text{Fe}(\text{p},\gamma)^{57}\text{Co}$	ave.	6028.2	0.8	6027.5	0.4	-0.9	1	29	29 $^{57}\text{Co}$		average	
		-1226.4	0.5	-1225.9	0.3	0.9	2				70Ob02	
		-1225.6	0.4			-0.6	2				71Le21	
$^{57}\text{Cu}^i(\text{p})^{56}\text{Ni}$		4650	50	4609	25	-0.8	2				76Vi02	
		4568	10			4.1	C				98Jo.A	
		4595	29			0.5	2				02Jo09	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{57}\text{Ti}(\beta^-)^{57}\text{V}$	11020	950	10500	270	-0.5	U					96Do23
$^{57}\text{Cr}(\beta^-)^{57}\text{Mn}$	5100	100	4961.5	1.8	-1.4	U			ANB		78Da04
$^{57}\text{Mn}(\beta^-)^{57}\text{Fe}$	2690	50	2695.6	1.5	0.1	U					63Va37
$^{57}\text{Co}(\epsilon)^{57}\text{Fe}$	810	30	836.3	0.5	0.9	U					71La02 *
$^{57}\text{Fe}(\text{p,n})^{57}\text{Co}$	-1619.4	2.0	-1618.6	0.5	0.4	-			Oak		64Jo11 Z
	-1618.2	2.0			-0.2	-			Can		70Kn03
ave.	-1618.8	1.4			0.1	1	10	10 $^{57}\text{Co}$			average
$^{57}\text{Fe}(\beta^+)^{57}\text{Co}^i$	-8122	7	-8108.2	0.3	2.0	U					71Be29 *
$^{57}\text{Ni}(\beta^+)^{57}\text{Co}$	3245	10	3261.7	0.6	1.7	U					50Fr10 *
	3235	10			2.7	U					51Ca28 *
	3246	10			1.6	U					58Ko60 *
$^{57}\text{Cu}(\beta^+)^{57}\text{Ni}$	8742	130	8774.9	0.4	0.3	U					84Sh28
$^{56}\text{Fe}(\text{n},\gamma)^{56}\text{Fe}$	Original error 0.0005 increased for calibration										GAu **
$^{56}\text{Fe}(\text{p},\gamma)^{56}\text{Co}^i$	T=1247.9 recalibrated to T=1248.5(0.6) keV										AHW **
$^{56}\text{Fe}(\text{p},\gamma)^{56}\text{Co}^i$	T=1247.1 recalibrated to T=1247.7(0.4) keV										AHW **
$^{57}\text{Co}(\epsilon)^{57}\text{Fe}$	IBE=674(30) to $5/2^-$ level at 136.4743 keV										Ens98c **
$^{57}\text{Fe}(\beta^+)^{57}\text{Co}^i$	CDE=8893(7) $Q=-8129(7)$ ; recalibration +7 keV for $^{54}\text{Fe}(\text{p,n})^{54}\text{Co}$ from Ame1961										MMC123**
$^{57}\text{Ni}(\beta^+)^{57}\text{Co}$	$E_{\beta^+}=845(10) 835(10) 849(10)$ respectively, to $13/2^-$ level at 1377.663 keV										Ens98c **
$^{58}\text{V}-u$	-43210	280	-43370	100	-0.4	2			TO3	1.5	90Tu01
	-43350	280			-0.1	2			TO5	1.5	94Se12
	-42700	400			-1.1	2			TO6	1.5	98Ba.A
	-43328	107			-0.3	2			GT1	1.5	04Ma.A
	-43457	134			0.6	2			LZ1	1.0	15Xu14
$^{58}\text{Cr}-u$	-55680	230	-55815.5	1.6	-0.4	U			TO3	1.5	90Tu01
	-55750	260			-0.2	U			TO5	1.5	94Se12
	-55490	270			-0.8	U			TO6	1.5	98Ba.A
$^{58}\text{Cr}-^{85}\text{Rb}_{.682}$	4343.9	1.6				2			MA8	1.0	17Mo.A
$^{58}\text{Mn}-^{39}\text{K}_{1.487}$	-5964.9	2.9				2			MA8	1.0	12Na15 *
$\text{C}_3 \text{H}_8 \text{N}-^{58}\text{Fe}$	132382	12	132400.5	0.4	0.4	U			R09	4.0	72De11
$\text{C}_3 \text{H}_6 \text{O}-^{58}\text{Fe}$	108576	13	108591.1	0.4	0.3	U			R09	4.0	72De11
$\text{C}_2 \text{H}_4 \text{N O}-^{58}\text{Fe}$	95999	13	96015.0	0.4	0.3	U			R09	4.0	72De11
$\text{C}_3 \text{H}_6 \text{O}-^{58}\text{Ni}$	106491	8	106523.0	0.4	1.0	U			R10	4.0	74De22
$\text{C}_3^{13}\text{C H}_9-^{58}\text{Ni}$	138424	14	138438.3	0.4	0.3	U			R10	4.0	74De22
$\text{C}_3 \text{H}_8 \text{N}-^{58}\text{Ni}$	130302	25	130332.5	0.4	0.3	U			R10	4.0	74De22
$\text{C}_2 \text{H}_4 \text{O N}-^{58}\text{Ni}$	93926	10	93947.0	0.4	0.5	U			R10	4.0	74De22
	93928	15			0.3	U			R10	4.0	74De22
$\text{C}_3 \text{H}_6 \text{O}-^{58}\text{Ni}$	106504	14	106523.0	0.4	0.3	U			R10	4.0	74De22
$^{58}\text{Ni}-^{58}\text{Fe}$	2059	32	2068.0	0.3	0.1	U			R09	4.0	72De11
$^{58}\text{Cu}-^{58}\text{Ni}$	9190.61	0.50	9190.6	0.5	0.0	1	90	90 $^{58}\text{Cu}$	JY1	1.0	10Ka26
$^{58}\text{Ni}(\text{p},^6\text{He})^{53}\text{Co}$	-27889	18	-27872.4	1.7	0.9	U			MSU		75Mu09 *
$^{58}\text{Ni}(\alpha,^8\text{He})^{54}\text{Ni}$	-50190	50	-50135	5	1.1	U			Tex		77Tr05
$^{58}\text{Fe}(\text{p},\alpha)^{55}\text{Mn}$	420	9	421.36	0.24	0.2	U			MIT		64Sp12
$^{58}\text{Ni}(\text{p},\alpha)^{55}\text{Co}$	-1341.0	2.9	-1334.8	0.4	2.1	U			BNL		73Go19
	-1335.1	0.9			0.3	1	18	12 $^{55}\text{Co}$	NDm		74Jo14
$^{58}\text{Ni}(\beta^+)^{58}\text{Ni}$	-17556	11	-17553.8	0.7	0.2	U			MSU		77Mu03 *
$^{58}\text{Fe}(\text{d},\alpha)^{56}\text{Mn}$	5470	12	5467.23	0.28	-0.2	U			Kop		67Ha.A
$^{56}\text{Fe}(\beta^+)^{56}\text{Co}$	6853	15	6882.3	1.1	2.0	U			MIT		72Ly01
$^{58}\text{Ni}(\text{d},\alpha)^{56}\text{Co}$	6522	12	6522.5	0.4	0.0	U			Kop		67Ha.A
	6506	10			1.6	U			MIT		68Be10
$^{58}\text{Ni}(\text{p},\text{t})^{56}\text{Ni}$	-13987	18	-13982.0	0.3	0.3	U			Bld		65Ho07
$^{58}\text{Ni}(\text{p},\text{t})^{56}\text{Ni}^j$	-23926	4				2					84Ka07 *
$^{57}\text{Fe}(\text{n},\gamma)^{58}\text{Fe}$	10044.60	0.3	10044.59	0.18	0.0	-			MMn		80Is02 Z
	10044.65	0.24			-0.2	-			Bdn		06Fi.A
$^{57}\text{Fe}(\text{d},\text{p})^{58}\text{Fe}$	7815	8	7820.02	0.18	0.6	U			MIT		64Sp12
	7824	12			-0.3	U			Kop		67Ha.A
$^{57}\text{Fe}(\text{n},\gamma)^{58}\text{Fe}$	ave.	10044.63	0.19	10044.59	0.18	-0.2	1	93	82 $^{58}\text{Fe}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{57}\text{Fe}(p,\gamma)^{58}\text{Co}$	6952	3	6954.3	1.1	0.8	1	14	14	$^{58}\text{Co}$	70Er03
$^{58}\text{Ni}(p,d)^{57}\text{Ni}$	-9971.2	7.	-9991.7	0.5	-2.9	U				79Ik04 *
$^{58}\text{Ni}(^3\text{He},\alpha)^{57}\text{Ni}$	8360.3	4.	8361.4	0.5	0.3	U	MSU			76Na23
	8384.8	15.			-1.6	U				79Fo09 *
$^{58}\text{Ni}(^7\text{Li},^8\text{He})^{57}\text{Cu}$	-29564	50	-29622.4	0.5	-1.2	U	MSU			85Sh03
	-29613	17			-0.6	U	Tex			86Ga19
$^{58}\text{Ni}(^{14}\text{N},^{15}\text{C})^{57}\text{Cu}$	-19900	40	-19929.6	0.9	-0.7	U	Ber			87St04
$^{58}\text{Mn}(\beta^-)^{58}\text{Fe}$	5890	100	6327.6	2.7	4.4	B				69Wa10 *
	5958	100			3.7	B				71Dy01 *
$^{58}\text{Fe}(t,^3\text{He})^{58}\text{Mn}$	-6318	15	-6309.0	2.7	0.6	U	LAI			77Fl03 *
$^{58}\text{Co}(\beta^+)^{58}\text{Fe}$	2305	6	2308.0	1.1	0.5	U				52Ch31 *
	2307	4			0.2	U				63Rh02 *
$^{58}\text{Fe}(^3\text{He},t)^{58}\text{Co}^j$	-8079	8				2				71Be29 *
$^{58}\text{Ni}(p,n)^{58}\text{Cu}$	-9351	5	-9343.4	0.4	1.5	U	Mar			64Ma.A
	-9352.6	3.4			2.7	U	Ric			66Bo20 Z
	-9346	10			0.3	U	Ric			66Ri09
	-9346.6	1.7			1.9	U	Yal			69Ov01 Z
	-9347.8	4.0			1.1	U	Har			76Fr13
$^{58}\text{Ni}(\pi^+, \pi^-)^{58}\text{Zn}$	-16908	50				2				86Se04
$^{58}\text{Mn}-^{39}\text{K}_{1.487}$	$D_M = -5887.8(2.9) \mu\text{u}$ for $^{58}\text{Mn}^m$ at 71.77 keV; $M - A = -55755.6(2.7) \text{ keV}$									Nub16b **
$^{58}\text{Ni}(p, ^6\text{He})^{53}\text{Co}$	$Q$ increased 1 for recalibration									AHW **
$^{58}\text{Ni}(^3\text{He}, ^6\text{He})^{55}\text{Ni}$	Averaged with reference See $^{46}\text{Ti}(^3\text{He}, ^6\text{He})$									75Mu09 **
$^{58}\text{Ni}(p,t)^{56}\text{Ni}^j$	Strongest of three fragments IT=9943(4); $Q$ rebuilt with Ame1977									MMC129**
$^{58}\text{Ni}(p,d)^{57}\text{Ni}$	$Q = -15210(7)$ for $^{57}\text{Ni}^i$ at 5238.8(0.7) keV, strongest fragment IT=5230(7); rebuilt with $Q_{gs} = -9975 \text{ keV}$ , average of 73Ed01 and 65Sh06									Nub16b **
*										73Ed01 **
$^{58}\text{Ni}(^3\text{He}, \alpha)^{57}\text{Ni}$	IT=5235(15); $Q = 3146(15)$ for $^{57}\text{Ni}^i$ at 5238.8(0.7) rebuilt with Ame1977									MMC129**
$^{58}\text{Mn}(\beta^-)^{58}\text{Fe}$	$Q_{\beta^-} = 6100(300)$ ; and 5930(100) from $^{58}\text{Mn}^m$ at 71.77 keV									Nub16b **
$^{58}\text{Mn}(\beta^-)^{58}\text{Fe}$	$Q_{\beta^-} = 6030(100)$ from $^{58}\text{Mn}^m$ at 71.77 keV									Nub16b **
$^{58}\text{Fe}(t, ^3\text{He})^{58}\text{Mn}$	And $Q = -6318(15) - 77(8)$ to $^{58}\text{Mn}^m$ at 71.77 keV									Nub16b **
$^{58}\text{Co}(\beta^+)^{58}\text{Fe}$	$E_{\beta^+} = 472(6) 474(4)$ respectively, to $2^+$ level at 810.7662 keV									En104 **
$^{58}\text{Fe}(^3\text{He}, t)^{58}\text{Co}^j$	Strongest of two fragments IT=5759(8); $Q$ rebuilt with Ame1964									71Be29 **
*	recalibration +7 keV for $^{54}\text{Fe}(p,n)^{54}\text{Co}$ from Ame1961									MMC123**
$^{59}\text{V}-u$	-38500	400	-40610	170	-3.5	B				TO3 1.5 90Tu01
	-40700	350			0.2	2				TO5 1.5 94Se12
	-39900	400			-1.2	2				TO6 1.5 98Ba.A
	-40677	129			0.3	2				GT1 1.5 04Ma.A
$^{59}\text{Cr}-u$	-51490	290	-51620	230	-0.3	2				TO3 1.5 90Tu01 *
	-51640	310			0.0	2				TO5 1.5 94Se12 *
	-51100	310			-1.1	2				TO6 1.5 98Ba.A *
	-52380	500			1.5	2				MT1 1.0 16Me07 *
$^{59}\text{Mn}-^{39}\text{K}_{1.513}$	-4696.8	2.5				2				MA8 1.0 12Na15
$^{59}\text{Fe}-^{85}\text{Rb}_{.694}$	-3908.1	1.0	-3908.4	0.4	-0.3	1	15	15	$^{59}\text{Fe}$	MA8 1.0 17Ma.A
$\text{C}_3 \text{H}_7 \text{O}-^{59}\text{Co}$	116467	12	116496.2	0.4	0.6	U				R10 4.0 74De22
$\text{C}_2^{13}\text{C} \text{H}_6 \text{O}-^{59}\text{Co}$	112011	25	112026.0	0.4	0.1	U				R10 4.0 74De22
$\text{C}_2 \text{H}_5 \text{O} \text{N}-^{59}\text{Co}$	103901	6	103920.1	0.4	0.8	U				R10 4.0 74De22
$^{59}\text{Zn}-u$	-50698	29	-50688.0	0.8	0.3	U				LZ1 1.0 11Tu09
$^{58}\text{Ni} \text{H}-^{59}\text{Co}$	9970	15	9973.16	0.22	0.1	U				R10 4.0 74De22
$^{59}\text{Zn}-^{58}\text{Cu}_{1.017}$	5722.4	1.3	5722.6	0.8	0.1	1	36	27	$^{59}\text{Zn}$	JY1 1.0 10Ka26
$^{59}\text{Zn}-^{59}\text{Cu}$	9815.22	0.72	9815.2	0.6	-0.1	1	81	73	$^{59}\text{Zn}$	JY1 1.0 10Ka26
$^{59}\text{Co}-^{58}\text{Ni}$	-2182	35	-2148.12	0.22	0.2	U				R10 4.0 74De22
$^{59}\text{Co}(p,\alpha)^{56}\text{Fe}$	3245	8	3241.4	0.3	-0.4	U				MIT 64Sp12
	3243	9			-0.2	U				Ald 66Br05
	3240.4	1.4			0.7	U				NDm 74Jo14
$^{59}\text{Co}(d,\alpha)^{57}\text{Fe}$	8667	15	8662.9	0.3	-0.3	U				Kop 67Ha.A
	8659.3	3.2			1.1	U				NDm 74Jo14



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference		
$^{59}\text{Ni}(p,t)^{57}\text{Ni}$	-12738.2	3.3	-12733.7	0.5	1.4	U	MSU		76Na23		
	-12738.4	5.0			0.9	U			78Na11 *		
$^{58}\text{Fe}(n,\gamma)^{59}\text{Fe}$	6581.15	0.30	6581.01	0.11	-0.5	-	Ptn		73Sp06 Z		
	6580.94	0.20			0.3	-			80Ve05 Z		
	6581.02	0.14			-0.1	-			Bdn	06Fi.A	
$^{58}\text{Fe}(d,p)^{59}\text{Fe}$	4357	8	4356.44	0.11	-0.1	U	MIT		64Sp12		
	4369	8			-1.6	U			Kop	67Ha.A	
$^{58}\text{Fe}(n,\gamma)^{59}\text{Fe}$	ave.	0.11	6581.01	0.11	0.0	1	99	85 $^{59}\text{Fe}$	average		
$^{58}\text{Fe}(p,\gamma)^{59}\text{Co}$	7359.7	2.0	7363.6	0.4	1.9	U				74Ke14 Z	
$^{58}\text{Fe}(^3\text{He},d)^{59}\text{Co}$	1871	20	1870.1	0.4	0.0	U	LAL		65B113		
$^{58}\text{Fe}(p,\gamma)^{59}\text{Co}-^{56}\text{Fe}(^57\text{Co})$	1336.5	0.7	1336.1	0.4	-0.6	1			41	28 $^{57}\text{Co}$	75Br29
$^{59}\text{Co}(\gamma,n)^{58}\text{Co}$	-10441	26	-10453.9	1.1	-0.5	U	Phi		60Ge01		
$^{59}\text{Co}(d,t)^{58}\text{Co}$	-4196.0	1.4	-4196.6	1.1	-0.5	1			62	61 $^{58}\text{Co}$	74Jo14
$^{58}\text{Ni}(n,\gamma)^{59}\text{Ni}$	8999.37	0.30	8999.28	0.05	-0.3	U	MMn		75Wi06 Z		
	8999.38	0.20			-0.5	U			77Is01 Z		
	8999.10	0.23			0.8	U			ILn	93Ha05 Z	
	8999.28	0.05			0.0	1			99	72 $^{59}\text{Ni}$	04Ra23
	8999.15	0.18			0.7	U			Bdn	06Fi.A	
$^{58}\text{Ni}(d,p)^{59}\text{Ni}$	6797	10	6774.71	0.05	-2.2	U	Kop		67Ha.A		
	6785	5			-2.1	U			MIT	70An25	
	6773.5	1.7			0.7	U			NDm	74Jo14	
$^{58}\text{Ni}(p,\gamma)^{59}\text{Cu}$	3418.5	0.5	3418.6	0.4	0.1	1	62	62 $^{59}\text{Cu}$	63Bo07 Z		
	3419	2			-0.2	U				70Fo09	
	3416.7	2.0			0.9	U				75K106 Z	
$^{58}\text{Ni}(p,\pi^-)^{59}\text{Zn}$	-144735	40	-144783.4	0.8	-1.2	U	ANB		83Sh31		
$^{59}\text{Mn}(\beta^-)^{59}\text{Fe}$	5200	100	5139.5	2.4	-0.6	U			77Pa18		
$^{59}\text{Fe}(\beta^-)^{59}\text{Co}$	1570	4	1564.9	0.4	-1.3	U				52Me53 *	
	1563	3			0.6	U				63Wo01 *	
$^{59}\text{Ni}(\epsilon)^{59}\text{Co}$	1074.5	1.3	1073.00	0.19	-1.2	U				76Be02 *	
$^{59}\text{Co}(p,n)^{59}\text{Ni}$	-1855.8	2.0	-1855.35	0.19	0.2	U	MIT		51Mc48 Z		
	-1854.3	4.0			-0.3	U				57Bu37 Z	
	-1861	5			1.1	U			Ric	57Ch30 Z	
	-1855.8	1.6			0.3	U			Oak	64Jo11 Z	
	-1855.33	0.20			-0.1	1			95	90 $^{59}\text{Co}$	98Bo30
$^{59}\text{Co}(^3\text{He},t)^{59}\text{Ni}^i$	-8436	8	-8433.5	2.1	0.3	U	PTB		71Be29 *		
$^{59}\text{Zn}(\beta^+)^{59}\text{Cu}$	9120	100	9142.8	0.6	0.2	U				81Ar13	
* $^{59}\text{Cr}-u$	Original -51220(240) $\mu\text{u}$ or $M-A=-47710(230)$ keV								GAu **		
* $^{59}\text{Cr}-u$	Original -51370(270) $\mu\text{u}$ or $M-A=-47850(250)$ keV								GAu **		
* $^{59}\text{Cr}-u$	$M-A=-47350(250)$ keV for mixture gs+m at 503.0 keV								Nub16b **		
* $^{59}\text{Cr}-u$	$M-A=-48540(440)$ keV for mixture gs+m at 503.0 keV								Nub16b **		
* $^{59}\text{Ni}(p,t)^{57}\text{Ni}$	Strongest of three IAS fragments, $Q=-17977.2(5.0)$ for $^{57}\text{Ni}^i$ at 5238.8								Nub16b **		
* $^{59}\text{Fe}(\beta^-)^{59}\text{Co}$	$E_{\beta^-}=475(3)$ to $3/2^-$ level at 1099.256 keV								Ens024 **		
* $^{59}\text{Fe}(\beta^-)^{59}\text{Co}$	$E_{\beta^-}=462(3), 273(3)$ to $3/2^-$ levels at 1099.256, 1291.605 keV								Ens024 **		
* $^{59}\text{Ni}(\epsilon)^{59}\text{Co}$	Authors add B(K)=8.3 of Ni, changed in 7.7 of Co								AHW **		
* $^{59}\text{Co}(^3\text{He},t)^{59}\text{Ni}^i$	Strongest fragment $Q=-8441(8)$ ; recalibration +5 keV for $^{58}\text{Ni}(p,n)^{58}\text{Cu}^i$ from Ame1961								MMC129**		
$^{60}\text{V}-u$	-33860	700	-35690	240	-1.7	U	TO3	1.5	90Tu01 *		
	-35560	600			-0.1	2			TO5	94Se12 *	
	-35180	520			-0.6	2			TO6	98Ba.A *	
	-35889	215			0.6	2			GT1	04Ma.A	
	-35510	430			-0.4	2			MT1	1.0	11Es06 *
$^{60}\text{Cr}-u$	-49680	240	-50100	210	-1.2	2	TO3	1.5	90Tu01		
	-50270	280			0.4	2			TO5	94Se12	
	-49910	280			-0.5	2			TO6	1.5	98Ba.A
	-50929	494			1.7	2			MT1	1.0	16Me07

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{60}\text{Mn}-u$	-56550	240	-56863.4	2.5	-0.9	U			TO3	1.5	90Tu01 *
	-56810	290			-0.1	U			TO5	1.5	94Se12 *
	-56530	280			-0.8	U			TO6	1.5	98Ba.A *
$^{60}\text{Mn}-^{39}\text{K}_{1.538}$	-1044.0	2.5				2			MA8	1.0	12Na15 *
$^{60}\text{Co}-u$	-66380	280	-66184.3	0.5	0.5	U			TO6	1.5	98Ba.A *
$\text{C}_3 \text{H}_8 \text{O}-^{60}\text{Ni}$	126796	14	126729.6	0.4	-1.2	U			R10	4.0	74De22
$\text{C}_2 \text{H}_6 \text{O N}-^{60}\text{Ni}$	114231	10	114153.6	0.4	-1.9	U			R10	4.0	74De22
$\text{C}_2^{13}\text{C H}_7 \text{O}-^{60}\text{Ni}$	122315	10	122259.4	0.4	-1.4	U			R10	4.0	74De22
$\text{C H}_2 \text{N O}_2-^{60}\text{Ni}$	77843	16	77768.1	0.4	-1.2	U			R10	4.0	74De22
$\text{C}_5-^{60}\text{Ni}$	69275	14	69214.7	0.4	-1.1	U			R10	4.0	74De22
$^{60}\text{Ni}-^{85}\text{Rb}_{.706}$	-6937.8	1.6	-6938.3	0.4	-0.3	U			MA8	1.0	07Gu09
$^{60}\text{Zn}-^{58}\text{Ni}_{1.034}$	8698.02	0.55	8698.1	0.4	0.1	1	65	65 $^{60}\text{Zn}$	JY1	1.0	10Ka26
$^{60}\text{Zn}-^{59}\text{Cu}_{1.017}$	3373.19	0.55	3373.2	0.4	-0.1	1	65	35 $^{60}\text{Zn}$	JY1	1.0	10Ka26
$^{60}\text{Ni}-^{58}\text{Ni H}$	-12513	30	-12381.56	0.08	1.1	U			R10	4.0	74De22
$^{60}\text{Ni}-^{59}\text{Co}$	-2503	40	-2408.40	0.22	0.6	U			R10	4.0	74De22
$^{60}\text{Ni}-^{58}\text{Ni}$	-4624	25	-4556.52	0.08	0.7	U			R10	4.0	74De22
	-4627	45			0.4	U			R10	4.0	74De22
$^{60}\text{Ni H}-^{59}\text{Co}$	5310	40	5416.63	0.22	0.7	U			R10	4.0	74De22
$^{60}\text{Ni}(\text{p},\alpha)^{57}\text{Co}$	-263.6	0.7	-263.5	0.4	0.1	1	37	33 $^{57}\text{Co}$	NDm		74Jo14
$^{58}\text{Fe}(\text{t},\text{p})^{60}\text{Fe}$	6907	15	6919	3	0.8	2			LAI		71Ca19
	6947	10			-2.8	2			MSU		76St11
	6913	4			1.4	2			LAI		78No05
$^{60}\text{Ni}(\text{d},\alpha)^{58}\text{Co}$	6084.5	2.2	6084.8	1.1	0.2	1	25	25 $^{58}\text{Co}$	NDm		74Jo14
$^{58}\text{Ni}(\text{t},\text{p})^{60}\text{Ni}$	11905	10	11905.21	0.07	0.0	U			Ald		71Da16
$^{60}\text{Ni}(\text{p},\text{t})^{58}\text{Ni}^i$	-20735	40				2					74Ko08 *
$^{60}\text{Ni}(\text{p},\text{t})^{58}\text{Ni}^j$	-26444	7				2					84Ka07 *
$^{58}\text{Ni}(\text{t},\text{p})^{60}\text{Cu}$	5770	12	5758.6	1.6	-0.9	U			CIT		67Mi02
	5746	20			0.6	U			MIT		68Yo01
$^{58}\text{Ni}(\text{t},\text{p})^{60}\text{Cu}^i$	3210	10	3218	5	0.8	1	26	26 $^{60}\text{Cu}^i$	MIT		68Yo01
$^{58}\text{Ni}(\text{t},\text{p})^{60}\text{Zn}$	818	18	805.5	0.4	-0.7	U			CIT		67Mi02
	821	13			-1.2	U			Oak		72Gr39
$^{58}\text{Ni}(\text{t},\text{p})^{60}\text{Zn}^j$	-6562	24				2					74Ev02 *
$^{59}\text{Co}(\text{n},\gamma)^{60}\text{Co}$	7491.88	0.08	7491.92	0.07	0.5	2			BNn		84Ko29 Z
	7492.05	0.15			-0.9	2			Bdn		06Fi.A
$^{59}\text{Co}(\text{d},\text{p})^{60}\text{Co}$	5267	11	5267.35	0.07	0.0	U			MIT		64Sp12
	5272	8			-0.6	U			Kop		67Ha.A
$^{59}\text{Co}(\text{p},\gamma)^{60}\text{Ni}^i$	-1594	4				2					67Ar01
$^{59}\text{Ni}(\text{n},\gamma)^{60}\text{Ni}$	11387.6	0.4	11387.73	0.05	0.3	U					75Wi06 Z
	11387.73	0.05			0.0	1	99	75 $^{60}\text{Ni}$	ORn		04Ra23
$^{60}\text{Ni}(\text{p},\text{d})^{59}\text{Ni}$	-9180	50	-9163.16	0.05	0.3	U			Pri		64Le10
$^{60}\text{Ni}(\text{d},\text{t})^{59}\text{Ni}$	-5130.2	2.1	-5130.50	0.05	-0.1	U			NDm		74Jo14
$^{60}\text{Ni}(\text{p},\text{d})^{59}\text{Ni}^i$	-16505.1	2.1				2					78Ik02 *
$^{60}\text{Mn}(\beta^-)^{60}\text{Fe}$	8234	86	8445	4	2.5	U			ANB		78No03 *
$^{60}\text{Co}(\beta^-)^{60}\text{Ni}$	2823.6	1.0	2822.81	0.21	-0.8	U					68Wo02 *
$^{60}\text{Cu}(\beta^+)^{60}\text{Ni}$	6250	40	6128.0	1.6	-3.1	B					54Nu26
$^{60}\text{Ni}(\text{p},\text{n})^{60}\text{Cu}$	-6912	20	-6910.3	1.6	0.1	U			ChR		58Go77
	-6909	10			-0.1	U			Ric		66Ri09
	-6910.3	1.6				2			Yal		69Ov01 Z
$^{60}\text{Ni}(\text{t},\text{p})^{60}\text{Cu}^i$	-8685	6	-8688	5	-0.5	1	74	74 $^{60}\text{Cu}^i$			71Be29 *
$^{60}\text{Zn}(\beta^+)^{60}\text{Cu}$	4166	64	4170.8	1.6	0.1	U					86Ka38
$^{60}\text{V}-u$	Original -33800(700) $\mu\text{u}$ or $M-A=-31500(650)$ keV										
$^{60}\text{V}-u$	Original -35500(600) $\mu\text{u}$ or $M-A=-33070(560)$ keV										
$^{60}\text{V}-u$	$M-A=-32700(470)$ keV for mixture gs+m+n at 0#150 and 203.7(0.7) keV										
$^{60}\text{V}-u$	$M-A=-33010(390)$ keV for mixture gs+m+n at 0#150 and 203.7(0.7) keV										
$^{60}\text{Mn}-u$	$M-A=-52540(230)$ keV for mixture gs+m at 271.90 keV										
$^{60}\text{Mn}-u$	$M-A=-52780(260)$ keV for mixture gs+m at 271.90 keV										
$^{60}\text{Mn}-u$	$M-A=-52520(250)$ keV for mixture gs+m at 271.90 keV										
$^{60}\text{Mn}-^{39}\text{K}_{1.538}$	$D_M=-752.1(2.5)$ $\mu\text{u}$ for $^{60}\text{Mn}^m$ at 271.90 keV; $M-A=-52695.9(2.4)$ keV										
$^{60}\text{Co}-u$	$M-A=-61800(260)$ keV for mixture gs+m at 58.59 keV										
$^{60}\text{Ni}(\text{p},\text{t})^{58}\text{Ni}^i$	IT=8830(40); $Q$ rebuilt with Ame1971										
$^{60}\text{Ni}(\text{p},\text{t})^{58}\text{Ni}^j$	IT=14537(7); $Q$ rebuilt with Ame1977										
$^{58}\text{Ni}(\text{t},\text{p})^{60}\text{Zn}^j$	IT=7380(30); $Q$ rebuilt with Ame1971										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>60</sup> Ni(p,d) <sup>59</sup> Ni <sup>i</sup>	Strongest fragment IT=7341; $Q$ rebuilt with Ame1977										MMC129**
* <sup>60</sup> Mn( $\beta^-$ ) <sup>60</sup> Fe	$E_{\beta^-}$ =5714(86) from <sup>60</sup> Mn <sup>m</sup> at 271.90 to (3 <sup>+</sup> ,4 <sup>+</sup> ) 2792.68 level										Ens13c **
* <sup>60</sup> Co( $\beta^-$ ) <sup>60</sup> Ni	$E_{\beta^-}$ =317.88(0.10) to 4 <sup>+</sup> level at 2505.753 keV										Ens13c **
* <sup>60</sup> Ni( <sup>3</sup> He,t) <sup>60</sup> Cu <sup>i</sup>	CDE=9454(6) $Q$ =-8690(6); recalibration +5 keV for <sup>58</sup> Ni(p,n) <sup>58</sup> Cu <sup>i</sup> from Ame1961										MMC123**
<sup>61</sup> V-u	-32750	960							MT1	1.0	11Es06
<sup>61</sup> Cr-u	-44500	400	-45600	110	-1.8	2			TO3	1.5	90Tu01
	-45910	300			0.7	2			TO5	1.5	94Se12
	-45120	280			-1.1	2			TO6	1.5	98Ba.A
	-45679	107			0.5	2			GT1	1.5	04Ma.A
	-45634	176			0.2	2			LZ1	1.0	15Xu14
<sup>61</sup> Mn-u	-46248	548			1.2	U			MT1	1.0	16Me07
	-55160	300	-55547.5	2.5	-0.9	U			TO3	1.5	90Tu01
	-55540	280			0.0	U			TO5	1.5	94Se12
	-55320	270			-0.6	U			TO6	1.5	98Ba.A
<sup>61</sup> Mn- <sup>39</sup> K <sub>1.564</sub>	1215.6	2.5				2		MA8	1.0	12Na15	
<sup>61</sup> Fe- <sup>39</sup> K <sub>1.564</sub>	-6490.7	2.8				2		MA8	1.0	12Na15	
C H <sub>3</sub> N O <sub>2</sub> - <sup>61</sup> Ni	85373	14	85323.4	0.4	-0.9	U			R10	4.0	74De22
C <sub>5</sub> H- <sup>61</sup> Ni	76810	10	76770.1	0.4	-1.0	U			R10	4.0	74De22
<sup>61</sup> Ga-u	-50654	59	-50600	40	0.9	1	48	48 <sup>61</sup> Ga	LZ1	1.0	11Tu09
<sup>60</sup> Ni H- <sup>61</sup> Ni	7539	14	7555.34	0.05	0.3	U			R10	4.0	74De22
<sup>61</sup> Ni- <sup>60</sup> Ni	339	60	269.69	0.05	-0.3	U			R10	4.0	74De22
<sup>61</sup> Ni- <sup>58</sup> Ni H	-12187	30	-12111.87	0.09	0.6	U			R10	4.0	74De22
<sup>61</sup> Ni- <sup>59</sup> Co	-2220	30	-2138.71	0.22	0.7	U			R10	4.0	74De22
<sup>58</sup> Ni( $\alpha$ ,n) <sup>61</sup> Zn	-9810	30	-9526	16	9.5	B			Oak		64St01
<sup>58</sup> Ni( <sup>6</sup> Li,t) <sup>61</sup> Zn	-4736	23	-4743	16	-0.3	R			LAI		78Wo01
<sup>59</sup> Co( <sup>3</sup> He,p) <sup>61</sup> Ni	9635	10	9634.44	0.21	-0.1	U			MIT		67Sp09
<sup>60</sup> Ni(n, $\gamma$ ) <sup>61</sup> Ni	7820.22	0.40	7820.10	0.05	-0.3	U					75Wi06 Z
	7819.96	0.20			0.7	U			MMn		77Is01 Z
	7820.02	0.20			0.4	U			ILn		93Ha05 Z
	7820.11	0.05			-0.1	-			ORn		04Ra23
	7820.06	0.16			0.3	-			Bdn		06Fi.A
<sup>60</sup> Ni(d,p) <sup>61</sup> Ni	5604	8	5595.54	0.05	-1.1	U			MIT		70An25
	5596.1	1.3			-0.4	U			NDm		74Jo14
<sup>60</sup> Ni(n, $\gamma$ ) <sup>61</sup> Ni	ave.	7820.11	0.05	7820.10	0.05	0.0	1	100	79 <sup>61</sup> Ni		average
<sup>61</sup> Ga <sup>i</sup> (p) <sup>60</sup> Zn	3110	30				2					87Ho.A
<sup>61</sup> Fe( $\beta^-$ ) <sup>61</sup> Co	3827	100	3977.6	2.7	1.5	U					67Eh02 *
	3887	100			0.9	U					67Gu06 *
<sup>61</sup> Co( $\beta^-$ ) <sup>61</sup> Ni	1290	40	1323.8	0.8	0.8	U					56Nu02
<sup>61</sup> Cu( $\beta^+$ ) <sup>61</sup> Ni	2227	5	2237.8	1.0	2.2	U					50Ow03
<sup>61</sup> Ni(p,n) <sup>61</sup> Cu	-3024.0	4.	-3020.2	1.0	1.0	U			Oak		64Jo11 Z
<sup>61</sup> Ni( <sup>3</sup> He,t) <sup>61</sup> Cu <sup>i</sup>	-8630	7				2					71Be29 *
<sup>61</sup> Zn( $\beta^+$ ) <sup>61</sup> Cu	5400	200	5635	16	1.2	U					59Cu86
<sup>61</sup> Ga( $\beta^+$ ) <sup>61</sup> Zn	9255	50	9210	40	-0.8	1	57	52 <sup>61</sup> Ga			02We07
* <sup>61</sup> Fe( $\beta^-$ ) <sup>61</sup> Co	$E_{\beta^-}$ =2800(100) 2860(100) respectively, to 3/2 <sup>-</sup> level at 1027.48 keV										Ens153 **
* <sup>61</sup> Ni( <sup>3</sup> He,t) <sup>61</sup> Cu <sup>i</sup>	Strongest fragment IT=6380(7); $Q$ rebuilt with Ame1964										MMC129**
*	recalibration +5 keV for <sup>58</sup> Ni(p,n) <sup>58</sup> Cu <sup>i</sup> from Ame1961										MMC123**
<sup>62</sup> Cr-u	-42400	600	-43900	160	-1.7	2			TO3	1.5	90Tu01
	-44200	400			0.5	2			TO5	1.5	94Se12
	-43100	350			-1.5	2			TO6	1.5	98Ba.A
	-44026	118			0.7	2			GT1	1.5	04Ma.A
	-43897	526			0.0	U			MT1	1.0	16Me07
<sup>62</sup> Mn-u	-51510	270	-52093	7	-1.4	U			TO3	1.5	90Tu01
	-52030	280			-0.1	U			TO5	1.5	94Se12
	-51180	280			-2.2	U			TO6	1.5	98Ba.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{62}\text{Mn}^m - ^{39}\text{K}_{1.590}$	5982.3	2.8				2			MA8	1.0	12Na15
$^{62}\text{Fe} - ^{39}\text{K}_{1.590}$	-5501.5	3.0				2			MA8	1.0	12Na15
$\text{C}_5 \text{H}_2 - ^{62}\text{Ni}$	87299	10	87305.2	0.5	0.2	U			R10	4.0	74De22
$\text{C H}_4 \text{N O}_2 - ^{62}\text{Ni}$	95859	12	95858.5	0.5	0.0	U			R10	4.0	74De22
$^{62}\text{Cu} - ^{62}\text{Ni}$	4250.05	0.51				2			JY1	1.0	06Er03
$^{62}\text{Zn} - ^{62}\text{Ni}$	5988.49	0.58	5988.6	0.5	0.2	1	68	$^{68}\text{Zn}$	JY1	1.0	06Er03
$^{62}\text{Ga} - ^{62}\text{Ni}$	15845.06	0.71	15844.9	0.5	-0.2	1	52	$^{52}\text{Ga}$	JY1	1.0	06Er03
$^{62}\text{Ga} - ^{62}\text{Zn}$	9856.21	0.45	9856.3	0.4	0.2	1	81	$^{48}\text{Ga}$	JY1	1.0	06Er03
$^{62}\text{Ni} - ^{61}\text{Ni}$	-2669	15	-2710.1	0.3	-0.7	U			R10	4.0	74De22
$^{62}\text{Ni} - ^{60}\text{Ni}$	-2333	30	-2440.4	0.3	-0.9	U			R10	4.0	74De22
$^{62}\text{Ni}(\text{p},\alpha)^{59}\text{Co}$	342	10	347.4	0.4	0.5	U			MIT		64Sp12
	343.3	0.7			5.9	B			NDm		74Jo14
$^{59}\text{Co}(\alpha,\text{p})^{62}\text{Ni}$	-346.5	2.3	-347.4	0.4	-0.4	U			NDm		74Jo14
$^{62}\text{Ni}(\text{t},\alpha)^{60}\text{Co}$	911	20	926	3	0.7	U			Hei		84Ha31
$^{62}\text{Ni}(\text{d},\alpha)^{60}\text{Co}$	5611.2	2.4	5614.8	0.4	1.5	U			NDm		74Jo14
$^{60}\text{Ni}(\text{t},\text{p})^{62}\text{Ni}$	9937	10	9934.0	0.3	-0.3	U			Ald		71Da16
$^{60}\text{Ni}(\text{t},\text{p})^{62}\text{Cu}$	5938	25	5956.6	0.6	0.7	U			MIT		67Sp09
$^{60}\text{Ni}(\text{t},\text{p})^{62}\text{Zn}$	3580	30	3554.8	0.5	-0.8	U			Oak		72Gr39
$^{62}\text{Ni}(\text{t},\alpha)^{61}\text{Fe}$	-7921	100	-7661.5	2.7	2.6	F			Ors		84De33 *
$^{62}\text{Ni}(\text{t},\alpha)^{61}\text{Co}$	8689	20	8676.6	0.7	-0.6	U			LAI		66B115
$^{61}\text{Ni}(\text{n},\gamma)^{62}\text{Ni}$	10596.2	1.5	10595.7	0.3	-0.3	-					70Fa06
	10595.8	0.7			-0.1	-					75Wi06 Z
	10595.6	0.4			0.3	-			Bdn		06Fi.A
$^{61}\text{Ni}(\text{d},\text{p})^{62}\text{Ni}$	8379	8	8371.2	0.3	-1.0	U			MIT		64Sp12
	8369	15			0.1	U			Ald		67Te02
$^{62}\text{Ni}(\text{d},\text{t})^{61}\text{Ni}$	-4340.6	1.3	-4338.5	0.3	1.6	-			NDm		74Jo14
$^{61}\text{Ni}(\text{n},\gamma)^{62}\text{Ni}$	ave. 10595.8	0.3	10595.7	0.3	-0.2	1	88	$^{67}\text{Ni}$			average
$^{62}\text{Mn}^m(\text{IT})^{62}\text{Mn}$	343	6				3					15Ga38 *
$^{62}\text{Fe}(\beta^-)^{62}\text{Co}$	3000	200	2546	19	-2.3	U					75Fr16
$^{62}\text{Co}(\beta^-)^{62}\text{Ni}$	5195	30	5322	19	4.2	C					57Ga15 *
$^{62}\text{Ni}(\text{t},\text{t})^{62}\text{Co}$	-5350	50	-5303	19	0.9	2					72Ba31
	-5296	20			-0.4	2			LAI		76Aj03
$^{62}\text{Cu}(\beta^+)^{62}\text{Ni}$	3932	10	3958.9	0.5	2.7	U					54Nu27
	3942	10			1.7	U					64Sa32
	3956	7			0.4	U					67An01
$^{62}\text{Ni}(\text{p},\text{n})^{62}\text{Cu}$	-4733	10	-4741.2	0.5	-0.8	U			Bar		61Ri02
	-4734.8	10.			-0.6	U			Ric		66Ri09
$^{62}\text{Ni}(\text{t},\text{He},\text{t})^{62}\text{Cu}^i$	-8591	6				2					71Be29 *
$^{62}\text{Zn}(\beta^+)^{62}\text{Cu}$	1682	10	1619.5	0.7	-6.3	B					50Ha65
	1697	10			-7.8	B					54Nu27
$^{62}\text{Ga}(\beta^+)^{62}\text{Zn}$	9171	26	9181.1	0.4	0.4	U			ANB		79Da04
$^{62}\text{Ni}(\text{t},\text{C},\text{O})^{61}\text{Fe}$	F : not unambiguously ground state transition										
$^{62}\text{Mn}^m(\text{IT})^{62}\text{Mn}$	$E_x = 418(2) - 72(+8-3)$										
$^{62}\text{Co}(\beta^-)^{62}\text{Ni}$	$E_{\beta^-} = 5217(30)$ from $^{62}\text{Co}^m$ at 22(5) keV										
$^{62}\text{Ni}(\text{t},\text{He},\text{t})^{62}\text{Cu}^i$	CDE=9360(6) $Q = -8596(6)$ ; recalibration +5 keV for $^{58}\text{Ni}(\text{p},\text{n})^{58}\text{Cu}^i$ from Ame1961										
	MMC124**										
$^{63}\text{Cr}-\text{u}$	-38819	462	-38660	380	0.2	2			GT1	1.5	04Ma.A
	-37870	700			-1.1	o			MT1	1.0	11Es06
	-38583	462			-0.2	2			MT1	1.0	16Me07
$^{63}\text{Mn}-\text{u}$	-49300	400	-50335	4	-1.7	U			TO3	1.5	90Tu01
	-50190	300			-0.3	U			TO5	1.5	94Se12
	-49600	290			-1.7	U			TO6	1.5	98Ba.A
	-50500	107			1.0	o			GT1	1.5	04Ma.A
	-50829	102			1.9	U			GT2	2.5	08Kn.A
$^{63}\text{Mn} - ^{39}\text{K}_{1.615}$	8278.7	4.0				2			MA8	1.0	12Na15

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{63}\text{Fe}-u$	-59190	240	-59727	5	-1.5	U			TO3	1.5	90Tu01
	-59570	290			-0.4	U			TO5	1.5	94Se12
	-58990	300			-1.6	U			TO6	1.5	98Ba.A
$^{63}\text{Fe}-^{39}\text{K}_{1,615}$	-1114.5	6.1	-1113	5	0.2	1	57	57 $^{63}\text{Fe}$	MA8	1.0	12Na15
$^{63}\text{Fe}-\text{H C}_2 \text{ F}_2$	-64354	10	-64359	5	-0.5	o			MS1	1.0	08B105
	-64353	10			-0.6	1	21	21 $^{63}\text{Fe}$	MS1	1.0	10Fe01
$^{63}\text{Fe}-\text{C }^{32}\text{S F}$	-30204	10	-30202	5	0.2	1	21	21 $^{63}\text{Fe}$	MS1	1.0	10Fe01
$\text{C}_5 \text{ H}_3 - ^{63}\text{Cu}$	93930	4	93877.9	0.5	-3.3	B			R10	4.0	74De22
$\text{C}_4 \text{ H N} - ^{63}\text{Cu}$	81347	10	81301.8	0.5	-1.1	U			R10	4.0	74De22
$\text{C}_4 \text{ }^{13}\text{C H}_2 - ^{63}\text{Cu}$	89466	14	89407.7	0.5	-1.0	U			R10	4.0	74De22
$\text{C}_2 \text{ H}_7 \text{ O}_2 - ^{63}\text{Cu}$	115064	16	115007.2	0.5	-0.9	U			R10	4.0	74De22
$^{13}\text{C C H}_8 \text{ O N} - ^{63}\text{Cu}$	134404	18	134346.5	0.5	-0.8	U			R10	4.0	74De22
$^{47}\text{Ti O} - ^{63}\text{Cu}$	17036	23	17075.1	0.5	0.4	U			R09	4.0	72De11
$^{63}\text{Ga} - ^{85}\text{Rb}_{,741}$	4658.0	1.4				2			MA8	1.0	07Gu09
$\text{C}_5 \text{ H}_2 - ^{63}\text{Ga}_{,984}$	75382.6	6.7	75384.6	1.4	0.3	U			MS1	1.0	07Sc24
$^{63}\text{Ge}-u$	-50372	40				2			LZ1	1.0	11Tu02
$^{63}\text{Cu} - ^{62}\text{Ni}$	1193	35	1252.37	0.06	0.4	U			R10	4.0	74De22
$^{63}\text{Cu} - ^{61}\text{Ni}$	-1449	30	-1457.7	0.3	-0.1	U			R10	4.0	74De22
$^{63}\text{Cu}(\text{p},\alpha)^{60}\text{Ni}$	3757	8	3757.4	0.3	0.1	U			MIT		64Sp12
	3780	10			-2.3	U			Min		67Jo03
	3754.9	1.5			1.7	U			NDm		76Jo01
$^{60}\text{Ni}(\alpha,n)^{63}\text{Zn}$	-7970	40	-7906.1	1.6	1.6	U			Oak		64St01
	-7910	20			0.2	U					67Bi04
$^{63}\text{Cu}(\text{d},\alpha)^{61}\text{Ni}$	9376	30	9353.0	0.3	-0.8	U					67Hj01
$^{62}\text{Ni}(\text{n},\gamma)^{63}\text{Ni}$	6838.04	0.20	6837.77	0.06	-1.4	-			MMn		77Is01 Z
	6837.88	0.18			-0.6	-			ILn		92Ha21 Z
	6837.89	0.14			-0.9	-			Bdn		06Fi.A
	6837.75	0.18			0.1	-			JAn		12Os04
$^{62}\text{Ni}(\text{d},\text{p})^{63}\text{Ni}$	4620	6	4613.20	0.06	-1.1	U			MIT		70An25
	4614.0	1.1			-0.7	U			NDm		74Jo14
$^{62}\text{Ni}(\text{n},\gamma)^{63}\text{Ni}$	ave. 6837.88	0.09	6837.77	0.06	-1.3	1	48	34 $^{63}\text{Ni}$			average
$^{62}\text{Ni}(\text{p},\gamma)^{63}\text{Cu}$	6119.2	1.5	6122.40	0.06	2.1	U					72Ki15
	6122.30	0.08			1.2	1	54	38 $^{63}\text{Cu}$	Utr		86De14 Z
$^{63}\text{Cu}(\gamma,\text{n})^{62}\text{Cu}$	-10833	17	-10863.6	0.5	-1.8	U			Phi		60Ge01
$^{63}\text{Co}(\beta^-)^{63}\text{Ni}$	3590	50	3661	19	1.4	1	14	14 $^{63}\text{Co}$			69Ki.A
$^{63}\text{Ni}(\beta^-)^{63}\text{Cu}$	65.87	0.15	66.977	0.015	7.4	B					66Hs01
	66.946	0.020			1.5	o					87He14
	66.945	0.004			7.9	F					92Ka29 *
	66.9459	0.0054			5.7	F					93Oh02 *
	66.980	0.015			-0.2	1	98	55 $^{63}\text{Ni}$			99Ho09
$^{63}\text{Zn}(\beta^+)^{63}\text{Cu}$	3352	20	3366.4	1.5	0.7	U					61Cu02
	3390	30			-0.8	U					61Va08
$^{63}\text{Cu}(\text{p},\text{n})^{63}\text{Zn}$	-4146.5	4.	-4148.7	1.5	-0.6	-			Ric		55Br16
	-4139.5	8.			-1.2	U			Oak		55Ki28 Z
	-4150.1	4.4			0.3	-			Tkm		63Ok01
ave. -4148.1	2.9			-0.2	1	28	27 $^{63}\text{Zn}$				average
$^{63}\text{Cu}(\text{}^3\text{He},\text{t})^{63}\text{Zn}^i$	-8875	6				2					71Be29 *
$^{63}\text{Ga}(\beta^+)^{63}\text{Zn}$	5520	100	5666.3	2.0	1.5	U					72Fi.A
$^{63}\text{Ni}(\beta^-)^{63}\text{Cu}$	F : excitation of atomic electron not taken into account										
$^{63}\text{Cu}(\text{}^3\text{He},\text{t})^{63}\text{Zn}^i$	CDE=9644(6) $Q = -8880(6)$ ; recalibration +5 keV for $^{58}\text{Ni}(\text{p},\text{n})^{58}\text{Cu}^i$ from Ame1961										
$^{64}\text{Cr}-u$	-35942	472				2			MT1	1.0	16Me07
$^{64}\text{Mn}-u$	-45340	350	-46151	4	-1.5	U			TO3	1.5	90Tu01 *
	-46340	350			0.4	U			TO5	1.5	94Se12 *
	-45664	306			-1.1	U			TO6	1.5	98Ba.A *
	-46280	129			0.7	U			GT1	1.5	04Ma.A *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{64}\text{Mn}-^{85}\text{Rb}_{.753}$	20271.7	3.8				2			MA8	1.0	12Na15
$^{64}\text{Fe}-u$	-58600	400	-59012	5	-0.7	U			TO3	1.5	90Tu01
	-59130	300			0.3	U			TO5	1.5	94Se12
	-58500	350			-1.0	U			TO6	1.5	98Ba.A
	-59012.2	5.3			0.0	o			MS1	1.0	08BI05
	62699.4	5.3				2			MS1	1.0	10Fe01
$\text{H C}_2 \text{ F}_2-^{64}\text{Fe}_{.984}$	62699.4	5.3				2			MS1	1.0	10Fe01
$^{64}\text{Co}^m-u$	-64075.5	4.5	-64075	4	0.1	o			MS1	1.0	08BI05
$\text{H C}_2 \text{ F}_2-^{64}\text{Co}^m_{.984}$	67681.6	4.6	67681	4	-0.1	1	87	$^{87}\text{Co}^m$	MS1	1.0	10Fe01
$^{64}\text{Co}^m-^{32}\text{S O}_2$	-25974	12	-25976	4	-0.1	1	13	$^{13}\text{Co}^m$	MS1	1.0	10Fe01
$\text{C}_5 \text{ H}_4-^{64}\text{Ni}$	103278	10	103333.8	0.5	1.4	U			R10	4.0	74De22
$\text{C}_4 \text{ }^{13}\text{C H}_3-^{64}\text{Ni}$	98809	12	98863.6	0.5	1.1	U			R10	4.0	74De22
$\text{C}_4 \text{ H}_2 \text{ N}-^{64}\text{Ni}$	90703	16	90757.7	0.5	0.9	U			R10	4.0	74De22
$^{64}\text{Ni}-^{85}\text{Rb}_{.753}$	-5609.2	1.4	-5611.3	0.5	-1.5	1	13	$^{13}\text{Ni}$	MA8	1.0	07Gu09
$^{64}\text{Zn}-^{85}\text{Rb}_{.753}$	-4430.1	8.4	-4435.9	0.7	-0.7	U			MA8	1.0	07Ke09
$^{64}\text{Ga}-^{85}\text{Rb}_{.753}$	3261.3	2.5	3262.7	1.5	0.6	1	38	$^{38}\text{Ga}$	MA8	1.0	07Gu09
$\text{C}_5 \text{ H}_2-^{64}\text{Ga}_{.969}$	76851.5	2.6	76851.7	1.5	0.1	1	33	$^{33}\text{Ga}$	MS1	1.0	07Sc24
$^{64}\text{Ge}-u$	-57090	690	-58310	4	-1.2	U			GA6	1.5	02Li24
$\text{H }^{32}\text{S O}_2-^{64}\text{Ge H}$	20210.5	4.0				2			MS1	1.0	07Sc24
$^{64}\text{Ge}-^{85}\text{Rb}_{.753}$	8070	43	8112	4	1.0	U			MS1	1.0	12Sc.A
$^{64}\text{Ga}-^{64}\text{Zn}$	7698.5	4.1	7698.6	1.6	0.0	1	15	$^{15}\text{Ga}$	CP1	1.0	07CI01
$^{64}\text{Ge}-^{64}\text{Zn}$	12517	33	12548	4	0.9	U			CP1	1.0	07CI01
$^{64}\text{Ni}-^{63}\text{Cu}$	-1523	30	-1630.90	0.22	-0.9	U			R10	4.0	74De22
$^{64}\text{Ga}-^{63}\text{Ga}$	-2730	150	-2453.8	2.1	0.7	U			CR1	2.5	89Sh10
$^{64}\text{Ni}-^{62}\text{Ni}$	-352	25	-378.53	0.23	-0.3	U			R10	4.0	74De22
$^{64}\text{Ni}(^3\text{He}, ^8\text{B})^{59}\text{Mn}$	-19610	30	-19564.0	2.6	1.5	U			MSU		76Ka24
$^{64}\text{Ni}(^3\text{He}, ^7\text{Be})^{60}\text{Fe}$	-6511	10	-6524	3	-1.3	R			MSU		76St11
$^{64}\text{Ni}(\alpha, ^7\text{Be})^{61}\text{Fe}$	-21523	20	-21522.5	2.7	0.0	U			Tex		77Co08
$^{64}\text{Ni}(^{14}\text{C}, ^{17}\text{O})^{61}\text{Fe}$	-4609	100	-4349.8	2.7	2.6	U			Ors		84Be.A *
$^{64}\text{Ni}(p, \alpha)^{61}\text{Co}$	663.2	0.7				2			NDm		74Jo14
$^{64}\text{Zn}(p, \alpha)^{61}\text{Cu}$	830	15	844.1	0.7	0.9	U					67Br10
	830	10			1.4	U			Min		67Jo03
	844.1	0.7				2			NDm		76Jo01
$^{64}\text{Zn}(^3\text{He}, ^6\text{He})^{61}\text{Zn}$	-12331	23	-12316	16	0.7	-			MSU		79We02
	ave.	-12320	16			0.3	1	95	$^{95}\text{Zn}$		average
$^{64}\text{Ni}(^{11}\text{B}, ^{13}\text{N})^{62}\text{Fe}$	-4930	70	-4898.6	2.8	0.4	U			Tex		77Co08
$^{64}\text{Ni}(^{14}\text{C}, ^{16}\text{O})^{62}\text{Fe}$	-501	40	-464.0	2.8	0.9	U			Ors		81Be40
$^{64}\text{Ni}(^{18}\text{O}, ^{20}\text{Ne})^{62}\text{Fe}$	-1915	50	-1961.8	2.8	-0.9	U			Can		76Hi14
	-1920	21			-2.0	U			Hei		77Bh03 *
	-1947	26			-0.6	U			Hei		84Ha31
$^{64}\text{Ni}(d, \alpha)^{62}\text{Co}$	5190	20	5036	19	-7.7	B					72Ba31
$^{62}\text{Ni}(t, p)^{64}\text{Ni}$	7999	20	8013.44	0.21	0.7	U			Ald		71Da16
$^{62}\text{Ni}(^3\text{He}, p)^{64}\text{Cu}$	6299	25	6320.47	0.11	0.9	U			MIT		67Sp09
$^{62}\text{Ni}(^3\text{He}, n)^{64}\text{Zn}$	6118	12	6117.6	0.6	0.0	U			Oak		72Gr39
$^{64}\text{Zn}(d, \alpha)^{62}\text{Cu}$	7508	15	7494.2	0.8	-0.9	U			MIT		67Sp09
$^{64}\text{Zn}(p, t)^{62}\text{Zn}$	-12493	10	-12496.9	0.8	-0.4	U			Bld		72Fa08
$^{64}\text{Ni}(^{14}\text{C}, ^{15}\text{O})^{63}\text{Fe}$	-11387	60	-11299	4	1.5	U			Ors		82De.A *
$^{64}\text{Ni}(^{34}\text{S}, ^{35}\text{Ar})^{63}\text{Fe}$	-17931	260	-18348	4	-1.6	U			Hei		83Wi.B
$^{64}\text{Ni}(t, \alpha)^{63}\text{Co}$	7266	20	7277	19	0.6	1	86	$^{86}\text{Co}$	LAl		66B115
	9657.32	0.4	9657.46	0.20	0.4	-					75Wi06 Z
$^{63}\text{Ni}(n, \gamma)^{64}\text{Ni}$	9657.58	0.24			-0.5	-			ILn		92Ha21
	ave.	9657.51	0.21			-0.2	1	98	$^{87}\text{Ni}$		average
	7916.07	0.12	7916.11	0.10	0.3	-			BNn		83De28 Z
$^{63}\text{Cu}(n, \gamma)^{64}\text{Cu}$	7915.52	0.08			7.4	B					02Bo11
	7916.14	0.16			-0.2	-			Bdn		06Fi.A
	5697	8	5691.54	0.10	-0.7	U			MIT		64Sp12
ave.	7916.10	0.10	7916.11	0.10	0.2	1	99	$^{90}\text{Cu}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{64}\text{Zn}(n,d)^{63}\text{Cu}$	-5520	50	-5488.7	0.6	0.6	U					65Wa14	
$^{64}\text{Zn}(d,t)^{63}\text{Zn}$	-5604.9	1.7	-5604.7	1.5	0.1	1	76	73 $^{63}\text{Zn}$	NDm		76Jo01	
$^{64}\text{Co}(\beta^-)^{64}\text{Ni}$	7000	500	7307	20	0.6	U					69Wa15	
	7000	400			0.8	U					74Ra31	
$^{64}\text{Ni}(t,^3\text{He})^{64}\text{Co}$	-7288	20				2			LAI		72FI17	
$^{64}\text{Cu}(\beta^+)^{64}\text{Ni}$	1673.4	1.0	1674.38	0.23	1.0	U					83Ch47	
$^{64}\text{Ni}(p,n)^{64}\text{Cu}$	-2458	6	-2456.72	0.23	0.2	U					61Va19	
	-2458.22	0.31				B			PTB		92Bo02	
$^{64}\text{Cu}(\beta^-)^{64}\text{Zn}$	577.8	1.0	579.5	0.6	1.7	1	42	32 $^{64}\text{Zn}$			83Ch47	
$^{64}\text{Ga}(\beta^+)^{64}\text{Zn}$	7072	30	7171.2	1.5	3.3	B					60Ja07	
$^{64}\text{Zn}(p,n)^{64}\text{Ga}$	-7951	4	-7953.5	1.5	-0.6	1	14	12 $^{64}\text{Ga}$	Tex		72Da.A	
$^{64}\text{Zn}(^3\text{He},t)^{64}\text{Ga}$	-7206	8	-7189.8	1.5	2.0	U			MSU		74Ro16 *	
$^{64}\text{Zn}(^3\text{He},t)^{64}\text{Ga}^i$	-9141	17	-9096.8	2.5	2.6	U			MIT		70Hi06	
	-9110	6				1	17	17 $^{64}\text{Ga}^i$			71Be29 *	
$^{64}\text{Ga}^i(\text{IT})^{64}\text{Ga}$	1905.1	2.3	1907.0	2.2	0.8	1	88	83 $^{64}\text{Ga}^i$			74Ro16	
$^{64}\text{Ge}(\beta^+)^{64}\text{Ga}$	4410	250	4517	4	0.4	U					73Da01 *	
* $^{64}\text{Mn}-u$	Original -45270(350) $\mu\text{u}$ or $M-A=-42170(330)$ keV										GAu	**
* $^{64}\text{Mn}-u$	Original -46270(350) $\mu\text{u}$ or $M-A=-43100(330)$ keV										GAu	**
* $^{64}\text{Mn}-u$	$M-A=-42430(280)$ keV for mixture gs+m at 174.1(0.5) ( $4^+$ ) keV										Nub16b	**
* $^{64}\text{Ni}(^{14}\text{C},^{17}\text{O})^{61}\text{Fe}$	Cited in reference and confirmed in PrvCom sep 88										84De33	**
* $^{64}\text{Ni}(^{18}\text{O},^{20}\text{Ne})^{62}\text{Fe}$	$Q-Q(^{62}\text{Ni}(^{18}\text{O},^{20}\text{Ne}))=-2843(20)$ , $Q(62)=923(4)$ keV										AHW	**
* $^{64}\text{Ni}(^{14}\text{C},^{15}\text{O})^{63}\text{Fe}$	Original -11743(60) reinterpreted as ( $3/2^-$ ) 356.2 level in $^{63}\text{Fe}$										GAu	**
* $^{64}\text{Zn}(^3\text{He},t)^{64}\text{Ga}$	$M-A=-58819(8)$ ; $Q$ rebuilt with Ame1971										GAu	**
* $^{64}\text{Zn}(^3\text{He},t)^{64}\text{Ga}^i$	CDE=9879(6) $Q=-9115(6)$ ; recalibration +5 keV for $^{58}\text{Ni}(p,n)^{58}\text{Cu}^i$ from Ame1961										MMC124**	
* $^{64}\text{Ge}(\beta^+)^{64}\text{Ga}$	$E_{\beta^+}=2960(250)$ to ( $1^+$ ) level at 427.03 keV										Ens073	**
$^{65}\text{Mn}-u$	-43900	600	-43980	4	-0.1	U			TO5	1.5	94Se12	
	-43500	500			-0.6	U			TO6	1.5	98Ba.A	
	-43790	330			-0.6	U			MT1	1.0	11Es06	
$^{65}\text{Mn}-^{85}\text{Rb}_{.765}$	23500.6	4.0				2			MA8	1.0	12Na15	
$^{65}\text{Fe}-u$	-54680	300	-54985	5	-0.7	U			TO3	1.5	90Tu01 *	
	-55270	320			0.6	U			TO5	1.5	94Se12 *	
	-54290	380			-1.2	U			TO6	1.5	98Ba.A *	
	-54985.9	5.4			0.2	o			MS1	1.0	08B105 *	
$\text{O}_2-^{65}\text{Fe}_{.492}$	16881.7	2.7				2			MS1	1.0	10Fe01 *	
$^{65}\text{Co}-u$	-63537.9	2.3	-63537.9	2.2	0.0	o			MS1	1.0	08B105	
$\text{O}_2-^{65}\text{Co}_{.492}$	21089.9	1.1				2			MS1	1.0	10Fe01	
$^{65}\text{Ni}-^{85}\text{Rb}_{.765}$	-2438.0	2.4	-2434.5	0.5	1.5	U			MA8	1.0	07Gu09	
$\text{C}_5 \text{H}_5-^{65}\text{Cu}$	111384	4	111335.7	0.7	-3.0	B			R10	4.0	74De22	
$\text{C}_4 \text{H}_3 \text{N}-^{65}\text{Cu}$	98800	8	98759.6	0.7	-1.3	U			R10	4.0	74De22	
$\text{C}_4 \text{ }^{13}\text{C} \text{H}_4-^{65}\text{Cu}$	106921	18	106865.5	0.7	-0.8	U			R10	4.0	74De22	
$^{49}\text{Ti} \text{O}-^{65}\text{Cu}$	15030	10	14989.8	0.7	-1.0	U			R09	4.0	72De11	
$^{65}\text{Cu}-^{85}\text{Rb}_{.765}$	-4730.6	1.2	-4729.7	0.7	0.8	1	34	34 $^{65}\text{Cu}$	MA8	1.0	07Gu09	
$^{65}\text{Ga}-^{85}\text{Rb}_{.765}$	215.4	1.5	215.2	0.9	-0.1	1	34	34 $^{65}\text{Ga}$	MA8	1.0	07Gu09	
$^{65}\text{Ge}-u$	-60080	270	-60631.9	2.3	-1.4	U			GA6	1.5	02Li24	
$\text{C}_5 \text{H}_2-^{65}\text{Ge}_{.939}$	72585.2	4.0	72583.4	2.2	-0.5	-			MS1	1.0	07Sc24 *	
$\text{C}_5 \text{H}_5-^{65}\text{Ge}_{.985}$	98847.2	4.2	98847.5	2.3	0.1	-			MS1	1.0	07Sc24 *	
$\text{C}_5 \text{H}_2-^{65}\text{Ge}_{.939}$	ave.	72584.2	72583.4	2.2	-0.3	1	57	57 $^{65}\text{Ge}$			average	
$^{65}\text{Ge} \text{H}-^{85}\text{Rb}_{.776}$	15634.4	6.2	15644.3	2.3	1.6	1	14	14 $^{65}\text{Ge}$	MS1	1.0	07Sc24 *	
$^{65}\text{Ge} \text{O} \text{H}-^{85}\text{Rb}_{.965}$	27237.1	4.3	27230.7	2.3	-1.5	1	29	29 $^{65}\text{Ge}$	MS1	1.0	12Sc.A	
$^{65}\text{As}-u$	-50389	91				2			LZ1	1.0	11Tu02	
$^{65}\text{Cu}-^{64}\text{Ni}$	-275	40	-176.9	0.7	0.6	U			R10	4.0	74De22	
$^{65}\text{Cu}-^{63}\text{Cu}$	-1784	10	-1807.7	0.7	-0.6	U			R10	4.0	74De22	
$^{65}\text{Cu}(p,\alpha)^{62}\text{Ni}$	4345	8	4346.7	0.7	0.2	U			MIT		64Sp12	
	4340	10			0.7	U			Min		67Jo03	
	4344.6	1.8			1.2	1	14	10 $^{65}\text{Cu}$	NDm		76Jo01	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{62}\text{Ni}(\alpha, n)^{65}\text{Zn}$	-6510	40	-6480.7	0.7	0.7	U			Oak		64St01
$^{65}\text{Cu}(d, \alpha)^{63}\text{Ni}$	9012	40	8959.9	0.7	-1.3	U					67Hj01
$^{63}\text{Cu}(t, p)^{65}\text{Cu}$	9351	25	9344.7	0.7	-0.3	U			Ald		66Bj02
$^{64}\text{Ni}(n, \gamma)^{65}\text{Ni}$	6097.86	0.20	6098.08	0.14	1.1	2			MMn		77Is01 Z
	6098.28	0.19			-1.0	2			Bdn		06Fi.A
$^{64}\text{Ni}(d, p)^{65}\text{Ni}$	3876	6	3873.51	0.14	-0.4	U			MIT		64Sp12
	3867	15			0.4	U			Ald		67Te02
	3870	5			0.7	U			MIT		70An25
$^{65}\text{Cu}(t, \alpha)^{64}\text{Ni}$	12352	11	12360.2	0.7	0.7	U					72He23
$^{65}\text{Cu}(\gamma, n)^{64}\text{Cu}$	-9896	28	-9910.4	0.7	-0.5	U			Phi		60Ge01
$^{65}\text{Cu}(d, t)^{64}\text{Cu}$	-3650	60	-3653.2	0.7	-0.1	U			ANL		60Ze02
$^{64}\text{Zn}(n, \gamma)^{65}\text{Zn}$	7979.3	0.8	7979.32	0.17	0.0	U					71Ot01 Z
	7979.2	0.5			0.2	U					75De.A Z
	7979.28	0.17			0.2	1	98	55 $^{65}\text{Zn}$	Bdn		06Fi.A
$^{64}\text{Zn}(d, p)^{65}\text{Zn}$	5758	10	5754.76	0.17	-0.3	U			ANL		67Vo05
$^{64}\text{Zn}(p, \gamma)^{65}\text{Ga}$	3942.0	1.0	3942.5	0.6	0.5	-					75We24 Z
	3943.0	1.0			-0.5	-					87Vi01
	ave.	3942.5	0.7		-0.1	1	83	66 $^{65}\text{Ga}$			average
$^{65}\text{Ge}(\epsilon p)^{64}\text{Zn}$	2300	100	2236.8	2.3	-0.6	U			ChR		81Ha44
$^{65}\text{As}^i(p)^{64}\text{Ge}$	3603	30	3576	17	-0.9	3					93Ba12
	3564	20			0.6	3					11Ro47 *
$^{65}\text{Ni}(\beta^-)^{65}\text{Cu}$	2140	10	2138.0	0.7	-0.2	U					64Fr04
$^{65}\text{Zn}(\beta^+)^{65}\text{Cu}$	1347	2	1351.6	0.4	2.3	U					49Ma57
	1347	2			2.3	U					53Ba82
	1347	3			1.5	U					53Pe14
	1349	3			0.9	U					53Sa26
	1342	4			2.4	U					53Yu04
	1346	4			1.4	U					56Av28
$^{65}\text{Cu}(p, n)^{65}\text{Zn}$	-2131.4	1.5	-2134.0	0.4	-1.7	U					56Ma14 Z
	-2135.8	2.5			0.7	U					57Be44 Z
	-2135.3	1.8			0.7	U			Tkm		63Ok01
	-2135.6	1.7			0.9	U			Oak		64Jo11 Z
	-2134.6	0.8			0.7	-			Yal		69Ov01 Z
	ave.	-2133.55	0.43			-1.0	-		PTB		89Sc24
$^{65}\text{Ga}(\beta^+)^{65}\text{Zn}$	3277	30	3254.5	0.7	-0.7	U					average
$^{65}\text{Ge}(\beta^+)^{65}\text{Ga}$	5220	400	6179.3	2.3	2.4	U					57Da07
* $^{65}\text{Fe}-u$	$M - A = -50740(250)$ keV for mixture gs+m at 393.7(0.2) keV										Nub16b **
* $^{65}\text{Fe}-u$	$M - A = -51290(280)$ keV for mixture gs+m at 393.7(0.2) keV										Nub16b **
* $^{65}\text{Fe}-u$	$M - A = -50370(330)$ keV for mixture gs+m at 393.7(0.2) keV and assuming ratio $R=0.13(6)$ , from half-life=430 ns and TOF=1 $\mu\text{s}$										Nub16b **
* $^{65}\text{Fe}-u$	$D_M = -54988.9(7.1)$ $\mu\text{u}$ to ground state, $D_M = -54557.0(8.4)$ to $^{65}\text{Fe}^m$ at 393.7 keV										GAu **
* $\text{O}_2 - ^{65}\text{Fe}_{.492}$	$D_M = 16883.6(3.6)$ $\mu\text{u}$ to ground state, $D_M = 16671.2(4.2)$ to $^{65}\text{Fe}^m$ at 393.7 keV										Nub16b **
* $\text{C}_5 \text{H}_2 - ^{65}\text{Ge}_{.939}$	For original doublet $\text{C}_5 \text{H}_2 - (^{65}\text{Ge} \text{H})_{0.939}$ , $D_M = 65237.5(4.0)$ $\mu\text{u}$										GAu **
* $\text{C}_5 \text{H}_5 - ^{65}\text{Ge}_{.985}$	For original doublet $\text{C}_5 \text{H}_5 - (^{65}\text{Ge} \text{H})_{0.985}$ , $D_M = 91139.5(4.2)$ $\mu\text{u}$										GAu **
* $^{65}\text{Ge} \text{H} - ^{85}\text{Rb}_{.776}$	Combining their 3 results yields a precision for $^{65}\text{Ge}$ of 2.4 keV, while their Table III gives 1.2 keV										GAu **
* $^{65}\text{As}^i(p)^{64}\text{Ge}$	And $E_p = 2620(30)$ to 901.7 level										GAu **
$^{66}\text{Mn}-u$	-39860	860	-39453	12	0.5	U			MT1	1.0	11Es06 *
$^{66}\text{Mn} - ^{85}\text{Rb}_{.776}$	28998	12				2			MA8	1.0	12Na15
$^{66}\text{Fe}-u$	-52300	700	-53750	4	-1.4	U			TO3	1.5	90Tu01
	-54020	350			0.5	U			TO5	1.5	94Se12
	-52800	300			-2.1	U			TO6	1.5	98Ba.A
	-53935	150			0.8	U			GT1	1.5	04Ma.A



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{66}\text{Fe}-^{28}\text{Si F}_2$	-27482.9	4.4				2			MS1	1.0	10Fe01
$^{66}\text{Co}-\text{u}$	-60470	300	-60557	15	-0.2	U			TO5	1.5	94Se12 *
	-59870	290			-1.6	U			TO6	1.5	98Ba.A *
$^{66}\text{Co}-\text{O C F}_2$	-52278	15	-52278	15	0.0	o			MS1	1.0	08B105
	-52278	15				2			MS1	1.0	10Fe01
$^{66}\text{Ni}-^{85}\text{Rb}_{.776}$	-2409.5	1.5				2			MA8	1.0	07Gu09
$^{66}\text{Cu}-^{85}\text{Rb}_{.776}$	-2680.6	2.2	-2680.0	0.7	0.3	1	10	10 $^{66}\text{Cu}$	MA8	1.0	07Gu09
$\text{C}_5 \text{H}_5-^{66}\text{Ge}_{.970}$	103278.9	2.5				2			MS1	1.0	07Sc24 *
$^{66}\text{As}-\text{u}$	-55290	730	-55851	6	-0.5	U			GA6	1.5	02Li24
$^{66}\text{As}-^{85}\text{Rb}_{.776}$	12607	32	12600	6	-0.2	U			MS1	1.0	07Sc24
$^{66}\text{As O}-^{85}\text{Rb}_{.965}$	24186.3	6.1				2			MS1	1.0	12Sc.A
$^{66}\text{Zn}(\text{p},\alpha)^{63}\text{Cu}$	1544.3	0.8	1544.6	0.7	0.4	1	74	66 $^{66}\text{Zn}$	NDm		76Jo01
$^{63}\text{Cu}(\alpha,\text{n})^{66}\text{Ga}$	-7670	30	-7502.5	1.1	5.6	B			Oak		64St01
$^{64}\text{Ni}(\text{t,p})^{66}\text{Ni}$	6559	25	6568.2	1.5	0.4	U			Ald		71Da16
$^{64}\text{Zn}(\text{t,p})^{66}\text{Zn}$	10582	15	10556.0	0.9	-1.7	U			Ald		72Hu06
$^{65}\text{Cu}(\text{n},\gamma)^{66}\text{Cu}$	7065.80	0.12	7065.93	0.09	1.1	-			BNn		83De29 Z
	7066.13	0.15			-1.3	-			Bdn		06Fi.A
$^{65}\text{Cu}(\text{d,p})^{66}\text{Cu}$	4837	8	4841.36	0.09	0.5	U			MIT		64Sp12
$^{65}\text{Cu}(\text{n},\gamma)^{66}\text{Cu}$	ave. 7065.93	0.09	7065.93	0.09	0.0	1	100	90 $^{66}\text{Cu}$			average
$^{66}\text{Zn}(\text{d,t})^{65}\text{Zn}$	-4770	60	-4801.2	0.9	-0.5	U			ANL		60Ze02
$^{66}\text{Co}(\beta^-)^{66}\text{Ni}$	9700	500	9598	14	-0.2	U					88Bo06
$^{66}\text{Ni}(\beta^-)^{66}\text{Cu}$	200	30	252.0	1.5	1.7	U					56Jo20
$^{66}\text{Cu}(\beta^-)^{66}\text{Zn}$	2650	30	2640.9	0.9	-0.3	U					51Fr19 *
	2650	30			-0.3	U					56Jo20
$^{66}\text{Ga}(\beta^+)^{66}\text{Zn}$	5175.0	3.0	5175.5	0.8	0.2	U					63Ca03 *
	5175.5	0.8				2					14Se12
$^{66}\text{Zn}(\beta^+)^{66}\text{Ga}^i$	-9044	6				2					71Be29 *
$^{66}\text{Ge}(\beta^+)^{66}\text{Ga}$	2490	50	2116.6	2.6	-7.5	F					69Ba31 *
	2420	30			-10.1	F					69Sa08 *
	2100	30			0.6	U					70De39 *
$^{66}\text{As}(\beta^+)^{66}\text{Ge}$	9550	50	9582	6	0.6	U			ANB		79Da.A
$^{66}\text{Mn}-\text{u}$	$M - A = -36900(790)$ keV for mixture gs+m at 464.5 keV ( $5^-$ )										Nub16b **
$^{66}\text{Co}-\text{u}$	Original $-60160(300)$ $\mu\text{u}$ or $M - A = -56040(280)$ keV										GAu **
$^{66}\text{Co}-\text{u}$	$M - A = -55480(270)$ keV for mixture gs+m+n at 175.1 and 642(5) keV										Nub16b **
*	and assuming for first isomer a ratio $R=0.5(0.2)$ to ground state,										GAu **
*	from half-life= $1.21 \mu\text{s}$ and TOF= $1 \mu\text{s}$										GAu **
$^{66}\text{C}_5 \text{H}_5-^{66}\text{Ge}_{.970}$	For original doublet $\text{C}_5 \text{H}_5-(^{66}\text{Ge H})_{0.970}$ , $D_M=95688.6(2.5)$ $\mu\text{u}$										GAu **
$^{66}\text{Cu}(\beta^-)^{66}\text{Zn}$	$E_{\beta^-} = 2630(30)$ 1640(30) to ground state and $2^+$ level at 1039.2279 keV										Ens104 **
$^{66}\text{Ga}(\beta^+)^{66}\text{Zn}$	Original $E_{\beta^+} = 4153(3)$ corrected for recoil										14Se12 **
$^{66}\text{Zn}(\beta^+)^{66}\text{Ga}^i$	CDE= $9813(6)$ $Q = -9049(6)$ ; recalibration +5 keV for $^{58}\text{Ni}(\text{p,n})^{58}\text{Cu}^i$ from Ame1961										MMC124**
$^{66}\text{Ge}(\beta^+)^{66}\text{Ga}$	$E_{\beta^+} = 1440(50)$ to 43.9 level; F : probably distorted by annihilation pile up										AHW **
$^{66}\text{Ge}(\beta^+)^{66}\text{Ga}$	$E_{\beta^+} = 1370(30)$ to 43.9 level; F : probably distorted by annihilation pile up										AHW **
$^{66}\text{Ge}(\beta^+)^{66}\text{Ga}$	$E_{\beta^+} = 1028(30)$ , 668(30), 558(50) to 43.812 $1^+$ , 381.859 $1^+$ , 536.618 $1^+$ level										Ens104 **
$^{67}\text{Mn}-\text{u}$	-36600	670	-35920#	320#	1.0	D			MT1	1.0	15Me.A *
$^{67}\text{Fe}-\text{u}$	-50190	500	-48960	290	1.6	2			TO5	1.5	94Se12 *
	-48450	380			-0.9	2			TO6	1.5	98Ba.A *
	-49641	440			1.0	2			GT1	1.5	04Ma.A
	-49580	300			2.1	o			MT1	1.0	11Es06 *
	-48510	460			-1.0	2			MT1	1.0	15Me.A
$^{67}\text{Co}-\text{u}$	-59390	300	-59390	7	0.0	U			TO5	1.5	94Se12
	-58730	350			-1.3	U			TO6	1.5	98Ba.A
$^{28}\text{Si F}_2-^{67}\text{Co}_{.985}$	32232.9	8.0	32232	7	-0.1	2			MS1	1.0	10Fe01
	32231	13			0.1	2			MS1	1.0	10Fe01 *
$^{67}\text{Ni}-\text{u}$	-68370	430	-68431	3	-0.1	U			TO5	1.5	94Se12 *
	-68090	470			-0.5	U			TO6	1.5	98Ba.A *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{67}\text{Ni}-^{85}\text{Rb}_{.788}$	1079.1	3.1				2			MA8	1.0	07Gu09
$^{67}\text{Cu}-^{85}\text{Rb}_{.788}$	-2760.0	1.3	-2760.8	1.0	-0.6	1	54	54 $^{67}\text{Cu}$	MA8	1.0	07Gu09
$^{67}\text{As}-u$	-60500	260	-60748.9	0.5	-0.6	U			GA6	1.5	02Li24
$^{67}\text{As}-^{86}\text{Kr}_{.779}$	8885.7	2.9	8885.4	0.5	-0.1	U			MS1	1.0	07Sc24
	8808	30			2.6	U			MS1	1.0	07Sc24
$^{67}\text{As}-^{85}\text{Rb}_{.788}$	8762.5	1.7	8760.8	0.5	-1.0	U			MS1	1.0	07Sc24
	8760.87	0.54			-0.1	1	77	77 $^{67}\text{As}$	MS1	1.0	12Sc.A
$^{67}\text{As O}-^{85}\text{Rb}_{.976}$	20258.7	1.0	20258.9	0.5	0.2	1	23	23 $^{67}\text{As}$	MS1	1.0	12Sc.A
$^{67}\text{Se}-u$	-50006	72				2			LZ1	1.0	11Tu02
$^{67}\text{Zn N}-^{66}\text{Zn }^{15}\text{N}$	4060.21	0.32	4058.88	0.25	-1.7	U			H30	2.5	77Ba10 *
$^{64}\text{Zn}(\alpha,n)^{67}\text{Ge}$	-9240	60	-8992	5	4.1	B			Oak		64St01
	-8987.5	12.			-0.4	2			ANL		78Mu05
	-8993	5			0.2	2					79Al04
$^{65}\text{Cu}(t,p)^{67}\text{Cu}$	7716	25	7716.7	1.1	0.0	U			Ald		66Bj02
$^{65}\text{Cu}(^3\text{He,p})^{67}\text{Zn}$	8185	40	8258.9	0.9	1.8	U			MIT		74Is01
$^{66}\text{Zn}(n,\gamma)^{67}\text{Zn}$	7052.5	0.6	7052.47	0.23	-0.1	-					71Ot01 Z
	7052.5	0.5			-0.1	-					75De.A Z
	7052.5	0.3			-0.1	-			Bdn		06Fi.A
$^{66}\text{Zn}(d,p)^{67}\text{Zn}$	4827	10	4827.90	0.23	0.1	U			ANL		67Vo05
	4820	5			1.6	U			MIT		74Is01
$^{67}\text{Zn}(d,t)^{66}\text{Zn}$	-800	60	-795.24	0.23	0.1	U			ANL		60Ze02
$^{66}\text{Zn}(n,\gamma)^{67}\text{Zn}$	ave. 7052.50	0.24	7052.47	0.23	-0.1	1	98	64 $^{67}\text{Zn}$			average
$^{67}\text{Ni}(\beta^-)^{67}\text{Cu}$	3830	90	3577	3	-2.8	U					75Re09
$^{67}\text{Cu}(\beta^-)^{67}\text{Zn}$	577	8	560.8	0.8	-2.0	U					53Ea11
	560.3	1.0			0.5	1	69	46 $^{67}\text{Cu}$	ANB		15Ch57
$^{67}\text{Zn}(p,n)^{67}\text{Ga}$	-1776	5	-1783.6	1.1	-1.5	U			Ric		57Ch30 Y
	-1783.3	1.4			-0.2	1	66	55 $^{67}\text{Ga}$	Oak		64Jo11 Z
$^{67}\text{Ge}(\beta^+)^{67}\text{Ga}$	4330	100	4221	5	-1.1	U					59Ri35 *
	4370	150			-1.0	U					69Ba07 *
$^{67}\text{As}(\beta^+)^{67}\text{Ge}$	6010	100	6071	5	0.6	U			ANB		80Mu12
$^{*67}\text{Mn}-u$	Trends from Mass Surface TMS suggest $^{67}\text{Mn}$ 630 less bound										GAu **
$^{*67}\text{Fe}-u$	Original -50000(500) $\mu\text{u}$ or $M-A=-46570(470)$ keV										GAu **
$^{*67}\text{Fe}-u$	$M-A=-44930(330)$ keV for mixture gs+m at 402(9) keV										Nub16b **
$^{*67}\text{Fe}-u$	$M-A=-45980(250)$ keV for mixture gs+m at 402(9) keV										Nub16b **
$^{*28}\text{Si F}_2-^{67}\text{Co}_{.985}$	$M-A=-54829(12)$ for $^{67}\text{Co}^m$ at 491.55(0.11) keV										10Fe01 **
$^{*67}\text{Ni}-u$	Original -67840(300) $\mu\text{u}$ or $M-A=-63190(280)$ keV										GAu **
$^{*67}\text{Ni}-u$	$M-A=-62930(330)$ keV for mixture gs+m at 1006.6(0.2) keV										Nub16b **
$^{*67}\text{Zn N}-^{66}\text{Zn }^{15}\text{N}$	Original 4060.21(0.25); plus syst 0.20 $\mu\text{u}$ estimated by evaluator										GAu **
$^{*67}\text{Ge}(\beta^+)^{67}\text{Ga}$	$E_{\beta^+}=3140(100)$ 3180(100) respectively, to $1/2^-$ level at 166.98 keV										Ens05c **
$^{68}\text{Fe}-u$	-46300	500	-46690	390	-0.5	2			TO6	1.5	98Ba.A
	-47330	460			1.4	o			MT1	1.0	11Es06
	-46830	460			0.3	2			MT1	1.0	15Me.A
$^{68}\text{Co}-u$	-55640	350	-55750	200	-0.2	o			TO5	1.5	94Se12
	-54750	300			-2.2	U			TO6	1.5	98Ba.A
	-55730	140			-0.1	2			GT2	2.5	08Kn.A *
	-55760	250			0.0	2			MT1	1.0	11Es06 *
$^{68}\text{Ni}-u$	-68030	930	-68131	3	-0.1	U			TO5	1.5	94Se12 *
	-67530	930			-0.4	U			TO6	1.5	98Ba.A *
$^{68}\text{Ni}-^{85}\text{Rb}_{.800}$	2437.0	3.2				2			MA8	1.0	07Gu09
$^{68}\text{Cu}-u$	-70570	440	-70389.1	1.7	0.3	U			TO6	1.5	98Ba.A *
$^{68}\text{Cu}-^{85}\text{Rb}_{.800}$	179.1	1.7				2			MA8	1.0	07Gu09 *
$^{68}\text{Ga}-^{85}\text{Rb}_{.800}$	-1484	37	-1451.6	1.5	0.9	U			MA8	1.0	07Gu09
$^{68}\text{Ge}-\text{C}_5\text{H}_8$	-134496.7	8.6	-134504.9	2.0	-1.0	U			CP1	1.0	04Cl03
	-134506.3	2.8			0.5	2			CP1	1.0	04Cl03
	-134503.5	2.9			-0.5	2			CP1	1.0	04Cl03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{68}\text{As}-u$	-63221	107	-63225.9	2.0	0.0	U			GT1	1.5	01Ha66	
$^{68}\text{As}-\text{C}_5\text{H}_8$	-125839	13	-125826.1	2.0	1.0	U			CP1	1.0	04Cl03	
	-125827.7	9.9			0.2	U			CP1	1.0	04Cl03	
	-125827.1	2.9			0.3	-			CP1	1.0	04Cl03	
	-125824.4	3.1			-0.6	-			CP1	1.0	04Cl03	
	ave.	-125825.8	2.1			-0.1	1	88	88 $^{68}\text{As}$			average
$\text{C F}_3-^{68}\text{As}_{1.015}$	59385.8	5.7	59383.7	2.0	-0.4	1	12	12 $^{68}\text{As}$	MS1	1.0	07Sc24	
$^{68}\text{Se}-u$	-56197	86	-58174.8	0.5	-23.0	F			CS1	1.0	01La31 *	
	-57560	1070			-0.4	U			GA6	1.5	02Li24	
	-57900	300			-0.9	U			CS1	1.0	08Go23	
$^{68}\text{Se}-\text{C}_5\text{H}_8$	-120801	31	-120775.0	0.5	0.8	U			CP1	1.0	04Cl03	
$\text{C F}_3-^{68}\text{Se}_{1.015}$	54256.87	0.54				2		MS1	1.0	09Sa12		
$^{68}\text{Zn }^{35}\text{Cl}-^{66}\text{Zn }^{37}\text{Cl}$	1757.9	1.0	1760.7	0.3	0.7	U			H18	4.0	64Ba03	
$^{68}\text{As}-^{68}\text{Ge}$	8698.8	9.9	8678.8	2.8	-2.0	U			CP1	1.0	04Cl03	
$^{68}\text{Se}-^{68}\text{Ge}$	13669	27	13729.9	2.1	2.3	U			CP1	1.0	04Cl03	
$^{65}\text{Cu}(\alpha,n)^{68}\text{Ga}$	-5800	40	-5824.1	1.5	-0.6	U			Oak		64St01	
$^{66}\text{Ni}(t,p)^{68}\text{Ni}-^{68}\text{Zn}()^{70}\text{Zn}$	-2110	21	-2100	4	0.5	U			Hei		77Bh03	
$^{66}\text{Zn}(t,p)^{68}\text{Zn}$	8758	15	8768.8	0.3	0.7	U			Ald		72Hu06	
$^{68}\text{Zn}(^{14}\text{C}, ^{15}\text{O})^{67}\text{Ni}$	-6052	150	-6100	3	-0.3	U			Ors		84De33	
	$^{67}\text{Zn}(n,\gamma)^{68}\text{Zn}$	10198.2	0.4	10198.10	0.19	-0.3	-					71Ot01 Z
	10198.06	0.22			0.2	-			Bdn		06Fi.A	
$^{68}\text{Zn}(d,t)^{67}\text{Zn}$	-3930	60	-3940.87	0.19	-0.2	U			ANL		60Ze02	
$^{67}\text{Zn}(n,\gamma)^{68}\text{Zn}$	ave.	10198.09	0.19	10198.10	0.19	0.0	1	100	99 $^{68}\text{Zn}$			average
	$^{68}\text{Cu}(\beta^-)^{68}\text{Zn}$	4580	60	4440.1	1.8	-2.3	U					64Ba13
	4590	50			-3.0	B					72Sw01	
$^{68}\text{Zn}(t, ^3\text{He})^{68}\text{Cu}$	-4410	20	-4421.5	1.8	-0.6	U			LAI		77Sh08	
$^{68}\text{Ga}(\beta^+)^{68}\text{Zn}$	2921.1	1.2				2					72Si03	
	2915	10	2921.1	1.2	0.6	U					85Bo58	
$^{68}\text{Zn}(p,n)^{68}\text{Ga}$	-3693	6	-3703.4	1.2	-1.7	U			Ric		55Br16 Z	
	-3703	5			-0.1	U			Ric		57Ch30 Z	
	-3707	5			0.7	U			Oak		64Jo11 Z	
$^{68}\text{As}(\beta^+)^{68}\text{Ge}$	8100	100	8084.3	2.6	-0.2	U			ANB		77Pa13	
	8073	54			0.2	U					02Cl.A *	
$^{68}\text{Se}(\beta^+)^{68}\text{As}$	4710	200	4705.1	1.9	0.0	U					04Wo16	
$^{68}\text{Co}-u$	$M - A = -51838(96)$ keV for mixture gs+m at 150#(150#) keV											
$^{68}\text{Co}-u$	$M - A = -51860(210)$ keV for mixture gs+m at 150#(150#) keV											
$^{68}\text{Ni}-u$	$M - A = -61950(280)$ keV for mixture gs+n at 2849.1 keV											
$^{68}\text{Ni}-u$	$M - A = -61480(280)$ keV for mixture gs+n at 2849.1 keV											
$^{68}\text{Cu}-u$	$M - A = -65380(350)$ keV for mixture gs+m at 721.26 keV											
$^{68}\text{Cu}-^{85}\text{Rb}_{.800}$	This result was first published in reference											
$^{68}\text{Cu}-^{85}\text{Rb}_{.800}$	Also 948.6(1.6) $\mu\text{u}$ for $^{68}\text{Cu}^m - ^{85}\text{Rb}_{.800}$ , yielding excit. of 716.7(2.2) keV											
$^{68}\text{Se}-u$	F : other results in same paper not trusted, see $^{80}\text{Y}$ and $^{80}\text{Zr}$											
$^{68}\text{As}(\beta^+)^{68}\text{Ge}$	From mass difference 8667(64) $\mu\text{u}$											
$^{69}\text{Fe}-u$	-42240	640	-41900#	430#	0.5	D			MT1	1.0	15Me.A *	
$^{69}\text{Co}-u$	-54800	400	-53980	150	1.4	o			TO5	1.5	94Se12	
	-53050	300			-2.1	2			TO6	1.5	98Ba.A	
	-54070	230			0.4	2			MT1	1.0	11Es06	
	-54117	223			0.6	2			LZ1	1.0	15Xu14	
$^{69}\text{Ni}-u$	-64600	400	-64390	4	0.4	U			TO5	1.5	94Se12 *	
	-64250	450			-0.2	U			TO6	1.5	98Ba.A *	
$^{69}\text{Ni}-^{85}\text{Rb}_{.812}$	7237.0	4.0				2			MA8	1.0	07Gu09	
$^{69}\text{Cu}-^{85}\text{Rb}_{.812}$	1056.0	1.5				2			MA8	1.0	07Gu09	
$^{69}\text{Zn}-u$	-73580	400	-73449.6	0.9	0.2	U			TO6	1.5	98Ba.A *	
$\text{C}_5\text{H}_9-^{69}\text{Ga}$	144852.7	2.4	144851.8	1.3	-0.2	U			M15	2.5	63Ri07	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{69}\text{Ga}-^{85}\text{Rb}_{.812}$	-2799.8	1.6	-2799.7	1.3	0.0	1	65	65 $^{69}\text{Ga}$	MA8	1.0	07Gu09
$\text{C F}_3-^{69}\text{Se}$	55794.7	1.6	55794.6	1.6	0.0	1	100	100 $^{69}\text{Se}$	MS1	1.0	07Sc24
$^{69}\text{Ga}(\text{p},\alpha)^{66}\text{Zn}$	4440	10	4435.4	1.4	-0.5	U			ANL		67Ka11
$^{66}\text{Zn}(\alpha,\text{n})^{69}\text{Ge}$	-7520	30	-7444.9	1.5	2.5	U			Oak		64St01
$^{67}\text{Zn}(\text{t},\text{p})^{69}\text{Zn}$	8168	20	8198.37	0.25	1.5	U			Ald		72Hu06
$^{68}\text{Zn}(\text{n},\gamma)^{69}\text{Zn}$	6482.3	0.8	6482.07	0.16	-0.3	U					71Ot01 Z
	6481.8	0.5			0.5	U					75De.A Z
	6482.07	0.16				2			Bdn		06Fi.A
$^{68}\text{Zn}(\text{d},\text{p})^{69}\text{Zn}$	4259	10	4257.50	0.16	-0.1	U			ANL		67Vo05
	4243	10			1.5	U			MIT		75Is04
$^{68}\text{Zn}(^3\text{He},\text{d})^{69}\text{Ga}$	1126	20	1116.2	1.4	-0.5	U					74Ri08
$^{69}\text{Se}(\varepsilon\text{p})^{68}\text{Ge}$	3390	50	3255.1	2.4	-2.7	U			ChR		76Ha29
	3370	70			-1.6	U			ChR		77Ma24
$^{69}\text{Br}(\text{p})^{68}\text{Se}$	789	37	640	40	-4.0	B			MSU		11Ro18 *
	641	42				3			MSU		14De41 *
$^{69}\text{Br}^i(\text{p})^{68}\text{Se}$	4131	50	4129	19	0.0	3					97Xu01 *
	4210.6	50.			-1.6	3			MSU		11Ro47 *
	4113	22			0.7	3			MSU		14De41 *
$^{69}\text{Cu}(\beta^-)^{69}\text{Zn}$	2480	70	2681.6	1.6	2.9	U					66Va12
$^{69}\text{Zn}(\beta^-)^{69}\text{Ga}$	897	5	910.0	1.4	2.6	U					53Du03
$^{69}\text{Ge}(\beta^+)^{69}\text{Ga}$	2225	15	2227.1	0.5	0.1	U					51Hu38 *
$^{69}\text{Ga}(\text{p},\text{n})^{69}\text{Ge}$	-3008.8	3.2	-3009.5	0.5	-0.2	U			Tkm		63Ok01
	-3006.0	4.			-0.9	U			Oak		64Jo11 Z
	-3009.50	0.55			0.0	1	100	100 $^{69}\text{Ge}$	PTB		92Bo.B Z
$^{69}\text{As}(\beta^+)^{69}\text{Ge}$	3972	50	3990	30	0.3	-					70Bo19
	4067	50			-1.6	-			ChR		77Ma24 *
	ave.	4020	40		-0.9	1	82	82 $^{69}\text{As}$			average
$^{69}\text{Se}(\beta^+)^{69}\text{As}$	6817	75	6680	30	-1.9	1	18	18 $^{69}\text{As}$	ChR		77Ma24 *
$^{69}\text{Fe}-\text{u}$	Trends from Mass Surface TMS suggest $^{69}\text{Fe}$ 320 less bound										
$^{69}\text{Ni}-\text{u}$	$M - A = -59940(330)$ keV for mixture gs+m+n at 321(2) and 2700.0 keV										
$^{69}\text{Ni}-\text{u}$	$M - A = -59620(380)$ keV for mixture gs+m+n at 321(2) and 2700.0 keV										
*	and assuming for second isomer a ratio $R=0.13(0.06)$ to ground state,										
*	from half-life=439 ns and TOF=1 $\mu\text{s}$										
$^{69}\text{Zn}-\text{u}$	$M - A = -68320(350)$ keV for mixture gs+m at 438.636 keV										
$^{69}\text{Br}(\text{p})^{68}\text{Se}$	Symmetrized from $Q_p=785(+40-34)$ keV										
$^{69}\text{Br}(\text{p})^{68}\text{Se}$	And $E_p=751(+132-82)$ in good agreement with previous item										
$^{69}\text{Br}^i(\text{p})^{68}\text{Se}$	Might be also a more intense peak around 3 MeV										
$^{69}\text{Br}^i(\text{p})^{68}\text{Se}$	$E_p=2970(50)$ to $(2^+)$ level at 1196.7 keV										
$^{69}\text{Br}^i(\text{p})^{68}\text{Se}$	A weaker peak around 4 MeV cannot be excluded, could feed lower $(2^+)$										
$^{69}\text{Br}^i(\text{p})^{68}\text{Se}$	Original $Q = 2939(22)$ corrected to $Q_p=2916(22)$ to $(2^+)$ level at 1196.7 keV										
$^{69}\text{Ge}(\beta^+)^{69}\text{Ga}$	$E_{\beta^+}=1215, 610$ to ground state $3/2^-, 574.220$ $5/2^-$ levels										
$^{69}\text{As}(\beta^+)^{69}\text{Ge}$	$E_{\beta^+}=2812(50)$ to $3/2^-$ level at 232.70 keV										
$^{69}\text{Se}(\beta^+)^{69}\text{As}$	$E_{\beta^+}=5006(75)$ to 789.46 $(1/2^-, 3/2^-)$ level, and others										
$^{70}\text{Co}-\text{u}$	-49000	600	-50060#	320#	-1.2	U			TO6	1.5	98Ba.A
	-50370	320			1.0	D			MT1	1.0	11Es06 *
$^{70}\text{Ni}-\text{u}$	-63980	350	-63568.7	2.3	0.8	U			TO5	1.5	94Se12 *
	-63020	350			-1.0	U			TO6	1.5	98Ba.A *
$^{70}\text{Cu}-^{85}\text{Rb}_{.824}$	5077.6	1.7	5077.3	1.2	-0.2	2			MA8	1.0	07Gu09 *
	5077.2	2.2			0.1	2			MA8	1.0	07Gu09 *
	5077.0	2.3			0.1	2			MA8	1.0	07Gu09 *
$^{70}\text{Ga}-^{85}\text{Rb}_{.824}$	-1293.0	2.3	-1292.8	1.3	0.1	1	31	31 $^{70}\text{Ga}$	MA8	1.0	07Gu09
$\text{C}_5 \text{H}_{10}-^{70}\text{Ge}$	154001.3	2.2	154001.6	0.9	0.1	U			M15	2.5	63Ri07
$\text{C}_4 \text{H}_6 \text{O}-^{70}\text{Ge}$	117616.1	1.8	117616.1	0.9	0.0	U			M15	2.5	63Ri07
$^{70}\text{Se}-\text{u}$	-66890	490	-66484.5	1.7	0.6	o			GA6	1.5	98Ch20
	-66635	75			1.3	U			GT1	1.5	01Ha66
	-66520	140			0.2	U			GA6	1.5	02Li24

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{70}\text{Se}-^{13}\text{C F}_3$	-65048.8	1.7		2			MS1	1.0	09Sa12	
$^{70}\text{Se}-^{85}\text{Rb}_{.824}$	6209	18	6200.8	1.7	-0.5	U	MA8	1.0	11He10	
$^{70}\text{Br}-^{13}\text{C F}_3$	-53772	16		2			MS1	1.0	09Sa12 *	
$^{70}\text{Ni}-^{72}\text{Ge}_{.972}$	12173.6	2.3		2			JY1	1.0	07Ra27	
$^{70}\text{Zn }^{35}\text{Cl}-^{68}\text{Zn }^{37}\text{Cl}$	3429.5	1.7	3425.0	2.2	-0.7	1	10	9 $^{70}\text{Zn}$	H18 4.0	64Ba03
$^{70}\text{Zn}(^3\text{He}, ^8\text{B})^{65}\text{Co}$	-18385	13	-18370	3	1.2	U			Pri	78Ko24
$^{70}\text{Zn}(\alpha, ^7\text{Be})^{67}\text{Ni}$	-19155	36	-19166	3	-0.3	U			Tex	78Co.A
	-19164	22			-0.1	U			Pri	78Ko28
$^{70}\text{Zn}(^{14}\text{C}, ^{17}\text{O})^{67}\text{Ni}$	-1661	100	-1993	3	-3.3	B			Ors	88Gi04
$^{70}\text{Ge}(\text{p}, \alpha)^{67}\text{Ga}$	1180.9	1.5	1181.2	1.2	0.2	1	60	45 $^{67}\text{Ga}$	NDm	76Jo01
$^{70}\text{Ge}(^3\text{He}, ^6\text{He})^{67}\text{Ge}$	-10572	30	-10565	5	0.2	U			MSU	78Pa11
$^{70}\text{Zn}(^{14}\text{C}, ^{16}\text{O})^{68}\text{Ni}$	1727	30	1656	4	-2.4	U			Ors	88Gi04
$^{70}\text{Zn}(^{18}\text{O}, ^{20}\text{Ne})^{68}\text{Ni}$	172	26	158	4	-0.5	U			Hei	84Ha31
$^{68}\text{Zn}(\text{t}, \text{p})^{70}\text{Zn}$	7196	15	7218.5	2.0	1.5	U			Ald	72Hu06
$^{70}\text{Ge}(\text{p}, \text{t})^{68}\text{Ge}$	-11251	13	-11243.9	2.1	0.5	U			ChR	72Hs01
	-11242	7			-0.3	U			Ors	77Gu02
$^{70}\text{Zn}(^{14}\text{C}, ^{15}\text{O})^{69}\text{Ni}$	-8936	150	-9422	4	-3.2	B			Ors	84De33
$^{70}\text{Zn}(\text{d}, ^3\text{He})^{69}\text{Cu}$	-5605	10	-5624.0	2.4	-1.9	U			ANL	78Ze04
	-5622	13			-0.2	U			Hei	84Ha31
$^{70}\text{Zn}(\text{t}, \alpha)^{69}\text{Cu}$	8682	20	8696.4	2.4	0.7	U			LAL	81Aj02
$^{69}\text{Ga}(\text{n}, \gamma)^{70}\text{Ga}$	7654.0	1.0	7653.65	0.17	-0.4	U				71Ar12 Z
	7651.6	1.0			2.0	F				71Ve03 *
	7653.65	0.17			0.0	1	100	64 $^{70}\text{Ga}$	Bdn	06Fi.A
$^{69}\text{Ga}(\text{d}, \text{p})^{70}\text{Ga}$	5430	10	5429.08	0.17	-0.1	U			Kop	71Ar12
$^{70}\text{Ge}(\text{d}, ^3\text{He})^{69}\text{Ga}$	-3030	7	-3029.6	1.5	0.1	U			Ors	78Ro14
$^{70}\text{Cu}(\beta^-)^{70}\text{Zn}$	6310	110	6588.4	2.2	2.5	U				75Re09 *
	5928	110			6.0	B				75Re09 *
$^{70}\text{Zn}(\text{t}, ^3\text{He})^{70}\text{Cu}$	-6559	20	-6569.8	2.2	-0.5	U			LAL	77Sh08
	-6602	20			1.6	U			LAL	87Aj.A
$^{70}\text{Zn}(\text{p}, \text{n})^{70}\text{Ga}$	-1436.3	2.0	-1436.9	1.6	-0.3	-			Nvl	59Go68 Z
	-1439.1	3.0			0.7	-			Oak	64Jo11 Z
	ave.	1.6			0.1	1	92	88 $^{70}\text{Zn}$		average
$^{70}\text{Ga}(\beta^-)^{70}\text{Ge}$	1650	10	1651.7	1.5	0.2	U				57Bu41
$^{70}\text{As}(\beta^+)^{70}\text{Ge}$	6220	50			2					63Bo14 *
$^{70}\text{Se}(\beta^+)^{70}\text{As}$	2780	200	2410	50	-1.8	F				75La02 *
	2736	85			-3.8	B				01To06
$^{70}\text{Br}(\beta^+)^{70}\text{Se}$	9970	170	10504	15	3.1	C			ANB	79Da.A
	9898	80			7.6	B				04Ka38 *
* $^{70}\text{Co}-\text{u}$	$M - A = -46820(280)$ keV for mixture gs+m at 200#200 keV									
* $^{70}\text{Co}-\text{u}$	Trends from Mass Surface TMS suggest $^{70}\text{Co}$ 290 less bound									
* $^{70}\text{Ni}-\text{u}$	Original $-63860(350)$ $\mu\text{u}$ or $M - A = -59490(330)$ keV									
* $^{70}\text{Ni}-\text{u}$	$M - A = -58590(330)$ keV for mixture gs+m at 2860(2) keV and									
*	assuming ratio $R=0.04(2)$ , from half-life= $232$ ns and $\text{TOF}=1$ $\mu\text{s}$									
* $^{70}\text{Cu}-^{85}\text{Rb}_{.824}$	The three results for $^{70}\text{Cu}$ were first published in reference									
* $^{70}\text{Cu}-^{85}\text{Rb}_{.824}$	$D_M = 5185.7(2.2)$ $\mu\text{u}$ for $^{70}\text{Cu}^m$ at 101.1(0.3) keV; $M - A = -62875.4(2.0)$ keV									
* $^{70}\text{Cu}-^{85}\text{Rb}_{.824}$	$D_M = 5337.4(2.3)$ $\mu\text{u}$ for $^{70}\text{Cu}^n$ at 242.6(0.5) keV; $M - A = -62734.1(2.2)$ keV									
* $^{70}\text{Br}-^{13}\text{C F}_3$	$D_M = -51311(16)$ $\mu\text{u}$ for $^{70}\text{Br}^m$ at 2292.3(0.8) keV									
* $^{69}\text{Ga}(\text{n}, \gamma)^{70}\text{Ga}$	F: $E(\gamma)$ systematically lower than for other authors; Z recalibrated									
* $^{70}\text{Cu}(\beta^-)^{70}\text{Zn}$	$E_{\beta^-} = 4550(120), 3370(170)$ to $4^+$ level at 1786.33, $5^-$ at 3037.61 keV									
* $^{70}\text{Cu}(\beta^-)^{70}\text{Zn}$	$E_{\beta^-} = 6170(110)$ from $^{70}\text{Cu}^n$ $1^+$ at 242.6 keV									
* $^{70}\text{As}(\beta^+)^{70}\text{Ge}$	$E_{\beta^+} = 2144(50)$ to $3^+$ level at 3046.427, $4^+$ at 3058.707 keV									
* $^{70}\text{Se}(\beta^+)^{70}\text{As}$	$E_{\beta^+} = 1500(200)$ to $1^+$ level at 81.49, $1^+$ at 234.70, $1^+$ at 458.12 keV									
* $^{70}\text{Se}(\beta^+)^{70}\text{As}$	F: author's half-life 20(2)m disagrees with Nubase 41.1(0.3)m									
* $^{70}\text{Br}(\beta^+)^{70}\text{Se}$	$Q_{\beta^+} = 12190(80)$ from $2292.3$ $^{70}\text{Br}^m$									

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{71}\text{Co}-u$	-47100	600	-47630	500	-0.6	2			TO6	1.5	98Ba.A
	-47870	600			0.4	2			MT1	1.0	11Es06
$^{71}\text{Ni}-u$	-60000	400	-59481.0	2.4	0.9	U			TO5	1.5	94Se12
	-58700	350			-1.5	U			TO6	1.5	98Ba.A
$^{71}\text{Cu}-^{85}\text{Rb}_{.835}$	6332.4	1.6				2			MA8	1.0	07Gu09
$^{71}\text{Zn}-u$	-72080	380	-72280.4	2.8	-0.4	U			TO6	1.5	98Ba.A *
$^{71}\text{Zn}^m-^{85}\text{Rb}_{.835}$	1544.3	2.6	1544.4	2.5	0.0	1	95	95 $^{71}\text{Zn}^m$	MA8	1.0	08Ba54
$\text{C}_5\text{H}_{11}-^{71}\text{Ga}$	161370.2	3.2	161372.8	0.9	0.3	U			M15	2.5	63Ri07
$^{71}\text{Ga}-^{85}\text{Rb}_{.835}$	-1641.6	3.0	-1641.9	0.9	-0.1	-			MA8	1.0	07Gu09
	-1640.2	1.3			-1.3	-			MA8	1.0	07Ke09
	ave.	-1640.4	1.2		-1.2	1	53	53 $^{71}\text{Ga}$			average
$^{71}\text{Se}-u$	-68160	340	-67791	3	0.7	o			GA6	1.5	98Ch20
	-67687	75			-0.9	U			GT1	1.5	01Ha66
	-67830	120			0.2	U			GA6	1.5	02Li24
$^{71}\text{Se}-^{85}\text{Rb}_{.835}$	5865.0	3.0				2			MA8	1.0	11He10
$^{71}\text{Br}-u$	-61260	610	-60658	6	0.7	U			GA6	1.5	02Li24
$^{71}\text{Br H}_2-\text{C}_4\text{H}_9\text{O}$	-110347.7	5.8	-110348	6	0.0	1	100	100 $^{71}\text{Br}$	MS1	1.0	09Sa12
$^{71}\text{Kr}-u$	-49727	151	-49730	140	0.0	1	84	84 $^{71}\text{Kr}$	LZ1	1.0	11Tu02
$^{71}\text{Ni}-^{72}\text{Ge}_{.986}$	17352.2	2.4				2			JY1	1.0	07Ra27
$^{68}\text{Zn}(\alpha,n)^{71}\text{Ge}$	-5630	40	-5747.0	1.1	-2.9	U			Oak		64St01
$^{70}\text{Zn}(^{18}\text{O},^{17}\text{F})^{71}\text{Cu}$	-9529	35	-9588.1	2.4	-1.7	U			Ber		89Bo.A
$^{70}\text{Zn}(d,p)^{71}\text{Zn}$	3609	10	3611	3	0.2	1	10	7 $^{71}\text{Zn}$	ANL		67Vo05
$^{70}\text{Zn}(^3\text{He},d)^{71}\text{Ga}$	2380	20	2369.9	2.1	-0.5	U					74Ri08
$^{71}\text{Ga}(\gamma,n)^{70}\text{Ga}$	-9240	60	-9300.3	1.4	-1.0	U			Phi		60Ge01
$^{71}\text{Ga}(d,t)^{70}\text{Ga}$	-3054	10	-3043.1	1.4	1.1	U			Kop		71Ar12
$^{70}\text{Ge}(n,\gamma)^{71}\text{Ge}$	7415.3	1.5	7415.94	0.11	0.4	U					70Or.A
	7415.1	2.			0.4	U					72Gr34
	7415.95	0.15			-0.1	-			MMn		91Is01 Z
	7415.93	0.15			0.1	-			Bdn		06Fi.A
$^{70}\text{Ge}(d,p)^{71}\text{Ge}$	5182	10	5191.37	0.11	0.9	U			Kyu		73Ka03
$^{70}\text{Ge}(n,\gamma)^{71}\text{Ge}$	ave.	7415.94	0.11	7415.94	0.11	0.0	1	100	85 $^{70}\text{Ge}$		average
$^{70}\text{Ge}(p,\gamma)^{71}\text{As}$	4619	5	4620	4	0.2	R					75Li14
$^{71}\text{Zn}^m(\text{IT})^{71}\text{Zn}$	157.7	1.3	157.7	1.3	0.0	1	98	93 $^{71}\text{Zn}$			Ens10c
$^{71}\text{Zn}(\beta^-)^{71}\text{Ga}$	2610	50	2810.4	2.8	4.0	B					61Th01 *
	2786	50			0.5	U					61Th01 *
	2796	50			0.3	U					64So01 *
$^{71}\text{Ge}(\epsilon)^{71}\text{Ga}$	233.0	0.5	232.64	0.22	-0.7	-			Hei		84Ha.A
	229.3	1.0			3.3	F					91Zl01 *
	232.1	0.5			1.1	-					93Di03 *
	232.71	0.29			-0.2	-					95Le19
	233.5	1.2			-0.7	U			TT1		13Fr13
$^{71}\text{Ga}(p,n)^{71}\text{Ge}$	-1018.4	2.0	-1014.98	0.22	1.7	U			Oak		64Jo11 Z
$^{71}\text{Ge}(\epsilon)^{71}\text{Ga}$	ave.	232.65	0.22	232.64	0.22	0.0	1	99	86 $^{71}\text{Ge}$		average
$^{71}\text{Ga}(^3\text{He},t)^{71}\text{Ge}-^{65}\text{Cu}(^65)\text{Zn}$	1122.0	0.9	1119.0	0.4	-3.3	B			Pri		84Ko10
$^{71}\text{As}(\beta^+)^{71}\text{Ge}$	1997	20	2013	4	0.8	U					53St31 *
	2010	10			0.3	2					54Th36 *
	2012	10			0.1	2					55Gr08 *
$^{71}\text{Se}(\beta^+)^{71}\text{As}$	4428	125	4747	5	2.5	U					73Sc17
	4762	35			-0.4	U					01To06
$^{71}\text{Kr}(\epsilon)^{71}\text{Br}$	10140	320	10180	130	0.1	1	16	16 $^{71}\text{Kr}$			97Oi01
$^{*71}\text{Zn}-u$	$M-A=-67060(350)$ keV for mixture gs+m at 157.7 keV										Nub16c **
$^{*71}\text{Zn}(\beta^-)^{71}\text{Ga}$	$E_{\beta^-}=1450(50)$ 1460(50) respectively, from $^{71}\text{Zn}^m$ at 157.7(1.3) to $9/2^+$ at 1493.74										Ens10c **
$^{*71}\text{Ge}(\epsilon)^{71}\text{Ga}$	F : sees 17 keV neutrino										AHW **
$^{*71}\text{Ge}(\epsilon)^{71}\text{Ga}$	Original error 0.1 increased for calibration uncertainty										GAu **
$^{*71}\text{As}(\beta^+)^{71}\text{Ge}$	$E_{\beta^+}=800(20)$ 813(10) 815(10) respectively, to $5/2^-$ level at 174.943 keV										Ens10c **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{72}\text{Ni}-u$	-58700	500	-58214.1	2.4	0.6	U			TO5	1.5	94Se12
	-57400	400			-1.4	U			TO6	1.5	98Ba.A
$^{72}\text{Cu}-u$	-64250	510	-64179.7	1.5	0.1	U			TO6	1.5	98Ba.A *
$^{72}\text{Cu}-^{85}\text{Rb}_{.847}$	10534.4	1.5				2			MA8	1.0	07Gu09
$^{72}\text{Zn}-^{85}\text{Rb}_{.847}$	1556.9	2.3				2			MA8	1.0	08Ba54
$^{72}\text{Ga}-^{85}\text{Rb}_{.847}$	1079.5	1.5	1081.5	0.9	1.4	1	34	34 $^{72}\text{Ga}$	MA8	1.0	07Gu09
$\text{C}_4 \text{H}_8 \text{O}-^{72}\text{Ge}$	135438.4	2.1	135439.05	0.08	0.1	U			M15	2.5	63Ri07
$^{72}\text{Ge}-u$	-77906	25	-77924.17	0.08	-0.7	U			MR1	1.0	15Wi.A
	-77927	22			0.1	U			MR1	1.0	15Wi.A
	-77898	20			-1.3	U			MR1	1.0	15Wi.A
$^{72}\text{Se}-^{85}\text{Rb}_{.847}$	1854.6	2.1				2			MA8	1.0	11He10
$^{72}\text{Br}^{27}\text{Al}-^{85}\text{Rb}_{1.165}$	20892.1	7.2	20898.0	1.1	0.8	U			MA8	1.0	11He10
$^{72}\text{Br}-^{85}\text{Rb}_{.847}$	11308.7	1.1				2			MS1	1.0	15Va05 *
$^{72}\text{Kr}-^{85}\text{Rb}_{.847}$	16806.5	8.6				2			MA8	1.0	06Ro11
$^{70}\text{Ge} \text{H}_2-^{72}\text{Ge}$	17821.3	1.7	17822.9	0.9	0.4	U			M15	2.5	63Ri07
$^{72}\text{Ge}^{35}\text{Cl}-^{70}\text{Ge}^{37}\text{Cl}$	779.8	5.9	777.2	0.9	-0.2	U			H40	2.5	85El01
$^{72}\text{Ni}-^{72}\text{Ge}$	19710.1	2.4				2			JY1	1.0	07Ra27
$^{70}\text{Zn}(\text{t,p})^{72}\text{Zn}$	6231	20	6241.6	2.9	0.5	U			Ald		72Hu06
$^{71}\text{Ga}(\text{n},\gamma)^{72}\text{Ga}$	6521.1	1.0	6520.47	0.19	-0.6	U					70Li04 Z
	6519.8	1.0			0.7	F					71Ve03 *
	6520.44	0.19			0.2	1	99	66 $^{72}\text{Ga}$	Bdn		06Fi.A
$^{72}\text{Ge}(\text{d},^3\text{He})^{71}\text{Ga}$	-4241	7	-4242.3	0.8	-0.2	U			Ors		78Ro14
$^{72}\text{Zn}(\beta^-)^{72}\text{Ga}$	422	20	442.8	2.3	1.0	U					63De11 *
	458	6			-2.5	B					63Th03 *
$^{72}\text{Ga}(\beta^-)^{72}\text{Ge}$	4000	20	3997.6	0.8	-0.1	U					55Jo09 *
	3984	10			1.4	U					60La04 *
$^{72}\text{As}(\beta^+)^{72}\text{Ge}$	4361	10	4356	4	-0.5	2					50Me55
	4345	10			1.1	2					68Vi05
$^{72}\text{Ge}(\text{p,n})^{72}\text{As}$	-5140	5	-5138	4	0.3	2			Kyu		76Ki12
$^{72}\text{Br}(\beta^+)^{72}\text{Se}$	8869	95	8806.4	2.2	-0.7	U					01To06
$^{72}\text{Kr}(\beta^+)^{72}\text{Br}$	5040	80	5121	8	1.0	U					73Sc17 *
$^{72}\text{Cu}-u$	$M-A=-59710(470)$ keV for mixture gs+m at 270.3 keV										Nub16b **
$^{72}\text{Br}-^{85}\text{Rb}_{.847}$	$D_M=11308.4(1.1)\text{uu}, 11417.2(1.2)$ for ground state and $^{72}\text{Br}^m$ at 100.76(0.15) keV										Nub16b **
*	respectively $M-A=-59062.0(1.0)$ and $-58960.7(1.1)$ keV										15Va05 **
$^{71}\text{Ga}(\text{n},\gamma)^{72}\text{Ga}$	F : $E(\gamma)$ systematically lower than for other authors; Z recalibrated										AHW **
$^{72}\text{Zn}(\beta^-)^{72}\text{Ga}$	$E_{\beta^-}=260(20) 296(6)$ respectively, to $1^+$ level at 161.53 keV										Ens102 **
$^{72}\text{Ga}(\beta^-)^{72}\text{Ge}$	$E_{\beta^-}=3166(20) 3150(10)$ respectively, to $2^+$ level at 834.01 keV										Ens102 **
$^{72}\text{Kr}(\beta^+)^{72}\text{Br}$	$E_{\beta^+}=3794(180), 3626(105), 3682(80), 3364(155)$ to 162.67, 309.84 $1^+$ ,										73Sc17 **
*	415.05 $1^+$ and 576.74 $1^+$ levels										Ens102 **
$^{73}\text{Ni}-u$	-52500	500	-53793.3	2.6	-1.7	U			TO6	1.5	98Ba.A
$^{73}\text{Cu}-u$	-62740	350	-63325.6	2.1	-1.1	U			TO6	1.5	98Ba.A
$^{73}\text{Cu}-^{85}\text{Rb}_{.859}$	12447.9	4.2	12447.0	2.1	-0.2	1	25	25 $^{73}\text{Cu}$	MA8	1.0	07Gu09
$^{73}\text{Zn}-u$	-70100	380	-70417.4	2.0	-0.6	U			TO6	1.5	98Ba.A *
$^{73}\text{Zn}-^{85}\text{Rb}_{.859}$	5355.2	2.0				2			MA8	1.0	08Ba54
$^{73}\text{Ga}-^{85}\text{Rb}_{.859}$	947.3	1.8				2			MA8	1.0	07Gu09
$\text{C}_4 \text{H}_9 \text{O}-^{73}\text{Ge}$	141878.4	2.1	141880.95	0.06	0.5	U			M15	2.5	63Ri07
$^{73}\text{Se}-^{85}\text{Rb}_{.859}$	2511	11	2528	8	1.5	1	52	52 $^{73}\text{Se}$	MA8	1.0	11He10 *
$^{73}\text{Br}-u$	-68428	97	-68328	8	0.7	U			GT1	1.5	01Ha66
$^{73}\text{Br}^{27}\text{Al}-^{85}\text{Rb}_{1.176}$	16945.3	7.8				2			MA8	1.0	11He10
$^{73}\text{Kr}-^{85}\text{Rb}_{.859}$	15061.8	7.1	15062	7	0.0	o			MA8	1.0	04Ro32 *
	15062.8	9.7			-0.1	2			MA8	1.0	06Ro11
	15060.7	10.3			0.1	2			MA8	1.0	06Ro11
$^{73}\text{Ni}-^{72}\text{Ge}_{1.014}$	25221.8	2.6				2			JY1	1.0	07Ra27
$^{73}\text{Cu}-^{72}\text{Ge}_{1.014}$	15689.2	2.4	15689.5	2.1	0.1	1	75	75 $^{73}\text{Cu}$	JY1	1.0	07Ra27

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{72}\text{Ge H}-^{73}\text{Ge}$	6443.9	1.3	6441.90	0.05	-0.6	U			M15	2.5	63Ri07
$^{73}\text{Br}-^{72}\text{Br}$	-4610	330	-4923	8	-0.4	U			CR1	2.5	89Sh10 *
	-4709	166			-0.9	U			CR2	1.5	91Sh19 *
$^{72}\text{Ge}(n,\gamma)^{73}\text{Ge}$	6783.4	0.9	6782.94	0.05	-0.5	U					72Gr34
	6780.9	2.			1.0	U					72Ha74
	6782.94	0.05			0.0	1	100	100 $^{72}\text{Ge}$	MMn		91Is01 Z
	6783.12	0.15			-1.2	U			Bdn		06Fi.A
$^{72}\text{Ge}(d,p)^{73}\text{Ge}$	4571	4	4558.37	0.05	-3.2	B			Ald		69He05
	4563	10			-0.5	U			Kop		72Ha74
	4541	10			1.7	U			Kyu		73Ka03
$^{72}\text{Ge}(^3\text{He},d)^{73}\text{As}$	160	4	162	4	0.6	1	93	93 $^{73}\text{As}$	Hei		76Sc13
$^{73}\text{Kr}(\epsilon p)^{72}\text{Se}$	3700	150	4027	7	2.2	U			ChR		81Ha44
$^{73}\text{Rb}^i(p)^{72}\text{Kr}$	3803	40			3						93Ba61
$^{73}\text{Ga}(\beta^-)^{73}\text{Ge}$	1554	40	1598.2	1.7	1.1	U					58Yt22 *
	1564	80			0.4	U					70Wa21 *
$^{73}\text{Ge}(p,n)^{73}\text{As}$	-1121.6	15.	-1127	4	-0.4	U			Oak		64Jo11 *
$^{73}\text{Se}(\beta^+)^{73}\text{As}$	2740	10	2725	7	-1.5	1	55	48 $^{73}\text{Se}$			56Ha10 *
$^{73}\text{Br}(\beta^+)^{73}\text{Se}$	4748	500	4580	10	-0.3	U					70Mu02 *
	4648	400			-0.2	U					74Ro11 *
	4688	140			-0.8	U					87He21 *
	4610	70			-0.4	U					01To06
$^{73}\text{Kr}(\beta^+)^{73}\text{Br}$	6790	350	7096	10	0.9	U					73Sc17 *
	6860	220			1.1	U					97Oi01
$^{*73}\text{Zn}-u$	$M - A = -65200(350)$ keV for mixture gs+m at 195.5 keV										Nub16c **
$^{*73}\text{Se}-^{85}\text{Rb}_{.859}$	$D_M = 2524.6(7.3)$ $\mu\text{u}$ for mixture gs+m at 25.71 keV; $M - A = -68230.0(6.8)$ keV										Nub16b **
$^{*73}\text{Kr}-^{85}\text{Rb}_{.859}$	Combined results of next two items										GAu **
$^{*73}\text{Br}-^{72}\text{Br}$	$D_M = 4660(330)$ $\mu\text{u}$ corrected for $^{72}\text{Br}$ gs+m mixture at 100.76 keV										Nub16b **
$^{*73}\text{Br}-^{72}\text{Br}$	From $^{72}\text{Br}/^{73}\text{Br} = 0.98635312(227)$										AHW **
$^{*73}\text{Ga}(\beta^-)^{73}\text{Ge}$	$E_{\beta^-} = 1190(40)$ $1200(80)$ respectively, to $3/2^-$ level at 364.02 keV										Ens043 **
$^{*73}\text{Ge}(p,n)^{73}\text{As}$	$T = 1205(15)$ to $5/2^-$ level at 67.039(0.008)keV										Ens043 **
$^{*73}\text{Se}(\beta^+)^{73}\text{As}$	$E_{\beta^+} = 1290(10)$ to $9/2^+$ level at 427.906 keV										Ens043 **
$^{*73}\text{Br}(\beta^+)^{73}\text{Se}$	$E_{\beta^+} = 3700(500)$ $3600(400)$ $3640(140)$ respectively, to $^{73}\text{Se}^m$ at 25.71 keV										Nub16b **
$^{*73}\text{Kr}(\beta^+)^{73}\text{Br}$	$E_{\beta^+} = 5589(350)$ to $3/2^-$ level at 178.08 keV										Ens043 **
$^{74}\text{Ni}-u$	-52830	1060	-52020#	210#	0.8	D			MT1	1.0	11Es06 *
$^{74}\text{Cu}-u$	-59400	400	-60125	7	-1.2	U			TO6	1.5	98Ba.A
$^{74}\text{Cu}-^{85}\text{Rb}_{.871}$	16706.0	6.6				2			MA8	1.0	07Gu09
$^{74}\text{Zn}-^{85}\text{Rb}_{.871}$	6238.4	2.7				2			MA8	1.0	08Ba54
$^{74}\text{Ga}-^{85}\text{Rb}_{.871}$	3776.9	22.6	3777	3	0.0	U			MA8	1.0	07Ke09 *
	3776.9	4.0			0.0	2			MA8	1.0	07Gu09
	3806.5	34.6			-0.9	U			MA8	1.0	07Ke09 *
	3776.8	5.4			0.0	2			TT1	1.0	15Ma30
$\text{C } ^{32}\text{S}_2-^{74}\text{Ge H}_2$	7314.0	1.4	7314.522	0.014	0.1	U			M15	2.5	63Ri07
$^{74}\text{Ge}-^{84}\text{Kr}$	9680.0337	0.0128	9680.034	0.013	0.0	1	100	100 $^{74}\text{Ge}$	FS1	1.0	10Mo03
$\text{C}_6 \text{H}_2-^{74}\text{Se}$	93173.8	3.8	93174.129	0.016	0.0	U			M15	2.5	63Ri07
$^{74}\text{Se}-^{84}\text{Kr}$	10978.2066	0.0128	10978.207	0.015	0.0	o			FS1	1.0	10Mo03 *
$^{74}\text{Se}-^{85}\text{Rb}_{.871}$	-691.4	7.3	-692.927	0.016	-0.2	U			MA8	1.0	11He10
$^{74}\text{Br } ^{27}\text{Al}-^{85}\text{Rb}_{1.188}$	16246.0	6.8	16242	6	-0.5	1	85	85 $^{74}\text{Br}$	MA8	1.0	11He10 *
$^{74}\text{Kr}-^{85}\text{Rb}_{.871}$	9915.0	2.2	9915.2	2.2	0.1	o			MA8	1.0	04Ro32 *
	9916.8	2.6			-0.6	-			MA8	1.0	06Ro11
	9909.7	4.4			1.2	-			MA8	1.0	06Ro11
	ave.	9915.0			0.1	1	93	93 $^{74}\text{Kr}$			average
$^{74}\text{Rb}-^{85}\text{Rb}_{.871}$	21109	19	21097	3	-0.6	U			MA8	1.0	07Ke09
	21097.9	4.3			-0.2	o			MA8	1.0	04Ke10 *
	21095.7	5.2			0.3	-			MA8	1.0	07Ke09
	21102.7	7.5			-0.8	-			MA8	1.0	07Ke09
	21096.4	6.5			0.1	-			TT1	1.0	15Ma30
	ave.	21097			-0.1	1	83	83 $^{74}\text{Rb}$			average



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{74}\text{Rb}-u$	-55765	125	-55734	3	0.2	U			P40	1.0	06Lu19
$^{74}\text{Ge } ^{35}\text{Cl}-^{72}\text{Ge } ^{37}\text{Cl}$	2047.5	1.1	2052.05	0.11	1.0	U			H18	4.0	64Ba03
	2047.74	0.71			2.4	U			H40	2.5	85E101
	2052.01	0.26			0.1	U			H44	1.5	91Hy01
$^{74}\text{Se } ^{35}\text{Cl}-^{72}\text{Ge } ^{37}\text{Cl}$	3347.9	4.7	3350.22	0.11	0.2	U			H40	2.5	85E101
$^{73}\text{Ge } \text{H}-^{74}\text{Ge}$	10105.1	1.7	10106.23	0.06	0.3	U			M15	2.5	63Ri07
$^{74}\text{Se}-^{74}\text{Ge}$	1298.5	8.5	1298.173	0.008	0.0	U			H40	2.5	85E101
	1298.7	3.7			-0.1	U			H40	2.5	85E101
	1298.096	0.053			1.5	U			JY1	1.0	10Ko15
	1298.1729	0.0080			0.0	1	100	100 $^{74}\text{Se}$	FS1	1.0	10Mo03
$^{74}\text{Br}-^{73}\text{Br}$	-1244	410	-1761	10	-0.5	U			CR1	2.5	89Sh10 *
$^{74}\text{Se}(\text{p,t})^{72}\text{Se}$	-11979	24	-12005.9	2.0	-1.1	U			Win		74De31 *
$^{74}\text{Ge}(\text{d},^{14}\text{C},^{15}\text{O})^{73}\text{Zn}$	-8018	150	-7664.8	1.9	2.4	U			Ors		84De33
$^{74}\text{Ge}(\text{d},^3\text{He})^{73}\text{Ga}$	-5515	7	-5518.6	1.7	-0.5	U			Ors		78Ro14
	-5509	13			-0.7	U			Hei		84Ha31
$^{73}\text{Ge}(\text{n},\gamma)^{74}\text{Ge}$	10200.2	0.6	10196.24	0.06	-6.6	B					70Ha60
	10198	2			-0.9	U					74Ch18
	10195.90	0.15			2.2	-			ILn		85Ho.A Z
	10196.32	0.14			-0.6	-					89Bu.A
	10196.31	0.07			-1.1	-			MMn		91Is01 Z
	10196.06	0.20			0.9	-			Bdn		06Fi.A
	ave.	10196.24	0.06			0.0	1	100	100 $^{73}\text{Ge}$		
$^{74}\text{Se}(\text{d},^3\text{He})^{73}\text{As}$	-3027	8	-3056	4	-3.6	B			Ors		83Ro08 *
$^{74}\text{Zn}(\beta^-)^{74}\text{Ga}$	2350	100	2293	4	-0.6	U					72Er05 *
$^{74}\text{Ga}(\beta^-)^{74}\text{Ge}$	5400	100	5372.8	3.0	-0.3	U					62Ei02 *
$^{74}\text{As}(\beta^+)^{74}\text{Ge}$	2558	4	2562.4	1.7	1.1	-					71Bo01 *
$^{74}\text{Ge}(\text{p,n})^{74}\text{As}$	-3343.5	5.6	-3344.7	1.7	-0.2	-			Tkm		63Ok01
	-3348.3	5.			0.7	-			Oak		64Jo11 Z
	-3346	5			0.3	-					70Fi03 Z
	-3347	3			0.8	-			Kyu		73Ki11
ave.	2562.9	1.9	2562.4	1.7	-0.3	1	82	82 $^{74}\text{As}$			average
$^{74}\text{As}(\beta^-)^{74}\text{Se}$	1351	4	1353.1	1.7	0.5	1	18	18 $^{74}\text{As}$			71Bo01 *
$^{74}\text{Br}(\beta^+)^{74}\text{Se}$	6857	100	6925	6	0.7	U					69La15 *
$^{74}\text{Se}(\text{p,n})^{74}\text{Br}$	-7689	15	-7707	6	-1.2	1	15	15 $^{74}\text{Br}$			75Lu02 *
$^{74}\text{Kr}(\beta^+)^{74}\text{Br}$	3000	200	2956	6	-0.2	U					74Ro11
$^{74}\text{Rb}(\beta^+)^{74}\text{Kr}$	3327	125			-3.0	B					75Sc07
	10000	1500	10416	3	0.3	U					76Da.D
	10413.8	7.0			0.3	1	24	17 $^{74}\text{Rb}$			03Pi08 *
* $^{74}\text{Ni}-u$	Trends from Mass Surface TMS suggest $^{74}\text{Ni}$ 750 less bound										GAu **
* $^{74}\text{Ga}-^{85}\text{Rb}_{.871}$	$D_M=3780.1(22.5) \mu\text{u}$ corrected $-3.0(1.7) \text{keV}$ for gs+m mixture $R<0.1$ at 59.571 keV										Nub16b **
* $^{74}\text{Ga}-^{85}\text{Rb}_{.871}$	$D_M=3809.7(34.6) \mu\text{u}$ corrected $-3.0(1.7) \text{keV}$ for gs+m mixture $R<0.1$ at 59.571 keV										Nub16b **
* $^{74}\text{Se}-^{84}\text{Kr}$	Not independent measurement, use $^{74}\text{Ge}-^{74}\text{Se}$ below										10Mo03 **
* $^{74}\text{Br } ^{27}\text{Al}-^{85}\text{Rb}_{1.188}$	$D_M=16253.4(5.3) \mu\text{u}$ for mixture gs+m at 13.58 keV; $M-A=-82474.9(4.9) \text{keV}$										Nub16b **
* $^{74}\text{Kr}-^{85}\text{Rb}_{.871}$	Combined results of next two items										GAu **
* $^{74}\text{Rb}-^{85}\text{Rb}_{.871}$	Combined results of next two items										GAu **
* $^{74}\text{Br}-^{73}\text{Br}$	$D_M=-1230(410) \mu\text{u}$ for $^{74}\text{Br}^m$ at 13.58 keV										Nub16b **
* $^{74}\text{Se}(\text{p,t})^{72}\text{Se}$	Original error 12; added systematic error 21 keV										GAu **
* $^{74}\text{Se}(\text{d},^3\text{He})^{73}\text{As}$	$Q=-3033(8)$ for $Q(^{76}\text{Se}(\text{d},^3\text{He}))=-4020.7(2.0)$ , now 4014.5 keV										AHW **
* $^{74}\text{Zn}(\beta^-)^{74}\text{Ga}$	$E_{\beta^-}=2100(100)$ to $1^+$ level at 251.787 keV										Ens067 **
* $^{74}\text{Ga}(\beta^-)^{74}\text{Ge}$	$E_{\beta^-}=2450(100)$ to $3^-$ level at 2949.48 keV										Ens067 **
* $^{74}\text{As}(\beta^+)^{74}\text{Ge}$	Original error increased: authors report $E(2^+)=593.1(1.5)$ while										AHW **
*	$E(2^+)=595.850(0.006) \text{keV}$ ; see also $^{84}\text{Rb}(\beta^+)$										Ens067 **
* $^{74}\text{As}(\beta^-)^{74}\text{Se}$	Original value 1350.1(0.7), error increased, see $^{84}\text{Rb}(\beta^+)$										AHW **
* $^{74}\text{Br}(\beta^+)^{74}\text{Se}$	$E_{\beta^+}=5200(100), 4500(100)$ to 634.76, 1363.21 levels										69La15 **
*	from $^{74}\text{Br}^m$ at 13.8(0.5) keV										93Do05 **
* $^{74}\text{Se}(\text{p,n})^{74}\text{Br}$	$T=7868(15)$ to $(2^-)$ level at 72.65 keV										Ens067 **
* $^{74}\text{Rb}(\beta^+)^{74}\text{Kr}$	Deduced from measured half-life and branching ratio										GAu **
* $^{74}\text{Rb}(\beta^+)^{74}\text{Kr}$	Original 10405(9) re-evaluated in reference										11To.A **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{75}\text{Cu}-u$	-58100	700	-58477.4	2.5	-0.4	U			TO6	1.5	98Ba.A	
$^{75}\text{Zn}-^{85}\text{Rb}_{.882}$	10641.7	2.1				2			MA8	1.0	08Ba54	
$^{75}\text{Ga}-^{85}\text{Rb}_{.882}$	4301.7	2.6				2			MA8	1.0	07Gu09	
$\text{C}_3 \text{H}_7 \text{O}_2 - ^{75}\text{As}$	123009.8	2.6	123009.9	0.9	0.0	U			M15	2.5	63Ri07	
$^{75}\text{As}-^{85}\text{Rb}_{.882}$	-601.3	7.6	-604.0	0.9	-0.4	U			MA8	1.0	02Ke.A	
$^{75}\text{Br}^{27}\text{Al}-^{85}\text{Rb}_{1.200}$	13201.3	4.6				2			MA8	1.0	11He10	
$^{75}\text{Kr}-^{85}\text{Rb}_{.882}$	8747.2	8.7				2			MA8	1.0	06Ro11	
$^{75}\text{Rb}-^{85}\text{Rb}_{.882}$	16371	8	16374.7	1.3	0.5	U			MA2	1.0	94Ot01	
	16374.7	1.7			0.0	2			MA8	1.0	07Ke09	
	16368	21			0.3	U			MA8	1.0	07Ke09	
	16374.6	1.9			0.0	2			TT1	1.0	15Ma30	
$^{75}\text{Cu}-^{72}\text{Ge}_{1.042}$	22719.6	2.5				2			JY1	1.0	07Ra27	
$^{75}\text{As}^{35}\text{Cl}-^{73}\text{Ge}^{37}\text{Cl}$	1079.6	5.0	1085.7	1.0	0.5	U			H40	2.5	85El01	
$^{74}\text{Ge}(n,\gamma)^{75}\text{Ge}$	6505.9	1.1	6505.84	0.05	-0.1	U					72Gr34	
	6505.5	2.			0.2	U					72Ha74	
	6505.81	0.30			0.1	U					89Bu.A *	
	6505.26	0.08			7.3	B		MMn			91Is01 Z	
	6505.45	0.14			2.8	C		Bdn			06Fi.A	
	6505.84	0.05				2					12Me04	
$^{74}\text{Ge}(d,p)^{75}\text{Ge}$	4265	15	4281.27	0.05	1.1	U			MIT		67Sp09	
	4282	10			-0.1	U			Kop		72Ha74	
	4268	10			1.3	U			Kyu		73Ka03	
$^{74}\text{Ge}(p,\gamma)^{75}\text{As}$	6901.6	5.	6900.7	0.9	-0.2	U					74Wa08	
$^{74}\text{Ge}(^3\text{He},d)^{75}\text{As}$	1414	4	1407.2	0.9	-1.7	U			Hei		76Sc13	
$^{75}\text{As}(\gamma,n)^{74}\text{As}$	-10259	31	-10245.5	1.9	0.4	U			Phi		60Ge01	
$^{74}\text{Se}(n,\gamma)^{75}\text{Se}$	8027.84	0.30	8027.60	0.07	-0.8	U			BNn		81En07 Z	
	8027.60	0.08			0.0	-			ILn		84To11 Z	
	8027.59	0.16			0.0	-			Bdn		06Fi.A	
	ave.	8027.60			0.0	1	100	100 $^{75}\text{Se}$			average	
$^{75}\text{Zn}(\beta^-)^{75}\text{Ga}$	6060	80	5906	3	-1.9	U			Stu		86Ek01	
$^{75}\text{Ga}(\beta^-)^{75}\text{Ge}$	3300	200	3392.4	2.4	0.5	U					60Mo01	
$^{75}\text{Ge}(\beta^-)^{75}\text{As}$	1188	20	1177.2	0.9	-0.5	U					55Sc09	
$^{75}\text{As}(p,n)^{75}\text{Se}$	-1647.2	2.0	-1647.1	0.9	0.1	-			Nvl		59Go68 Z	
	-1647.3	1.1			0.3	-			Oak		64Jo11 Z	
	-1643	5			-0.8	U					70Fi03	
	ave.	-1647.3			0.3	1	85	85 $^{75}\text{As}$			average	
$^{75}\text{Br}(\beta^+)^{75}\text{Se}$	3010	20	3062	4	2.6	U					52Fu04 *	
	3030	50			0.6	U					61Ba43 *	
	3050	20			0.6	U					69Ra24 *	
$^{75}\text{Kr}(\beta^+)^{75}\text{Br}$	4400	200	4783	9	1.9	U					74Ro12 *	
$^{75}\text{Sr}(\epsilon)^{75}\text{Rb}$	10600	220				3					03Hu01	
$^{74}\text{Ge}(n,\gamma)^{75}\text{Ge}$	Original error 0.03 keV increased										GAu	**
$^{75}\text{Br}(\beta^+)^{75}\text{Se}$	$E_{\beta^+}=1700(20) 1720(50) 1740(20)$ respectively, to $3/2^-$ level at 286.5714 keV										Ens139	**
$^{75}\text{Kr}(\beta^+)^{75}\text{Br}$	$E_{\beta^+}=3200(200)$ to $132.46 (5/2)^+, 154.61 (3/2)^+$ levels										Ens139	**
$^{76}\text{Cu}-^{85}\text{Rb}_{.894}$	24135.0	7.2				2			MA8	1.0	07Gu09	
$^{76}\text{Zn}-^{85}\text{Rb}_{.894}$	11975.5	2.0	11974.9	1.6	-0.3	1	61	61 $^{76}\text{Zn}$	MA8	1.0	08Ba54	
$^{76}\text{Zn}-^{88}\text{Rb}_{.864}$	9737.4	2.5	9738.3	1.6	0.4	1	39	39 $^{76}\text{Zn}$	JY1	1.0	08Ha23	
$^{76}\text{Ga}-^{85}\text{Rb}_{.894}$	7687.6	2.1				2			MA8	1.0	07Gu09	
$\text{C}^{32}\text{S}_2-^{76}\text{Ge}$	22741.6	1.5	22739.622	0.019	-0.5	U			M15	2.5	63Ri07	
$^{76}\text{Ge}-u$	-78597.242	0.096	-78597.273	0.019	-0.3	U			ST2	1.0	01Do08	
$\text{C}_6 \text{H}_4-^{76}\text{Se}$	112100	8	112086.424	0.017	-0.7	U			M15	2.5	63Ri07	
$^{76}\text{Se}-u$	-80786.205	0.081	-80786.296	0.017	-1.1	U			ST2	1.0	01Do08	
$^{76}\text{Kr}-^{85}\text{Rb}_{.894}$	4774.3	4.7	4771	4	-0.8	1	84	84 $^{76}\text{Kr}$	MA8	1.0	06Ro11	
$^{76}\text{Rb}-^{85}\text{Rb}_{.894}$	13931	8	13933.0	1.0	0.3	U			MA2	1.0	94Ot01	
	13932.2	2.0			0.4	2			MA8	1.0	07Ke09	
	13923	15			0.7	U			MA8	1.0	07Ke09	
	13935.3	1.6			-1.4	2			MA8	1.0	07Ke09	
	13931.0	1.7			1.2	2			TT1	1.0	15Ma30	
	13925.2	4.5			1.7	U			TT1	1.0	15Ma30	
	13933.5	3.4			-0.1	o			TT1	1.0	15Ma30 *	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{76}\text{Sr } ^{19}\text{F}-^{85}\text{Rb}_{1.118}$	38785	37				2			MA8	1.0	05Si34
$^{76}\text{Sr}-\text{u}$	-58813	107	-58240	40	5.4	F			CS1	1.0	01La31 *
$^{76}\text{Ge}-^{84}\text{Kr}$	9904.9983	0.0175	9904.998	0.019	0.0	o			FS1	1.0	10Mo03 *
$^{76}\text{Se}-^{84}\text{Kr}$	7715.9762	0.0169	7715.976	0.017	0.0	1	100	100 $^{76}\text{Se}$	FS1	1.0	10Mo03
$^{74}\text{Ge } \text{H}_2-^{76}\text{Ge}$	15425.0	1.7	15425.100	0.023	0.0	U			M15	2.5	63Ri07
$^{76}\text{Ge } ^{35}\text{Cl}-^{74}\text{Ge } ^{37}\text{Cl}$	3175.7	1.5	3175.07	0.07	-0.1	U			H18	4.0	64Ba03
	3170.41	0.74			2.5	U			H40	2.5	85E101
	3174.61	0.41			0.8	U			H44	1.5	91Hy01
$^{76}\text{Se } ^{35}\text{Cl}-^{74}\text{Ge } ^{37}\text{Cl}$	986.30	0.65	986.05	0.07	-0.3	U			H44	1.5	91Hy01
$^{76}\text{Ge}-^{76}\text{Se}$	2190.92	0.59	2189.022	0.008	-1.3	U			H40	2.5	85E101
	2188.60	0.42			0.7	U			H44	1.5	91Hy01
	2188.963	0.054			1.1	U			ST2	1.0	01Do08
	2188.98	0.16			0.3	U			JY1	1.0	08Ra09
	2189.0221	0.008			0.0	1	100	100 $^{76}\text{Ge}$	FS1	1.0	10Mo03
$^{75}\text{Rb}-^{76}\text{Rb}_{.493} \ ^{74}\text{Rb}_{.507}$	-1140	170	-1081.1	2.0	0.1	U			P20	2.5	82Au01
$^{76}\text{Ge}(^{14}\text{C}, ^{17}\text{O})^{73}\text{Zn}$	-3779	40	-3790.8	1.9	-0.3	U			Ors		84Be10 *
$^{76}\text{Ge}(^{14}\text{C}, ^{16}\text{O})^{74}\text{Zn}$	163	40	300.7	2.5	3.4	B			Ors		84Be10
$^{76}\text{Ge}(^{18}\text{O}, ^{20}\text{Ne})^{74}\text{Zn}$	-1219	21	-1197.1	2.5	1.0	U			Hei		84Ha31
$^{76}\text{Ge}(^{14}\text{C}, ^{15}\text{O})^{75}\text{Zn}$	-10354	150	-10489.7	2.0	-0.9	U			Ors		84De33
$^{76}\text{Ge}(\text{d}, ^3\text{He})^{75}\text{Ga}$	-6545	7	-6543.8	2.4	0.2	U			Ors		78Ro14
	-6536	22			-0.4	U			Hei		84Ha31
$^{75}\text{As}(\text{n}, \gamma)^{76}\text{As}$	7329	2	7328.50	0.07	-0.3	U					68Jo11
	7328.421	0.075			1.0	2			ILn		90Ho10 Z
	7328.81	0.15			-2.1	2			Bdn		06Fi.A
$^{75}\text{As}(\text{d}, \text{p})^{76}\text{As}$	5105	5	5103.93	0.07	-0.2	U					76Mo32
$^{75}\text{Se}(\text{n}, \gamma)^{76}\text{Se}$	11154.15	0.30	11153.79	0.07	-1.2	U			ILn		83To20 Z
$^{76}\text{Zn}(\beta^-)^{76}\text{Ga}$	4160	80	3993.6	2.4	-2.1	U			Stu		86Ek01
$^{76}\text{Ga}(\beta^-)^{76}\text{Ge}$	6770	150	6916.2	2.0	1.0	o			Stu		77A117
	7010	90			-1.0	U			Stu		86Ek01
$^{76}\text{Ge}(\text{p}, \text{n})^{76}\text{As}$	-1705	5	-1703.9	0.9	0.2	U					70Fi03
$^{76}\text{As}(\beta^-)^{76}\text{Se}$	2970	2	2960.6	0.9	-4.7	B					69Na11
$^{76}\text{Br}(\beta^+)^{76}\text{Se}$	5002	20	4963	9	-2.0	2					71Dz08
$^{76}\text{Br}(\text{n}, \text{p})^{76}\text{Se}$	5730	15	5745	9	1.0	2			ILL		78An14
$^{76}\text{Se}(\text{p}, \text{n})^{76}\text{Br}$	-5738.6	15.	-5745	9	-0.4	2					75Lu02
$^{76}\text{Rb}(\beta^+)^{76}\text{Kr}$	8793	570	8535	4	-0.5	U					75We23
	8063	44			10.7	F					82Mo10 *
	8094	162			2.7	F			BNL		83Li11 *
	8250	150			1.9	U			IRS		93A103
* $^{76}\text{Rb}-^{85}\text{Rb}_{.894}$	no accuracy check was performed for this $12^+$ charge state										15Ma30 **
* $^{76}\text{Sr}-\text{u}$	F : other results in same paper not trusted, see $^{80}\text{Y}$										GAu **
* $^{76}\text{Ge}-^{84}\text{Kr}$	Not independent measurement, use $^{76}\text{Ge}-^{76}\text{Se}$ below										10Mo03 **
* $^{76}\text{Ge}(^{14}\text{C}, ^{17}\text{O})^{73}\text{Zn}$	$Q = -3974(40) \text{ MeV} - A = -65410(40) \text{ MeV}$ to $^{73}\text{Zn}^m$ at 195.5 keV										GAu **
* $^{76}\text{Rb}(\beta^+)^{76}\text{Kr}$	$E_{\beta^+} = 4558(30) \text{ keV}$ to level $(1^+, 2^+)$ at 2570.95 keV and corrections by authors										Ens953 **
* $^{76}\text{Rb}(\beta^+)^{76}\text{Kr}$	F : 29.6% feeding above 2570.95 level from reference										84Mo22 **
$^{77}\text{Cu}-\text{u}$	-51850	540	-52200#	160#	-0.6	D			OR1	1.0	06Ha62 *
$^{77}\text{Zn}-\text{u}$	-62790	780	-63112.8	2.1	-0.3	U			TO6	1.5	98Ba.A *
	-63380	260			0.4	U			GT2	2.5	08Kn.A *
$^{77}\text{Zn}-^{88}\text{Rb}_{.875}$	14485.7	4.5	14486.1	2.1	0.1	1	22	22 $^{77}\text{Zn}$	JY1	1.0	08Ha23
$\text{C}_6 \text{H}_5-^{77}\text{Se}$	119211.9	4.2	119211.01	0.07	-0.1	U			M15	2.5	63Ri07
$^{77}\text{Rb}-^{39}\text{K}_{1.974}$	1912	27	2045.0	1.4	4.9	B			MA2	1.0	87Bo59
$^{77}\text{Zn}-^{85}\text{Rb}_{.906}$	16805.8	2.4	16805.7	2.1	0.0	1	78	78 $^{77}\text{Zn}$	MA8	1.0	08Ba54
$^{77}\text{Ga}-^{85}\text{Rb}_{.906}$	9072.8	2.6				2			MA8	1.0	07Gu09
$^{77}\text{Kr}-^{85}\text{Rb}_{.906}$	4588.5	2.1				2			MA8	1.0	06Ro11
$^{77}\text{Rb}-^{85}\text{Rb}_{.906}$	10327	8	10320.1	1.4	-0.9	U			MA2	1.0	94O101
	10320.1	1.4				2			MA8	1.0	07Ke09

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{77}\text{Sr } ^{19}\text{F}-^{85}\text{Rb}_{1,129}$	35938.0	8.5				2			MA8	1.0	05Si34	
$^{75}\text{Rb}-^{77}\text{Rb}_{,325} \ ^{74}\text{Rb}_{,676}$	-1340	380	-1053.6	2.4	0.3	U			P20	2.5	82Au01	
$^{76}\text{Rb}-^{77}\text{Rb}_{,494} \ ^{75}\text{Rb}_{,507}$	525	30	557.1	1.3	0.4	U			P20	2.5	82Au01	
$^{76}\text{Ge}(n,\gamma)^{77}\text{Ge}$	6072.5	1.0	6071.29	0.05	-1.2	U					72Gr34 Z	
	6071.7	1.2			-0.3	U					72Ha74 Z	
	6072.3	0.4			-2.5	U			Bdn		06Fi.A	
	6071.29	0.05				2					12Me04	
$^{76}\text{Ge}(d,p)^{77}\text{Ge}$	3839	10	3846.72	0.05	0.8	U			Kop		72Ha74	
	3823	12			2.0	U			Kyu		73Ka03	
$^{76}\text{Ge}(^3\text{He,d})^{77}\text{As}$	2497	3	2498.9	1.7	0.6	1	32	32 $^{77}\text{As}$	Hei		76Sc13	
$^{76}\text{Se}(n,\gamma)^{77}\text{Se}$	7418.87	0.20	7418.86	0.06	-0.1	-			BNn		81En07	
	7418.85	0.07			0.1	-			ILn		85To10 Z	
	7418.85	0.15			0.0	-			Bdn		06Fi.A	
$^{76}\text{Se}(d,p)^{77}\text{Se}$	5192	10	5194.29	0.06	0.2	U			Ald		63Ma27	
$^{76}\text{Se}(n,\gamma)^{77}\text{Se}$	ave.	7418.85	7418.86	0.06	0.1	1	99	99 $^{77}\text{Se}$			average	
$^{77}\text{Sr}(ep)^{76}\text{Kr}$	3850	200	3921	9	0.4	U			ChR		76Ha29	
$^{77}\text{Zn}(\beta^-)^{77}\text{Ga}$	7270	120	7203	3	-0.6	U			Stu		86Ek01	
$^{77}\text{Ga}(\beta^-)^{77}\text{Ge}$	5340	60	5220.5	2.4	-2.0	U			Stu		77Al17	
	5690	300			-1.6	U			Stu		86Ek01	
$^{77}\text{Ge}(\beta^-)^{77}\text{As}$	2670	100	2703.5	1.7	0.3	U					52Sm13 *	
$^{77}\text{As}(\beta^-)^{77}\text{Se}$	700	7	683.2	1.7	-2.4	U					51Ca04	
	679	4			1.0	1	18	18 $^{77}\text{As}$			51Je01	
$^{77}\text{Br}(\beta^+)^{77}\text{Se}$	1358	20	1364.7	2.8	0.3	U					51Ca28	
$^{77}\text{Se}(p,n)^{77}\text{Br}$	-2147	4	-2147.0	2.8	0.0	2			Oak		58Jo01	
	-2147.0	4.			0.0	2			Tkm		63Ok01	
$^{77}\text{Kr}(\beta^+)^{77}\text{Br}$	3012	30	3065	3	1.8	U					55Th01 *	
	3027	40			1.0	U					73Ba22 *	
	3300	100			-2.3	U					74Ro11 *	
	2760	42			7.3	B					82Mo10 *	
$^{77}\text{Rb}(\beta^+)^{77}\text{Kr}$	5180	390	5339.0	2.4	0.4	U					75We23	
	5272	26			2.6	U					82Mo10	
	5113	69			3.3	B			BNL		83Li11	
	5320	70			0.3	U			IRS		93Al03	
$^{77}\text{Sr}(\beta^+)^{77}\text{Rb}$	6986	227	7027	8	0.2	U			BNL		83Li11	
* $^{77}\text{Cu}-u$	Trends from Mass Surface TMS suggest $^{77}\text{Cu}$ 320 more bound										GAu	**
* $^{77}\text{Zn}-u$	$M - A = -58100(700)$ keV for mixture gs+m at 772.440 keV										Nub16b	**
* $^{77}\text{Zn}-u$	$M - A = -58648(95)$ keV for mixture gs+m at 772.440 keV ( $1/2^-$ )										Nub16b	**
* $^{77}\text{Ge}(\beta^-)^{77}\text{As}$	$E_{\beta^-} = 2196(100)$ to $9/2^+$ $^{77}\text{As}^m$ at 475.48 keV										Nub16c	**
* $^{77}\text{Kr}(\beta^+)^{77}\text{Br}$	Error not in 55Th01, estimated by evaluator										AHW	**
* $^{77}\text{Kr}(\beta^+)^{77}\text{Br}$	$E_{\beta^+} = 1860(30)$ 1875(40) respectively, to $5/2^+$ level at 129.64 keV										Ens126	**
* $^{77}\text{Kr}(\beta^+)^{77}\text{Br}$	$E_{\beta^+} = 2000(100)$ 1528(36) respectively, to $(3/2)^+$ level at 276.22 keV										Ens126	**
$^{78}\text{Cu}-u$	-47770	540				2			OR1	1.0	06Ha62	
$^{78}\text{Zn}-^{88}\text{Rb}_{,886}$	16863.8	2.9	16863.6	2.1	-0.1	1	52	52 $^{78}\text{Zn}$	JY1	1.0	08Ha23	
$^{78}\text{Ga}-^{88}\text{Rb}_{,886}$	10184.3	3.3	10183.2	2.0	-0.3	1	38	38 $^{78}\text{Ga}$	JY1	1.0	08Ha23	
$\text{C}_6 \text{H}_6-^{78}\text{Se}$	129642.6	2.2	129640.95	0.19	-0.3	U			M15	2.5	63Ri07	
$\text{C}_6 \text{H}_6-^{78}\text{Kr}$	126548.3	3.6	126583.9	0.3	4.0	B			M15	2.5	63Ri07	
	126554	17			1.2	U			R11	1.5	78Di09	
	126560	7			2.3	U			R11	1.5	78Di09	
$\text{C}_5 \text{N H}_4-^{78}\text{Kr}$	113994	20	114007.8	0.3	0.5	U			R11	1.5	78Di09	
$^{78}\text{Kr}-^{86}\text{Kr}_{,907}$	1441.2	1.0	1442.5	0.3	1.3	1	11	11 $^{78}\text{Kr}$	MS1	1.0	06Ri15	
$^{78}\text{Zn}-^{85}\text{Rb}_{,918}$	19266.0	3.0	19266.2	2.1	0.1	1	48	48 $^{78}\text{Zn}$	MA8	1.0	08Ba54	
$^{78}\text{Ga}-^{85}\text{Rb}_{,918}$	12585.2	2.6	12585.9	2.0	0.3	1	62	62 $^{78}\text{Ga}$	MA8	1.0	07Gu09	
$^{78}\text{Kr}-^{85}\text{Rb}_{,918}$	1342.3	1.4	1343.4	0.3	0.8	U			MA8	1.0	06Ro11	
	1338.9	2.2			2.0	U			MA8	1.0	06Ro11	
$^{78}\text{Rb}-^{85}\text{Rb}_{,918}$	9118	8	9119	3	0.1	2			MA2	1.0	94Ot01	
	9121.3	7.5			-0.3	2			TT1	1.0	12Ga15 *	
	9118.3	4.5			0.1	2			TT1	1.0	12Ga15 *	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{78}\text{Sr}-^{85}\text{Rb}_{.918}$	13157	8				2			MA2	1.0	94Ot01
$^{78}\text{Se } ^{35}\text{Cl}_2-^{74}\text{Ge } ^{37}\text{Cl}_2$	2030.4	2.2	2031.70	0.24	0.4	U			H44	1.5	91Hy01
$^{78}\text{Se } ^{35}\text{Cl}-^{76}\text{Ge } ^{37}\text{Cl}$	-1147.60	0.92	-1143.37	0.20	1.8	U			H40	2.5	85El01
	-1143.57	0.72			0.2	U			H44	1.5	91Hy01
$^{78}\text{Se } ^{35}\text{Cl}-^{76}\text{Se } ^{37}\text{Cl}$	1042.03	1.35	1045.65	0.20	1.1	U			H40	2.5	85El01
	1044.58	0.45			1.6	U			H44	1.5	91Hy01
$^{76}\text{Se } \text{H}_2-^{78}\text{Kr}$	14440	25	14497.4	0.3	1.5	U			R11	1.5	78Di09
$^{77}\text{Se } \text{H}-^{78}\text{Kr}$	7367	26	7372.8	0.3	0.1	U			R11	1.5	78Di09
$^{78}\text{Kr}-^{78}\text{Se}$	3074	16	3057.10	0.28	-0.7	U			R11	1.5	78Di09
	3098	20			-1.4	U			R11	1.5	78Di09
	3057.18	0.29			-0.3	1	92	$89 ^{78}\text{Kr}$	MS1	1.0	13Bu17
$^{78}\text{Se } \text{H}-^{78}\text{Kr}$	4724	33	4767.93	0.28	0.9	U			R11	1.5	78Di09
$^{76}\text{Rb}-^{78}\text{Rb}_{.325}^{x} \ ^{75}\text{Rb}_{.676}$	-130	40	-69	4	0.6	U			P20	2.5	82Au01
$^{77}\text{Rb}-^{78}\text{Rb}_{.494}^{x} \ ^{76}\text{Rb}_{.507}$	-1192	19	-1138	6	1.1	U			P20	2.5	82Au01
$^{78}\text{Kr}(\alpha, ^8\text{He})^{74}\text{Kr}$	-41080	75	-41031.2	2.0	0.7	U			Tex		82Mo23 *
$^{78}\text{Se}(\text{p}, \alpha)^{75}\text{As}$	870.9	2.3	872.3	0.9	0.6	1	15	$15 ^{75}\text{As}$	NDm		82Zu04
$^{78}\text{Kr}(\beta^-, ^6\text{He})^{75}\text{Kr}$	-12581	14	-12516	8	4.7	B					87Mo06
$^{76}\text{Ge}(\text{t}, \text{p})^{78}\text{Ge}$	6310	5	6310	4	0.0	2			LAI		78Ar12
	6310	5			0.0	2			Phi		81St18
$^{78}\text{Se}(\text{p}, \text{t})^{76}\text{Se}$	-9433.7	4.3	-9434.83	0.18	-0.3	U			NDm		82Zu04
$^{78}\text{Kr}(\alpha, ^6\text{He})^{76}\text{Kr}$	-20351	10	-20332	4	1.9	o			Tex		82Mo23 *
$^{78}\text{Kr}(\text{p}, \text{t})^{76}\text{Kr}$	-12840	15	-12825	4	1.0	U			Tky		81Ma30
$^{78}\text{Se}(\text{d}, ^3\text{He})^{77}\text{As}$	-4904	4	-4905.1	1.7	-0.3	1	18	$18 ^{77}\text{As}$	Ors		83Ro08 *
$^{77}\text{Se}(\text{n}, \gamma)^{78}\text{Se}$	10497.7	0.3	10497.77	0.17	0.2	-			BNn		81En07 Z
	10497.75	0.21			0.1	-			Bdn		06Fi.A
$^{78}\text{Se}(\text{p}, \text{d})^{77}\text{Se}$	-8271.9	4.0	-8273.21	0.17	-0.3	U			NDm		82Zu04
$^{77}\text{Se}(\text{n}, \gamma)^{78}\text{Se}$	ave. 10497.73	0.17	10497.77	0.17	0.2	1	96	$95 ^{78}\text{Se}$			average
$^{78}\text{Kr}(\text{d}, \text{t})^{77}\text{Kr}$	-5804	7	-5822.9	2.0	-2.7	U					87Mo06
$^{78}\text{Zn}(\beta^-)^{78}\text{Ga}$	6440	140	6222.7	2.7	-1.6	o			Stu		86Ek01
	6364	90			-1.6	U			Stu		00Me.A
$^{78}\text{Ga}(\beta^-)^{78}\text{Ge}$	8140	160	8156	4	0.1	o			Stu		77Al17
	8200	80			-0.5	o			Stu		86Ek01
	8054	43			2.4	U			Stu		00Me.A
$^{78}\text{Ge}(\beta^-)^{78}\text{As}$	967	30	955	10	-0.4	R					65Fr04 *
	987	20			-1.6	R					65Kv01 *
$^{78}\text{As}(\beta^-)^{78}\text{Se}$	4270	100	4209	10	-0.6	U					70Mc01
	4310	100			-1.0	U					71Mo20 *
$^{78}\text{Br}(\beta^+)^{78}\text{Se}$	3542	50	3574	4	0.6	U			Bar		61Ri02
$^{78}\text{Se}(\text{p}, \text{n})^{78}\text{Br}$	-4344	10	-4356	4	-1.2	2			Bar		61Ri02
	-4370	10			1.4	2			LAI		61Sc11
	-4355.5	7.4			-0.1	2			Tkm		63Ok01 Z
	-4356	5			0.0	2					70Fi03 Z
$^{78}\text{Rb}(\beta^+)^{78}\text{Kr}$	7085	370	7243	3	0.4	U					81Ba40 *
	7240	50			0.1	U					93Al03 *
	7185	50			1.2	U			IRS		93Al03 *
$^{78}\text{Rb}^{\text{v}}(\text{IT})^{78}\text{Rb}$	74	12				3					82Au01 *
$^{78}\text{Rb}-^{85}\text{Rb}_{.918}$	Correction for $e^-$ binding= $+97\text{eV}$ is negligible										
$^{78}\text{Rb}-^{85}\text{Rb}_{.918}$	$D_M=9237.6(4.5) \mu\text{u } M-A=-66824.9(4.2)\text{keV}$ corrected for $e^-$ binding= $+97\text{eV}$										
$^{78}\text{Rb}-^{85}\text{Rb}_{.918}$	from $^{78}\text{Rb}^{\text{v}}$ at $111.19(0.22) \text{ keV}$										
$^{78}\text{Kr}(\alpha, ^8\text{He})^{74}\text{Kr}$	Original $-41120(75)$ for 4 events included 1 background event										
$^{78}\text{Kr}(\alpha, ^6\text{He})^{76}\text{Kr}$	Replaced by calibration free $^{80}\text{Kr}(\alpha, ^6\text{He})^{78}\text{Kr}-^{78}\text{Kr}(\text{t})^{76}\text{Kr}$										
$^{78}\text{Se}(\text{d}, ^3\text{He})^{77}\text{As}$	Original value $-4910(4)$ corrected, see $^{74}\text{Se}(\text{d}, ^3\text{He})$										
$^{78}\text{Ge}(\beta^-)^{78}\text{As}$	$E_{\beta^-} = 690(30) \text{ 710(20)}$ respectively, to $1^+$ level at $277.3 \text{ keV}$										
$^{78}\text{As}(\beta^-)^{78}\text{Se}$	$E_{\beta^-} = 3000(100)$ to $2^+$ level at $1308.644 \text{ keV}$										
$^{78}\text{Rb}(\beta^+)^{78}\text{Kr}$	$E_{\beta^+} = 3410(370)$ from $^{78}\text{Rb}^{\text{v}} 4^{(-)}$ at $111.19(0.22)$ to $(4)^-$ level at $2764.10 \text{ keV}$										
$^{78}\text{Rb}(\beta^+)^{78}\text{Kr}$	$Q_{\beta^+} = 7180(80)$ ; and $7300(50)$ from $^{78}\text{Rb}^{\text{v}}$ at $111.19(0.22) \text{ keV}$										
$^{78}\text{Rb}^{\text{v}}(\text{IT})^{78}\text{Rb}$	Corrected; using $^{78}\text{Rb}^{\text{v}}(\text{IT}) = 111.2 \text{ keV}$										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{79}\text{Cu}-u$	-46700	540	-44810#	320#	3.5	D			OR1	1.0	06Ha62 *
$^{79}\text{Zn}-^{88}\text{Rb}_{.898}$	22278.1	2.9	22276.7	2.4	-0.5	1	68	68 $^{79}\text{Zn}$	JY1	1.0	08Ha23
$^{79}\text{Ga}-^{88}\text{Rb}_{.898}$	12490.9	2.0	12490.9	2.0	0.0	1	100	100 $^{79}\text{Ga}$	JY1	1.0	08Ha23
$^{79}\text{Ga}-u$	-67064	129	-67147.7	2.0	-0.3	U			GT2	2.5	08Su19
$\text{C}_6 \text{H}_7-^{79}\text{Br}$	136444.3	2.4	136437.6	1.1	-1.1	U			M15	2.5	63Ri07
	136444	15			-0.3	U			R11	1.5	78Di09
	136449	12			-0.6	U			R11	1.5	78Di09
$\text{C}_5 \text{}^{13}\text{C} \text{H}_6-^{79}\text{Br}$	131976	16	131967.4	1.1	-0.4	U			R11	1.5	78Di09
	131974	17			-0.3	U			R11	1.5	78Di09
$\text{C}_5 \text{N} \text{H}_5-^{79}\text{Br}$	123870	7	123861.6	1.1	-0.8	U			R11	1.5	78Di09
	123871	14			-0.4	U			R11	1.5	78Di09
$\text{C}_5 \text{O} \text{H}_3-^{79}\text{Br}$	100061	15	100052.1	1.1	-0.4	U			R11	1.5	78Di09
	100057	20			-0.2	U			R11	1.5	78Di09
$\text{C}_4 \text{N} \text{O} \text{H}-^{79}\text{Br}$	87489	20	87476.1	1.1	-0.4	U			R11	1.5	78Di09
$^{79}\text{Kr}-u$	-79981	52	-79917	4	1.2	U			GS2	1.0	05Li24 *
$^{79}\text{Y}-u$	-62070	85				2			LZ1	1.0	16Xi.A
$^{79}\text{Zn}-^{85}\text{Rb}_{.929}$	24582.4	4.2	24585.4	2.4	0.7	1	32	32 $^{79}\text{Zn}$	MA8	1.0	08Ba54
$^{79}\text{Rb}-^{85}\text{Rb}_{.929}$	5934	8	5937.2	2.3	0.4	U			MA2	1.0	94Ot01
	5937.2	2.3				2			MA8	1.0	07Ke09
$^{79}\text{Sr}-^{85}\text{Rb}_{.929}$	11655	9				2			MA2	1.0	94Ot01
$^{77}\text{Se} \text{H}_2-^{79}\text{Br}$	17239	8	17226.6	1.1	-1.0	U			R11	1.5	78Di09
$^{78}\text{Se} \text{H}-^{79}\text{Br}$	6806	8	6796.7	1.1	-0.8	U			R11	1.5	78Di09
$^{79}\text{Br}-^{78}\text{Kr}$	-2072	30	-2028.7	1.1	1.0	U			R11	1.5	78Di09
$^{77}\text{Rb}-^{79}\text{Rb}_{.487} \text{}^{75}\text{Rb}_{.513}$	-1010	40	-996.2	1.8	0.1	U			P20	2.5	82Au01
$^{77}\text{Rb}-^{79}\text{Rb}_{.325} \text{}^{76}\text{Rb}_{.675}$	-1060	40	-996.1	1.6	0.6	U			P20	2.5	82Au01
	-990	70			0.0	U			P20	2.5	82Au01
$^{78}\text{Rb}^+-^{79}\text{Rb}_{.494} \text{}^{77}\text{Rb}_{.506}$	940	40	919	12	-0.2	U			P20	2.5	82Au01
$^{78}\text{Se}(n,\gamma)^{79}\text{Se}$	6962.6	0.3	6962.83	0.13	0.8	2					79Br.A Z
	6962.2	0.3			2.1	2			BNn		81En07 Z
	6963.11	0.17			-1.6	2			Bdn		06Fi.A
$^{78}\text{Se}(d,p)^{79}\text{Se}$	4756	6	4738.27	0.13	-3.0	B			MIT		64Sp12
$^{78}\text{Kr}(d,p)^{79}\text{Kr}$	5980	50	6111	3	2.6	U			Yal		56B110
$^{78}\text{Kr}(^3\text{He},d)^{79}\text{Rb}$	-1585	10	-1579.8	2.2	0.5	U			Phi		87St11
$^{79}\text{Zn}(\beta^-)^{79}\text{Ga}$	8550	240	9115.4	2.9	2.4	U			Stu		86Ek01
$^{79}\text{Ga}(\beta^-)^{79}\text{Ge}$	6770	80	6980	40	2.6	o			Stu		77A117
	7000	80			-0.3	o			Stu		86Ek01
	6979	40			0.0	1	86	86 $^{79}\text{Ge}$	Stu		00Me.A
$^{79}\text{Ge}(\beta^-)^{79}\text{As}$	4300	200	4110	40	-1.0	U					70Ka04
	4110	100			0.0	1	14	14 $^{79}\text{Ge}$	Stu		81Al20
$^{79}\text{As}(\beta^-)^{79}\text{Se}$	2230	50	2281	5	1.0	U					61Ku09 *
$^{79}\text{Se}(\beta^-)^{79}\text{Br}$	160	5	150.6	1.0	-1.9	U					49Pa.A
$^{79}\text{Kr}(\beta^+)^{79}\text{Br}$	1612	10	1626	3	1.4	4					52Be55
	1620	5			1.2	4					54Th39
	1635	5			-1.8	4					64Bo25
$^{79}\text{Rb}(\beta^+)^{79}\text{Kr}$	3530	50	3639	4	2.2	U					71Li02 *
	3720	90			-0.9	U					72Br31 *
	3650	70			-0.2	U			IRS		93Al03
$^{79}\text{Sr}(\beta^+)^{79}\text{Rb}$	5259	78	5326	9	0.9	U			BNL		81Li12
	5059	67			4.0	B			Ors		82De36
$^{79}\text{Y}(\beta^+)^{79}\text{Sr}$	7120	450	7660	80	1.2	U					92Mu12
* $^{79}\text{Cu}-u$	Trends from Mass Surface TMS suggest $^{79}\text{Cu}$ 1760 less bound										
* $^{79}\text{Kr}-u$	$M-A=-74437(30)$ keV for mixture gs+m at 129.77 7/2 <sup>+</sup> keV										
* $^{79}\text{As}(\beta^-)^{79}\text{Se}$	$E_{\beta^-}=1700(50)$ to 527.93 3/2 <sup>-</sup> level, and other $E_{\beta^-}$										
* $^{79}\text{Rb}(\beta^+)^{79}\text{Kr}$	$E_{\beta^+}=1825(50)$ 2010(90) respectively, to 3/2 <sup>+</sup> level at 688.17 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{80}\text{Ga}-\text{u}$	-63441	129	-63579	3	-0.4	U			GT2	2.5	08Su19
$\text{C}_6 \text{H}_8-^{80}\text{Se}$	146068.5	2.9	146078.5	1.0	1.4	U			M15	2.5	63Ri07
$\text{C}_6 \text{H}_8-^{80}\text{Kr}$	146225.7	4.6	146222.2	0.7	-0.3	U			M15	2.5	63Ri07
	146235	18			-0.5	U			R11	1.5	78Di09
	146215	16			0.3	U			R11	1.5	78Di09
$\text{C}_5 \text{O H}_4-^{80}\text{Kr}$	109834	20	109836.7	0.7	0.1	U			R11	1.5	78Di09
$^{80}\text{Y}-\text{u}$	-65720	190	-65645	7	0.4	U			1.0	1.0	98Is06
	-66664	86			11.8	F			CS1	1.0	01La31 *
	-65600	200			-0.2	U			CS1	1.0	08Go23
$^{80}\text{Y O}-^{96}\text{Mo}$	24594.6	6.7				2			JY1	1.0	06Ka48
$^{80}\text{Zr}-\text{u}$	-59600	1600	-58360#	320#	0.8	D			1.0	1.0	98Is06 *
	-59740	161			8.6	F			CS1	1.0	01La31 *
$^{80}\text{Zn}-^{88}\text{Rb}_{.909}$	25165.2	7.3	25167.1	2.8	0.3	1	14	14 $^{80}\text{Zn}$	JY1	1.0	08Ha23
$^{80}\text{Ga}-^{88}\text{Rb}_{.909}$	17034.9	3.1				2			JY1	1.0	08Ha23
$^{80}\text{Ge}-^{88}\text{Rb}_{.909}$	5964.9	2.2				2			JY1	1.0	08Ha23
$^{80}\text{Kr}-^{86}\text{Kr}_{.930}$	-488.9	1.1	-489.8	0.7	-0.8	1	46	46 $^{80}\text{Kr}$	MS1	1.0	06Ri15
$^{80}\text{Zn}-^{85}\text{Rb}_{.941}$	27559.1	3.0	27558.8	2.8	-0.1	1	86	86 $^{80}\text{Zn}$	MA8	1.0	08Ba54
$^{80}\text{Kr}-^{85}\text{Rb}_{.941}$	-614.5	1.7	-616.1	0.7	-0.9	1	19	19 $^{80}\text{Kr}$	MA8	1.0	06Ro11
	-627.1	9.6			1.1	U			MA8	1.0	10Na13 *
$^{80}\text{Rb}-^{85}\text{Rb}_{.941}$	5528	8	5522.3	2.0	-0.7	U			MA2	1.0	94Ot01
	5522.3	2.0				2			MA8	1.0	07Ke09
$^{80}\text{Sr}-^{85}\text{Rb}_{.941}$	7531	8	7523	4	-1.0	2			MA2	1.0	94Ot01
	7513	14			0.7	U			MA8	1.0	05Si34
	7521.3	4.2			0.5	2			SH1	1.0	11Ha08
$^{80}\text{Se } ^{35}\text{Cl}-^{78}\text{Se } ^{37}\text{Cl}$	2164.8	1.4	2162.7	1.0	-0.4	U			H18	4.0	64Ba03
	2160.8	9.2			0.1	U			H40	2.5	85Ei01
$^{80}\text{As}-^{80}\text{Kr}$	6096.5	3.5				2			MS1	1.0	07Bo50
$^{80}\text{Kr}-^{79}\text{Br}$	-1955	28	-1959.6	1.2	-0.1	U			R11	1.5	78Di09
$^{80}\text{Kr}-^{78}\text{Kr}$	-4046	30	-3988.3	0.8	1.3	U			R11	1.5	78Di09
$^{79}\text{Rb}-^{80}\text{Rb}_{.658} \text{ } ^{77}\text{Rb}_{.342}$	-1218	27	-1139.5	2.5	1.2	U			P20	2.5	82Au01
$^{79}\text{Rb}-^{80}\text{Rb}_{.494} \text{ } ^{78}\text{Rb}_{.506}$	-1313	24	-1316	7	-0.1	U			P20	2.5	82Au01
$^{80}\text{Se}(\text{p},\alpha)^{77}\text{As}$	1020.0	2.8	1020.9	1.8	0.3	1	40	32 $^{77}\text{As}$	NDm		82Zu04
$^{80}\text{Kr}(\alpha,^3\text{He},^6\text{He})^{77}\text{Kr}$	-10398	24	-10384.8	2.1	0.6	U					87Mo06
$^{80}\text{Se}(\text{d},\alpha)^{78}\text{As}$	5755	12	5768	10	1.1	2			Phi		77Mo13
$^{80}\text{Se}(\text{p,t})^{78}\text{Se}$	-8395.1	3.0	-8394.4	1.0	0.2	-			NDm		82Zu04
ave.	-8394.1	2.1			-0.1	1	21	20 $^{80}\text{Se}$			average
$^{80}\text{Kr}(\alpha,^6\text{He})^{78}\text{Kr}-^{78}\text{Kr}(\text{)}^{76}\text{Kr}$	1432	10	1449	4	1.7	1	17	16 $^{76}\text{Kr}$			82Mo23
$^{78}\text{Kr}(\alpha,^3\text{He},\text{n})^{80}\text{Sr}$	2990	30	2993	3	0.1	U					79A119
$^{80}\text{Se}(\text{d},^3\text{He})^{79}\text{As}$	-5921	7	-5919	5	0.3	-			Ors		83Ro08 *
	-5921	13			0.2	-			Hei		83Wi14
$^{80}\text{Se}(\text{t},\alpha)^{79}\text{As}$	8407	10	8401	5	-0.6	-			Phi		83Mo09
$^{80}\text{Se}(\text{d},^3\text{He})^{79}\text{As}$	ave. -5919	5	-5919	5	0.0	1	100	100 $^{79}\text{As}$			average
$^{80}\text{Se}(\text{p,d})^{79}\text{Se}$	-7687.6	3.0	-7688.7	1.0	-0.4	R			NDm		82Zu04
$^{79}\text{Br}(\text{n},\gamma)^{80}\text{Br}$	7892.11	0.20	7892.28	0.13	0.8	3			ILn		78Do06 Z
	7892.41	0.18			-0.7	3			Bdn		06Fi.A
$^{79}\text{Br}(\text{d,p})^{80}\text{Br}$	5640	20	5667.71	0.13	1.4	U			Mtr		72Ch33
$^{80}\text{Zn}(\beta^-)^{80}\text{Ga}$	7540	200	7575	4	0.2	U			Stu		86Ek01
	7150	150			2.8	U			Trs		86Gi07
$^{80}\text{Ga}(\beta^-)^{80}\text{Ge}$	10000	300	10312	4	1.0	o			Stu		81Al20
	10380	120			-0.6	U			Stu		86Ek01
$^{80}\text{Ge}(\beta^-)^{80}\text{As}$	2640	70	2679	4	0.6	U			Stu		77A117
	2630	20			2.5	U			Trs		86Gi07
$^{80}\text{As}(\beta^-)^{80}\text{Se}$	6000	200	5545	3	-2.3	U					59Me68
	5470	90			0.8	U			Trs		86Gi07
$^{80}\text{Se}(\text{t},^3\text{He})^{80}\text{As}$	-5560	25	-5526	3	1.3	U			LAl		79Aj02

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{80}\text{Br}(\beta^+)^{80}\text{Se}$	1884	10	1870.5	0.3	-1.4	U					54Li19
	1872	7			-0.2	U					69Ka06
$^{80}\text{Se}(\text{p,n})^{80}\text{Br}$	-2655.2	2.8	-2652.8	0.3	0.9	U			Tkm		63Ok01
	-2652.5	3.0			-0.1	U			Oak		64Jo11 Z
	-2653.2	5.			0.1	U					70Fi03
	-2652.81	0.31				2			PTB		92Bo02 Z
$^{80}\text{Br}(\beta^-)^{80}\text{Kr}$	1970	30	2004.4	1.2	1.1	U					52Fu04
	2040	20			-1.8	U					54Li19
	1997	10			0.7	U					69Ka06
$^{80}\text{Rb}(\beta^+)^{80}\text{Kr}$	5120	500	5717.9	2.0	1.2	U					61Ho13
	5500	350			0.6	U					75We23 *
	5650	100			0.7	U			IRS		93Al03
$^{80}\text{Kr}(\text{p,n})^{80}\text{Rb}$	-6484.0	20.	-6500.2	2.0	-0.8	U					72Ja.A
$^{80}\text{Y}(\beta^+)^{80}\text{Sr}$	6952	152	9163	7	14.5	F			BNL		81Li12 *
	6934	242			9.2	F			Ors		82De36 *
	6200	600			4.9	F					96Sh27 *
* $^{80}\text{Y}-\text{u}$	F : below lower limit $M > -65890(90) \mu\text{u} - 61376(83) \text{keV}$ determined in reference										
* $^{80}\text{Zr}-\text{u}$	Trends from Mass Surface TMS suggest $^{80}\text{Zr}$ 1160 less bound										
* $^{80}\text{Zr}-\text{u}$	F : other results in same paper not trusted, see $^{80}\text{Y}$ and $^{68}\text{Se}$										
* $^{80}\text{Kr}-^{85}\text{Rb}_{.941}$	Only one measurement										
* $^{80}\text{Se}(\text{d},^3\text{He})^{79}\text{As}$	Originally -5927(7), see $^{74}\text{Se}(\text{d},^3\text{He})$										
* $^{80}\text{Rb}(\beta^+)^{80}\text{Kr}$	$E_{\beta^+} = 3860(350) \text{ to } 2^+ \text{ level at } 616.60 \text{ keV}$										
* $^{80}\text{Y}(\beta^+)^{80}\text{Sr}$	F : below lower limit $Q_{\beta^-} > 8929(23) \text{ keV}$ determined in reference										
$^{81}\text{Ge}-\text{u}$	-71710	240	-71167.1	2.2	0.9	U			GT2	2.5	08Kn.A *
$\text{C}_6 \text{H}_9-^{81}\text{Br}$	154135.3	3.8	154137.1	1.0	0.2	U			M15	2.5	63Ri07
	154143	17			-0.2	U			R11	1.5	78Di09
	154134	10			0.2	U			R11	1.5	78Di09
$\text{C}_5 \text{N H}_7-^{81}\text{Br}$	141561	10	141561.0	1.0	0.0	U			R11	1.5	78Di09
	141553	18			0.3	U			R11	1.5	78Di09
$\text{C}_5 \text{O H}_5-^{81}\text{Br}$	117742	12	117751.6	1.0	0.5	U			R11	1.5	78Di09
$\text{C}_4 \text{O}_2 \text{H}-^{81}\text{Br}$	81356	20	81366.1	1.0	0.3	U			R11	1.5	78Di09
$\text{C}_4^{13}\text{C O H}_4-^{81}\text{Br}$	113275	14	113281.4	1.0	0.3	U			R11	1.5	78Di09
$^{81}\text{Rb}-\text{u}$	-80958	41	-81006	5	-1.2	U			GS2	1.0	05Li24 *
$^{81}\text{Y O}-^{97}\text{Mo}$	18352.0	5.8				2			JY1	1.0	06Ka48
$^{81}\text{Zr}-\text{u}$	-61686	101				2			LZ1	1.0	16Xi.A
$^{81}\text{Ga}-^{88}\text{Rb}_{.920}$	19723.5	3.5				2			JY1	1.0	08Ha23
$^{81}\text{Ge}-^{88}\text{Rb}_{.920}$	10422.6	2.2				2			JY1	1.0	08Ha23
$^{81}\text{As}-^{88}\text{Rb}_{.920}$	3721.9	3.3	3721.9	2.8	0.0	1	74	74 $^{81}\text{As}$	JY1	1.0	08Ha23
$^{81}\text{Zn}-^{85}\text{Rb}_{.953}$	34467.0	5.4				2			MA8	1.0	08Ba54
$^{81}\text{Rb}-^{85}\text{Rb}_{.953}$	3063	8	3058	5	-0.6	-			MA2	1.0	94Ot01
	3055.4	9.2			0.3	-			SH1	1.0	11Ha08 *
	ave.	6			-0.2	1	76	76 $^{81}\text{Rb}$			average
$^{81}\text{Sr}-^{85}\text{Rb}_{.953}$	7278	8	7276	3	-0.3	2			MA2	1.0	94Ot01
	7272	12			0.3	U			MA8	1.0	05Si34
	7275.3	3.7			0.1	2			SH1	1.0	11Ha08
$^{81}\text{Se}-^{80}\text{Kr}_{1.013}$	2704.2	2.4	2702.1	1.2	-0.9	1	26	18 $^{81}\text{Se}$	MS1	1.0	07Bo50 *
$^{80}\text{Se H}-^{81}\text{Br}$	8023	8	8058.6	1.5	3.0	B			R11	1.5	78Di09
$^{80}\text{Kr H}-^{81}\text{Br}$	7922	18	7914.9	1.3	-0.3	U			R11	1.5	78Di09
$^{81}\text{Br}-\text{H }^{79}\text{Br}$	-9865	13	-9874.4	1.5	-0.3	U			M15	2.5	63Ri07
$^{81}\text{Br}-^{80}\text{Kr}$	-91	32	-89.8	1.3	0.0	U			R11	1.5	78Di09
$^{81}\text{Br}-^{79}\text{Br}$	-2020	32	-2049.4	1.5	-0.6	U			R11	1.5	78Di09
	-2014	35			-0.7	U			R11	1.5	78Di09
$^{79}\text{Rb}-^{81}\text{Rb}_{.325}$ $^{78}\text{Rb}_{.675}$	-1130	30	-1148	9	-0.2	U			P20	2.5	82Au01 Y
$^{80}\text{Rb}-^{81}\text{Rb}_{.494}$ $^{79}\text{Rb}_{.506}$	927	29	926	3	0.0	U			P20	2.5	82Au01 Y
$^{80}\text{Se}(\text{n},\gamma)^{81}\text{Se}$	6700.9	0.5	6700.8	0.3	-0.1	-			BNn		81En07 Z
	6700.9	0.5			-0.1	-			Bdn		06Fi.A



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{80}\text{Se}(d,p)^{81}\text{Se}$	4490	6	4476.3	0.3	-2.3	U			MIT		64Sp12
	4477.5	3.0			-0.4	U		NDm			82Zu04
$^{80}\text{Se}(n,\gamma)^{81}\text{Se}$	ave. 6700.9	0.4	6700.8	0.3	-0.1	1	97	$^{71}\text{Se}$			average
$^{81}\text{Br}(\gamma,n)^{80}\text{Br}$	-10130	35	-10159.4	1.4	-0.8	U			Phi		60Ge01
$^{80}\text{Kr}(d,p)^{81}\text{Kr}$	5660	15	5649.6	1.2	-0.7	U			Tex		75Ch11 *
	5646	4			0.9	1	10	$^{71}\text{Kr}$	Oak		86Bu18
$^{80}\text{Kr}(^3\text{He},d)^{81}\text{Rb}$	-637	10	-641	5	-0.4	1	24	$^{24}\text{Rb}$	Phi		87St11
$^{81}\text{Zr}(\epsilon p)^{80}\text{Sr}$	4700	200	5560	90	4.3	B					99Hu05
$^{81}\text{Ga}(\beta^-)^{81}\text{Ge}$	8320	150	8664	4	2.3	U			Stu		81Al20
$^{81}\text{Ge}(\beta^-)^{81}\text{As}$	6230	120	6242	3	0.1	U			Stu		81Al20 *
$^{81}\text{As}(\beta^-)^{81}\text{Se}$	3800	200	3855.7	2.8	0.3	U					60Mo01
	3730	100			1.3	U			Stu		77Al17
$^{81}\text{Se}(\beta^-)^{81}\text{Br}$	1600	50	1588.0	1.4	-0.2	U					60Ku06
	1560	50			0.6	U					67Yt03
$^{81}\text{Kr}(\epsilon)^{81}\text{Br}$	280.7	0.5	280.9	0.5	0.3	1	89	$^{84}\text{Kr}$			88Ax01 *
$^{81}\text{Br}(p,n)^{81}\text{Kr}$	-1062	4	-1063.2	0.5	-0.3	U					84Fi.A
$^{81}\text{Br}(^3\text{He},t)^{81}\text{Kr}$	-296	6	-299.4	0.5	-0.6	U					84Bu23
$^{81}\text{Br}(^3\text{He},t)^{81}\text{Kr}-^{51}\text{V}()^{51}\text{Cr}$	470.6	1.8	471.6	0.5	0.6	U			Pri		82Ko06 *
$^{81}\text{Rb}(\beta^+)^{81}\text{Kr}$	2260	30	2240	5	-0.7	U					75Va24 *
	2290	50			-1.0	U					77Li14
$^{81}\text{Sr}(\beta^+)^{81}\text{Rb}$	3990	30	3929	6	-2.0	U					73Br32 *
$^{81}\text{Y}(\beta^+)^{81}\text{Sr}$	5408	86	5815	6	4.7	B			BNL		81Li12
	5620	89			2.2	U			Ors		82De36
$^{81}\text{Zr}(\beta^+)^{81}\text{Y}$	7160	290	8250	90	3.8	B			Ors		82De36
$^{81}\text{Ge}-u$	$M-A=-66454(93)$ keV for mixture gs+m at 679.14 keV										Nub16b **
$^{81}\text{Rb}-u$	$M-A=-75369(29)$ keV for mixture gs+m at 86.31 keV										Nub16b **
$^{81}\text{Rb}-^{85}\text{Rb}_{.953}$	$D_M=3148.1(9.2)$ keV for $^{81}\text{Rb}^m$ at 86.31(0.07) keV; $M-A=-75373.1(8.6)$ keV										Nub16b **
$^{81}\text{Se}-^{80}\text{Kr}_{1.013}$	$D_M=2814.8(2.4)$ $\mu$ u for $^{81}\text{Se}^m$ at 103.00(0.06)keV; $M-A=-76283.2(2.4)$ keV										Nub16b **
$^{80}\text{Kr}(d,p)^{81}\text{Kr}$	Original value 5610(15) reinterpreted as going to 49.57 level										76Me08 **
$^{81}\text{Ge}(\beta^-)^{81}\text{As}$	$Q_{\beta^-}=6230(120)$ ; and 6930(280) from $^{81}\text{Ge}^m$ at 679.14 keV										Nub16b **
$^{81}\text{Kr}(\epsilon)^{81}\text{Br}$	$LM=0.42(0.05)$ , $Q(\epsilon)=4.7(0.5)$ to $5/2^-$ level at 275.985 keV										Ens08a **
$^{81}\text{Br}(^3\text{He},t)^{81}\text{Kr}-^{51}\text{V}()^{51}\text{Cr}$	$Q-Q$ to 456.89(0.03) level=13.7(1.8) keV										GAu **
$^{81}\text{Rb}(\beta^+)^{81}\text{Kr}$	$E_{\beta^+}=1050(30)$ to $1/2^-$ level at 190.64 keV										Ens08a **
$^{81}\text{Sr}(\beta^+)^{81}\text{Rb}$	$E_{\beta^+}=2684(30)$ to 301.241 $3/2^-$ level, and other $E_{\beta^+}$										Ens08a **
$^{82}\text{Ga}-u$	-56812	268	-56823.5	2.6	0.0	U			GT1	1.5	04Ma.A
$^{82}\text{Ge}-u$	-70400	129	-70226.0	2.4	0.5	U			GT2	2.5	08Su19
$\text{C}_6 \text{H}_{10}-^{82}\text{Se}$	161545.0	4.6	161550.8	0.5	0.5	U			M15	2.5	63Ri07
$\text{C}_6 \text{H}_{10}-^{82}\text{Kr}$	164769.8	3.4	164769.167	0.006	-0.1	U			M15	2.5	63Ri07
	164787	14			-0.8	U			R11	1.5	78Di09
	164784	16			-0.6	U			R11	1.5	78Di09
$\text{C}_5 \text{N H}_8-^{82}\text{Kr}$	152200	25	152193.107	0.006	-0.2	U			R11	1.5	78Di09
$\text{C}_5 \text{O H}_6-^{82}\text{Kr}$	128396	20	128383.658	0.006	-0.4	U			R11	1.5	78Di09
$^{82}\text{Rb}-u$	-81775	39	-81791	3	-0.4	U			GS2	1.0	05Li24 *
$^{82}\text{Sr}-u$	-81604	63	-81600	6	0.1	U			GS2	1.0	05Li24
$^{82}\text{Y O}-^{98}\text{Mo}$	16441.2	5.9				2			JY1	1.0	06Ka48
$^{82}\text{Zr}-u$	-68311	12				2			LZ1	1.0	16Xi.A
$^{82}\text{Ga}-^{88}\text{Rb}_{.932}$	25830.4	2.6				2			JY1	1.0	08Ha23
$^{82}\text{Ge}-^{88}\text{Rb}_{.932}$	12427.9	2.4				2			JY1	1.0	08Ha23
$^{82}\text{As}-^{88}\text{Rb}_{.932}$	7392.6	4.0				2			JY1	1.0	08Ha23 *
$^{82}\text{Kr}-^{86}\text{Kr}_{.953}$	-1329.4	1.1	-1330.771	0.005	-1.2	U			MS1	1.0	06Ri15
	-1330.7684	0.0094			-0.3	1	26	$^{25}\text{Kr}$	FS1	1.0	13Ho22
$^{82}\text{Zn}-^{85}\text{Rb}_{.965}$	39697.0	3.3				2			MA8	1.0	13Wo06
$^{82}\text{Kr}-^{85}\text{Rb}_{.965}$	-1394.9	2.6	-1395.944	0.006	-0.4	U			MA8	1.0	06Ro11
$^{82}\text{Rb}^m-^{85}\text{Rb}_{.965}$	3407	9	3406.0	2.8	-0.1	U			MA2	1.0	94O01
	3406.0	2.8				2			MA8	1.0	05Gu37

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{82}\text{Sr}-^{85}\text{Rb}_{.965}$	3517	8	3523	6	0.7	1	65	$^{65}\text{ }^{82}\text{Sr}$	MA2	1.0	94Ot01
$^{82}\text{Se } ^{35}\text{Cl}-^{80}\text{Se } ^{37}\text{Cl}$	3128.92	0.63	3127.9	1.0	-0.7	1	41	$^{37}\text{ }^{80}\text{Se}$	H40	2.5	85El01
$^{80}\text{Se H}_2-^{82}\text{Kr}$	18665	18	18690.7	1.0	1.0	U			R11	1.5	78Di09
	18671	19			0.7	U			R11	1.5	78Di09
$^{82}\text{Kr}-^{84}\text{Kr}_{.976}$	-140.6277	0.0051	-140.627	0.005	0.1	1	78	$^{75}\text{ }^{82}\text{Kr}$	FS1	1.0	13Ho22
$^{81}\text{Br H}-^{82}\text{Se}$	7419	8	7413.7	1.2	-0.4	U			R11	1.5	78Di09
$^{81}\text{Br H}-^{82}\text{Kr}$	10662	20	10632.1	1.0	-1.0	U			R11	1.5	78Di09
$^{82}\text{Se}-^{82}\text{Kr}$	3222	16	3218.4	0.5	-0.2	U			R11	1.5	78Di09
	3218	22			0.0	U			R11	1.5	78Di09
	3216.1	1.6			1.0	U			H45	1.5	93Nx01
	3218.36	0.52			0.0	1	93	$^{93}\text{ }^{82}\text{Se}$	MS1	1.0	13Li01
$^{82}\text{Kr}-^{78}\text{Se H}_3$	-27269	35	-27303.19	0.19	-0.7	U			R11	1.5	78Di09
$^{80}\text{Se H}_3-^{82}\text{Kr}$	26466	32	26515.7	1.0	1.0	U			R11	1.5	78Di09
$^{82}\text{Kr}-^{81}\text{Br}$	-2805	32	-2807.1	1.0	0.0	U			R11	1.5	78Di09
$^{82}\text{Se H}-^{82}\text{Kr}$	11082	40	11043.4	0.5	-0.6	U			R11	1.5	78Di09
$^{79}\text{Rb}-^{82}\text{Rb}_{.241}$ $^{78}\text{Rb}_{.760}^x$	-1536	29	-1627	10	-1.3	U			P20	2.5	82Au01 Y
$^{81}\text{Rb}-^{82}\text{Rb}_{.741}$ $^{78}\text{Rb}_{.260}^x$	-1680	40	-1618	6	0.6	U			P20	2.5	82Au01 Y
$^{80}\text{Rb}-^{82}\text{Rb}_{.325}$ $^{79}\text{Rb}_{.675}$	440	40	377.6	2.6	-0.6	U			P20	2.5	82Au01 Y
$^{82}\text{Kr}(^3\text{He}, ^6\text{He})^{79}\text{Kr}$	-8822	31	-8810	3	0.4	U					87Mo06
$^{82}\text{Se}(^{14}\text{C}, ^{16}\text{O})^{80}\text{Ge}$	-449	60	-301.7	2.1	2.5	U			Ors		83Be.C
$^{82}\text{Se}(^{18}\text{O}, ^{20}\text{Ne})^{80}\text{Ge}$	-2020	40	-1799.5	2.1	5.5	B			Hei		83Wi14 *
$^{82}\text{Se}(p,t)^{80}\text{Se}$	-7496.1	3.0	-7495.3	0.9	0.3	1	10	$^{9}\text{ }^{80}\text{Se}$	NDm		82Zu04
$^{82}\text{Se}(d, ^3\text{He})^{81}\text{As}$	-6864	10	-6856.1	2.7	0.8	-			Ors		83Ro08 *
	-6861	18			0.3	U			Hei		83Wi14
$^{82}\text{Se}(t, \alpha)^{81}\text{As}$	7467	6	7464.3	2.7	-0.4	-			Phi		82Mo04
$^{82}\text{Se}(d, ^3\text{He})^{81}\text{As}$	ave. -6856	5	-6856.1	2.7	0.0	1	27	$^{26}\text{ }^{81}\text{As}$			average
$^{82}\text{Se}(p,d)^{81}\text{Se}$	-7051.8	2.8	-7051.6	1.0	0.1	1	12	$^{11}\text{ }^{81}\text{Se}$	NDm		82Zu04
$^{81}\text{Br}(n, \gamma)^{82}\text{Br}$	7592.80	0.20	7592.94	0.12	0.7	-			ILn		78Do06 Z
	7593.02	0.15			-0.5	-			Bdn		06Fi.A
$^{81}\text{Br}(d,p)^{82}\text{Br}$	5400	20	5368.38	0.12	-1.6	U			Mtr		72Ch33
$^{81}\text{Br}(n, \gamma)^{82}\text{Br}$	ave. 7592.94	0.12	7592.94	0.12	0.0	1	100	$^{94}\text{ }^{81}\text{Br}$			average
$^{82}\text{Ge}(\beta^-)^{82}\text{As}$	4700	140	4690	4	-0.1	U			Stu		81Al20
$^{82}\text{As}(\beta^-)^{82}\text{Se}$	7270	200	7488	4	1.1	U					70Va31 *
	6360	200			5.6	B					70Ka04 *
	7740	30			-8.4	C			Stu		00Me.A *
	7531	21			-2.0	U			Stu		04Ga44 *
$^{82}\text{Se}(t, ^3\text{He})^{82}\text{As}$	-7500	25	-7470	4	1.2	U			LAl		79Aj02
$^{82}\text{Br}(\beta^-)^{82}\text{Kr}$	3092.9	1.0	3093.1	1.0	0.2	1	94	$^{94}\text{ }^{82}\text{Br}$			56Wa24 *
$^{82}\text{Rb}(\beta^+)^{82}\text{Kr}$	4400	15	4404	3	0.3	U					69Be74 *
	4420	60			-0.3	U			IRS		93Al03 *
$^{82}\text{Kr}(p,n)^{82}\text{Rb}$	-5161	20	-5186	3	-1.3	U					72Ja.A
$^{82}\text{Rb}^m(\text{IT})^{82}\text{Rb}$	69.0	1.5				3					Ens035
$^{82}\text{Y}(\beta^+)^{82}\text{Sr}$	7868	185	7946	8	0.4	U			BNL		81Li12
	7793	123			1.2	U			Ors		82De36
$^{82}\text{Zr}(\beta^+)^{82}\text{Y}$	4000	500	4433	12	0.9	F			Ors		82De36 *
$^{*82}\text{Rb}-u$	$M-A=-76138(30)$ keV for mixture gs+m at 69.0(1.5) keV										Nub16b **
$^{*82}\text{As}-^{88}\text{Rb}_{.932}$	$D_M=7395.1(4.6)$ 7532.5(4.0) $\mu\text{u}$ to ground state and $^{82}\text{As}^m$ at 132.1(0.2) keV										Nub16b **
$^{*82}\text{Se}(^{18}\text{O}, ^{20}\text{Ne})^{80}\text{Ge}$	Recalibrated to $^{64}\text{Ni}(^{62}\text{Fe})=-1938(15)$ keV										AHW **
$^{*82}\text{Se}(d, ^3\text{He})^{81}\text{As}$	Originally -6870(10), see $^{74}\text{Se}(d, ^3\text{He})$										AHW **
$^{*82}\text{As}(\beta^-)^{82}\text{Se}$	$E_{\beta^-}=7200(20)$ to ground state (80%) and 654.75 $2^+$ level (10%)										Ens035 **
$^{*82}\text{As}(\beta^-)^{82}\text{Se}$	$E_{\beta^-}=3600(200)$ from $^{82}\text{As}^m$ at 132.1 to $5^-$ level at 2893.70 and higher levels										Ens035 **
$^{*82}\text{As}(\beta^-)^{82}\text{Se}$	and $E_{\beta^-}=7625(22)$ from $^{82}\text{As}^m$ at 132.1 keV										00Me.A **
$^{*82}\text{As}(\beta^-)^{82}\text{Se}$	Average of 3 branches; and 7677(17) average of 2 branches from $^{82}\text{As}^m$ at 132.1										04Ga44 **
$^{*82}\text{Br}(\beta^-)^{82}\text{Kr}$	$E_{\beta^-}=444(1)$ to $4^-$ level at 2648.360 keV										Ens035 **
$^{*82}\text{Rb}(\beta^+)^{82}\text{Kr}$	$E_{\beta^+}=3350(60)$ ; and 800(15) from $^{82}\text{Rb}^m$ at 69.0(1.5) to $4^-$ level at 2648.360										Ens035 **
$^{*82}\text{Rb}(\beta^+)^{82}\text{Kr}$	$Q_{\beta^+}=4360(100)$ ; and 4510(60) of $^{82}\text{Rb}^m$ at 69.0(1.5) keV										Nub16b **
$^{*82}\text{Zr}(\beta^+)^{82}\text{Y}$	F : for 2.5(0.1) m activity, but Ensdf adopts 32(5) s										Ens035 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{83}\text{Ge}-u$	-65626	268	-65460.9	2.6	0.4	o			GT1	1.5	04Ma.A
	-65270	320			-0.6	U			OR1	1.0	06Ha62
	-65276	129			-0.6	U			GT2	2.5	08Su19
$^{83}\text{As}-u$	-74677	129	-74793	3	-0.4	U			GT2	2.5	08Su19
	$\text{C}_6 \text{H}_{11}-^{83}\text{Kr}$	171946.8	3.4	171948.836	0.010	0.2	U		M15	2.5	63Ri07
$\text{C}_5 \text{N} \text{H}_9-^{83}\text{Kr}$	171948	16			0.0	U			R11	1.5	78Di09
	159344	25	159372.776	0.010	0.8	U			R11	1.5	78Di09
	159360	19			0.4	U			R11	1.5	78Di09
$\text{C}_5 \text{O} \text{H}_7-^{83}\text{Kr}$	135543	25	135563.327	0.010	0.5	U			R11	1.5	78Di09
$^{83}\text{Y} \text{O}-^{98}\text{Mo}_{1,010}$	12941	20				2			JY1	1.0	06Ka48 *
$^{83}\text{Zr} \text{O}-^{98}\text{Mo}_{1,010}$	19697.9	6.9				2			JY1	1.0	06Ka48
$^{83}\text{Nb}-u$	-61789	162				2			LZ1	1.0	16Xi.A
$^{83}\text{Ga}-^{88}\text{Rb}_{,943}$	30749.7	2.8				2			JY1	1.0	08Ha23
$^{83}\text{Ge}-^{88}\text{Rb}_{,943}$	18168.5	2.6				2			JY1	1.0	08Ha23
$^{83}\text{As}-^{88}\text{Rb}_{,943}$	8836.3	3.0				2			JY1	1.0	08Ha23
$^{80}\text{Se} \text{H}_3-^{83}\text{Kr}$	25825	25	25870.4	1.0	1.2	U			R11	1.5	78Di09
$^{83}\text{Kr}-^{86}\text{Kr}_{,965}$	386.6	1.1	387.261	0.009	0.6	U			MS1	1.0	06Ri15
$^{83}\text{Rb}-^{85}\text{Rb}_{,976}$	1207	8	1207.4	2.5	0.1	U			MA2	1.0	94Ot01
	1207.4	2.5			0.0	1	100	100 $^{83}\text{Rb}$	MA8	1.0	07Ke09
$^{82}\text{Se} \text{H}-^{83}\text{Kr}$	10380	18	10398.1	0.5	0.7	U			R11	1.5	78Di09
	10368	16			1.3	U			R11	1.5	78Di09
$^{82}\text{Kr} \text{H}-^{83}\text{Kr}$	7160	18	7179.669	0.010	0.7	U			R11	1.5	78Di09
$^{83}\text{Kr}-^{84}\text{Kr}_{,988}$	1566.7601	0.0089	1566.760	0.009	0.0	1	100	100 $^{83}\text{Kr}$	FS1	1.0	13Ho22
$^{83}\text{Sr}-^{83}\text{Rb}$	2447	9	2440	7	-0.8	1	59	59 $^{83}\text{Sr}$	MA2	1.0	94Ot01
$^{83}\text{Kr}-^{80}\text{Se} \text{H}_2$	-18022	36	-18045.3	1.0	-0.4	U			R11	1.5	78Di09
$^{85}\text{Rb}-^{83}\text{Kr} \text{H}$	-10211	45	-10161.813	0.010	0.7	U			R11	1.5	78Di09
$^{83}\text{Kr}-^{82}\text{Se}$	-2572	35	-2573.0	0.5	0.0	U			R11	1.5	78Di09
$^{83}\text{Kr}-^{82}\text{Kr}$	648	12	645.363	0.010	-0.1	U			M15	2.5	63Ri07
$^{81}\text{Rb}-^{83}\text{Rb}_{,488} \text{ } ^{79}\text{Rb}_{,513}$	-529	26	-548	5	-0.3	U			P20	2.5	82Au01 Y
$^{81}\text{Rb}-^{83}\text{Rb}_{,325} \text{ } ^{80}\text{Rb}_{,675}$	-1054	27	-1040	5	0.2	U			P20	2.5	82Au01 Y
$^{82}\text{Rb}-^{83}\text{Rb}_{,659} \text{ } ^{80}\text{Rb}_{,342}$	627	24	604	3	-0.4	U			P20	2.5	82Au01 Y
$^{82}\text{Rb}-^{83}\text{Rb}_{,494} \text{ } ^{81}\text{Rb}_{,506}$	1098	23	1054	4	-0.8	U			P21	2.5	82Au01 Y
$^{82}\text{Ge}(\text{d,p})^{83}\text{Ge}$	1470	70	1408	3	-0.9	U			NDm		05Th03
$^{82}\text{Se}(\text{d,p})^{83}\text{Se}$	3593.4	3.0				2			NDm		78Mo12
$^{82}\text{Se}(\text{ } ^3\text{He,d})^{83}\text{Br}$	3207.4	5.6	3215	4	1.4	1	46	46 $^{83}\text{Br}$	NDm		83Zu01
$^{82}\text{Kr}(\text{ } ^3\text{He,d})^{83}\text{Rb}$	288	10	274.3	2.3	-1.4	U			Phi		87St11
$^{83}\text{Zr}(\text{ep})^{82}\text{Sr}$	2750	100	2809	9	0.6	U					83Ha06
$^{83}\text{As}(\beta^-)^{83}\text{Se}$	5460	220	5671	4	1.0	U			Stu		77A117
$^{83}\text{Se}(\beta^-)^{83}\text{Br}$	3610	40	3673	5	1.6	U					67Ma35
	3681	20			-0.4	U					68Sc10 *
	982	10	977	4	-0.5	-					51Du03 *
$^{83}\text{Br}(\beta^-)^{83}\text{Kr}$	967	15			0.7	U					63Pa09 *
	966	6			1.8	-					69Ph03 *
	ave.	970	5		1.3	1	54	54 $^{83}\text{Br}$			average
$^{83}\text{Rb}(\epsilon)^{83}\text{Kr}$	750	20	920.0	2.3	8.5	B					70Go45 *
$^{83}\text{Sr}(\beta^+)^{83}\text{Rb}$	2264	10	2273	6	0.9	1	41	41 $^{83}\text{Sr}$			68Et01 *
$^{83}\text{Y}(\beta^+)^{83}\text{Sr}$	4509	85	4592	20	1.0	U					81Li12 *
	4455	50			2.7	B			BNL		82De36 *
$^{83}\text{Zr}(\beta^+)^{83}\text{Y}$	5868	85	6294	20	5.0	B			Ors		82De36 *
$^{83}\text{Nb}(\beta^+)^{83}\text{Zr}$	7500	300	8360	150	2.9	B					88Ku14
$^{*83}\text{Y} \text{O}-^{98}\text{Mo}_{1,010}$	$D_M=12973.8(5.9) \mu\text{V}$ for mixture gs+m at 62.04(0.10) keV; $M-A=-72172.9(5.8) \text{keV}$										
$^{*83}\text{Se}(\beta^-)^{83}\text{Br}$	$Q_{\beta^-}=3910(20) \text{ from } ^{83}\text{Se}^m \text{ at } 228.92 \text{ keV}$										
$^{*83}\text{Br}(\beta^-)^{83}\text{Kr}$	$E_{\beta^-}=940(10) \text{ } 925(15) \text{ } 924(6) \text{ respectively, to } ^{83}\text{Kr}^n \text{ at } 41.5575 \text{ keV}$										
$^{*83}\text{Rb}(\epsilon)^{83}\text{Kr}$	$\text{LK}=0.132(0.002) \text{ to } 5/2^- \text{ level at } 561.9585, \text{ recalculated } Q$										
$^{*83}\text{Sr}(\beta^+)^{83}\text{Rb}$	$E_{\beta^+}=1227(8) \text{ } 24\% \text{ to ground state, } 20\% \text{ to } ^{83}\text{Rb}^m \text{ at } 42.0780, \text{ and other } E_{\beta^+}$										
$^{*83}\text{Y}(\beta^+)^{83}\text{Sr}$	$E_{\beta^+}=2868(85) \text{ from } ^{83}\text{Y}^m \text{ at } 62.04 \text{ keV to } (3/2^-) \text{ level at } 681.11 \text{ keV}$										
$^{*83}\text{Y}(\beta^+)^{83}\text{Sr}$	$E_{\beta^+}=3353(50) \text{ to } 9/2^+ \text{ level at } 35.47 \text{ keV}$										
	and $E_{\beta^+}=2941(84) \text{ from } ^{83}\text{Y}^m \text{ at } 62.04 \text{ to } (3/2^-) \text{ level at } 681.11 \text{ keV}$										
$^{*83}\text{Zr}(\beta^+)^{83}\text{Y}$	$Q_{\beta^+}=5806(85) \text{ to } ^{83}\text{Y}^m \text{ at } 62.04 \text{ keV}$										
$^{*83}\text{Zr}(\beta^+)^{83}\text{Y}$	Recalculated value 5802(50) of reference not accepted										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{84}\text{Ge}-u$	-62270	430	-62425	3	-0.4	U			OR1	1.0	06Ha62	
$^{84}\text{As}-u$	-70530	320	-70697	3	-0.5	U			OR1	1.0	06Ha62 *	
	-70710	140			0.0	U			GT2	2.5	08Su19 *	
$\text{C}_6 \text{H}_{12}-^{84}\text{Kr}$	182399.4	2.5	182402.658	0.004	0.5	U			M15	2.5	63Ri07	
	182392	6			1.2	U			R11	1.5	78Di09	
$\text{C}_5 \text{N} \text{H}_{10}-^{84}\text{Kr}$	169819	18	169826.598	0.004	0.3	U			R11	1.5	78Di09	
	169819	13			0.4	U			R11	1.5	78Di09	
$\text{C}_5 \text{O} \text{H}_8-^{84}\text{Kr}$	146010	20	146017.149	0.004	0.2	U			R11	1.5	78Di09	
$\text{C}_4 \text{ }^{13}\text{C} \text{O} \text{H}_7-^{84}\text{Kr}$	141543	18	141546.952	0.004	0.1	U			R11	1.5	78Di09	
$^{84}\text{Kr}-\text{N}_6$	-106946.3154	0.0152	-106946.298	0.004	1.1	o			FS1	1.0	05Sh38 *	
	-106946.2971	0.0086			-0.1	1	22	21	$^{84}\text{Kr}$	FS1	1.0	09Re03
$^{84}\text{Kr} \text{H}-^{85}\text{Rb}$	7515	18	7533.023	0.004	0.7	U			R11	1.5	78Di09	
$\text{C}_6 \text{H}_{12}-^{84}\text{Sr}$	180470.8	2.6	180481.3	1.3	1.6	U			M15	2.5	63Ri07	
$^{84}\text{Y} \text{O}-^{97}\text{Mo}_{1.031}$	12483.5	5.1	12482	5	-0.2	1	82	82	$^{84}\text{Y}$	JY1	1.0	08We10
$^{84}\text{Zr} \text{O}-^{98}\text{Mo}_{1.020}$	14728.6	5.9				2			JY1	1.0	06Ka48	
$^{84}\text{Nb}-u$	-65721	14				2			LZ1	1.0	16Xi.A	
$^{84}\text{Ge}-^{88}\text{Rb}_{.955}$	22268.7	3.4				2			JY1	1.0	08Ha23	
$^{84}\text{As}-^{88}\text{Rb}_{.955}$	13996.9	3.4				2			JY1	1.0	08Ha23	
$^{84}\text{Se}-^{88}\text{Rb}_{.955}$	3160.4	2.1	3160.4	2.1	0.0	1	100	100	$^{84}\text{Se}$	JY1	1.0	08Ha23
$^{82}\text{Se} \text{H}_2-^{84}\text{Kr}$	20834	16	20851.9	0.5	0.7	U			R11	1.5	78Di09	
$^{84}\text{Kr}-^{86}\text{Kr}_{.977}$	-1168.3	1.0	-1168.8534	0.0022	-0.6	U			MS1	1.0	06Ri15	
$^{83}\text{Kr} \text{H}-^{84}\text{Kr}$	10465	16	10453.822	0.009	-0.5	U			R11	1.5	78Di09	
$^{84}\text{Kr}-^{85}\text{Rb}_{.988}$	-1349.4	1.5	-1350.534	0.004	-0.8	U			MA8	1.0	06De36	
$^{84}\text{Rb}-^{85}\text{Rb}_{.988}$	1536	8	1527.0	2.4	-1.1	U			MA2	1.0	94Oi01	
$^{84}\text{Sr}-^{85}\text{Rb}_{.988}$	570.9	1.5	570.9	1.3	0.0	-			MA8	1.0	07Ke09	
	572.1	4.3			-0.3	-			SH1	1.0	11Ha08	
ave.	571.0	1.4			-0.1	1	89	89	$^{84}\text{Sr}$		average	
$^{84}\text{Kr}-^{80}\text{Se} \text{H}_3$	-28505	48	-28499.2	1.0	0.1	U			R11	1.5	78Di09	
$^{84}\text{Kr}-^{83}\text{Kr}$	-2628	12	-2628.790	0.009	0.0	U			M15	2.5	63Ri07	
	-2646	30			0.4	U			R11	1.5	78Di09	
$^{84}\text{Kr}-^{40}\text{Ar}_2$	-13268.5136	0.0171	-13268.519	0.006	-0.3	1	13	7	$^{40}\text{Ar}$	FS1	1.0	05Sh38 *
$\text{C}_2 \text{O}_4-^{84}\text{Kr}$	68160.7359	0.0205	68160.750	0.004	0.7	o			FS1	1.0	05Sh38 *	
	68160.7516	0.0131			-0.1	1	10	9	$^{84}\text{Kr}$	FS1	1.0	09Re03
$^{82}\text{Se}(\text{t,p})^{84}\text{Se}$	6016	15	6014.7	2.0	-0.1	U			LAl		74Kn02	
$^{84}\text{Sr}(\text{p,t})^{82}\text{Sr}$	-12310	10	-12300	6	1.0	1	36	35	$^{82}\text{Sr}$	Oak		73Ba56
	-12295	24			-0.2	U			Win		74De31 *	
$^{83}\text{Kr}(\text{n},\gamma)^{84}\text{Kr}$	10519.5	1.8	10520.019	0.008	0.3	U					72Ma42 Z	
	10520.6	0.3			-1.9	U			Bdn		06Fi.A	
$^{84}\text{Sr}(\text{d,t})^{83}\text{Sr}$	-5720	30	-5666	7	1.8	U					70Be24 *	
$^{84}\text{As}(\beta^-)^{84}\text{Se}$	7195	200	10094	4	14.5	F			Trs		94Gi07 *	
	9120	880			1.1	U					96WaZX	
$^{84}\text{Se}(\beta^-)^{84}\text{Br}$	1818	50	1835	26	0.3	1	27	26	$^{84}\text{Br}$		68Re12 *	
	1808	100			0.3	U					70Ei02 *	
$^{84}\text{Br}(\beta^-)^{84}\text{Kr}$	4650	30	4656	26	0.2	1	74	74	$^{84}\text{Br}$		70Ha21 *	
$^{84}\text{Br}^m(\beta^-)^{84}\text{Kr}$	4970	100				2					70Ha21 *	
$^{84}\text{Rb}(\beta^+)^{84}\text{Kr}$	2679	3	2680.4	2.2	0.5	-					64La03	
	2682	5			-0.3	-					71Bo01 *	
$^{84}\text{Rb}(\text{n,p})^{84}\text{Kr}$	3450	10	3462.7	2.2	1.3	U			ILL		76An05	
$^{84}\text{Rb}(\beta^+)^{84}\text{Kr}$	ave. 2679.8	2.6	2680.4	2.2	0.2	1	73	73	$^{84}\text{Rb}$		average	
$^{84}\text{Rb}(\beta^-)^{84}\text{Sr}$	892	4	890.6	2.3	-0.3	1	34	27	$^{84}\text{Rb}$		71Bo01 *	
$^{84}\text{Y}(\beta^+)^{84}\text{Sr}$	6950	30	6755	4	-6.5	F					70Va.A *	
	6750	10			0.5	1	19	18	$^{84}\text{Y}$		70Re.A *	
	6423	135			2.5	U			BNL		81Li12 *	
	6408	124			2.8	U			Ors		82De36 *	
$^{84}\text{Nb}(\beta^+)^{84}\text{Zr}$	7200	300	10203	14	10.0	B					96Sh27	
* $^{84}\text{As}-u$	Erroneously reported as -75700(300) keV in the publication											
* $^{84}\text{As}-u$	$M - A = -65869(119)$ keV for mixture gs+m at 0#(100#) keV											
* $^{84}\text{Kr}-\text{N}_6$	Corrected in reference of same group											
* $^{84}\text{Kr}-^{40}\text{Ar}_2$	Corrected in reference of same group											
* $\text{C}_2 \text{O}_4-^{84}\text{Kr}$	Corrected in reference of same group											
* $^{84}\text{Sr}(\text{p,t})^{82}\text{Sr}$	Original error 12; added systematic error 21 keV											
* $^{84}\text{Sr}(\text{d,t})^{83}\text{Sr}$	$Q = -5755(30)$ to $9/2^+$ level at 35.47 keV											
	Ens153 **											

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
* <sup>84</sup> As( $\beta^-$ ) <sup>84</sup> Se	F: observed ( $\beta^-$ n) decay implies $Q_{\beta^-} > 8681(15)$ keV										93Ru01	**
* <sup>84</sup> Se( $\beta^-$ ) <sup>84</sup> Br	$E_{\beta^-} = 1410(50)$ 1400(100) respectively, to $1^+$ level at 408.2 keV										Ens09a	**
* <sup>84</sup> Br( $\beta^-$ ) <sup>84</sup> Kr	$E_{\beta^-} = 4626(15), 3810(50), 2700(50)$ to ground state, 881.615 $2^+$ , 1897.784 $0^+$										Ens09a	**
* <sup>84</sup> Br <sup>m</sup> ( $\beta^-$ ) <sup>84</sup> Kr	$E_{\beta^-} = 2200(100)$ to $5^-$ level at 2770.94 keV										Ens09a	**
* <sup>84</sup> Rb( $\beta^+$ ) <sup>84</sup> Kr	Original error increased: authors report $E(2^+) = 877.2(1.5)$ while $E(2^+) = 881.615(0.003)$ keV; see also <sup>74</sup> As( $\beta^+$ )										AHW	**
* <sup>84</sup> Rb( $\beta^-$ ) <sup>84</sup> Sr	Originally 891.8(2.0), error increased, see <sup>84</sup> Rb( $\beta^+$ )										Ens09a	**
* <sup>84</sup> Y( $\beta^+$ ) <sup>84</sup> Sr	F: possibly added with $e^+e^-$										AHW	**
* <sup>84</sup> Y( $\beta^+$ ) <sup>84</sup> Sr	$E_{\beta^+} = 1641(10)$ and 2242(17) to levels at 4062 and 3511 keV										70Re.A	**
* <sup>84</sup> Y( $\beta^+$ ) <sup>84</sup> Sr	$Q_{\beta^+} = 6409(170)$ , and 6499(135) from <sup>84</sup> Y <sup>m</sup> at 67 keV										00Do10	**
* <sup>84</sup> Y( $\beta^+$ ) <sup>84</sup> Sr	$Q_{\beta^+} = 6475(124)$ from <sup>84</sup> Y <sup>m</sup> at 67 keV										00Do10	**
<sup>85</sup> Ge-u	-57220	540	-57030	4	0.4	U			OR1	1.0	06Ha62	
<sup>85</sup> As-u	-68095	225	-67836	3	0.8	o			GT1	1.5	04Ma.A	
	-67887	129			0.2	U			GT2	2.5	08Su19	
C <sub>6</sub> H <sub>13</sub> - <sup>85</sup> Rb	189927.6	3.9	189935.682	0.005	0.8	U			M15	2.5	63Ri07	
	189930	15			0.3	U			R11	1.5	78Di09	
C <sub>4</sub> N O H <sub>7</sub> - <sup>85</sup> Rb	140985	18	140974.112	0.005	-0.4	U			R11	1.5	78Di09	
<sup>85</sup> Rb- <sup>39</sup> K <sub>2,179</sub>	-9124.6	2.7	-9126.700	0.012	-0.8	U			MA8	1.0	09Na.A	
<sup>85</sup> Rb- <sup>120</sup> Sn <sub>708</sub>	-18970.8	2.2	-18969.2	0.7	0.7	U			JY1	1.0	11Ha48	
<sup>85</sup> Y-u	-83559	31	-83567	20	-0.3	2			GS2	1.0	05Li24 *	
<sup>85</sup> Zr O- <sup>98</sup> Mo <sub>1,031</sub>	13886.7	6.9				2			JY1	1.0	06Ka48	
<sup>85</sup> Nb O- <sup>98</sup> Mo <sub>1,031</sub>	21246	26	21289	4	1.7	U			JY1	1.0	06Ka48 *	
<sup>85</sup> Ge- <sup>88</sup> Rb <sub>966</sub>	28638.8	4.0				2			JY1	1.0	08Ha23	
<sup>85</sup> As- <sup>88</sup> Rb <sub>966</sub>	17832.8	3.3				2			JY1	1.0	08Ha23	
<sup>85</sup> Se- <sup>88</sup> Rb <sub>966</sub>	7929.9	2.8				2			JY1	1.0	08Ha23	
<sup>85</sup> Br- <sup>88</sup> Rb <sub>966</sub>	1314.9	3.3				2			JY1	1.0	07Ra23	
<sup>85</sup> Nb- <sup>85</sup> Rb	17056.1	4.4				2			SH1	1.0	11Ha08 *	
<sup>85</sup> Mo- <sup>85</sup> Rb	26471	17				2			SH1	1.0	11Ha08	
C <sub>6</sub> H <sub>14</sub> - <sup>85</sup> Rb	197760.706	0.014	197760.714	0.005	0.6	U			MI2	1.0	99Br47	
<sup>85</sup> Rb-C <sub>6</sub> H <sub>12</sub>	-182110.662	0.024	-182110.649	0.005	0.5	U			MI2	1.0	99Br47	
<sup>85</sup> Rb- <sup>84</sup> Kr	300	32	292.009	0.004	-0.2	U			R11	1.5	78Di09	
	292.0121	0.0064			-0.5	1	39	34 <sup>85</sup> Rb	FS1	1.0	10Mo30	
<sup>83</sup> Rb- <sup>85</sup> Rb <sub>488</sub> <sup>81</sup> Rb <sub>512</sub>	-351	22	-339	3	0.2	U			P21	2.5	82Au01 Y	
<sup>84</sup> Kr(d,p) <sup>85</sup> Kr	4895	8	4887.7	2.0	-0.9	U			MIT		63Ho.A	
<sup>85</sup> Rb( $\gamma$ ,n) <sup>84</sup> Rb	-10650	80	-10479.7	2.2	2.1	U			Phi		60Ge01	
<sup>85</sup> Rb(p,d) <sup>84</sup> Rb	-8275	6	-8255.1	2.2	3.3	B			Bld		78Sh11	
<sup>84</sup> Sr(d,p) <sup>85</sup> Sr	6303	8	6300	3	-0.3	1	14	12 <sup>85</sup> Sr			71Mo02	
<sup>85</sup> Mo( $\epsilon$ p) <sup>84</sup> Zr	5100	200	6623	17	7.6	B					99Hu05	
<sup>85</sup> Se( $\beta^-$ ) <sup>85</sup> Br	6185	90	6162	4	-0.3	o			Bwg		87Gr.A	
	6182	23			-0.9	U			Bwg		92Gr.A	
<sup>85</sup> Br( $\beta^-$ ) <sup>85</sup> Kr	2870	19	2905	4	1.8	U			Stu		79Al05	
<sup>85</sup> Kr( $\beta^-$ ) <sup>85</sup> Rb	687	2				2					70Wo08	
<sup>85</sup> Sr( $\epsilon$ ) <sup>85</sup> Rb	1007	30	1064.1	2.8	1.9	U					69Mc05	
<sup>85</sup> Rb(p,n) <sup>85</sup> Sr	-1890	30	-1846.4	2.8	1.5	U			BNL		58El44	
<sup>85</sup> Rb( <sup>3</sup> He,t) <sup>85</sup> Sr	-1083	3	-1082.6	2.8	0.1	1	88	88 <sup>85</sup> Sr	Pri		82Ko06	
<sup>85</sup> Y( $\beta^+$ ) <sup>85</sup> Sr	3255	25	3261	19	0.2	R					63Do07 *	
<sup>85</sup> Zr( $\beta^+$ ) <sup>85</sup> Y	4693	99	4667	20	-0.3	U			Ors		82De36	
<sup>85</sup> Nb( $\beta^+$ ) <sup>85</sup> Zr	6000	200	6896	8	4.5	F					88Ku14 *	
* <sup>85</sup> Y-u	$M - A = -77824(28)$ keV for mixture gs+m at 19.68 keV										Nub16c	**
* <sup>85</sup> Nb O- <sup>98</sup> Mo <sub>1,031</sub>	$D_M = 21292.2(6.9)$ $\mu$ u for mixture gs+m at 150#80 keV; $M - A = -66274.8(6.6)$ keV										Nub16b	**
* <sup>85</sup> Nb- <sup>85</sup> Rb	Misprint in publication 1.000200869 (not 1.00200869)										GAu	**
* <sup>85</sup> Y( $\beta^+$ ) <sup>85</sup> Sr	$E_{\beta^+} = 1540(20)$ to $3/2^-$ level at 743.25 keV										Ens148	**
*	and $E_{\beta^+} = 2240(10)$ from <sup>85</sup> Y <sup>m</sup> at 19.68 (conflicting $\rightarrow$ outer error used)										Nub16b	**
* <sup>85</sup> Nb( $\beta^+$ ) <sup>85</sup> Zr	F: see discussion of this result in reference										06Ka48	**
*	$Q_{\beta^+} = 6100(200)$ in text p.268 and in Table 1										GAu	**

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{86}\text{Ge}-u$	-54750	540	-53030	470	3.2	B			OR1	1.0	06Ha62
	-53033	188				2			GT3	2.5	16Kn03
$^{86}\text{As}-u$	-63586	247	-63298	4	0.8	o			GT1	1.5	04Ma.A
	-63189	129			-0.3	U			GT2	2.5	08Su19
$^{86}\text{Se}-u$	-75702	128	-75688.3	2.7	0.0	U			GT2	2.5	08Su19
$\text{C}_6 \text{H}_{14}-^{86}\text{Kr}$	198936.7	2.7	198939.825	0.004	0.5	U			M15	2.5	63Ri07
	198933	15			0.3	U			R11	1.5	78Di09
$\text{C}_5 \text{N} \text{H}_{12}-^{86}\text{Kr}$	186366	20	186363.765	0.004	-0.1	U			R11	1.5	78Di09
$^{86}\text{Kr}-u$	-89389.271	0.110	-89389.374	0.004	-0.9	U			ST2	1.0	02Bf02
$^{86}\text{Kr}-^{120}\text{Sn}_{.717}$	-19269.6	2.2	-19268.1	0.7	0.7	U			JY1	1.0	11Ha48
$\text{C}_6 \text{H}_{14}-^{86}\text{Sr}$	200264.9	3.6	200289.725	0.006	2.8	U			M15	2.5	63Ri07
$^{86}\text{Y}-u$	-85019	75	-85114	15	-1.3	U			GS2	1.0	05Li24 *
$^{86}\text{Zr} \text{O}-^{98}\text{Mo}_{1.041}$	9692.8	6.9	9686	4	-0.9	1	31	31 $^{86}\text{Zr}$	JY1	1.0	06Ka48
$^{86}\text{Nb} \text{O}-^{98}\text{Mo}_{1.041}$	19171.0	5.9				2			JY1	1.0	06Ka48
$^{86}\text{As}-^{88}\text{Rb}_{.977}$	23346.2	3.7				2			JY1	1.0	08Ha23
$^{86}\text{Se}-^{88}\text{Rb}_{.977}$	10956.4	2.7				2			JY1	1.0	08Ha23
$^{86}\text{Br}-^{88}\text{Rb}_{.977}$	5450.1	3.3				2			JY1	1.0	07Ra23
$^{86}\text{Kr}-^{84}\text{Kr}_{1.024}$	1236.9544	0.0042	1236.9565	0.0023	0.5	1	30	20 $^{84}\text{Kr}$	FS1	1.0	12Ra34
$^{86}\text{Sr}-^{84}\text{Kr}_{1.024}$	-112.9463	0.0054	-112.944	0.004	0.5	1	59	54 $^{86}\text{Sr}$	FS1	1.0	12Ra34
$^{86}\text{Kr}-^{85}\text{Rb}_{1.012}$	-120.3	3.6	-120.591	0.004	-0.1	U			MA8	1.0	06Ro11
	-119.1	1.6			-0.9	U			MA8	1.0	06De36
$^{86}\text{Rb}-^{85}\text{Rb}_{1.012}$	-54	88			-0.8	U			MA9	1.0	10Na13 *
	441	9	436.23	0.21	-0.5	U			MA2	1.0	94Oo1
$^{86}\text{Sr}^{19}\text{F}-^{85}\text{Rb}_{1.235}$	16608	13	16603.565	0.006	-0.3	U			MA8	1.0	05Si34
$^{86}\text{Zr}-^{85}\text{Rb}_{1.012}$	5562.7	4.6	5566	4	0.6	1	69	69 $^{86}\text{Zr}$	SH1	1.0	11Ha08
$^{86}\text{Mo}-^{85}\text{Rb}_{1.012}$	20443.6	4.0				2			SH1	1.0	11Ha08
$^{86}\text{Sr}-^{86}\text{Kr}$	-1349.8965	0.0060	-1349.900	0.004	-0.6	1	49	46 $^{86}\text{Sr}$	FS1	1.0	12Ra34
$^{86}\text{Kr}-^{85}\text{Rb}$	-1206	42	-1179.111	0.004	0.4	U			R11	1.5	78Di09
	-1179.1083	0.0071			-0.4	-			FS1	1.0	10Mo30 *
ave.	-1179.1109	0.0059			-0.1	-			FS1	1.0	10Mo30 *
	-1179.110	0.005			-0.3	1	69	66 $^{85}\text{Rb}$	average		
$^{86}\text{Kr}-\text{N}_6$	-107833.3986	0.0074	-107833.400	0.004	-0.3	1	28	27 $^{86}\text{Kr}$	FS1	1.0	09Re03
$\text{C}_2 \text{O}_4-^{86}\text{Kr}$	69047.8337	0.0155	69047.852	0.004	1.2	o			FS1	1.0	05Sh38 *
	69047.8440	0.0113			0.7	1	12	12 $^{86}\text{Kr}$	FS1	1.0	09Re03
$^{86}\text{Kr}-^{84}\text{Kr}$	-908	32	-887.1024	0.0023	0.4	U			R11	1.5	78Di09
	-887.1041	0.0125			0.1	o			FS1	1.0	05Sh38 *
ave.	-887.1080	0.0069			0.8	-			FS1	1.0	09Re03 *
	-887.0954	0.0060			-1.2	-			FS1	1.0	10Mo30 *
$^{86}\text{Sr}(\text{p,t})^{84}\text{Sr}$	-11535	10	-11534.4	1.2	0.1	U			Oak		73Ba56
$^{85}\text{Rb}(\text{n},\gamma)^{86}\text{Rb}$	8651.1	1.0	8650.98	0.20	-0.1	U					69Da15 Z
	8651.3	1.5			-0.2	U					70Or.A
$^{85}\text{Rb}(\text{d,p})^{86}\text{Rb}$	8650.98	0.20				2			Bdn		06Fi.A
	6433	10	6426.41	0.20	-0.7	U			Tal		69Da15
$^{86}\text{Se}(\beta^-)^{86}\text{Br}$	5095	100	5129	4	0.3	o			Bwg		87Gr.A
	5099	11			2.7	C			Bwg		92Gr.A
$^{86}\text{Br}(\beta^-)^{86}\text{Kr}$	7620	60	7633	3	0.2	U			Stu		79Al05
	7626	11			0.7	U			Bwg		92Gr.A
$^{86}\text{Rb}(\beta^-)^{86}\text{Sr}$	1774	5	1776.10	0.20	0.4	U					64Da16
	1770	3			2.0	U					66An10
ave.	1779.2	2.5			-1.2	U					75Be21
	1775	3			0.4	U					75Ra09
$^{86}\text{Y}(\beta^+)^{86}\text{Sr}$	5220	20	5240	14	1.0	2					62Ya01 *
	5260	20			-1.0	2					65Va02 *
$^{86}\text{Nb}(\beta^+)^{86}\text{Zr}$	7978	80	8835	7	10.7	F					82De43 *
$^{86}\text{Mo}(\beta^+)^{86}\text{Nb}$	5019	430	5024	7	0.0	U					94Sh07 *
$^{86}\text{Y}-u$	$M - A = -79086(29)$ keV for mixture gs+m at 218.21 keV										
$^{86}\text{Kr}-^{85}\text{Rb}_{1.012}$	Typo in original paper, ratio should read 1.011 763 90(1 03)										
$^{86}\text{Kr}-^{85}\text{Rb}$	Different charge states : $3^+$ and $2^+$										
$\text{C}_2 \text{O}_4-^{86}\text{Kr}$	Corrected in reference of same group										
$^{86}\text{Kr}-^{84}\text{Kr}$	Corrected in reference of same group										
$^{86}\text{Kr}-^{84}\text{Kr}$	Different charge states in these two results: $3^+$ and $2^+$										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* $^{86}\text{Y}(\beta^+)^{86}\text{Sr}$	$E_{\beta^+}=2019(20)$		$1960(20)$ respectively, to $2229.81$ $4^+$ level, and other $E_{\beta^+}$								Ens14c **
* $^{86}\text{Nb}(\beta^+)^{86}\text{Zr}$	F : see discussion of this result in reference										
* $^{86}\text{Mo}(\beta^+)^{86}\text{Nb}$	$E_{\beta^+}=3900(400)$		to $(1^+)$ level at $97.1$ keV								94Sh07 **
$^{87}\text{Se}-u$	-71357	128	-71311.4	2.4	0.1	U			GT2	2.5	08Su19
$^{87}\text{Kr}-u$	-86622	30	-86645.24	0.26	-0.8	U			GS2	1.0	05Li24
$\text{C}_4 \text{H}_7 \text{O}_2 - ^{87}\text{Rb}$	135417.8	2.7	135423.934	0.006	0.9	U			M15	2.5	63Ri07
$\text{C}_5 \text{}^{13}\text{C} \text{H}_{14} - ^{87}\text{Rb}$	203767	15	203724.755	0.006	-1.9	U			R11	1.5	78Di09
$\text{C}_4 \text{O} \text{N} \text{H}_9 - ^{87}\text{Rb}$	159277	15	159233.383	0.006	-1.9	U			R11	1.5	78Di09
$\text{C}_3 \text{}^{13}\text{C} \text{O} \text{N} \text{H}_8 - ^{87}\text{Rb}$	154809	25	154763.186	0.006	-1.2	U			R11	1.5	78Di09
$\text{C}_4 \text{H}_7 \text{O}_2 - ^{87}\text{Sr}$	135722.2	3.5	135726.969	0.005	0.5	U			M15	2.5	63Ri07
$^{87}\text{Y}-u$	-89153	30	-89123.9	1.2	1.0	U			GS2	1.0	03Li.A *
$^{87}\text{Zr}-u$	-85222	30	-85183	4	1.3	U			GS2	1.0	05Li24
$^{87}\text{Zr} \text{O} - ^{97}\text{Mo}_{1.062}$	9543.3	5.2	9542	4	-0.2	1	73	$^{87}\text{Zr}$	JY1	1.0	08We10
$^{87}\text{Nb}_{1.069} - \text{C}_7 \text{H}_9$	-155224	30	-155205	8	0.6	U			CP1	1.0	11Fa10
$^{87}\text{Nb} \text{O} - ^{98}\text{Mo}_{1.051}$	15027.9	7.3				2			JY1	1.0	06Ka48 *
$^{87}\text{Mo}_{1.069} - \text{C}_7 \text{H}_9$	-147186.1	4.8	-147184	3	0.5	1	47	$^{87}\text{Mo}$	CP1	1.0	11Fa10
$^{87}\text{Sr} - ^{84}\text{Kr}_{1.036}$	565.8516	0.0059	565.850	0.004	-0.2	1	47	$^{87}\text{Sr}$	FS1	1.0	12Ra34
$^{87}\text{Kr} - ^{85}\text{Rb}_{1.024}$	3683.0	2.9	3682.07	0.26	-0.3	U			MA8	1.0	06De36
	3684.1	4.7			-0.4	U			MA8	1.0	10Na13
$^{87}\text{Rb} - ^{85}\text{Rb}_{1.024}$	-490	9	-492.156	0.007	-0.2	U			MA2	1.0	94O101
	-493.0	2.7			0.3	U			MA8	1.0	06De36
	-492.33	0.80			0.2	U			MA8	1.0	07Ke09
	-492.4	1.4			0.2	U			MA8	1.0	09Na.A
	-492.04	0.87			-0.1	U			MA8	1.0	11He10
$^{87}\text{Sr} - ^{85}\text{Rb}_{1.024}$	-780	9	-795.191	0.005	-1.7	U			MA2	1.0	94O101
$^{87}\text{Mo} - ^{85}\text{Rb}_{1.024}$	18525.6	4.2	18524	3	-0.5	1	53	$^{87}\text{Mo}$	SH1	1.0	11Ha08
$^{87}\text{Tc} - ^{85}\text{Rb}_{1.024}$	28394.5	4.5				2			SH1	1.0	11Ha08 *
$^{87}\text{As} - ^{88}\text{Rb}_{.989}$	28000.6	3.2				2			JY1	1.0	08Ha23
$^{87}\text{Se} - ^{88}\text{Rb}_{.989}$	16397.5	2.4				2			JY1	1.0	08Ha23
$^{87}\text{Br} - ^{88}\text{Rb}_{.989}$	8382.9	3.4				2			JY1	1.0	07Ra23
$^{86}\text{Kr} \text{H} - ^{87}\text{Rb}$	9309	16	9255.127	0.006	-2.2	U			R11	1.5	78Di09
$^{87}\text{Sr} - ^{86}\text{Kr}_{1.012}$	-660.4616	0.0050	-660.461	0.004	0.2	1	62	$^{87}\text{Sr}$	FS1	1.0	12Ra34
$\text{C}_6 \text{H}_{16} - ^{87}\text{Rb}$	216019.966	0.023	216019.985	0.007	0.8	U			MI2	1.0	99Br47
$^{87}\text{Rb} - \text{C}_6 \text{H}_{14}$	-200369.931	0.015	-200369.920	0.007	0.7	1	19	$^{87}\text{Rb}$	MI2	1.0	99Br47
$^{87}\text{Rb} - ^{86}\text{Kr}$	-1477	30	-1430.095	0.006	1.0	U			R11	1.5	78Di09
	-1430.0932	0.0059			-0.3	1	87	$^{87}\text{Rb}$	FS1	1.0	10Mo30
$^{87}\text{Sr} - ^{86}\text{Sr}$	-382	12	-383.230	0.006	0.0	U			M15	2.5	63Ri07
$^{87}\text{Rb} - ^{85}\text{Rb}$	-2620	35	-2609.206	0.007	0.2	U			R11	1.5	78Di09
$^{85}\text{Rb} - ^{87}\text{Rb}_{.489} \text{}^{83}\text{Rb}_{.512}$	-310	30	-314.8	1.2	-0.1	U			P21	2.5	82Au01
$^{84}\text{Rb} - ^{87}\text{Rb}_{.241} \text{}^{83}\text{Rb}_{.759}$	850	72	643.7	2.8	-1.1	U			P21	2.5	82Au01 *
$^{87}\text{Sr}(\text{p,t})^{85}\text{Sr}$	-11440	10	-11437.6	2.8	0.2	U			Oak		73Ba56
$^{87}\text{Br}(\beta^- \text{n})^{86}\text{Kr}$	1335	25	1303	3	-1.3	U					84Kr.B
$^{86}\text{Kr}(\text{n},\gamma)^{87}\text{Kr}$	5515.04	0.6	5515.17	0.25	0.2	2					77Je03 Z
	5515.20	0.27			-0.1	2			Bdn		06Fi.A
$^{86}\text{Kr}(\text{d,p})^{87}\text{Kr}$	3286	8	3290.61	0.25	0.6	U			MIT		63Ho.A
$^{87}\text{Rb}(\gamma,\text{n})^{86}\text{Rb}$	-9990	70	-9922.11	0.20	1.0	U			Phi		60Ge01
$^{87}\text{Rb}(\text{d,t})^{86}\text{Rb}$	-3659	15	-3664.89	0.20	-0.4	U			Tal		69Da15
$^{86}\text{Sr}(\text{n},\gamma)^{87}\text{Sr}$	8428.12	0.17	8428.294	0.005	1.0	U			ILn		86Wi16 Z
	8428.17	0.17			0.7	U			Bdn		06Fi.A
$^{86}\text{Sr}(\text{d,p})^{87}\text{Sr}$	6203	8	6203.728	0.005	0.1	U					71Mo02
$^{86}\text{Sr}(\text{p},\gamma)^{87}\text{Y}$	5785.4	3.3	5784.3	1.1	-0.3	R					71Um03
$^{86}\text{Sr}(\text{}^3\text{He,d})^{87}\text{Y}$	346	15	290.8	1.1	-3.7	B			ANL		71Ma11
$^{87}\text{Mo}(\epsilon\text{p})^{86}\text{Zr}$	3700	300	3795	5	0.3	U					83Ha06
$^{87}\text{Se}(\beta^-)^{87}\text{Br}$	7250	150	7466	4	1.4	o			Bwg		87Gr.A
	7275	35			5.4	B			Bwg		92Gr.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{87}\text{Br}(\beta^-)^{87}\text{Kr}$	6830	120	6818	3	-0.1	U			Stu		79A105
	6750	150			0.5	o			Bwg		87Gr.B
	6855	25			-1.5	U			Bwg		92Gr.A
$^{87}\text{Kr}(\beta^-)^{87}\text{Rb}$	3888	7	3888.27	0.25	0.0	U					73Wo01
$^{87}\text{Rb}(\beta^-)^{87}\text{Sr}$	272	3	282.275	0.006	3.4	B					59Fl40
	274	3			2.8	B					61Be41
$^{87}\text{Rb}(^3\text{He,t})^{87}\text{Sr}-^{81}\text{Br}(^{81}\text{Kr})$	564.0	1.5	563.1	0.5	-0.6	1	10	9 $^{81}\text{Kr}$	Pri		82Ko06
$^{87}\text{Y}(\beta^+)^{87}\text{Sr}$	2190	50	1861.7	1.1	-6.6	B					67Mi13 *
	1791	40			1.8	U					69Zo04 *
$^{87}\text{Sr}(\text{p,n})^{87}\text{Y}$	-2644.2	1.2	-2644.0	1.1	0.1	2					71Um03 Z
$^{87}\text{Zr}(\beta^+)^{87}\text{Y}$	3663	40	3671	4	0.2	U					65Ba48 *
$^{87}\text{Nb}(\beta^+)^{87}\text{Zr}$	5165	60	5473	8	5.1	B					82De43 *
$^{87}\text{Mo}(\beta^+)^{87}\text{Nb}$	6382	308	6990	7	2.0	U					82De43 *
	6589	300			1.3	U					91Mi15 *
$^{*87}\text{Y}-\text{u}$	$M-A=-82665(28)$ keV for $^{87}\text{Y}^m$ at 380.82 keV										Nub16b **
$^{*87}\text{Nb O}-^{98}\text{Mo}_{1.051}$	$D_M=15030.0(6.9)$ $\mu\text{u}$ for mixture gs+m at 3.9(0.1) keV; $M-A=-73870.2(6.7)$ keV										Nub16b **
$^{*87}\text{Tc}-^{85}\text{Rb}_{1.024}$	Most probably the high-spin isomer										11Ha08 **
$^{*84}\text{Rb}-^{87}\text{Rb}_{.241}$ $^{83}\text{Rb}_{.759}$	$D_M=1080(40)$ keV corrected $-230(60)$ for mixture gs+m at 463.59 keV										Nub16b **
$^{*87}\text{Y}(\beta^+)^{87}\text{Sr}$	$E_{\beta^+}=780(50)$ to $^{87}\text{Sr}^m$ at 388.5287 keV										Nub16b **
$^{*87}\text{Y}(\beta^+)^{87}\text{Sr}$	$E_{\beta^+}=1150(40)$ from $^{87}\text{Y}^m$ at 380.82 keV										Nub16b **
$^{*87}\text{Zr}(\beta^+)^{87}\text{Y}$	$E_{\beta^+}=2260(40)$ to $^{87}\text{Y}^m$ at 380.82 keV										Nub16b **
$^{*87}\text{Nb}(\beta^+)^{87}\text{Zr}$	$Q_{\beta^+}=5169(60)$ from $^{87}\text{Nb}^m$ at 3.9(0.1) keV										Nub16b **
$^{*87}\text{Mo}(\beta^+)^{87}\text{Nb}$	$Q_{\beta^+}=6378(308)$ to $^{87}\text{Nb}^m$ at 3.9(0.1) keV										Nub16b **
$^{*87}\text{Mo}(\beta^+)^{87}\text{Nb}$	$E_{\beta^+}=5300(300)$ to $(7/2)^+$ level at 266.9 keV										Ens15a **
$^{88}\text{Se}-\text{u}$	-68555	129	-68583	4	-0.1	U			GT2	2.5	08Su19
$^{88}\text{Br}-\text{u}$	-75832	100	-75917	3	-0.3	o			GT2	2.5	08Kn.A
	-75823	129			-0.3	U			GT2	2.5	08Su19
$\text{C}_4 \text{H}_8 \text{O}_2 -^{88}\text{Sr}$	146789.1	4.7	146817.242	0.006	2.4	U			M15	2.5	63Ri07
$^{88}\text{Y}-\text{u}$	-90500	31	-90498.7	1.6	0.0	U			GS2	1.0	05Li24
$^{88}\text{Zr O}-^{98}\text{Mo}_{1.061}$	5502.3	6.9	5502	6	0.0	1	71	71 $^{88}\text{Zr}$	JY1	1.0	06Ka48
$^{88}\text{Nb O}-^{98}\text{Mo}_{1.061}$	13452	78	13510	60	0.7	1	66	66 $^{88}\text{Nb}$	JY1	1.0	06Ka48 *
$^{88}\text{Sr}-^{84}\text{Kr}_{1.048}$	-1637.3606	0.0069	-1637.366	0.005	-0.8	1	46	42 $^{88}\text{Sr}$	FS1	1.0	12Ra34
$^{88}\text{Kr}-^{85}\text{Rb}_{1.035}$	5745.5	2.8				2			MA8	1.0	06De36
$^{88}\text{Rb}-^{85}\text{Rb}_{1.035}$	2615	9	2613.21	0.17	-0.2	U			MA4	1.0	02Ra23
$^{88}\text{Sr}-^{85}\text{Rb}_{1.035}$	-3108	20	-3090.126	0.006	0.9	U			MA8	1.0	07Ke09
	-3088	11			-0.2	U			MA8	1.0	05Si34
$^{88}\text{Mo}-^{85}\text{Rb}_{1.035}$	13265.4	4.1				2			JY1	1.0	08We10
$^{88}\text{Tc}-^{85}\text{Rb}_{1.035}$	25080	160				2			JY1	1.0	08We10 *
$^{88}\text{Sr}-^{86}\text{Kr}_{1.023}$	-2942.4188	0.0059	-2942.415	0.005	0.7	1	61	58 $^{88}\text{Sr}$	FS1	1.0	12Ra34
$^{88}\text{Se}-^{88}\text{Rb}$	20101.9	3.6				2			JY1	1.0	08Ha23
$^{88}\text{Br}-^{88}\text{Rb}$	12767.7	3.4				2			JY1	1.0	07Ra23
$^{88}\text{Sr}-^{87}\text{Sr}$	-3260	12	-3265.241	0.006	-0.2	U			M15	2.5	63Ri07
$^{86}\text{Kr}(\text{t,p})^{88}\text{Kr}$	4091	15	4086.5	2.6	-0.3	U			LAl		76Fl02
$^{88}\text{Sr}(\text{p,t})^{86}\text{Sr}$	-11060	10	-11059.368	0.006	0.1	U			Oak		73Ba56
$^{87}\text{Rb}(\text{n},\gamma)^{88}\text{Rb}$	6082.52	0.16	6082.52	0.16	0.0	1	99	99 $^{88}\text{Rb}$	Bdn		06Fi.A
$^{87}\text{Rb}(\text{d,p})^{88}\text{Rb}$	3858	15	3857.96	0.16	0.0	U			Oak		71Ra17
	3837	20			1.0	U					71To05
$^{87}\text{Sr}(\text{n},\gamma)^{88}\text{Sr}$	11112.63	0.22	11112.869	0.006	1.1	U			ILn		87Wi15 Z
	11112.64	0.22			1.0	U			Bdn		06Fi.A
$^{87}\text{Sr}(\text{d,p})^{88}\text{Sr}$	8865	5	8888.303	0.005	4.7	B			MIT		68Co20
$^{88}\text{Se}(\beta^-)^{88}\text{Br}$	6854	31	6832	5	-0.7	U			Bwg		92Gr.A
$^{88}\text{Br}(\beta^-)^{88}\text{Kr}$	8970	130	8975	4	0.0	U			Stu		79A105
	8880	200			0.5	o			Bwg		87Gr.B
	8960	36			0.4	U			Bwg		92Gr.A



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{88}\text{Kr}(\beta^-)^{88}\text{Rb}$	2930	30	2917.7	2.6	-0.4	U			Trs		78Wo15
$^{88}\text{Rb}(\beta^-)^{88}\text{Sr}$	5310	60	5312.62	0.16	0.0	U			Trs		78Wo15
	5318	9			-0.6	U			Gsn		80De02 *
	5313	5			-0.1	U			Trs		82Br23
$^{88}\text{Y}(\beta^+)^{88}\text{Sr}$	3622.6	1.5				2					79An36 *
$^{88}\text{Sr}(\text{p,n})^{88}\text{Y}$	-4402	6	-4404.9	1.5	-0.5	U			Bar		61Sh23
	-4406	10			0.1	U			Tal		62Ne08
$^{88}\text{Nb}(\beta^+)^{88}\text{Zr}$	7550	100	7460	60	-0.9	1	35	34	$^{88}\text{Nb}$		84Ox01
$^{88}\text{Nb}^m(\beta^+)^{88}\text{Zr}$	7590	100				2					84Ox01
$^{88}\text{Tc}(\beta^+)^{88}\text{Mo}$	8600	1300	11010	150	1.9	U					96Od01
	7800	600			5.3	B					96Sh27
$^{88}\text{Nb O}-^{98}\text{Mo}_{1.061}$	$D_M=13527.5(6.9) \mu\text{u}$ for mixture gs+m at 140(120) keV; $M-A=-76150.9(6.7) \text{ keV}$										Nub16b **
$^{88}\text{Tc}-^{85}\text{Rb}_{1.035}$	Most probably the high-spin isomer										08We10 **
$^{88}\text{Tc}-^{85}\text{Rb}_{1.035}$	$D_M=25082.4(4.1) \mu\text{u}$ for mixture gs+m at 0#300 keV; $M-A=-61679.1(3.8) \text{ keV}$										Nub16b **
$^{88}\text{Rb}(\beta^-)^{88}\text{Sr}$	Original error 4 corrected in reference										94Ha.A **
$^{88}\text{Y}(\beta^+)^{88}\text{Sr}$	$E_{\beta^+}=764.6(1.5) \text{ to } 2^+ \text{ level at } 1836.090 \text{ keV}$										Ens141 **
$^{89}\text{Se}-\text{u}$	-63285	225	-63331	4	-0.1	o			GT1	1.5	04Ma.A
	-63291	129			-0.1	U			GT2	2.5	08Su19
$\text{C}_7 \text{H}_5-^{89}\text{Y}$	133247.0	3.4	133284.0	1.7	4.3	B			M15	2.5	63Ri07
$^{89}\text{Nb}-\text{u}$	-86588	34	-86555	25	1.0	-			GS2	1.0	05Li24 *
	ave.	-86582	29		1.0	1	78	78	$^{89}\text{Nb}$		average
$^{89}\text{Kr}-^{85}\text{Rb}_{1.047}$	10191.6	2.3				2			MA8	1.0	06De36
$^{89}\text{Rb}-^{85}\text{Rb}_{1.047}$	4628	9	4634	6	0.7	1	42	42	$^{89}\text{Rb}$	1.0	02Ra23
$^{89}\text{Mo}-^{85}\text{Rb}_{1.047}$	11824.3	4.2				2			JY1	1.0	08We10
$^{89}\text{Tc}-^{85}\text{Rb}_{1.047}$	20007	17	20005	4	-0.1	U			SH1	1.0	08We10
	20004.8	4.1				2			JY1	1.0	08We10
$^{89}\text{Se}-^{88}\text{Rb}_{1.011}$	26329.0	4.0				2			JY1	1.0	08Ha23
$^{89}\text{Br}-^{88}\text{Rb}_{1.011}$	16364.5	3.5				2			JY1	1.0	07Ra23
$^{88}\text{Rb}-^{89}\text{Rb}_{.494} \text{ } ^{87}\text{Rb}_{.506}$	545	23	563.4	2.7	0.3	U			P21	2.5	82Au01
$^{89}\text{Y}(\text{d},\alpha)^{87}\text{Sr}$	7889	15	7882.5	1.6	-0.4	U			Mtr		72Br13
$^{88}\text{Sr}(\text{n},\gamma)^{89}\text{Sr}$	6358.70	0.13	6358.72	0.09	0.1	-			ILn		89Wi05 Z
	6358.73	0.13			-0.1	-			Bdn		06Fi.A
$^{88}\text{Sr}(\text{d,p})^{89}\text{Sr}$	4133	5	4134.15	0.09	0.2	U			MIT		67Sp09
$^{88}\text{Sr}(\text{n},\gamma)^{89}\text{Sr}$	ave.	6358.71	0.09	6358.72	0.09	0.0	1	100	$^{89}\text{Sr}$		average
$^{88}\text{Sr}(\text{p},\gamma)^{89}\text{Y}$	7078	4	7075.7	1.6	-0.6	1	16	16	$^{89}\text{Y}$		75Be.B Z
$^{89}\text{Y}(\gamma,\text{n})^{88}\text{Y}$	-11540	40	-11480.7	2.2	1.5	U			Phi		63Ge02
$^{89}\text{Br}(\beta^-)^{89}\text{Kr}$	8140	140	8262	4	0.9	U			Stu		81Ho17
	8120	120			1.2	o			Bwg		87Gr.B
	8155	30			3.6	C			Bwg		92Gr.A
$^{89}\text{Kr}(\beta^-)^{89}\text{Rb}$	5150	30	5177	6	0.9	U					67Ki01
	5191	60			-0.2	U			Trs		78Wo15 *
	5140	120			0.3	U			Stu		81Ho17 *
$^{89}\text{Rb}(\beta^-)^{89}\text{Sr}$	4486	12	4497	5	0.9	-					66Ki06
	4491	15			0.4	o			Gsn		80Bl.A
	4510	9			-1.5	-			Gsn		80De02 *
	ave.	4501	7		-0.7	1	57	57	$^{89}\text{Rb}$		average
$^{89}\text{Sr}(\beta^-)^{89}\text{Y}$	1463	5	1499.3	1.6	7.3	B					49La06
	1488	4			2.8	B					70Wo05
$^{89}\text{Zr}(\beta^+)^{89}\text{Y}$	2841	10	2832.8	2.8	-0.8	U					51Hy24 *
	2832	10			0.1	U					53Sh48 *
	2828	7			0.7	-					60Ha26 *
$^{89}\text{Y}(\text{p,n})^{89}\text{Zr}$	-3612.8	4.	-3615.1	2.8	-0.6	-			Tkm		63Ok01 Z
	-3619.4	6.			0.7	-			Oak		64Jo11 Z
$^{89}\text{Zr}(\beta^+)^{89}\text{Y}$	ave.	2832	3	2832.8	2.8	0.4	1	85	$^{89}\text{Zr}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{89}\text{Nb}(\beta^+)^{89}\text{Zr}$	3870	100	4250	24	3.8	B					55Ma13
	4340	50			-1.8	1	23	22 $^{89}\text{Nb}$			74Vo08
$^{89}\text{Mo}(\beta^+)^{89}\text{Nb}$	5970	300	5610	24	-1.2	U					64Bu12
$^{89}\text{Tc}(\beta^+)^{89}\text{Mo}$	7510	210	7620	5	0.5	U					91He04 *
$^{89}\text{Nb}-u$	$M - A = -80656(28)$ keV for mixture gs+m at 0#30 keV										Nub16b **
$^{89}\text{Kr}(\beta^-)^{89}\text{Rb}$	$E_{\beta^-} = 4970(60)$ to 220.948 level										GAu **
$^{89}\text{Kr}(\beta^-)^{89}\text{Rb}$	Splitting Table 3a into 3 groups yields 4610(120) 4867(152) 5135(123) keV										GAu **
$^{89}\text{Rb}(\beta^-)^{89}\text{Sr}$	Original error 8 corrected in reference										94Ha.A **
$^{89}\text{Zr}(\beta^+)^{89}\text{Y}$	$E_{\beta^+} = 910(10)$ 901(10) 897(7) respectively, to $^{89}\text{Y}^m$ at 908.97 keV										Nub16b **
$^{89}\text{Tc}(\beta^+)^{89}\text{Mo}$	$E_{\beta^+} = 6370(210)$ to 118.8 level; no Fermi-Kurie plot										91He04 **
$^{90}\text{Se}-u$	-59904	236				2			GT1	1.5	04Ma.A
$\text{C}_4 \text{H}_{10} \text{O}_2 - ^{90}\text{Zr}$	163377	6	163380.80	0.13	0.3	U			M15	2.5	63Ri07
$^{90}\text{Zr}-u$	-95301.36	0.23	-95301.24	0.13	0.5	1	30	30 $^{90}\text{Zr}$	MS1	1.0	15Gu09
$^{90}\text{Nb}-u$	-88872	50	-88741	4	2.6	U			GS2	1.0	05Li24 *
$^{90}\text{Mo}-\text{C}_7 \text{H}_6$	-133018.9	4.7	-133019	4	0.0	1	63	63 $^{90}\text{Mo}$	CP1	1.0	11Fa10
$^{90}\text{Mo}_{1.033}-\text{C}_7 \text{H}_9$	-159318	23	-159334	4	-0.7	U			CP1	1.0	11Fa10
$^{90}\text{Tc}-\text{C}_7 \text{H}_6$	-122880.3	6.7	-122876.3	1.1	0.6	U			CP1	1.0	11Fa10
$^{90}\text{Tc}_{1.033}-\text{C}_7 \text{H}_9$	-148835	22	-148856.9	1.1	-1.0	U			CP1	1.0	11Fa10
	-148854.2	8.5			-0.3	U			CP1	1.0	11Fa10
$^{90}\text{Ru}_{1.033}-\text{C}_7 \text{H}_9$	-142382	11	-142380	4	0.2	1	14	14 $^{90}\text{Ru}$	CP1	1.0	11Fa10
$^{90}\text{Kr}-^{85}\text{Rb}_{1.059}$	12942.6	2.0				2			MA8	1.0	06De36
$^{90}\text{Rb}-^{85}\text{Rb}_{1.059}$	8211	9	8213	7	0.3	1	60	60 $^{90}\text{Rb}$	MA4	1.0	02Ra23 *
$^{90}\text{Tc}-^{85}\text{Rb}_{1.059}$	17489.2	8.0	17488.6	1.1	-0.1	U			SH1	1.0	08We10
	17489.8	4.2			-0.3	U			JY1	1.0	08We10
$^{90}\text{Ru}-^{85}\text{Rb}_{1.059}$	23775	11	23759	4	-1.5	-			SH1	1.0	08We10
	23756.6	4.7			0.5	-			JY1	1.0	08We10
ave.	23759	4			-0.1	1	86	86 $^{90}\text{Ru}$			average
$^{90}\text{Tc}-^{86}\text{Kr}_{1.047}$	17664.6	1.1				2			JY1	1.0	12Ka12
$^{90}\text{Tc}^m - ^{86}\text{Kr}_{1.047}$	17819.2	1.4				2			JY1	1.0	12Ka12
$^{90}\text{Zr}-^{87}\text{Rb}_{1.034}$	-1393.79	0.16	-1393.91	0.13	-0.7	1	62	62 $^{90}\text{Zr}$	MS1	1.0	15Gu09
$^{90}\text{Br}-^{88}\text{Rb}_{1.023}$	22017.0	3.6				2			JY1	1.0	07Ra23
$^{89}\text{Rb}-^{90}\text{Rb}_{91}^{85}\text{Rb}_{209}$	-1826	24	-1818	12	0.1	U			P21	2.5	82Au01
$^{90}\text{Zr}(\alpha, ^8\text{He})^{86}\text{Zr}$	-40136	30	-39988	4	4.9	B			INS		90Ka01
$^{90}\text{Zr}(^3\text{He}, ^6\text{He})^{87}\text{Zr}$	-12083	8	-12086	4	-0.4	1	27	27 $^{87}\text{Zr}$	MSU		78Pa11
$^{90}\text{Zr}(p,t)^{88}\text{Zr}$	-12805	10	-12805	5	0.0	1	29	29 $^{88}\text{Zr}$	Oak		71Ba43
$^{89}\text{Y}(n,\gamma)^{90}\text{Y}$	6857.1	1.0	6857.03	0.10	-0.1	U			ORn		81Ra07
	6857.26	0.30			-0.8	-					83De17
	6856.98	0.17			0.3	-			ILn		93Mi04 Z
	6857.01	0.14			0.1	-			Bdn		06Fi.A
$^{89}\text{Y}(d,p)^{90}\text{Y}$	4635	5	4632.46	0.10	-0.5	U					64Wa14
$^{89}\text{Y}(n,\gamma)^{90}\text{Y}$	ave.	6857.03	0.10	6857.03	0.10	0.0	1	100	63 $^{89}\text{Y}$		average
$^{89}\text{Y}(p,\gamma)^{90}\text{Zr}$	8351	4	8353.2	1.6	0.5	1	16	16 $^{89}\text{Y}$			75Be.B
$^{90}\text{Zr}(\gamma,n)^{89}\text{Zr}$	-11940	50	-11968	3	-0.6	U			Phi		63Ge02
$^{90}\text{Zr}(p,d)^{89}\text{Zr}$	-9728	10	-9744	3	-1.6	U			Oak		71Ba43
$^{90}\text{Zr}(d,t)^{89}\text{Zr}$	-5719.2	7.1	-5711	3	1.1	1	19	19 $^{89}\text{Zr}$	SPa		79Bo37
$^{90}\text{Zr}(^3\text{He}, \alpha)^{89}\text{Zr}$	8580	50	8609	3	0.6	U			Phi		67Fo04
$^{90}\text{Br}(\beta^-)^{90}\text{Kr}$	9800	400	10959	4	2.9	U			Stu		81Ho17
	10280	110			6.2	C			Bwg		87Gr.B
	10350	75			8.1	C			Bwg		92Gr.A
$^{90}\text{Kr}(\beta^-)^{90}\text{Rb}$	4410	30	4405	7	-0.2	U					70Ma11
	4390	40			0.4	U			Trs		78Wo15
	4380	25			1.0	U			Bwg		87Gr.A
$^{90}\text{Rb}^i(\text{IT})^{90}\text{Rb}$	71	12				2					82Au01
$^{90}\text{Rb}(\beta^-)^{90}\text{Sr}$	6550	60	6584	7	0.6	U			Trs		78Wo15
	6560	150			0.2	U			Bwg		78St02
	6585	15			-0.1	o			Gsn		80Bl.A
	6578	15			0.4	o			Gsn		80De02
	6587	10			-0.3	1	43	40 $^{90}\text{Rb}$	Gsn		92Pr03
$^{90}\text{Sr}(\beta^-)^{90}\text{Y}$	546	2	545.9	1.4	0.0	-					64Da16
	546	2			0.0	-					83Ha35

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item		Input value		Adjusted value	$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{90}\text{Sr}(\beta^-)^{90}\text{Y}$	ave.	546.0	1.4	545.9	1.4	0.0	1	99	$^{90}\text{Sr}$		average
$^{90}\text{Y}(\beta^-)^{90}\text{Zr}$		2271	2	2278.5	1.6	3.7	B				61Ni02
		2284	5			-1.1	-				64Da16
		2273	5			1.1	-				64La13
		2280	5			-0.3	-				66Ri01
		2278	8			0.1	U		Gsn		80Bl.A
		2279.5	2.9			-0.4	-				83Ha35
	ave.	2279.2	2.0			-0.4	1	62	$^{62}\text{Y}$		average
$^{90}\text{Nb}(\beta^+)^{90}\text{Zr}$		6111	4	6111	3	0.0	1	69	$^{69}\text{Nb}$		68Pe01 *
$^{90}\text{Mo}(\beta^+)^{90}\text{Nb}$		2489	4	2489	3	0.0	1	69	$^{37}\text{Mo}$		66Pe10 *
$^{90}\text{Tc}(\beta^+)^{90}\text{Mo}$		8920	410	9448	4	1.3	U				74Ia01 *
		9270	300			0.6	U				81Ox01 *
		8726	300			2.4	U				81Ox01 *
$^{90}\text{Nb}-u$		$M-A=-82721(29)$ keV for mixture gs+n at 122.370 keV									
$^{90}\text{Rb}-^{85}\text{Rb}_{1.059}$		$D_M=8326(9)$ $\mu\text{u}$ for $^{90}\text{Rb}^m$ at 106.90 keV; $M-A=-79260(9)$ keV									
$^{90}\text{Nb}(\beta^+)^{90}\text{Zr}$		$E_{\beta^+}=1500(4)$ to $6^+$ level at 3589.419 keV									
$^{90}\text{Mo}(\beta^+)^{90}\text{Nb}$		$E_{\beta^+}=1085(4)$ to $1^+$ level at 382.01 keV									
$^{90}\text{Tc}(\beta^+)^{90}\text{Mo}$		$E_{\beta^+}=7900(400)$ from $^{90}\text{Tc}^m$ at 144.1 to ground state (85%) and 947.97 (15%) level									
$^{90}\text{Tc}(\beta^+)^{90}\text{Mo}$		$E_{\beta^+}=5300(300)$ to 2946.82 level									
$^{90}\text{Tc}(\beta^+)^{90}\text{Mo}$		$E_{\beta^+}=6900(300)$ from $^{90}\text{Tc}^m$ at 144.0(1.7) to $2^+$ level a 947.97 keV									
$^{91}\text{Se}-u$		-54300	186				2				16Kn03
$^{91}\text{Rb}-u$		-83532	21	-83463	8	1.3	U		GT3	2.5	89Al33
$\text{C}_7 \text{H}_7-^{91}\text{Zr}$		149143.1	4.4	149135.00	0.11	-0.7	U		M15	2.5	63Ri07
$^{91}\text{Zr}-u$		-94359.85	0.25	-94359.78	0.11	0.3	1	20	$^{20}\text{Zr}$	1.0	15Gu09
$^{91}\text{Nb}-u$		-93064	46	-93010	3	1.2	U		GS2	1.0	05Li24 *
$^{91}\text{Mo}-\text{C}_7 \text{H}_7$		-143031.3	8.3	-143030	7	0.2	1	65	$^{65}\text{Mo}$	1.0	11Fa10
$^{91}\text{Tc}-\text{C}_7 \text{H}_7$		-136340.9	6.7	-136350.3	2.5	-1.4	-		CP1	1.0	11Fa10
		-136353.6	4.6			0.7	-		CP1	1.0	11Fa10
	ave.	-136350	4			-0.2	1	45	$^{45}\text{Tc}$		average
$^{91}\text{Ru}_{1.011}-\text{C}_7 \text{H}_8$		-136730	620	-136664.6	2.4	0.1	U		CP1	1.0	11Fa10
$^{91}\text{Ru}-\text{C}_7 \text{H}_7$		-128035.6	3.9	-128033.7	2.4	0.5	1	37	$^{37}\text{Ru}$	1.0	11Fa10
$^{91}\text{Ru}_{1.022}-\text{C}_7 \text{H}_9$		-145260	23	-145295.4	2.4	-1.5	U		CP1	1.0	11Fa10
$^{91}\text{Kr}-^{85}\text{Rb}_{1.071}$		18279.5	2.4				2		MA8	1.0	06De36
$^{91}\text{Rb}-^{85}\text{Rb}_{1.071}$		11003	10	11010	8	0.7	1	70	$^{70}\text{Rb}$	1.0	02Ra23
$^{91}\text{Sr}-^{85}\text{Rb}_{1.071}$		4702	9	4669	6	-3.7	B		MA4	1.0	02Ra23
$^{91}\text{Tc}-^{85}\text{Rb}_{1.071}$		12898.3	5.4	12898.2	2.5	0.0	1	22	$^{22}\text{Tc}$	1.0	08We10
$^{91}\text{Ru}-^{85}\text{Rb}_{1.071}$		21223	11	21214.7	2.4	-0.8	-		SH1	1.0	08We10
		21215.5	4.2			-0.2	-		JY1	1.0	08We10
	ave.	21216	4			-0.4	1	37	$^{37}\text{Ru}$		average
$^{91}\text{Zr}-^{87}\text{Rb}_{1.046}$		637.32	0.19	637.39	0.11	0.4	1	35	$^{35}\text{Zr}$	1.0	15Gu09
$^{91}\text{Br}-^{88}\text{Rb}_{1.034}$		26098.3	3.8				2		JY1	1.0	07Ra23
$^{91}\text{Tc}-^{94}\text{Mo}_{.968}$		10303.0	4.4	10304.1	2.5	0.2	1	33	$^{33}\text{Tc}$	1.0	08We10
$^{91}\text{Ru}-^{94}\text{Mo}_{.968}$		18620.9	4.7	18620.6	2.4	-0.1	1	26	$^{26}\text{Ru}$	1.0	08We10
$^{91}\text{Zr}-^{90}\text{Zr}$		942	12	941.47	0.16	0.0	U		M15	2.5	63Ri07
$^{90}\text{Rb}^+-^{91}\text{Rb}_{.824} \text{ } ^{85}\text{Rb}_{.176}$		-686	24	-770	15	-1.4	U		P21	2.5	82Au01
$^{91}\text{Zr}(\text{p,t})^{89}\text{Zr}$		-10677	10	-10681	3	-0.4	U		Oak		71Ba43
$^{90}\text{Zr}(\text{n},\gamma)^{91}\text{Zr}$		7194.4	0.5	7194.35	0.15	-0.1	-				81Lo.A Z
		7192.7	0.8			2.1	-		Bdn		06Fi.A
$^{90}\text{Zr}(\text{d,p})^{91}\text{Zr}$		4959	20	4969.78	0.15	0.5	U		Pit		64Co11
		4969	8			0.1	U		MIT		72Gr12
		4970.3	2.2			-0.2	U		SPa		79Bo37
$^{91}\text{Zr}(\text{p,d})^{90}\text{Zr}$		-4977	10	-4969.78	0.15	0.7	U		Oak		71Ba43
$^{91}\text{Zr}(\text{d,t})^{90}\text{Zr}$		-932	20	-937.12	0.15	-0.3	U		Pit		64Co11
		-940.3	3.7			0.9	U		SPa		79Bo37
$^{90}\text{Zr}(\text{n},\gamma)^{91}\text{Zr}$	ave.	7193.9	0.4	7194.35	0.15	1.0	1	13	$^{7}\text{Zr}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{90}\text{Zr}(p,\gamma)^{91}\text{Nb}$	5167	5	5154.4	2.9	-2.5	B					71Ra08
	5167	4			-3.1	C					75Be.B Z
$^{90}\text{Zr}(^3\text{He,d})^{91}\text{Nb}$	-227	20	-339.0	2.9	-5.6	B			Hei		70Kn05
$^{90}\text{Zr}(\alpha,t)^{91}\text{Nb}$	-14643	27	-14659.4	2.9	-0.6	U			Brk		71Zi03
$^{91}\text{Ru}^m(\epsilon p)^{90}\text{Mo}$	4300	500				2					83Ha06
$^{91}\text{Br}(\beta^-)^{91}\text{Kr}$	9790	100	9867	4	0.8	U					89Gr03
	9805	50			1.2	U			Bwg		92Gr.A
$^{91}\text{Kr}(\beta^-)^{91}\text{Rb}$	6420	80	6771	8	4.4	B			Trs		78Wo15
	6450	80			4.0	B			Bwg		89Gr03
$^{91}\text{Rb}(\beta^-)^{91}\text{Sr}$	5830	45	5907	9	1.7	U			Trs		78Wo15 *
	5927	24			-0.8	o			Gsn		80De02 *
	5920	28			-0.5	-			McG		83Ia02 *
	5930	22			-1.1	-			Gsn		92Pr03 *
ave.	5926	17			-1.1	1	26	18 $^{91}\text{Rb}$			average
$^{91}\text{Sr}(\beta^-)^{91}\text{Y}$	2669	10	2699	5	3.0	B					53Am08 *
	2684	10			1.5	-					73Ha11 *
	2704	8			-0.6	-			Gsn		80De02 *
	2709	15			-0.6	-			McG		83Ia02 *
ave.	2698	6			0.2	1	83	81 $^{91}\text{Sr}$			average
$^{91}\text{Y}(\beta^-)^{91}\text{Zr}$	1545	5	1544.3	1.8	-0.1	-					64La13
	1544	2			0.1	-					75Ra08
ave.	1544.1	1.9			0.1	1	98	98 $^{91}\text{Y}$			average
$^{91}\text{Zr}(p,n)^{91}\text{Nb}$	-2045	6	-2039.9	2.9	0.8	-					70Ki01
	-2038.8	3.4			-0.3	-			Oak		71Ma47
	ave.	-2040.3	3.0			0.1	1	98	98 $^{91}\text{Nb}$	Kyu	
$^{91}\text{Mo}(\beta^+)^{91}\text{Nb}$	4460	30	4429	7	-1.0	-					56Sm96
	4435	23			-0.3	-					93Os06
	ave.	4444	18			-0.8	1	14	11 $^{91}\text{Mo}$		
$^{91}\text{Tc}(\beta^+)^{91}\text{Mo}$	6220	200	6222	7	0.0	U					74Ia01
$^{91}\text{Nb}-u$	$M - A = -86636(30)$ keV for mixture gs+m at 104.60 keV										Nub16b **
$^{91}\text{Rb}(\beta^-)^{91}\text{Sr}$	$E_{\beta^-} = 5760(40)$ to $^{91}\text{Sr}$ ground state <8% and 93.628 keV $(3/2)^+$ level 25%										Ens13a **
$^{91}\text{Rb}(\beta^-)^{91}\text{Sr}$	Original error 8 corrected to 13 keV in reference										94Ha.A **
$^{91}\text{Rb}(\beta^-)^{91}\text{Sr}$	$E_{\beta^-} = 5857$ to mixture $^{91}\text{Sr}$ ground state <8% and 93.628 $(3/2)^+$ level 25%										Ens13a **
$^{91}\text{Rb}(\beta^-)^{91}\text{Sr}$	$E_{\beta^-} = 5850(20)$ and $E_{\beta^-} = 5860(10)$ respectively, to $^{91}\text{Sr}$ ground state <8% and 93.628 level 25%										Ens13a **
$^{91}\text{Sr}(\beta^-)^{91}\text{Y}$	$E_{\beta^-} = 2665(10), 2030(20), 1359(10), 1093(10)$ to										53Am08 **
*	ground state, $3/2^-$ level at 653.02 keV, $(5/2)^+$ at 1305.39, $(5/2)^-$ at 1545.90										Ens13a **
$^{91}\text{Sr}(\beta^-)^{91}\text{Y}$	Original error 4 increased: in disagreement with other results										AHW **
$^{91}\text{Sr}(\beta^-)^{91}\text{Y}$	Original error 3 corrected in reference										94Ha.A **
$^{92}\text{Br}-u$	-60711	103	-60368	7	1.3	U			GT2	2.5	08Kn.A
$^{92}\text{Rb}-u$	-80323	32	-80272	7	0.6	U			Pb1	2.5	89Al33
$C_7 \text{H}_8 - ^{92}\text{Zr}$	157569.4	3.8	157564.94	0.11	-0.5	U			M15	2.5	63Ri07
$^{92}\text{Zr}-u$	-94964.66	0.18	-94964.68	0.11	-0.1	1	37	37 $^{92}\text{Zr}$	MS1	1.0	15Gu09
$^{92}\text{Nb}-u$	-92851	56	-92811.4	1.9	0.7	U			GS2	1.0	05Li24 *
$C_7 \text{H}_8 - ^{92}\text{Mo}$	155790.0	3.2	155793.10	0.17	0.4	U			M15	2.5	63Ri07
$^{92}\text{Mo}_{1.011} - C_7 \text{H}_9$	-164641.3	7.0	-164643.26	0.17	-0.3	U			CP1	1.0	11Fa10
$^{92}\text{Mo}-u$	-93193.14	0.47	-93192.84	0.17	0.6	1	13	13 $^{92}\text{Mo}$	MS1	1.0	15Gu09
$^{92}\text{Tc}-C_7 \text{H}_8$	-147328	13	-147330	3	-0.2	U			CP1	1.0	11Fa10
$^{92}\text{Tc}_{.989} - C_7 \text{H}_7$	-138569	10	-138573	3	-0.4	-			CP1	1.0	11Fa10
$^{92}\text{Tc}_{1.011} - C_7 \text{H}_9$	-156087.6	6.1	-156088	3	0.0	-			CP1	1.0	11Fa10
$^{92}\text{Tc}_{.989} - C_7 \text{H}_7$	ave. -138572	5	-138573	3	-0.2	1	40	40 $^{92}\text{Tc}$			average
$^{92}\text{Ru}-C_7 \text{H}_8$	-142352	18	-142365.9	2.9	-0.8	o			CP1	1.0	08Fa11
	-142352	18			-0.8	U			CP1	1.0	11Fa10
	-142377	10			1.1	o			CP1	1.0	08Fa11
	-142378	10			1.2	U			CP1	1.0	11Fa10

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{92}\text{Ru}_{1.011}-\text{C}_7\text{H}_9$	-151074.5	5.6	-151068.3	2.9	1.1	o			CP1	1.0	08Fa11
	-151074.5	5.6			1.1	1	28	28 $^{92}\text{Ru}$	CP1	1.0	11Fa10
$^{92}\text{Rh}_{1.011}-\text{C}_7\text{H}_9$	-138825	23	-138802	5	1.0	U			CP1	1.0	11Fa10
	-138818	17			1.0	U			CP1	1.0	11Fa10
									MA8	1.0	06De36
$^{92}\text{Kr}-^{85}\text{Rb}_{1.082}$	21616.6	2.9				2			MA8	1.0	06De36
$^{92}\text{Rb}-^{85}\text{Rb}_{1.082}$	15176	9	15172	7	-0.4	1	53	53 $^{92}\text{Rb}$	MA4	1.0	02Ra23
$^{92}\text{Sr}-^{85}\text{Rb}_{1.082}$	6482	9	6482	4	0.0	-			MA4	1.0	02Ra23
	6484.0	4.3			-0.5	-			MA8	1.0	05Gu37
	ave.	6484	4		-0.5	1	90	90 $^{92}\text{Sr}$			average
$^{92}\text{Mo}-^{85}\text{Rb}_{1.082}$	2251.43	0.85	2250.66	0.17	-0.9	U			JY1	1.0	12Ka13
$^{92}\text{Tc}-^{85}\text{Rb}_{1.082}$	10728	12	10713	3	-1.2	U			SH1	1.0	08We10
	10712.5	4.3			0.2	1	60	60 $^{92}\text{Tc}$	JY1	1.0	08We10
	15684.3	5.7	15677.9	2.9	-1.1	-			SH1	1.0	08We10
$^{92}\text{Ru}-^{85}\text{Rb}_{1.082}$	15677.9	4.3			0.0	-			JY1	1.0	08We10
	ave.	15680	3		-0.7	1	72	72 $^{92}\text{Ru}$			average
$^{92}\text{Rh}-^{85}\text{Rb}_{1.082}$	27841	37	27811	5	-0.8	U			SH1	1.0	08We10
	27811.2	4.7				2			JY1	1.0	08We10
$^{92}\text{Zr}-^{87}\text{Rb}_{1.057}$	1031.51	0.21	1031.50	0.11	0.0	1	27	27 $^{92}\text{Zr}$	MS1	1.0	15Gu09
$^{92}\text{Mo}-^{87}\text{Rb}_{1.057}$	2803.38	0.18	2803.34	0.17	-0.2	1	87	87 $^{92}\text{Mo}$	MS1	1.0	15Gu09
$^{92}\text{Br}-^{88}\text{Rb}_{1.045}$	32306.8	7.2				2			JY1	1.0	07Ra23
$^{92}\text{Zr}-^{35}\text{Cl}-^{90}\text{Zr}-^{37}\text{Cl}$	3285	2	3286.67	0.18	0.2	U			H13	4.0	63Ba20
$^{92}\text{Zr}-^{91}\text{Zr}$	-603	12	-604.90	0.10	-0.1	U			M15	2.5	63Ri07
$^{88}\text{Rb}-^{92}\text{Rb}_{.410}$	-3258	22	-3309.2	2.5	-0.9	U			P21	2.5	82Au01
$^{89}\text{Rb}-^{92}\text{Rb}_{.553}$	-3457	24	-3470	6	-0.2	U			P21	2.5	82Au01
$^{91}\text{Rb}-^{92}\text{Rb}_{.848}$	-1703	25	-1766	9	-1.0	U			P21	2.5	82Au01
$^{90}\text{Rb}^x-^{92}\text{Rb}_{.699}$	-2059	24	-2131	14	-1.2	U			P21	2.5	82Au01
$^{90}\text{Rb}^x-^{92}\text{Rb}_{.326}$	209	24	157	14	-0.9	U			P21	2.5	82Au01
$^{92}\text{Mo}(\alpha,^8\text{He})^{88}\text{Mo}$	-43278	20	-43307	4	-1.4	U			INS		90Ka01
$^{92}\text{Mo}(\text{p},\alpha)^{89}\text{Nb}$	-1306	50	-1319	24	-0.3	R			ANL		75Se.A
$^{92}\text{Mo}(\alpha,^3\text{He},^6\text{He})^{89}\text{Mo}$	-14465	15	-14455	4	0.7	U			MSU		80Pa02
$^{92}\text{Zr}(\text{p,t})^{90}\text{Zr}$	-7372	14	-7347.33	0.15	1.8	U			Bld		66St15
	-7350	10			0.3	U			Oak		71Ba43
	-14330	30	-14297	3	1.1	U			VUn		76Ka08
$^{92}\text{Mo}(\text{p,t})^{90}\text{Mo}$	785	15	808	8	1.5	1	26	15 $^{92}\text{Rb}$			84Kr.B
$^{92}\text{Rb}(\beta^-)^{91}\text{Sr}$	8634.91	0.20	8634.78	0.09	-0.7	-			ILn		79Br.25
$^{91}\text{Zr}(\text{n},\gamma)^{92}\text{Zr}$	8634.64	0.15			0.9	-					81Su.A
	8635.00	0.24			-0.9	-			Bdn		06Fi.A
	6470	30	6410.21	0.09	-2.0	U					62Ma06
$^{91}\text{Zr}(\text{d,p})^{92}\text{Zr}$	6395	20			0.8	U			Pit		64Co11
	6410.9	4.3			-0.2	U			SFa		79Bo37
	-6410	11	-6410.21	0.09	0.0	U			Bld		66St15
$^{92}\text{Zr}(\text{p,d})^{91}\text{Zr}$	-6410	10			0.0	U		Oak		71Ba43	
$^{92}\text{Zr}(\text{d,t})^{91}\text{Zr}$	-2363	25	-2377.55	0.09	-0.6	U			Pit		64Co11
$^{91}\text{Zr}(\text{n},\gamma)^{92}\text{Zr}$	ave.	8634.79	0.11	8634.78	0.09	-0.1	1	75	39 $^{91}\text{Zr}$		average
		-10446	15	-10447	6	0.0	-		Tex		73Ko03
		-10432	25			-0.6	-		Grn		73Mo03
	ave.	-10442	13			-0.3	1	24	24 $^{91}\text{Mo}$		average
$^{92}\text{Br}(\beta^-)^{92}\text{Kr}$	12155	100	12537	7	3.8	B			Bwg		89Gr03
	12220	55			5.8	C			Bwg		92Gr.A
$^{92}\text{Kr}(\beta^-)^{92}\text{Rb}$	6160	80	6003	7	-2.0	U			Trs		78Wo15
	6045	80			-0.5	o			Bwg		89Gr03
	5987	10			1.6	o			Bwg		92Gr.A
	5993	27			0.4	U			Bwg		92Gr06
	8080	160	8095	6	0.1	U			Trs		78Wo15
$^{92}\text{Rb}(\beta^-)^{92}\text{Sr}$	8091	15			0.3	o			Gsn		80Bl.A
	8111	15			-1.1	o			Gsn		80De02
	8080	30			0.5	-			McG		83Ia02
	8095	25			0.0	o			Bwg		87Gr.A
	8096	16			-0.1	-			Bwg		92Gr.A
	8107	15			-0.8	-			Gsn		92Pr03
	ave.	8099	10			-0.4	1	39	32 $^{92}\text{Rb}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{92}\text{Sr}(\beta^-)^{92}\text{Y}$	1929	50	1949	9	0.4	U					57He39 *
	1930	30			0.6	-			Trs		78Wo15 *
	1920	20			1.5	-			McG		83Ia02
$^{92}\text{Y}(\beta^-)^{92}\text{Zr}$	ave.	1923	17		1.6	1	32	29 $^{92}\text{Y}$			average
		3640	20	3643	9	0.1	-				62Bu16
		3630	15			0.8	-		McG		83Ia02
$^{92}\text{Nb}(\beta^+)^{92}\text{Zr}$	ave.	3634	12		0.7	1	58	58 $^{92}\text{Y}$			average
		2005	6	2005.7	1.8	0.1	U				59We30 *
		2008	6			-0.4	U				62Bu16 *
$^{92}\text{Zr}(\text{p,n})^{92}\text{Nb}$		-2790.7	2.3	-2788.1	1.8	1.1	-		Kyu		74Ku01
		-2792	5			0.8	-				75Ke12
	ave.	-2790.9	2.1			1.4	1	73	73 $^{92}\text{Nb}$		average
$^{92}\text{Tc}(\beta^+)^{92}\text{Mo}$		7880	100	7883	3	0.0	U				64Va05
$^{92}\text{Mo}(\text{p,n})^{92}\text{Tc}$		-8672	50	-8665	3	0.1	U		Tal		66Mo06 *
$^{92}\text{Mo}(^3\text{He,t})^{92}\text{Tc}$		-7882	30	-7901	3	-0.6	U		ChR		73Ha02
$^{92}\text{Nb}-\text{u}$	$M - A = -86422(34)$ keV for mixture gs+m at 135.5 keV										Nub16b **
$^{92}\text{Sr}(\beta^-)^{92}\text{Y}$	$E_{\beta^-} = 545(50)$ 546(30) respectively, to $1^+$ level at 1383.91 keV										Ens12a **
$^{92}\text{Nb}(\beta^+)^{92}\text{Zr}$	$\text{p}^+ = 56(6) \times 10^{-5}$ , $60(6) \times 10^{-5}$ respectively, to $2^+$ level at 934.51 keV										Ens12a **
*	recalculated $Q_{\beta^+} = 2140(6)$ 2143(6) respectively, from $^{92}\text{Nb}^m$ at 135.5 keV										AHW **
$^{92}\text{Mo}(\text{p,n})^{92}\text{Tc}$	$T = 9040(50)$ to $(4^+)$ level at 270.09 keV										Ens12a **
$^{93}\text{Br}-\text{u}$	-56866	322	-56780	460	0.2	o			GT1	1.5	04Ma.A
	-56780	185				2			GT3	2.5	16Kn03
$^{93}\text{Rb}-\text{u}$	-78036	21	-77961	8	1.4	U			Pb1	2.5	89A133
	-77868	100			-0.4	U			GT2	2.5	08Kn.A
$\text{C}_7 \text{H}_9 - ^{93}\text{Nb}$	164046.9	3.5	164052.1	1.6	0.6	U			M15	2.5	63Ri07
$^{93}\text{Mo}-\text{u}$	-93194	30	-93191.23	0.19	0.1	U			GS2	1.0	05Li24 *
$^{93}\text{Tc}-\text{u}$	-89729	31	-89754.9	1.1	-0.8	U			GS2	1.0	05Li24
$^{93}\text{Tc}-\text{C}_7 \text{H}_9$	-160189.5	7.7	-160180.1	1.1	1.2	U			CP1	1.0	11Fa10
	-160170	22			-0.5	U			CP1	1.0	11Fa10
	-160189.4	8.5			1.1	U			CP1	1.0	11Fa10
$^{93}\text{Tc}_{989} - \text{C}_7 \text{H}_8$	-151270	190	-151367.8	1.1	-0.5	U			CP1	1.0	11Fa10
$^{93}\text{Ru}-\text{C}_7 \text{H}_9$	-153318.2	6.4	-153320.8	2.2	-0.4	-			CP1	1.0	11Fa10
	-153307	23			-0.6	U			CP1	1.0	11Fa10
	-153324.0	4.8			0.7	-			CP1	1.0	11Fa10
	-153321.9	3.5			0.3	-			CP1	1.0	11Fa10
	ave.	-153321.9	2.6			0.4	1	73	73 $^{93}\text{Ru}$		average
$^{93}\text{Rh}-\text{C}_7 \text{H}_9$	-144485	25	-144512.5	2.8	-1.1	o			CP1	1.0	08Fa11
	-144485	26			-1.1	U			CP1	1.0	11Fa10
	-144527.7	5.3			2.9	U			CP1	1.0	08Fa11
	-144527.7	5.2			2.9	U			CP1	1.0	11Fa10
$^{93}\text{Kr}-^{85}\text{Rb}_{1.094}$	-144512.9	3.8			0.1	1	55	55 $^{93}\text{Rh}$	CP1	1.0	11Fa10
	27649.2	2.7				2			MA8	1.0	06De36
$^{93}\text{Rb}-^{85}\text{Rb}_{1.094}$	18549	10	18541	8	-0.8	1	71	71 $^{93}\text{Rb}$	MA4	1.0	02Ra23
$^{93}\text{Sr}-^{85}\text{Rb}_{1.094}$	10526	10	10526	8	0.0	1	66	66 $^{93}\text{Sr}$	MA4	1.0	02Ra23
$^{93}\text{Ru}-^{85}\text{Rb}_{1.094}$	13609.4	4.3	13606.5	2.2	-0.7	1	27	27 $^{93}\text{Ru}$	JY1	1.0	08We10
$^{93}\text{Rh}-^{85}\text{Rb}_{1.094}$	22428	12	22414.8	2.8	-1.1	-			SH1	1.0	08We10
	22413.5	4.5			0.3	-			JY1	1.0	08We10
	ave.	22415	4			-0.1	1	45	45 $^{93}\text{Rh}$		average
$^{91}\text{Rb}-^{93}\text{Rb}_{.489} \ ^{89}\text{Rb}_{.511}$	-471	9	-479	9	-0.4	1	15	12 $^{91}\text{Rb}$	P31	2.5	86Au02
$^{91}\text{Rb}-^{93}\text{Rb}_{.326} \ ^{90}\text{Rb}_{.674}$	-656	23	-627	12	0.5	U			P21	2.5	82Au01
$^{92}\text{Rb}-^{93}\text{Rb}_{.495} \ ^{91}\text{Rb}_{.505}$	465	23	436	8	-0.5	U			P21	2.5	82Au01
$^{93}\text{Rb}(\beta^- \text{n})^{92}\text{Sr}$	2220	30	2176	9	-1.5	U					84Kr.B
$^{92}\text{Sr}(\text{n},\gamma)^{93}\text{Sr}$	5230	6	5290	8	10.0	B					80Kr07
$^{92}\text{Zr}(\text{n},\gamma)^{93}\text{Zr}$	6733.7	1.1	6734.3	0.4	0.6	-					72Gr23 Z
	6734.0	0.7			0.5	-					79Ke.D Z
	6735.3	0.7			-1.4	-			Bdn		06Fi.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{92}\text{Zr}(d,p)^{93}\text{Zr}$		4493	20	4509.8	0.4	0.8	U		Pit		64Co11	
$^{92}\text{Zr}(n,\gamma)^{93}\text{Zr}$	ave.	6734.5	0.5	6734.3	0.4	-0.4	1	98	$^{98}\text{Zr}$		average	
$^{93}\text{Nb}(\gamma,n)^{92}\text{Nb}$		-8780	60	-8830.9	2.0	-0.8	U		Phi		60Ge01	
		-8825	3			-2.0	1	44	$^{27}\text{Nb}$	McM	79Ba06	
$^{93}\text{Nb}(d,t)^{92}\text{Nb}$		-2581	20	-2573.6	2.0	0.4	U		Pit		64Co11	
		-2571	10			-0.3	U		Tal		64Sh04	
$^{92}\text{Mo}(n,\gamma)^{93}\text{Mo}$		8067.4	1.5	8069.81	0.09	1.6	U				73Wa17	
		8066	2			1.9	U				77Ri04	
		8069.81	0.09				2		MMn		91Is02	
		8070.0	0.3			-0.6	U		Bdn		06Fi.A	
$^{92}\text{Mo}(d,p)^{93}\text{Mo}$		5853	20	5845.24	0.09	-0.4	U		Pit		64Co11	
$^{92}\text{Mo}(p,\gamma)^{93}\text{Tc}$		4081	5	4086.5	1.0	1.1	U				75Be.B	
		4086.5	1.0				2				83Ay01	
$^{92}\text{Mo}(^3\text{He},d)^{93}\text{Tc}$		-1411	4	-1407.0	1.0	1.0	U		Hei		83Wi.A	
$^{93}\text{Kr}(\beta^-)^{93}\text{Rb}$		8700	500	8484	8	-0.4	U		Trs		78Wo15	
		8600	100			-1.2	U		Bwg		87Gr.A	
$^{93}\text{Rb}(\beta^-)^{93}\text{Sr}$		7560	120	7466	9	-0.8	U		Trs		78Wo15	
		7488	15			-1.5	o		Gsn		80Bl.A	
		7485	15			-1.3	o		Gsn		80De02	
		7440	30			0.9	-		McG		83Ia02	
		7455	35			0.3	-		Bwg		87Gr.A	
		7456	15			0.7	-		Gsn		92Pr03	
	ave.	7453	13			1.0	1	50	$^{26}\text{Rb}$		average	
$^{93}\text{Sr}(\beta^-)^{93}\text{Y}$		4130	100	4141	12	0.1	U		Bwg		78St02	
		4110	20			1.6	1	34	$^{24}\text{Y}$	McG	83Ia02	
$^{93}\text{Y}(\beta^-)^{93}\text{Zr}$		2890	20	2895	10	0.2	-				59Kn38	
		2880	15			1.0	-		McG		83Ia02	
	ave.	2884	12			0.9	1	76	$^{76}\text{Y}$		average	
$^{93}\text{Zr}(\beta^-)^{93}\text{Nb}$		93.8	2.	90.8	1.5	-1.5	1	55	$^{53}\text{Nb}$		53Gl.A	
$^{93}\text{Mo}(\epsilon)^{93}\text{Nb}$		158	15	405.8	1.5	16.5	B				64Ho08	
$^{93}\text{Nb}(p,n)^{93}\text{Mo}$		-1188	10	-1188.1	1.5	0.0	U				68Fi01	
		-1190	5			0.4	U				75Ch05	
$^{93}\text{Tc}(\beta^+)^{93}\text{Mo}$		3185.1	5.	3201.0	1.0	3.2	B				51Bo48	
		3192.1	3.			3.0	B				74An24	
$^{93}\text{Ru}(\beta^+)^{93}\text{Tc}$		6337	85	6389.4	2.3	0.6	U				83Ay01	
$^{93}\text{Mo}-u$	$M - A = -84385(28)$ keV for $^{93}\text{Mo}^m$ at 2424.95 keV										Ens115	**
$^{93}\text{Zr}(\beta^-)^{93}\text{Nb}$	$E_{\beta^-} = 63(2)$ to $1/2^-$ level at 30.77 keV										Ens115	**
$^{93}\text{Mo}(\epsilon)^{93}\text{Nb}$	$L/K = 0.36(0.04)$ to $1/2^-$ level at 30.82 keV, recalculated $Q$										Ens115	**
$^{93}\text{Tc}(\beta^+)^{93}\text{Mo}$	$E_{\beta^+} = 800(5)$ 807(3) respectively, to $7/2^+$ level 1363.048 keV										Ens115	**
$^{94}\text{Br}-u$		-50242	429	-50890#	320#	-0.6	D		GT3	2.5	16Kn03	*
$^{94}\text{Kr}-u$		-66238	247	-65860	13	1.0	U		GT1	1.5	04Ma.A	
$^{94}\text{Kr}-^{85}\text{Rb}_{1.106}$		31701	13				2		MA8	1.0	06De36	
		31665	24	31701	13	1.5	U		MA8	1.0	10Na13	
		31649	97			0.5	U		MA9	1.0	10Na13	*
$^{94}\text{Rb}-u$		-73602	54	-73605.2	2.2	0.0	F		Pb1	2.5	89Al33	*
$^{94}\text{Rb}-^{85}\text{Rb}_{1.106}$		23958	10	23955.4	2.2	-0.3	U		MA4	1.0	02Ra23	
		23955.6	2.6			-0.1	1	70	$^{70}\text{Rb}$	1.0	12Si10	*
$^{94}\text{Sr}-^{85}\text{Rb}_{1.106}$		12924	10	12916.2	1.8	-0.8	U		MA4	1.0	02Ra23	
		12916.0	1.8			0.1	1	98	$^{98}\text{Sr}$	1.0	12Si10	*
$C_7\text{H}_{10}-^{94}\text{Zr}$		171929.4	3.9	171937.80	0.18	0.9	U		M15	2.5	63Ri07	
$^{94}\text{Zr}-u$		-93687.34	0.20	-93687.48	0.18	-0.7	1	77	$^{77}\text{Zr}$	1.0	15Gu09	
$^{94}\text{Mo}-^{85}\text{Rb}_{1.106}$		2645.6	1.0	2644.14	0.15	-1.5	U		JY1	1.0	12Ka13	
$C_7\text{H}_{10}-^{94}\text{Mo}$		173159.6	3.2	173166.73	0.15	0.9	U		M15	2.5	63Ri07	
$^{94}\text{Mo}-u$		-94916.31	0.42	-94916.41	0.15	-0.2	1	13	$^{13}\text{Mo}$	1.0	15Gu09	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{94}\text{Tc}-u$	-90362	39	-90348	4	0.4	U			GS2	1.0	05Li24 *
$^{94}\text{Ru}-^{85}\text{Rb}_{1.106}$	8891	25	8903	3	0.5	U			SH1	1.0	08We10
	8907.1	4.5			-0.8	1	56	56 $^{94}\text{Ru}$	JY1	1.0	08We10
$^{94}\text{Ru}-\text{C}_7 \text{H}_{10}$	-166912.2	5.1	-166907	3	0.9	1	44	44 $^{94}\text{Ru}$	CP1	1.0	11Fa10
$^{94}\text{Rh}-^{85}\text{Rb}_{1.106}$	19291.2	4.6	19291	4	0.0	1	62	62 $^{94}\text{Rh}$	JY1	1.0	08We10
$^{94}\text{Rh}-\text{C}_7 \text{H}_{10}$	-156520.2	5.9	-156520	4	0.1	1	38	38 $^{94}\text{Rh}$	CP1	1.0	11Fa10
$^{94}\text{Rh}_{989}-\text{C}_7 \text{H}_9$	-147834	30	-147834	4	0.0	U			CP1	1.0	11Fa10
$^{94}\text{Kr}-^{86}\text{Kr}_{1.093}$	31710	110	31843	13	1.2	U			MA9	1.0	10Na13
$^{94}\text{Zr}-^{87}\text{Rb}_{1.080}$	4397.13	0.37	4397.55	0.18	1.1	1	23	23 $^{94}\text{Zr}$	MS1	1.0	15Gu09
$^{94}\text{Mo}-^{87}\text{Rb}_{1.080}$	3168.68	0.35	3168.62	0.15	-0.2	1	19	19 $^{94}\text{Mo}$	MS1	1.0	15Gu09
$^{94}\text{Rb}-^{88}\text{Rb}_{1.068}$	21109.1	4.0	21109.8	2.2	0.2	1	30	30 $^{94}\text{Rb}$	JY1	1.0	07Ra23
$^{94}\text{Zr}^{35}\text{Cl}-^{92}\text{Zr}^{37}\text{Cl}$	4235.0	2.	4227.31	0.22	-1.0	U			H13	4.0	63Ba20
$^{94}\text{Mo}^{35}\text{Cl}-^{92}\text{Mo}^{37}\text{Cl}$	1234.0	2.	1226.55	0.24	-0.9	U			H11	4.0	63Bi12
$^{94}\text{Pd}-^{94}\text{Mo}$	23952.7	4.6				2			JY1	1.0	08We10
$^{92}\text{Rb}-^{94}\text{Rb}_{.587} \text{ } ^{89}\text{Rb}_{.413}$	-764	24	-779	7	-0.2	U			P21	2.5	82Au01 Y
$^{92}\text{Rb}-^{94}\text{Rb}_{.489} \text{ } ^{90}\text{Rb}_{.511}$	-717	23	-726	9	-0.2	U			P21	2.5	82Au01 Y
$^{93}\text{Rb}-^{94}\text{Rb}_{.742} \text{ } ^{90}\text{Rb}_{.258}$	-1296	25	-1289	9	0.1	U			P21	2.5	82Au01 Y
$^{93}\text{Rb}-^{94}\text{Rb}_{.495} \text{ } ^{92}\text{Rb}_{.505}$	-840	40	-921	8	-0.8	F			P31	2.5	86Au02 *
$^{94}\text{Zr}(d,\alpha)^{92}\text{Y}$	8278	25	8258	9	-0.8	1	13	13 $^{92}\text{Y}$	Gm		74Gi09
$^{94}\text{Zr}(p,t)^{92}\text{Zr}$	-6466	12	-6471.13	0.19	-0.4	U			Bld		66St15
	-6470	10			-0.1	U			Oak		71Ba43
$^{94}\text{Ag}^n(2p)^{92}\text{Rh}$	3449	100	2500#	300#	-9.5	F					06Mu03 *
$^{94}\text{Rb}(\beta^-n)^{93}\text{Sr}$	3580	80	3452	8	-1.6	U					84Kr.B
$^{94}\text{Zr}(p,d)^{93}\text{Zr}$	-5983	15	-5994.0	0.5	-0.7	U			Bld		66St15
	-6000	10			0.6	U			Oak		71Ba43
$^{94}\text{Zr}(d,t)^{93}\text{Zr}$	-1969	20	-1961.4	0.5	0.4	U			Pit		64Co11
	-1960.2	2.4			-0.5	U			SPa		79Bo37
$^{93}\text{Nb}(n,\gamma)^{94}\text{Nb}$	7229.13	0.12	7227.54	0.08	-13.2	C					84Bo.C
	7227.51	0.09			0.3	-			MMn		88Ke09 Z
	7227.63	0.15			-0.6	-			Bdn		06Fi.A
ave.	7227.54	0.08			0.0	1	100	69 $^{94}\text{Nb}$			average
$^{94}\text{Mo}(d,t)^{93}\text{Mo}$	-3441	20	-3421.08	0.23	1.0	U			Pit		64Co11
$^{94}\text{Ag}^n(p)^{93}\text{Pd}$	5780	30	5790	17	0.3	4					05Mu15 *
	5794	20			-0.2	4					09Ce04 *
$^{94}\text{Rb}(\beta^-)^{94}\text{Sr}$	10304	20	10282.9	2.6	-1.1	o			Gsn		80Bl.A
	10322	100			-0.4	o			Gsn		80De02 *
	10353	140			-0.5	U			Trs		82Br23 *
	10335	45			-1.2	U			Bwg		82Pa24 *
	10312	20			-1.5	U			Gsn		92Pr03 *
$^{94}\text{Sr}(\beta^-)^{94}\text{Y}$	3512	10	3506	6	-0.6	1	41	40 $^{94}\text{Y}$	Gsn		80De02 *
$^{94}\text{Y}(\beta^-)^{94}\text{Zr}$	4920	9	4918	6	-0.2	1	50	50 $^{94}\text{Y}$	Gsn		80De02 *
$^{94}\text{Nb}(\beta^-)^{94}\text{Mo}$	2043.3	6.	2045.0	1.5	0.3	-					66Sn02 *
	2046.3	3.			-0.4	-					68Ho10 *
ave.	2045.7	2.7			-0.3	1	31	31 $^{94}\text{Nb}$			average
$^{94}\text{Tc}(\beta^+)^{94}\text{Mo}$	4261	5	4256	4	-1.1	2					64Ha29 *
$^{94}\text{Mo}(p,n)^{94}\text{Tc}$	-5027.8	7.	-5038	4	-1.5	2					73Mc04 *
$^{94}\text{Rh}(\beta^+)^{94}\text{Ru}$	9930	400	9676	5	-0.6	U					80Ox01 *
	9750	320			-0.2	U					06Ba55
$^{94}\text{Pd}(\beta^+)^{94}\text{Rh}$	6700	320	6805	5	0.3	U					06Ba55
$^{94}\text{Ag}^n(\beta^+)^{94}\text{Pd}$	17700	500	20180#	300#	5.0	D					04Mu30 *
* $^{94}\text{Br}-u$	Trends from Mass Surface TMS suggest $^{94}\text{Br}$ 600 more bound GAu **										
* $^{94}\text{Kr}-^{85}\text{Rb}_{1.106}$	Typo in original paper, ratio should read 1.006 255 64(1 14) GAu **										
* $^{94}\text{Rb}-u$	F : possibly isomeric mixture 92Al.B **										
* $^{94}\text{Rb}-^{85}\text{Rb}_{1.106}$	$D_M=23956.4(2.6) \mu\text{u}$ $M-A=-68561.8(2.4)\text{keV}$ corrected for $e^-$ binding= $-775\text{eV}$ 12Si10 **										
* $^{94}\text{Sr}-^{85}\text{Rb}_{1.106}$	$D_M=12916.7(1.8) \mu\text{u}$ $M-A=-78845.2(1.7)\text{keV}$ corrected for $e^-$ binding= $-625\text{eV}$ 12Si10 **										
* $^{94}\text{Tc}-u$	$M-A=-84133(29) \text{keV}$ for mixture gs+m at 76(3) keV Nub16b **										
* $^{93}\text{Rb}-^{94}\text{Rb}_{.495} \text{ } ^{92}\text{Rb}_{.505}$	F : rejection based on line-shape analysis 86Au02 **										
* $^{94}\text{Ag}^n(2p)^{92}\text{Rh}$	$Q_{2p}=1900(100) \text{ to } (11^+) \text{ level at } 1548.6 \text{ keV}$ Ens12a **										
* $^{94}\text{Ag}^n(2p)^{92}\text{Rh}$	F : no evidence from He-jet experiment 09Ce04 **										
* $^{94}\text{Ag}^n(2p)^{92}\text{Rh}$	F : possibly misidentified $^{92}\text{Rh}$ $\gamma$ rays 09Je05 **										
* $^{94}\text{Ag}^n(p)^{93}\text{Pd}$	$E_p=790(30), 1010(30) \text{ to } (33/2^+) \text{ at } 4995.6, (33/2^-, 35/2^-) \text{ at } 4752.7$ Ens115 **										
* $^{94}\text{Ag}^n(p)^{93}\text{Pd}$	$E_p=790(20) \text{ to level } (33/2^+) \text{ at } 4995.6 \text{ keV}$ Ens115 **										



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>94</sup> Rb( $\beta^-$ ) <sup>94</sup> Sr	Original value 10304(30) corrected in reference										94Ha.A **
* <sup>94</sup> Rb( $\beta^-$ ) <sup>94</sup> Sr	Original error 100 keV increased by 100 in reference for lower level feeding										94Ha.A **
* <sup>94</sup> Rb( $\beta^-$ ) <sup>94</sup> Sr	As corrected in reference										87Gr.A **
* <sup>94</sup> Rb( $\beta^-$ ) <sup>94</sup> Sr	$E_{\beta^-}$ =9475(20) to 2 <sup>+</sup> level at 836.91 keV										Ens116 **
* <sup>94</sup> Sr( $\beta^-$ ) <sup>94</sup> Y	Original error 6 corrected in reference										94Ha.A **
* <sup>94</sup> Y( $\beta^-$ ) <sup>94</sup> Zr	Original error 5 corrected in reference										94Ha.A **
* <sup>94</sup> Nb( $\beta^-$ ) <sup>94</sup> Mo	$E_{\beta^-}$ =470(6) 473(3) respectively, to level 4 <sup>+</sup> at 1573.76 keV										Ens069 **
* <sup>94</sup> Tc( $\beta^+$ ) <sup>94</sup> Mo	$E_{\beta^+}$ =816(5) to 6 <sup>+</sup> level at 2423.45 keV										Ens069 **
* <sup>94</sup> Mo(p,n) <sup>94</sup> Tc	T=5158(7) to <sup>94</sup> Tc <sup>m</sup> at 76(3) keV										Nub16b **
* <sup>94</sup> Rh( $\beta^+$ ) <sup>94</sup> Ru	$E_{\beta^+}$ =6400(400) to (3,4,5) level at 2503.2 keV										Ens069 **
* <sup>94</sup> Ag <sup>n</sup> ( $\beta^+$ ) <sup>94</sup> Pd	$Q_{\beta^+}$ larger than 17.7 MeV, uncertainty not given										04Mu30 **
* <sup>94</sup> Ag <sup>n</sup> ( $\beta^+$ ) <sup>94</sup> Pd	Trends from Mass Surface TMS suggest <sup>94</sup> Ag <sup>n</sup> 2480 less bound										GAu **
<sup>95</sup> Kr-u	-60183	150	-60289	20	-0.5	U			GT1	1.5	04Ma.A
<sup>95</sup> Kr- <sup>85</sup> Rb <sub>1.118</sub>	38330	20				2			MA8	1.0	06De36
<sup>95</sup> Rb-u	-70618	86	-70737	22	-0.6	U			Pb1	2.5	89Al33
<sup>95</sup> Sr- <sup>85</sup> Rb <sub>1.118</sub>	17987	10	17975	6	-1.2	1	39	39 <sup>95</sup> Sr	MA4	1.0	02Ra23
<sup>95</sup> Mo- <sup>85</sup> Rb <sub>1.118</sub>	4457.6	1.0	4456.52	0.13	-1.1	U			JY1	1.0	12Ka13
C <sub>7</sub> H <sub>11</sub> - <sup>95</sup> Mo	180236.5	3.5	180237.91	0.13	0.2	U			M15	2.5	63Ri07
<sup>95</sup> Mo-u	-94161.96	0.38	-94162.56	0.13	-1.6	1	12	12 <sup>95</sup> Mo	MS1	1.0	15Gu09
<sup>95</sup> Tc-u	-92417	32	-92348	5	2.2	U			GS2	1.0	05Li24 *
<sup>95</sup> Rh- <sup>85</sup> Rb <sub>1.118</sub>	14515.1	4.5	14517	4	0.4	1	86	86 <sup>95</sup> Rh	JY1	1.0	08We10
<sup>95</sup> Rh <sub>989</sub> -C <sub>7</sub> H <sub>10</sub>	-161416	11	-161427	4	-1.0	1	14	14 <sup>95</sup> Rh	CP1	1.0	11Fa10
<sup>95</sup> Mo- <sup>87</sup> Rb <sub>1.092</sub>	5012.33	0.44	5012.30	0.13	-0.1	U			MS1	1.0	15Gu09
<sup>95</sup> Sr- <sup>97</sup> Zr <sub>979</sub>	6529	10	6529	6	0.0	1	39	39 <sup>95</sup> Sr	JY1	1.0	06Ha03
<sup>95</sup> Y- <sup>97</sup> Zr <sub>979</sub>	-32.4	6.7	-9	7	3.6	B			JY1	1.0	07Ha32
<sup>95</sup> Pd- <sup>94</sup> Mo <sub>1.011</sub>	20848.9	4.7	20849	3	0.0	2			JY1	1.0	08We10
	20849.1	4.5			0.0	2			JY1	1.0	08We10 *
<sup>95</sup> Mo- <sup>94</sup> Mo	757	12	753.85	0.10	-0.1	U			M15	2.5	63Ri07
<sup>93</sup> Rb- <sup>95</sup> Rb <sub>653</sub> <sup>89</sup> Rb <sub>348</sub>	-1323	25	-1157	15	2.7	U			P21	2.5	82Au01
<sup>93</sup> Rb- <sup>95</sup> Rb <sub>587</sub> <sup>90</sup> Rb <sub>413</sub>	-1376	24	-1193	15	3.0	B			P21	2.5	82Au01
<sup>94</sup> Rb- <sup>95</sup> Rb <sub>792</sub> <sup>90</sup> Rb <sub>209</sub>	-16	28	196	16	3.0	B			P21	2.5	82Au01 Y
<sup>92</sup> Rb- <sup>95</sup> Rb <sub>242</sub> <sup>91</sup> Rb <sub>758</sub>	80	23	104	10	0.4	U			P21	2.5	82Au01
<sup>93</sup> Rb- <sup>95</sup> Rb <sub>489</sub> <sup>91</sup> Rb <sub>511</sub>	-654	12	-671	13	-0.6	F			P31	2.5	86Au02 *
<sup>94</sup> Rb- <sup>95</sup> Rb <sub>660</sub> <sup>92</sup> Rb <sub>341</sub>	433	15	423	14	-0.3	1	13	13 <sup>95</sup> Rb	P31	2.5	86Au02
	462	28			-0.6	U			P31	2.5	86Au02
<sup>95</sup> Mo(n, $\alpha$ ) <sup>92</sup> Zr	6405	30	6393.57	0.16	-0.4	U			ILL		75Em04
<sup>95</sup> Rb( $\beta^-$ -n) <sup>94</sup> Sr	5120	100	4883	20	-2.4	U					84Kr.B
<sup>94</sup> Zr(n, $\gamma$ ) <sup>95</sup> Zr	6461.6	1.0	6461.9	0.9	0.3	-					79Ke.D Z
	6357.8	0.3			347.1	F			Bdn		06Fi.A *
	6460.3	0.5			3.3	C			Bdn		08Fi.A *
<sup>94</sup> Zr(d,p) <sup>95</sup> Zr	4223	20	4237.4	0.9	0.7	U			Pit		64Co11
	4237.4	2.0			0.0	-			SPa		79Bo37
<sup>94</sup> Zr(n, $\gamma$ ) <sup>95</sup> Zr	ave.	6461.7	0.9	6461.9	0.9	0.3	1	92	91 <sup>95</sup> Zr		average
<sup>94</sup> Mo(n, $\gamma$ ) <sup>95</sup> Mo		7367	2	7369.11	0.09	1.1	U				77Ri04
		7369.10	0.10			0.1	1	89	68 <sup>94</sup> Mo	MMn	91Is02 Z
		7368.4	0.5			1.4	U		Bdn		06Fi.A
<sup>94</sup> Mo(d,p) <sup>95</sup> Mo		5137	20	5144.54	0.09	0.4	U		Pit		64Co11
<sup>95</sup> Pd(ep) <sup>94</sup> Ru		5116	300	5329	4	0.7	U				82Ku15 *
<sup>95</sup> Rb( $\beta^-$ ) <sup>95</sup> Sr		9224	30	9228	20	0.1	o		Gsn		80Bl.A
		9276	100			-0.5	o		Gsn		80De02 *
		9300	30			-2.4	C		Gsn		84Bl.A
		9280	45			-1.2	-		Bwg		87Gr.A
		9272	35			-1.3	-		Gsn		92Pr03
	ave.	9275	28			-1.7	1	53	51 <sup>95</sup> Rb		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{95}\text{Sr}(\beta^-)^{95}\text{Y}$	6110	150	6089	7	-0.1	U					70Ma.A	
	6060	100			0.3	U			Bwg		78St02	
	6082	10			0.7	1	52	32	$^{95}\text{Y}$	Gsn	84Bl.A	
	6052	25			1.5	U					90Ma03	
$^{95}\text{Y}(\beta^-)^{95}\text{Zr}$	4445	9	4451	7	0.7	1	57	56	$^{95}\text{Y}$	Gsn	80De02 *	
$^{95}\text{Zr}(\beta^-)^{95}\text{Nb}$	1125	8	1126.3	1.0	0.2	U					54Za05	
	1119	5			1.5	U					55Dr43	
	1122.7	3.			1.2	1	11	8	$^{95}\text{Zr}$		74An22 *	
$^{95}\text{Nb}(\beta^-)^{95}\text{Mo}$	925.5	0.5	925.6	0.5	0.2	1	98	97	$^{95}\text{Nb}$		63La06 *	
$^{95}\text{Tc}(\beta^+)^{95}\text{Mo}$	1683	10	1691	5	0.8	-					65Cr04 *	
	1693	6			-0.4	-					74An05 *	
$^{95}\text{Mo}(\text{p,n})^{95}\text{Tc}$	-2440	30	-2473	5	-1.1	U					57Le27	
	-2490	6			2.9	B			Oak		70Ki01	
$^{95}\text{Tc}(\beta^+)^{95}\text{Mo}$	ave.	1690	1691	5	0.0	1	97	97	$^{95}\text{Tc}$		average	
$^{95}\text{Ru}(\beta^+)^{95}\text{Tc}$	2558	30	2564	11	0.2	1	12	10	$^{95}\text{Ru}$		68Pi03 *	
$^{95}\text{Rh}(\beta^+)^{95}\text{Ru}$	5110	150	5117	10	0.0	U					75We03 *	
$^{95}\text{Tc}-\text{u}$	$M - A = -86066(28)$ keV for mixture gs+m at 38.91 keV										Nub16b **	
$^{95}\text{Pd}-^{94}\text{Mo}_{1.011}$	$D_M = 22862.1(4.5)$ $\mu\text{u}$ for $^{95}\text{Pd}^m$ at 1875.13 keV; $M - A = -68090.2(4.4)$ keV										Nub16b **	
$^{93}\text{Rb}-^{95}\text{Rb}_{.489}$ $^{91}\text{Rb}_{.511}$	F : Rejected by authors										86Au02 **	
$^{94}\text{Zr}(\text{n},\gamma)^{95}\text{Zr}$	F : value from 06Fi.A retracted										08Fi.A **	
$^{94}\text{Zr}(\text{n},\gamma)^{95}\text{Zr}$	Weak evidence										08Fi.A **	
$^{95}\text{Pd}(\text{e}p)^{94}\text{Ru}$	$E_p = 4300(300)$ from $^{95}\text{Pd}^m$ at 1875.13 to $^{94}\text{Ru}^m$ at 2644.1 keV										Nub16b **	
*	same $E_p$ ; both from figures										82No06 **	
$^{95}\text{Rb}(\beta^-)^{95}\text{Sr}$	$E_{\beta^-} = 8595(100)$ to $(3/2^+, 5/2^+)$ level at 680.70, corrected in reference										92Pr03 **	
$^{95}\text{Y}(\beta^-)^{95}\text{Zr}$	Original error 5 corrected in reference										94Ha.A **	
*	$Q_{\beta^-} = 4417(10)$ given by same group, not used										84Bl.A **	
$^{95}\text{Zr}(\beta^-)^{95}\text{Nb}$	$E_{\beta^-} = 887(3)$ to $1/2^-$ level at 235.69 keV										Ens10a **	
$^{95}\text{Nb}(\beta^-)^{95}\text{Mo}$	$E_{\beta^-} = 159.7(0.5)$ to $7/2^+$ level at 765.803 keV										Ens10a **	
$^{95}\text{Tc}(\beta^+)^{95}\text{Mo}$	$E_{\beta^+} = 700(10)$ $710(6)$ respectively, from $^{95}\text{Tc}^m$ at 38.91 keV										Nub16b **	
$^{95}\text{Ru}(\beta^+)^{95}\text{Tc}$	$E_{\beta^+} = 1200(30)$ to $7/2^+$ level at 336.413 keV										Ens10a **	
$^{95}\text{Rh}(\beta^+)^{95}\text{Ru}$	$E_{\beta^+} = 3150(150)$ to $7/2^+$ level at 941.79 keV										Ens10a **	
$^{96}\text{Kr}-^{85}\text{Rb}_{1.129}$	42606	22							MA8	1.0	10Na13	
$^{96}\text{Rb}-\text{u}$	-65508	43	-65867	4	-3.3	F			Pb1	2.5	89Al33 *	
$^{96}\text{Zr}-^{87}\text{Rb}_{1.103}$	8451.49	0.34	8451.50	0.12	0.0	1	13	13	$^{96}\text{Zr}$	MS1	1.0	15Gu09
$\text{C}_7 \text{H}_{12}-^{96}\text{Zr}$	185628	6	185622.77	0.12	-0.3	U			M15	2.5	63Ri07	
$^{96}\text{Zr}-\text{u}$	-91691	43	-91722.38	0.12	-0.7	U			JY0	1.0	04Ri12	
	-91722.60	0.17			1.3	1	52	52	$^{96}\text{Zr}$	MS1	1.0	15Gu09
$^{96}\text{Mo}-^{85}\text{Rb}_{1.129}$	4265.7	1.1	4264.16	0.13	-1.4	U			JY1	1.0	12Ka13	
$^{96}\text{Mo}-^{87}\text{Rb}_{1.103}$	4848.96	0.43	4848.65	0.13	-0.7	U			MS1	1.0	15Gu09	
$\text{C}_7 \text{H}_{12}-^{96}\text{Mo}$	189226.9	3.0	189225.61	0.13	-0.2	U			M15	2.5	63Ri07	
$^{96}\text{Mo}-\text{u}$	-95324.94	0.47	-95325.23	0.13	-0.6	U			MS1	1.0	15Gu09	
$^{96}\text{Tc}-\text{u}$	-92192	32	-92133	6	1.8	U			GS2	1.0	05Li24 *	
$\text{C}_7 \text{H}_{12}-^{96}\text{Ru}$	186304.6	3.8	186311.47	0.18	0.7	U			M16	2.5	63Da10	
$^{96}\text{Rb}-^{88}\text{Rb}_{1.091}$	30887.7	3.6	30888	4	0.1	1	100	100	$^{96}\text{Rb}$	JY1	1.0	07Ra23
$^{96}\text{Zr}-^{35}\text{Cl}-^{94}\text{Zr}-^{37}\text{Cl}$	4929	3	4915.21	0.22	-1.1	U			H13	4.0	63Ba20	
$^{96}\text{Mo}-^{35}\text{Cl}-^{94}\text{Mo}-^{37}\text{Cl}$	2539	2	2541.29	0.13	0.3	U			H11	4.0	63Bi12	
$^{96}\text{Pd}-^{94}\text{Mo}_{1.021}$	15123.4	4.5				2			JY1	1.0	08We10	
$^{96}\text{Sr}-^{97}\text{Zr}_{.990}$	9868	10	9865	9	-0.3	1	83	83	$^{96}\text{Sr}$	JY1	1.0	06Ha03
$^{96}\text{Y}-^{97}\text{Zr}_{.990}$	4053.7	6.8	4055	7	0.2	1	92	92	$^{96}\text{Y}$	JY1	1.0	07Ha32
$^{96}\text{Y}^m-^{97}\text{Zr}_{.990}$	5708.1	6.7				2			JY1	1.0	07Ha32	
$^{96}\text{Mo}-^{97}\text{Mo}_{.990}$	-2280.5	5.8	-2281.96	0.17	-0.3	U			JY1	1.0	06Ka48	
$^{96}\text{Zr}-^{96}\text{Nb}$	176.02	0.13	176.03	0.11	0.1	1	68	63	$^{96}\text{Nb}$	JY1	1.0	16Al03
$^{96}\text{Zr}-^{96}\text{Mo}$	3602.919	0.092	3602.85	0.08	-0.8	1	75	46	$^{96}\text{Mo}$	JY1	1.0	16Al03
$^{96}\text{Nb}-^{96}\text{Mo}$	3426.80	0.17	3426.82	0.11	0.1	1	46	37	$^{96}\text{Nb}$	JY1	1.0	16Al03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{96}\text{Ru}-^{96}\text{Mo}$	2914.14	0.13	2914.14	0.13	0.0	1	100	100 $^{96}\text{Ru}$	SH1	1.0	11El04
$^{96}\text{Mo}-^{95}\text{Mo}$	-1161	12	-1162.67	0.05	-0.1	U			M15	2.5	63Ri07
$^{93}\text{Rb}-^{96}\text{Rb}_{.554}$ $^{89}\text{Rb}_{.448}$	-2210	27	-2022	8	2.8	U			P21	2.5	82Au01
$^{95}\text{Rb}-^{96}\text{Rb}_{.848}$ $^{89}\text{Rb}_{.152}$	-1590	30	-1443	20	2.0	U			P21	2.5	82Au01
$^{94}\text{Rb}-^{96}\text{Rb}_{.699}$ $^{89}\text{Rb}_{.302}$	-1250	30	-999	4	3.3	B			P21	2.5	82Au01 Y
$^{94}\text{Rb}-^{96}\text{Rb}_{.588}$ $^{91}\text{Rb}_{.413}$	-380	25	-378	4	0.0	U			P21	2.5	82Au01
$^{95}\text{Rb}-^{96}\text{Rb}_{.742}$ $^{92}\text{Rb}_{.258}$	-1116	27	-1075	20	0.6	U			P21	2.5	82Au01
	-1143	16			1.7	1	26	26 $^{95}\text{Rb}$	P31	2.5	86Au02
$^{96}\text{Zr}(d,\alpha)^{94}\text{Y}$	7609	20	7623	6	0.7	1	10	10 $^{94}\text{Y}$	Grn		74Gi09
$^{96}\text{Zr}(p,t)^{94}\text{Zr}$	-5825	10	-5830.36	0.20	-0.5	U			Oak		71Ba43
$^{96}\text{Ru}(p,t)^{94}\text{Ru}$	-11165	10	-11158	3	0.7	U			Oak		71Ba01
$^{96}\text{Zr}(t,\alpha)^{95}\text{Y}$	8294	20	8295	7	0.0	1	11	11 $^{95}\text{Y}$	LAl		83Fl06
$^{96}\text{Zr}(p,d)^{95}\text{Zr}$	-5440	20	-5625.7	0.9	-9.3	B			Bld		67St24
	-5630	10			0.4	U			Oak		71Ba43
$^{96}\text{Zr}(d,t)^{95}\text{Zr}$	-1603	20	-1593.0	0.9	0.5	U			Pit		64Co11
	-1595.8	2.8			1.0	U			SPa		79Bo37
$^{96}\text{Mo}(t,\alpha)^{95}\text{Nb}$	10524	20	10516.3	0.5	-0.4	U			LAl		83Fl06
$^{95}\text{Mo}(n,\gamma)^{96}\text{Mo}$	9154.2	0.5	9154.33	0.05	0.3	U					70He27
	9154.32	0.05			0.3	1	96	67 $^{95}\text{Mo}$	MMn		91Is02 Z
	9153.90	0.20			2.2	U			Bdn		06Fi.A
$^{96}\text{Mo}(d,t)^{95}\text{Mo}$	-2923	20	-2897.11	0.05	1.3	U			Pit		64Co11
$^{96}\text{Ru}(p,d)^{95}\text{Ru}$	-8470	10	-8469	10	0.1	1	90	90 $^{95}\text{Ru}$	Oak		71Ba01
$^{96}\text{Ag}(\epsilon p)^{95}\text{Rh}$	6540	90				2					03Ba39 *
$^{96}\text{Rb}(\beta^-)^{96}\text{Sr}$	10800	220	11570	9	3.5	B					79Pe17
	11303	250			1.1	o			Gsn		80De02
	11547	100			0.2	U			Trs		82Br23
	11553	45			0.4	U			Gsn		84Bl.A
	11590	80			-0.3	U			Bwg		87Gr.A
	11709	40			-3.5	B			Gsn		92Pr03 *
$^{96}\text{Sr}(\beta^-)^{96}\text{Y}$	5332	30	5412	10	2.7	F					79Pe17 *
	5413	22			-0.1	-			Gsn		80De02 *
	5345	50			1.3	U			Bwg		87Gr.A
	5354	40			1.4	-					90Ma03
	ave.	5399	19		0.6	1	25	17 $^{96}\text{Sr}$			average
$^{96}\text{Y}(\beta^-)^{96}\text{Zr}$	7120	50	7103	6	-0.3	U			Gsn		80De02 *
	7030	70			1.0	U			Bwg		87Gr.A
	7067	30			1.2	U					90Ma03 *
$^{96}\text{Y}^m(\beta^-)^{96}\text{Zr}$	8030	150	8643	6	4.1	C			Bwg		87Gr.A
	8600	200			0.2	U					88St.A
	8237	21			19.3	C			Bwg		92Gr.A
$^{96}\text{Nb}(\beta^-)^{96}\text{Mo}$	3186.8	3.2	3192.06	0.11	1.6	U					68An03 *
$^{96}\text{Mo}(p,n)^{96}\text{Tc}$	-3760	10	-3756	5	0.4	2					74Do09
	-3754	6			-0.3	2					78Ke10
$^{96}\text{Rh}(\beta^+)^{96}\text{Ru}$	6472	200	6393	10	-0.4	U					75Gu01 *
$^{96}\text{Ru}(p,n)^{96}\text{Rh}$	-7175	10				2					70As08 Z
$^{96}\text{Pd}(\beta^+)^{96}\text{Rh}$	3450	150	3504	11	0.4	U					85Ry02 *
$^{96}\text{Rb}-u$	F : possibly isomeric mixture										
$^{96}\text{Tc}-u$	$M - A = -85860(28)$ keV for mixture gs+m at 34.23 keV										
$^{96}\text{Ag}(\epsilon p)^{95}\text{Rh}$	Original 6430(60) corrected by -110 keV for mixture of two $\beta^-$ -decaying $^{96}\text{Ag}$ states, to two isomeric states in $^{95}\text{Rh}$										
$^{96}\text{Rb}(\beta^-)^{96}\text{Sr}$	$E_{\beta^-} = 10894(40)$ to $2^+$ level at 814.93 keV										
$^{96}\text{Sr}(\beta^-)^{96}\text{Y}$	$E_{\beta^-} = 4400(30)$ to $1^+$ level at 931.70 keV, and other $E_{\beta^-}$										
$^{96}\text{Sr}(\beta^-)^{96}\text{Y}$	F : all other results of reference are strongly conflicting										
$^{96}\text{Sr}(\beta^-)^{96}\text{Y}$	Original error 20 corrected in reference										
$^{96}\text{Y}(\beta^-)^{96}\text{Zr}$	$Q_{\beta^-} = 5362(10)$ given by same group, not used										
$^{96}\text{Y}(\beta^-)^{96}\text{Zr}$	$Q_{\beta^-} = 7079(15)$ given by same group, not used										
$^{96}\text{Y}(\beta^-)^{96}\text{Zr}$	$E_{\beta^-} = 5326(36)$ to $2^+$ level at 1750.497 keV, and other $E_{\beta^-}$										
$^{96}\text{Nb}(\beta^-)^{96}\text{Mo}$	$E_{\beta^-} = 748.4(3.1)$ to $5^+$ level at 2438.477 keV										
$^{96}\text{Rh}(\beta^+)^{96}\text{Ru}$	$E_{\beta^+} = 3300(200)$ to $6^+$ level at 2149.74 keV										
$^{96}\text{Pd}(\beta^+)^{96}\text{Rh}$	$p^+ = 0.257(0.03)$ to $1^+$ level at 1274.78 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{97}\text{Kr}-^{86}\text{Kr}_{1.128}$	49920	140					MA9	1.0	10Na13
$^{97}\text{Rb}-u$	-62512	64	-62822.9	2.1	-1.9	U	Pb1	2.5	89Al33
$^{97}\text{Rb}-^{88}\text{Rb}_{1.102}$	34908.0	5.7	34907.3	2.1	-0.1	1	13 $^{97}\text{Rb}$	1.0	07Ra23
$^{97}\text{Rb}-^{85}\text{Rb}_{1.141}$	37824.9	2.2	37825.0	2.1	0.1	1	87 $^{97}\text{Rb}$	1.0	12Si10 *
$^{97}\text{Sr}-^{85}\text{Rb}_{1.141}$	27022.9	3.9	27023	4	-0.1	1	87 $^{97}\text{Sr}$	1.0	12Si10 *
$^{97}\text{Sr}-u$	-73599	99	-73625	4	-0.1	U	GT2	2.5	08Kn.A
$^{97}\text{Mo}-^{85}\text{Rb}_{1.141}$	6666.9	1.2	6664.82	0.18	-1.7	U	JY1	1.0	12Ka13
$^{97}\text{Mo}-^{87}\text{Rb}_{1.115}$	7280.86	0.39	7280.61	0.18	-0.6	1	21 $^{97}\text{Mo}$	1.0	15Gu09
$\text{C}_5 \text{H}_5 \text{O}_2 - ^{97}\text{Mo}$	122937.6	2.3	122937.50	0.18	0.0	U	M15	2.5	63Ri07
$^{97}\text{Mo}-u$	-93982.95	0.36	-93983.10	0.18	-0.4	1	24 $^{97}\text{Mo}$	1.0	15Gu09
$^{97}\text{Ru}-u$	-92471	30	-92454.2	3.0	0.6	U	GS2	1.0	05Li24
$^{97}\text{Pd}-^{85}\text{Rb}_{1.141}$	17119.9	5.2				2	JY1	1.0	09El08
$^{97}\text{Mo}-^{35}\text{Cl}-^{95}\text{Mo}-^{37}\text{Cl}$	3138	2	3129.57	0.19	-1.1	U	H11	4.0	63Bi12
$^{97}\text{Sr}-^{97}\text{Zr}$	15416	10	15417	4	0.1	1	13 $^{97}\text{Sr}$	1.0	06Ha03
$^{97}\text{Y}-^{97}\text{Zr}$	7322.9	7.2				2	JY1	1.0	07Ha32 *
$^{97}\text{Mo}-^{96}\text{Mo}$	1346	12	1342.13	0.18	-0.1	U	M15	2.5	63Ri07
$^{94}\text{Rb}-^{97}\text{Rb}_{.485} \ ^{91}\text{Rb}_{.516}$	-21	25	-65	5	-0.7	U	P21	2.5	82Au01 Y
$^{96}\text{Rb}-^{97}\text{Rb}_{.792} \ ^{92}\text{Rb}_{.209}$	650	30	620	4	-0.4	U	P21	2.5	82Au01
$^{95}\text{Rb}-^{97}\text{Rb}_{.490} \ ^{93}\text{Rb}_{.511}$	-165	25	-108	21	0.9	1	11 $^{95}\text{Rb}$	2.5	82Au01
$^{96}\text{Rb}-^{97}\text{Rb}_{.742} \ ^{93}\text{Rb}_{.258}$	848	19	803	4	-1.0	U	P31	2.5	86Au02
$^{96}\text{Zr}(n,\gamma)^{97}\text{Zr}$	5574	5	5575.1	0.4	0.2	U			77Ba33
	5575.1	0.4			0.1	1	99 $^{97}\text{Zr}$		06Fi.A
$^{96}\text{Zr}(d,p)^{97}\text{Zr}$	3338	20	3350.6	0.4	0.6	U	Pit		64Co11
$^{96}\text{Mo}(n,\gamma)^{97}\text{Mo}$	6821.1	1.0	6821.13	0.16	0.0	U			73De39
	6820	2			0.6	U			77Ri04
	6821.15	0.25			-0.1	-	MMn		91Is02 Z
	6821.5	0.4			-0.9	-	Bdn		06Fi.A
$^{96}\text{Mo}(d,p)^{97}\text{Mo}$	4582	20	4596.57	0.16	0.7	U	Pit		64Co11
$^{96}\text{Mo}(n,\gamma)^{97}\text{Mo}$	ave. 6821.25	0.21	6821.13	0.16	-0.5	1	59 $^{97}\text{Mo}$	44	average
$^{96}\text{Mo}(^3\text{He},d)^{97}\text{Tc}$	229	8	225	4	-0.5	-	ANL		74Co27
	220	8			0.6	-	Pit		74Co27
	ave. 225	6			0.1	1	53 $^{97}\text{Tc}$	53	average
$^{96}\text{Ru}(d,p)^{97}\text{Ru}$	5886	3	5886.9	2.8	0.3	2	Can		77Ho02
	5892	7			-0.7	2	ANL		77Me04
$^{97}\text{Rb}(\beta^-)^{97}\text{Sr}$	10020	50	10062	4	0.8	U	Gsn		80De02
	10450	30			-12.9	C	Gsn		84Bl.A
	10440	60			-6.3	C	Bwg		87Gr.A
	10462	40			-10.0	B	Gsn		92Pr03
$^{97}\text{Sr}(\beta^-)^{97}\text{Y}$	7452	40	7540	8	2.2	U	Gsn		84Bl.A
	7420	80			1.5	o	Bwg		87Gr.A
	7480	18			3.3	C	Bwg		92Gr.A
$^{97}\text{Y}(\beta^-)^{97}\text{Zr}$	6702	25	6821	7	4.8	C	Gsn		84Bl.A
	6640	70			2.6	o	Bwg		87Gr.A *
	6689	13			10.2	C	Bwg		92Gr.A *
$^{97}\text{Zr}(\beta^-)^{97}\text{Nb}$	2657.3	6.	2663	4	1.0	1	50 $^{97}\text{Nb}$	50	74Ra.A *
$^{97}\text{Nb}(\beta^-)^{97}\text{Mo}$	1933.1	6.	1939	4	1.0	1	50 $^{97}\text{Nb}$	50	74Ra.A *
$^{97}\text{Mo}(p,n)^{97}\text{Tc}$	-1128	9	-1103	4	2.8	B	Oak		70Ki01
	-1102	6			-0.1	1	47 $^{97}\text{Tc}$	47	74Co27
$^{97}\text{Ru}(\beta^+)^{97}\text{Tc}$	1150	100	1104	5	-0.5	U			70Ho01 *
$^{97}\text{Rh}(\beta^+)^{97}\text{Ru}$	3533	50	3520	40	-0.2	3			62Ba28 *
	3513	50			0.2	3			62Ch21 *
$^{97}\text{Pd}(\beta^+)^{97}\text{Rh}$	4790	300	4790	40	0.0	U			80Go11 *
$^{97}\text{Ag}(\beta^+)^{97}\text{Pd}$	6980	110				3			99Hu10
$*^{97}\text{Rb}-^{85}\text{Rb}_{1.141}$	$D_M=37825.7(2.2) \mu\text{u}$ $M-A=-58518.5(2.1)\text{keV}$ corrected for $e^-$ binding= $-703\text{eV}$								
$*^{97}\text{Sr}-^{85}\text{Rb}_{1.141}$	$D_M=27023.5(3.9) \mu\text{u}$ $M-A=-68580.7(3.7)\text{keV}$ corrected for $e^-$ binding= $-553\text{eV}$								
$*^{97}\text{Y}-^{97}\text{Zr}$	$D_M=8039.5(7.2) \mu\text{u}$ for $^{97}\text{Y}^m$ at 667.52(0.23) keV; $M-A=-75453.9(6.7) \text{keV}$								
$*^{97}\text{Y}(\beta^-)^{97}\text{Zr}$	$E_{\beta^-}=6645(70)$ ; and 7280(150) from $^{97}\text{Y}^m$ at 667.52 keV								
$*^{97}\text{Y}(\beta^-)^{97}\text{Zr}$	$E_{\beta^-}=6688(13)$ ; and 7361(26) from $^{97}\text{Y}^m$ at 667.52 keV								
$*^{97}\text{Zr}(\beta^-)^{97}\text{Nb}$	$E_{\beta^-}=1914(2)$ to $1/2^-$ level at 743.35 keV; error increased								
$*^{97}\text{Nb}(\beta^-)^{97}\text{Mo}$	$E_{\beta^-}=1275(2)$ to $7/2^+$ level at 658.13 keV; error increased								
$*^{97}\text{Ru}(\beta^+)^{97}\text{Tc}$	$p^+ < 0.0001$ (see fig 1), decay to $7/2^+$ level at 970.03 keV								
$*^{97}\text{Rh}(\beta^+)^{97}\text{Ru}$	$E_{\beta^+}=2090(50)$ 2070(50) respectively, to $7/2^+$ level at 421.54 keV								

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{97}\text{Pd}(\beta^+)^{97}\text{Rh}$	$E_{\beta^+}=3500(300)$ to $7/2^+$ level at 265.36 keV										Ens104 **
$^{98}\text{Rb}-^{85}\text{Rb}_{1.153}$	43331	32	43339	17	0.2	1	29	29 $^{98}\text{Rb}$	TT1	1.0	12Si10 *
$^{98}\text{Rb}-u$	-58359	29	-58368	17	-0.3	-			MA8	1.0	13Ma81
	-58370	29			0.1	-			TT1	1.0	16Kl04 *
ave.	-58365	21			-0.2	1	71	71 $^{98}\text{Rb}$			average
$^{98}\text{Rb}^m-u$	-58289	22				2			TT1	1.0	12Si10 *
$^{98}\text{Sr}-^{85}\text{Rb}_{1.153}$	30396.7	4.3	30398	3	0.4	-			TT1	1.0	12Si10 *
	30405.3	7.2			-1.0	-			TT1	1.0	16Kl04 *
ave.	30399	4			-0.2	1	88	88 $^{98}\text{Sr}$			average
$^{98}\text{Zr}-u$	-87247	43	-87265	9	-0.4	U			JY0	1.0	04Ri12
$^{98}\text{Mo}-^{85}\text{Rb}_{1.153}$	7104.1	5.7	7110.04	0.19	1.0	U			MA8	1.0	11He10
	7111.6	1.3			-1.2	U			JY1	1.0	12Ka13
$^{98}\text{Mo}-^{87}\text{Rb}_{1.126}$	7665.96	0.64	7666.33	0.19	0.6	U			MS1	1.0	15Gu09
$\text{C}_5 \text{H}_6 \text{O}_2-^{98}\text{Mo}$	131375.4	2.8	131375.82	0.19	0.1	U			M15	2.5	63Ri07
$^{98}\text{Mo}-u$	-94596.21	0.53	-94596.39	0.19	-0.3	1	12	12 $^{98}\text{Mo}$	MS1	1.0	15Gu09
$\text{C}_7 \text{H}_{14}-^{98}\text{Ru}$	204263.5	2.9	204264	7	0.0	1	92	92 $^{98}\text{Ru}$	M16	2.5	63Da10
$^{98}\text{Rh}-u$	-89302	46	-89292	13	0.2	U			GS2	1.0	05Li24 *
$^{98}\text{Pd}-^{85}\text{Rb}_{1.153}$	14404.5	5.1	14405	5	0.1	1	100	100 $^{98}\text{Pd}$	JY1	1.0	09Ei08
$^{98}\text{Ag}-^{85}\text{Rb}_{1.153}$	23283	40	23270	40	-0.4	1	78	78 $^{98}\text{Ag}$	MA8	1.0	11He10
$^{98}\text{Mo } ^{35}\text{Cl}-^{96}\text{Mo } ^{37}\text{Cl}$	3690	2	3678.94	0.20	-1.4	U			H11	4.0	63Bi12
$^{98}\text{Sr}-^{97}\text{Zr}_{1.010}$	18620	10	18625	3	0.5	1	12	12 $^{98}\text{Sr}$	JY1	1.0	06Ha03
$^{98}\text{Y}-^{97}\text{Zr}_{1.010}$	12321.4	8.5				2			JY1	1.0	07Ha32
$^{98}\text{Zr}-^{97}\text{Zr}_{1.010}$	2668	10	2668	9	0.0	1	82	82 $^{98}\text{Zr}$	JY1	1.0	06Ha03
$^{98}\text{Mo}-^{97}\text{Mo}_{1.010}$	327.9	5.8	326.53	0.07	-0.2	U			JY1	1.0	06Ka48
$^{98}\text{Mo}-^{97}\text{Mo}$	-614	12	-613.30	0.07	0.0	U			M15	2.5	63Ri07
$^{94}\text{Rb}-^{98}\text{Rb}_{.411} \ ^{91}\text{Rb}_{.590}$	-290	40	-347	8	-0.6	U			P21	2.5	82Au01 Y
$^{97}\text{Rb}-^{98}\text{Rb}_{.792} \ ^{93}\text{Rb}_{.209}$	-250	60	-281	13	-0.2	U			P21	2.5	82Au01
$^{96}\text{Rb}-^{98}\text{Rb}_{.490} \ ^{94}\text{Rb}_{.511}$	330	30	322	9	-0.1	U			P21	2.5	82Au01 Y
$^{97}\text{Rb}-^{98}\text{Rb}_{.660} \ ^{95}\text{Rb}_{.340}$	-300	50	-232	13	0.5	U			P21	2.5	82Au01
	-232	27			0.0	U			P31	2.5	86Au02
$^{96}\text{Zr}(t,p)^{98}\text{Zr}$	3508	20	3509	8	0.0	1	18	18 $^{98}\text{Zr}$	LAL		69B101
$^{96}\text{Zr}(^3\text{He},p)^{98}\text{Nb}$	5728	5				2			Phi		75Me13
$^{96}\text{Ru}(^{16}\text{O},^{14}\text{C})^{98}\text{Pd}$	-12529	20	-12516	5	0.6	U			BNL		82Th01
$^{98}\text{Mo}(t,\alpha)^{97}\text{Nb}$	10019	20	10015	4	-0.2	U			LAL		83F106
$^{97}\text{Mo}(n,\gamma)^{98}\text{Mo}$	8642.4	0.5	8642.60	0.06	0.4	U					71He10
	8642.60	0.07			0.0	-			MMn		91Is02 Z
	8642.57	0.18			0.2	-			Bdn		06Fi.A
$^{98}\text{Mo}(d,t)^{97}\text{Mo}$	-2379	20	-2385.37	0.06	-0.3	U			Pit		64Co11
$^{97}\text{Mo}(n,\gamma)^{98}\text{Mo}$	ave. 8642.60	0.07	8642.60	0.06	0.0	1	98	87 $^{98}\text{Mo}$			average
$^{97}\text{Mo}(^3\text{He},d)^{98}\text{Tc}$	680	8	683	3	0.4	-			ANL		74Co27
	686	10			-0.3	-			McM		76Ma16
ave.	682	6			0.1	1	29	29 $^{98}\text{Tc}$			average
$^{98}\text{Rb}(\beta^-)^{98}\text{Sr}$	11200	110	12054	16	7.8	B					79Pe17
	12343	150			-1.9	U			Trs		82Br23
	12519	60			-7.8	C			Gsn		84Bl.A
	12270	30			-7.2	C			McG		84Ia.A
	12440	75			-5.1	C			Bwg		87Gr.A
	12380	65			-5.0	B			Gsn		92Pr03
$^{98}\text{Rb}^m(\beta^-)^{98}\text{Sr}$	12710	120	12127	21	-4.9	C			Bwg		87Gr.A
$^{98}\text{Sr}(\beta^-)^{98}\text{Y}$	5730	40	5872	9	3.5	C					79Pe17
	5821	10			5.1	C			Gsn		84Bl.A
	5815	40			1.4	U			Bwg		87Gr.A
$^{98}\text{Y}(\beta^-)^{98}\text{Zr}$	8974	100	8992	12	0.2	U					79Pe17 *
	8780	30			7.1	C			Gsn		84Bl.A
	8840	55			2.8	C			Bwg		87Gr.A
	8963	41			0.7	U					88Ma.A
	8830	17			9.5	C			Bwg		92Gr.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{98}\text{Y}^m(\beta^-)^{98}\text{Zr}$	9780	200	9233	27	-2.7	o			Bwg		87Gr.A
	9233	27				2			Bwg		92Gr.A
$^{98}\text{Zr}(\beta^-)^{98}\text{Nb}$	2300	200	2238	10	-0.3	U					76He10
$^{98}\text{Nb}(\beta^-)^{98}\text{Mo}$	4300	200	4591	5	1.5	U					66Gu05
	4300	200			1.5	U					67Hu07
	4800	200			-1.0	U					76He10
	4580	100			0.1	U			Bwg		78St02
$^{98}\text{Mo}(\text{p,n})^{98}\text{Tc}$	-2458	10	-2466	3	-0.8	1	11	11 $^{98}\text{Tc}$	ANL		74Co27
$^{98}\text{Tc}(\beta^-)^{98}\text{Ru}$	1795	22	1793	7	-0.1	1	11	8 $^{98}\text{Ru}$			73Ok.A *
$^{98}\text{Rh}(\beta^+)^{98}\text{Ru}$	5120	100	5050	10	-0.7	U					72Ba37 *
	5151	50			-2.0	U					94Ba06 *
$^{98}\text{Ru}(\text{p,n})^{98}\text{Rh}$	-5832	10				2					70As08 Z
$^{98}\text{Ag}(\beta^+)^{98}\text{Pd}$	8420	150	8250	30	-1.1	U					79Ve.A *
	8200	70			0.8	1	22	22 $^{98}\text{Ag}$			00Hu17
$^{98}\text{Cd}(\beta^+)^{98}\text{Ag}$	5330	140	5430	40	0.7	U					92Pl01
$^{98}\text{Cd}(\epsilon)^{98}\text{Ag}$	5430	40				2					01St.A
$^{*98}\text{Rb}-^{85}\text{Rb}_{1.153}$	Original $D_M=43394.0(3.7) \mu\text{u}$ $M-A=-54317.7(3.4)\text{keV}$ corrected for gs+m mixture										16K104 **
$^{*98}\text{Rb}-^{85}\text{Rb}_{1.153}$	Original $M-A=-54309.4(4.0)$ corrected for isom mixture $E_x=80(30)$ $R=0.65(0.15)$										16K104 **
$^{*98}\text{Rb}-\text{u}$	Original $M-A=-54319.5(5.5)$ corrected for isom mixture $E_x=80(30)$ $R=0.65(0.15)$										16K104 **
$^{*98}\text{Rb}^m-\text{u}$	Data re-analysis										16K104 **
$^{*98}\text{Sr}-^{85}\text{Rb}_{1.153}$	$D_M=30397.3(4.3) \mu\text{u}$ $M-A=-66424.0(4.0)\text{keV}$ corrected for $e^-$ binding= $-529\text{eV}$										12Si10 **
$^{*98}\text{Sr}-^{85}\text{Rb}_{1.153}$	Former measured using $15^+$ ion; latter with $10^+$ ion										16K104 **
$^{*98}\text{Rh}-\text{u}$	$M-A=-83154(30) \text{keV}$ for mixture gs+m at 60#50 keV										Nub16b **
$^{*98}\text{Y}(\beta^-)^{98}\text{Zr}$	$E_{\beta^-}=4810(100)$ to $1^-$ level at 4164.60 keV										Ens035 **
$^{*98}\text{Tc}(\beta^-)^{98}\text{Ru}$	$E_{\beta^-}=397(22)$ to $4^+$ level at 1397.82 keV										Ens035 **
$^{*98}\text{Rh}(\beta^+)^{98}\text{Ru}$	$E_{\beta^+}=3450(100)$ to $2^+$ level at 652.44 keV, and others										Ens035 **
$^{*98}\text{Rh}(\beta^+)^{98}\text{Ru}$	$E_{\beta^+}=3476(50)$ to $2^+$ level at 652.44 keV										Ens035 **
$^{*98}\text{Ag}(\beta^+)^{98}\text{Pd}$	$Q_{\beta^+}=6880(150)$ to $4^+$ level at 1541.40 keV										Ens035 **
$^{99}\text{Rb}-^{85}\text{Rb}_{1.165}$	47885.1	4.8	47884	4	-0.2	2			MA8	1.0	13Ma81
	47880	10			0.4	2			TT1	1.0	16K104
$^{99}\text{Sr}-^{85}\text{Rb}_{1.165}$	35663	21	35645	5	-0.8	o			TT1	1.0	12Si10 *
	35644.5	7.0			0.1	1	53	53 $^{99}\text{Sr}$	TT1	1.0	16K104
$^{99}\text{Zr}-\text{u}$	-83323	19	-83329	11	-0.3	1	35	35 $^{99}\text{Zr}$	JY0	1.0	04Ri12
$\text{C7 H}_{15}-^{99}\text{Ru}$	211442.8	3.0	211445.2	0.4	0.3	U			M16	2.5	63Da10
$^{99}\text{Ag}-^{85}\text{Rb}_{1.165}$	20401.0	8.5	20411	7	1.1	2			SH1	1.0	07Ma92
	20427	11			-1.5	2			MA8	1.0	11He10
$^{99}\text{Cd}-^{85}\text{Rb}_{1.165}$	27690.8	1.7				2			MA8	1.0	09Br09
$^{99}\text{Pd}-^{96}\text{Mo}_{1.031}$	10052.8	5.5	10054	5	0.1	1	94	94 $^{99}\text{Pd}$	JY1	1.0	09El08
$^{99}\text{Sr}-^{97}\text{Zr}_{1.021}$	23794.1	7.4	23793	5	-0.1	1	47	47 $^{99}\text{Sr}$	JY1	1.0	06Ha03
$^{99}\text{Y}-^{97}\text{Zr}_{1.021}$	15066.8	7.1				2			JY1	1.0	07Ha32
$^{99}\text{Zr}-^{97}\text{Zr}_{1.021}$	7580	14	7583	11	0.2	1	65	65 $^{99}\text{Zr}$	JY1	1.0	06Ha03
$^{99}\text{Ru}-^{98}\text{Ru}$	652	11	644	7	-0.3	U			M16	2.5	63Da10
$^{97}\text{Rb}-^{99}\text{Rb}_{.653} \text{ } ^{93}\text{Rb}_{.348}$	100	100	135	4	0.1	U			P21	2.5	82Au01
$^{98}\text{Rb}-^{99}\text{Rb}_{.742} \text{ } ^{95}\text{Rb}_{.258}$	690	180	563	17	-0.3	U			P21	2.5	82Au01
$^{97}\text{Rb}-^{99}\text{Rb}_{.490} \text{ } ^{95}\text{Rb}_{.511}$	350	60	201	11	-1.0	U			P31	2.5	86Au02
$^{99}\text{Ru}(\text{n},\alpha)^{96}\text{Mo}$	6822	5	6815.9	0.4	-1.2	U					01Wa50
$^{96}\text{Ru}(\text{}^{16}\text{O}, \text{}^{13}\text{C})^{99}\text{Pd}$	-11723	20	-11760	5	-1.8	U			BNL		82Th01
	$^{98}\text{Mo}(\text{n},\gamma)^{99}\text{Mo}$	5927.7	1.	5925.44	0.15	-2.3	U				73De39
	5927	1			-1.6	U					74Er.A
	5924.6	0.6			1.4	U					76Ch02
	5923	2			1.2	U					77Ri04
	5925.42	0.15			0.2	1	100	99 $^{99}\text{Mo}$	MMn		91Is02 Z
	5927.7	0.5			-4.5	C			Bdn		06Fi.A
$^{98}\text{Mo}(\text{d,p})^{99}\text{Mo}$	3687	20	3700.88	0.15	0.7	U			Pit		64Co11

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{98}\text{Mo}(\beta^+)^{99}\text{Tc}$	1010	20	1007.4	0.9	-0.1	U			McM		77Ch06
$^{99}\text{Tc}(\text{p,d})^{98}\text{Tc}$	-6740	5	-6742	3	-0.5	-					76SI06
	-6755	9			1.4	-			Bld		77Em02
ave.	-6744	4			0.3	1	59	57	$^{98}\text{Tc}$		average
$^{99}\text{Rb}(\beta^-)^{99}\text{Sr}$	11340	120	11400	6	0.5	U			McG		84Ia.A
	10960	130			3.4	C			Bwg		87Gr.A
$^{99}\text{Sr}(\beta^-)^{99}\text{Y}$	8030	80	8128	8	1.2	U			McG		84Ia.A
	8360	75			-3.1	C			Bwg		87Gr.A
$^{99}\text{Y}(\beta^-)^{99}\text{Zr}$	7605	60	6971	12	-10.6	C			Bwg		87Gr.A
	7568	14			-42.7	C			Bwg		92Gr.A
$^{99}\text{Zr}(\beta^-)^{99}\text{Nb}$	4550	35	4715	16	4.7	C			Bwg		87Gr.A
	4559	15			10.4	C			Bwg		92Gr.A
$^{99}\text{Nb}(\beta^-)^{99}\text{Mo}$	3740	200	3635	12	-0.5	U					70Ei02 *
$^{99}\text{Mo}(\beta^-)^{99}\text{Tc}$	1356.7	1.0	1357.8	0.9	1.1	1	79	78	$^{99}\text{Tc}$		71Na01 *
$^{99}\text{Tc}(\beta^-)^{99}\text{Ru}$	292	3	297.5	0.9	1.8	U					51Ta05
	290	4			1.9	U					52Fe16
	293.5	2.0			2.0	1	22	20	$^{99}\text{Tc}$		80AI02 *
$^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$	2038	10	2044	7	0.6	-					52Sc11 *
	2053	10			-0.9	-					59To25 *
	2170	30			-4.2	B					74An23 *
ave.	2046	7			-0.2	1	90	89	$^{99}\text{Rh}$		average
$^{99}\text{Pd}(\beta^+)^{99}\text{Rh}$	3410	20	3399	8	-0.6	1	16	11	$^{99}\text{Rh}$		69Ph01 *
$^{99}\text{Ag}(\beta^+)^{99}\text{Pd}$	5430	150	5470	8	0.3	U					81Hu03
$^{99}\text{Sr}-^{85}\text{Rb}_{1.165}$	Original $D_M=35661.6(4.4) \mu\text{u}$ $M-A=-62506.3(4.1)\text{keV}$ re-evaluated										
$^{99}\text{Nb}(\beta^-)^{99}\text{Mo}$	$E_{\beta^-}=3500(200)$ to $7/2^+$ level at 235.508 keV										
$^{99}\text{Mo}(\beta^-)^{99}\text{Tc}$	$E_{\beta^-}=1214(1)$ to $1/2^-$ level at 142.6832 keV										
$^{99}\text{Tc}(\beta^-)^{99}\text{Ru}$	$E_{\beta^+}=434.8(2.6), 346.7(2.0)$ from $^{99}\text{Tc}^m$ at 142.6832 to ground state, 89.57 level										
$^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$	$E_{\beta^+}=740(10)$ from $^{99}\text{Rh}^m$ at 64.5 to 340.90 $7/2^+$ level										
$^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$	$E_{\beta^+}=1030(10), 710(10), 590(10), 420(20)$ keV										
*	to ground state, 321.99 $3/2^+$ , 442.59 $3/2^+$ , 618.13 $1/2^+$ levels										
$^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$	$Q_{\beta^+}$ larger than 2059.34 keV because that state populated via $\beta$ decay										
$^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$	Ref. proposed that $E_{\beta^+}=1030(10)$ keV decays to 89.57 $3/2^+$ level										
$^{99}\text{Rh}(\beta^+)^{99}\text{Ru}$	$E_{\beta^+}=1100(50), 680(30)$ and $540(30)$ to 89.57, 442.59 and 618.13 levels										
$^{99}\text{Pd}(\beta^+)^{99}\text{Rh}$	$E_{\beta^+}=2180(20), 1930(20), 1510(20)$ keV										
*	to 200.7 $7/2^+$ , 464.4 $(5/2, 7/2)^+$ , 874.5 $5/2^+$ levels above $(1/2^-)$ ground state										
$^{100}\text{Rb}-^{85}\text{Rb}_{1.176}$	54087	21							MA8	1.0	13Ma81
	54150	140	54087	21	-0.5	U			TT1	1.0	16KI04
$^{100}\text{Sr}-^{85}\text{Rb}_{1.176}$	39520	12	39515	8	-0.4	1	41	41	$^{100}\text{Sr}$	1.0	16KI04
$^{100}\text{Y}-\text{u}$	-72270	110	-72279	12	0.0	o			GT2	2.5	08Kn.A *
	-72290	140			0.0	U			GT2	2.5	08Su19 *
$^{100}\text{Zr}-\text{u}$	-82016	18	-81995	9	1.2	1	24	24	$^{100}\text{Zr}$	1.0	04Ri12
$^{100}\text{Mo}-^{85}\text{Rb}_{1.176}$	11216	27	11203.2	0.3	-0.5	U			MA8	1.0	11He10
	11205.5	1.4			-1.6	U			JY1	1.0	12Ka13
$^{100}\text{Mo}-^{87}\text{Rb}_{1.149}$	11820.25	0.57	11819.6	0.3	-1.2	1	32	32	$^{100}\text{Mo}$	1.0	15Gu09
$\text{C}_7 \text{H}_{16}-^{100}\text{Mo}$	217730.3	4.2	217732.5	0.3	0.2	U			M15	2.5	63Ri07
$^{100}\text{Mo}-\text{u}$	-92532.51	0.40	-92532.0	0.3	1.2	1	66	66	$^{100}\text{Mo}$	1.0	15Gu09
$\text{C}_7 \text{H}_{16}-^{100}\text{Ru}$	220983.8	3.7	220990.1	0.4	0.7	U			M16	2.5	63Da10
$^{100}\text{Rh}-\text{u}$	-91855	46	-91886	19	-0.7	1	18	18	$^{100}\text{Rh}$	1.0	05Li24 *
$^{100}\text{Ag}-^{85}\text{Rb}_{1.176}$	19849.9	7.1	19851	5	0.1	2			JY1	1.0	09EI08 *
	19851.8	8.2			-0.1	2			MA8	1.0	11He10 *
$^{100}\text{Cd}-\text{u}$	-79636	214	-79651.2	1.8	-0.1	U			CS1	1.0	96Ch32
$^{100}\text{Cd}-^{85}\text{Rb}_{1.176}$	24084.1	1.8	24084.1	1.8	0.0	1	100	100	$^{100}\text{Cd}$	1.0	09Br09
$^{100}\text{In}-\text{u}$	-69405	322	-69040	200	1.1	1	37	37	$^{100}\text{In}$	1.0	96Ch32
$^{100}\text{Sn}-\text{u}$	-62020	1020	-61500	320	0.5	U			CS1	1.0	96Ch32

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{100}\text{Sr}-^{97}\text{Zr}_{1.031}$	27579	10	27583	8	0.4	1	59	$^{100}\text{Sr}$	JY1	1.0	06Ha03
$^{100}\text{Y}-^{97}\text{Zr}_{1.031}$	19524	12				2			JY1	1.0	07Ha32
$^{100}\text{Y}^m-^{97}\text{Zr}_{1.031}$	19679	12				2			JY1	1.0	07Ha32
$^{100}\text{Zr}-^{97}\text{Zr}_{1.031}$	9815	10	9808	9	-0.7	1	76	$^{100}\text{Zr}$	JY1	1.0	06Ha03
$^{100}\text{Nb}^m-^{97}\text{Zr}_{1.031}$	6472.6	2.1				2			JY1	1.0	07Ha32
$^{100}\text{Mo}-^{35}\text{Cl}-^{98}\text{Mo}-^{37}\text{Cl}$	5019	2	5014.5	0.4	-0.6	U			H11	4.0	63Bi12
$^{100}\text{Nb}-^{100}\text{Nb}^m$	-335.7	8.3				3			JY1	1.0	07Ha32
$^{100}\text{Mo}-^{100}\text{Ru}$	3257.55	0.18	3257.52	0.18	-0.1	1	99	$^{100}\text{Ru}$	JY1	1.0	08Ra09
$^{100}\text{Ru}-^{99}\text{Ru}$	-1718	11	-1719.826	0.028	-0.1	U			M16	2.5	63Da10
$^{96}\text{Ru}(^{16}\text{O}, ^{12}\text{C})^{100}\text{Pd}$	-5599	26	-5605	18	-0.2	1	46	$^{100}\text{Pd}$	BNL		82Th01
$^{100}\text{Mo}(d, ^3\text{He})^{99}\text{Nb}$	-5639	15	-5653	12	-0.9	2			Tex		74Bi08
$^{100}\text{Mo}(t, \alpha)^{99}\text{Nb}$	8642	20	8667	12	1.3	2			LAI		83FI06
$^{100}\text{Mo}(d, t)^{99}\text{Mo}$	-2038	20	-2037.0	0.4	0.0	U			Pit		64Co11
$^{99}\text{Tc}(n, \gamma)^{100}\text{Tc}$	6764.4	1.				2					79Pi08
$^{99}\text{Ru}(n, \gamma)^{100}\text{Ru}$	6765.20	0.04	6764.4	1.0	-20.0	C					04Fu.A
	9673.2	0.7	9673.324	0.026	0.2	U					74Ri03
	9672.65	0.06			11.2	B			ILn		88Co18 Z
	9673.39	0.05			-1.3	-			MMn		91Is02 Z
	9673.30	0.03			0.8	-			ILn		00Ge01
	9673.41	0.19			-0.5	U			Bdn		06Fi.A
ave.	9673.324	0.026			0.0	1	100	$^{99}\text{Ru}$			average
$^{100}\text{Sr}(\beta^-)^{100}\text{Y}$	7460	140	7506	13	0.3	U			McG		84Ia.A *
$^{100}\text{Y}(\beta^-)^{100}\text{Zr}$	7075	100			4.3	C			Bwg		87Gr.A
	7920	100	9050	14	11.3	C			McG		84Ia.A *
	9310	70			-3.7	C			Bwg		87Gr.A
$^{100}\text{Zr}(\beta^-)^{100}\text{Nb}$	3335	25	3420	11	3.4	C			Bwg		87Gr.A
$^{100}\text{Nb}(\beta^-)^{100}\text{Mo}$	6245	25	6396	8	6.0	C			Bwg		87Gr.A
$^{100}\text{Nb}^m(\beta^-)^{100}\text{Mo}$	6745	75	6708.3	2.0	-0.5	U			Bwg		87Gr.A
$^{100}\text{Mo}(t, ^3\text{He})^{100}\text{Nb}^m$	-6690	30	-6689.7	2.0	0.0	U			LAI		79Aj03
$^{100}\text{Rh}(\beta^+)^{100}\text{Ru}$	3630	20	3636	18	0.3	1	82	$^{100}\text{Rh}$			53Ma64
$^{100}\text{Ag}(\beta^+)^{100}\text{Pd}$	7075	90	7075	18	0.0	U					79Ve.A *
	7022	200			0.3	U					80Ha20 *
$^{100}\text{Cd}(\beta^+)^{100}\text{Ag}$	3890	70	3943	5	0.8	U					89Ry02
$^{100}\text{In}(\beta^+)^{100}\text{Cd}$	10900	930	9880	180	-1.1	U			Lvp		95Sz01 *
	10080	230			-0.9	1	63	$^{100}\text{In}$			02PI03
$^{100}\text{Sn}(\beta^+)^{100}\text{In}$	7390	660	7030	240	-0.5	o			GSI		97Su06 *
	7840	660			-1.2	o			GSI		02Fa13 *
	7030	240			2				GSI		12Hi07 *
* $^{100}\text{Y}-u$	$M-A=-67245(93)$ keV for mixture gs+m at 144(16) keV										
* $^{100}\text{Y}-u$	$M-A=-67264(119)$ keV for mixture gs+m at 144(16) keV										
* $^{100}\text{Rh}-u$	$M-A=-85508(29)$ keV for mixture gs+m at 107.6 keV										
* $^{100}\text{Ag}-^{85}\text{Rb}_{1.176}$	$D_M=19858.2(5.2)$ $\mu\text{u}$ for mixture gs+m at 15.52 keV; $M-A=-78131.0(4.9)$ keV										
* $^{100}\text{Ag}-^{85}\text{Rb}_{1.176}$	$D_M=19860.2(6.6)$ $\mu\text{u}$ for mixture gs+m at 15.52 keV; $M-A=-78129.1(6.2)$ keV										
* $^{100}\text{Sr}(\beta^-)^{100}\text{Y}$	$E_{\beta^-}=7450(140)$ assumed by evaluator to $1^+$ level at 10.70 keV										
* $^{100}\text{Y}(\beta^-)^{100}\text{Zr}$	Not unambiguously ground state transition										
* $^{100}\text{Ag}(\beta^+)^{100}\text{Pd}$	From $5^+$ ground state to high spin level at 2920.4 keV										
* $^{100}\text{Ag}(\beta^+)^{100}\text{Pd}$	$E_{\beta^+}=5350(200)$ from $^{100}\text{Ag}^m 2^+$ at 15.52 to $2^+$ level at 665.50 keV										
* $^{100}\text{In}(\beta^+)^{100}\text{Cd}$	From lower and upper limits 9300-12500										
* $^{100}\text{Sn}(\beta^+)^{100}\text{In}$	$Q_{\beta^+}=7200(+800-500)$ ; also $E_{\beta^+}=3400(+700-300)$ keV										
* $^{100}\text{Sn}(\beta^+)^{100}\text{In}$	$E_{\beta^+}=3800(+700-300)$ to 2760(430) level										
* $^{100}\text{Sn}(\beta^+)^{100}\text{In}$	$E_{\beta^+}=3290(200)$ to $1^+$ level at 2721+x, with $x < 80$ keV										
$^{101}\text{Sr}-^{85}\text{Rb}_{1.188}$	45410	22	45400	9	-0.5	2			MA8	1.0	16De.A
	45398	10			0.2	2			TT1	1.0	16KI04
$^{101}\text{Zr}-u$	-78573	20	-78547	9	1.3	1	20	$^{101}\text{Zr}$	JY0	1.0	04Ri12



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_8 H_5 -^{101}Ru$	133549.5	2.2	133552.1	0.4	0.5	U			M16	2.5	63Da10
$^{101}Rh-u$	-93821	58	-93841	6	-0.3	U			GS2	1.0	05Li24 *
$^{101}Pd-u$	-91816	30	-91715	5	3.4	C			GS2	1.0	05Li24
$^{101}Ag-^{85}Rb_{1.188}$	17470.5	7.2	17478	5	1.0	2			SH1	1.0	07Ma92
	17485.6	7.5			-1.0	2			MA8	1.0	11He10
$^{101}Cd-^{85}Rb_{1.188}$	23367	11	23380.0	1.6	1.2	U			SH1	1.0	07Ma92
	23380.0	1.6				2			MA8	1.0	09Br09
$^{101}Pd-^{96}Mo_{1.052}$	8567.4	5.1	8567	5	-0.1	1	93	93 $^{101}Pd$	JY1	1.0	09El08
$^{101}Cd-^{96}Mo_{1.052}$	18872.7	5.5	18868.4	1.6	-0.8	U			JY1	1.0	09El08
$^{101}Y-^{97}Zr_{1.041}$	22847.5	7.6				2			JY1	1.0	07Ha32
$^{101}Zr-^{97}Zr_{1.041}$	14153	10	14146	9	-0.7	1	80	80 $^{101}Zr$	JY1	1.0	06Ha03
$^{101}Nb-^{102}Ru_{.990}$	10009.6	4.0				2			JY1	1.0	07Ha32
$^{101}Ru-^{100}Ru$	1368	11	1362.62	0.25	-0.2	U			M16	2.5	63Da10
$^{100}Mo(n,\gamma)^{101}Mo$	5398.4	0.5	5398.24	0.07	-0.3	U			ILn		75Ka.A
	5399.6	0.7			-1.9	U			ORn		79We07 Z
	5398.23	0.08			0.1	2			ILn		90Se17 Z
	5398.27	0.13			-0.2	2			Bdn		06Fi.A
$^{100}Mo(d,p)^{101}Mo$	3161	6	3173.68	0.07	2.1	U					72S125
$^{100}Ru(n,\gamma)^{101}Ru$	6802.0	0.7	6802.04	0.24	0.1	-					82Ba69
	6802.04	0.25			0.0	-			Bdn		06Fi.A
$^{100}Ru(d,p)^{101}Ru$	4581	4	4577.48	0.24	-0.9	U					77Ho02
$^{100}Ru(n,\gamma)^{101}Ru$	ave. 6802.04	0.24	6802.04	0.24	0.0	1	100	99 $^{101}Ru$			average
$^{101}Sn(\epsilon p)^{100}Cd$	6600	300				2					10St.A
$^{101}Rb(\beta^-)^{101}Sr$	11810	110	12480#	200#	6.1	D			Bwg		92Ba28 *
$^{101}Sr(\beta^-)^{101}Y$	9505	80	9736	11	2.9	B			Bwg		92Ba28
$^{101}Y(\beta^-)^{101}Zr$	8545	90	8105	11	-4.9	B			Bwg		92Ba28
$^{101}Zr(\beta^-)^{101}Nb$	5520	45	5726	9	4.6	B			Bwg		87Gr18
	5485	25			9.6	C			Bwg		92Gr.A
$^{101}Nb(\beta^-)^{101}Mo$	4575	30	4628	4	1.8	U			Bwg		87Gr.A
	4590	30			1.3	U			Bwg		87Gr18
	4569	18			3.3	C			Bwg		92Gr.A
$^{101}Mo(\beta^-)^{101}Tc$	2836	40	2825	24	-0.3	R					57Ok.A *
$^{101}Tc(\beta^-)^{101}Ru$	1620	30	1614	24	-0.2	2					71Ar23 *
$^{101}Pd(\beta^+)^{101}Rh$	1980	4	1980	4	0.1	1	95	88 $^{101}Rh$			71Ib01 *
$^{101}Ag(\beta^+)^{101}Pd$	4100	200	4098	7	0.0	U					72We.A
	4350	200			-1.3	U					78Ha11
	4180	150			-0.5	U					79Ve.A *
$^{101}Cd(\beta^+)^{101}Ag$	5530	130	5498	5	-0.2	U					70Be.A *
	5350	200			0.7	U					72We.A
* $^{101}Rh-u$	$M-A=-87315(29)$ keV for mixture gs+m at 157.32 keV										Nub16b **
* $^{101}Rb(\beta^-)^{101}Sr$	Trends from Mass Surface TMS suggest $^{101}Rb$ 670 less bound										GAu **
* $^{101}Mo(\beta^-)^{101}Tc$	$E_{\beta^-}=2230(40)$ to $(1/2^+, 3/2^+)$ level at 606.47 keV										Ens06a **
* $^{101}Tc(\beta^-)^{101}Ru$	$E_{\beta^-}=1320(30)$ to 306.858 $7/2^+$ and 1070(30) to 545.115 $7/2^+$ levels										Ens06a **
* $^{101}Pd(\beta^+)^{101}Rh$	$E_{\beta^+}=776(4)$ to $7/2^+$ level at 181.78 keV										Ens06a **
* $^{101}Ag(\beta^+)^{101}Pd$	$E_{\beta^+}=2895(150)$ to $7/2^+$ level at 261.0 keV, and others										Ens06a **
* $^{101}Cd(\beta^+)^{101}Ag$	Measured $E_{\beta^+}$ may go to excited state										70Be.A **
$^{102}Sr-^{85}Rb_{1.200}$	49857	72				2			MA8	1.0	16De.A
$^{102}Y-^{120}Sn_{.850}$	17456.3	4.3				2			JY1	1.0	11Ha48 *
$^{102}Zr-u$	-76780	43	-76853	9	-1.7	U			JY0	1.0	04Ri12
$C_8 H_6-^{102}Ru$	142604.8	3.2	142609.9	0.4	0.6	U			M16	2.5	63Da10
$C_8 H_6-^{102}Pd$	141324	19	141318.1	0.6	-0.1	U			M16	2.5	63Da10
	141346	18			-1.0	U			R12	1.5	83De51
$C_7 N H_4-^{102}Pd$	128775	19	128742.1	0.6	-1.2	U			R12	1.5	83De51
$^{102}Pd-u$	-94370.9	15.6	-94367.9	0.6	0.2	U			GS4	1.0	14Ya.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{102}\text{Ag}-u$	-88315	30	-88295	9	0.7	U			GS2	1.0	05Li24 *
$^{102}\text{Cd}-^{85}\text{Rb}_{1,200}$	20320.9	7.3	20334.1	1.8	1.8	U			SH1	1.0	07Ma92
	20334.2	1.9			0.0	1	88	88 $^{102}\text{Cd}$	MA8	1.0	09Br09
$^{102}\text{In}-^{85}\text{Rb}_{1,200}$	29959	13	29958	5	-0.1	1	14	14 $^{102}\text{In}$	SH1	1.0	07Ma92
$^{102}\text{Cd}-^{96}\text{Mo}_{1,063}$	15811.9	5.2	15812.5	1.8	0.1	1	12	12 $^{102}\text{Cd}$	JY1	1.0	09El08
$^{102}\text{In}-^{96}\text{Mo}_{1,063}$	25436.5	5.3	25437	5	0.0	1	86	86 $^{102}\text{In}$	JY1	1.0	09El08
$^{102}\text{Zr}-^{97}\text{Zr}_{1,052}$	16822.0	9.8	16820	9	-0.2	1	92	92 $^{102}\text{Zr}$	JY1	1.0	06Ha03
$^{102}\text{Nb}-^{97}\text{Zr}_{1,052}$	11756.4	2.7	11756.5	2.7	0.0	1	99	99 $^{102}\text{Nb}$	JY1	1.0	07Ha32
$^{102}\text{Mo}-^{97}\text{Zr}_{1,052}$	3961.0	9.8	3961	9	0.0	1	83	83 $^{102}\text{Mo}$	JY1	1.0	06Ha03
$^{102}\text{Nb}^m-^{102}\text{Nb}$	100.2	7.9	101	8	0.1	1	95	94 $^{102}\text{Nb}^m$	JY1	1.0	07Ha32
$^{102}\text{Pd}-^{102}\text{Ru}$	1291.76	0.39	1291.8	0.4	0.0	1	100	100 $^{102}\text{Pd}$	SH1	1.0	11Go23
$^{102}\text{Ru}-^{101}\text{Ru}$	-1233	11	-1232.78	0.05	0.0	U			M16	2.5	63Da10
$^{100}\text{Mo}(\text{t,p})^{102}\text{Mo}$	5034	20	5034	8	0.0	1	17	17 $^{102}\text{Mo}$	LAl		72Ca10
$^{100}\text{Mo}(\text{}^3\text{He,p})^{102}\text{Tc}$	6054	20	6022	9	-1.6	1	21	21 $^{102}\text{Tc}$	Pri		82De03
$^{102}\text{Pd}(\text{p,t})^{100}\text{Pd}$	-10356	24	-10351	18	0.2	1	54	54 $^{100}\text{Pd}$	Win		74De31 *
$^{101}\text{Ru}(\text{n},\gamma)^{102}\text{Ru}$	9220.4	0.9	9219.64	0.05	-0.8	U					74Ri03
	9219.64	0.05			0.0	1	100	99 $^{102}\text{Ru}$	MMn		91Is02 Z
	9219.63	0.19			0.1	U			Bdn		06Fi.A
$^{102}\text{In}(\text{e}\text{p})^{101}\text{Ag}$	3420	310	3351	7	-0.2	o			Lvp		91Re.A *
$^{102}\text{Sr}(\beta^-)^{102}\text{Y}$	8815	70	9010	70	2.8	B			Bwg		92Ba28
$^{102}\text{Y}(\beta^-)^{102}\text{Zr}$	9850	70	10415	10	8.1	B			Bwg		92Ba28
$^{102}\text{Zr}(\beta^-)^{102}\text{Nb}^m$	4605	30	4622	11	0.6	1	14	8 $^{102}\text{Zr}$	Bwg		87Gr18
$^{102}\text{Nb}(\beta^-)^{102}\text{Mo}$	7300	50	7262	9	-0.8	o			Bwg		87Gr.A
	7335	40			-1.8	U			Bwg		87Gr18
$^{102}\text{Nb}^m(\beta^-)^{102}\text{Mo}$	7215	40	7356	11	3.5	C			Bwg		87Gr.A
	7210	35			4.2	B			Bwg		87Gr18
$^{102}\text{Tc}(\beta^-)^{102}\text{Ru}$	4420	100	4534	9	1.1	U					69B116
$^{102}\text{Rh}(\beta^+)^{102}\text{Ru}$	2317	10	2323	6	0.6	2					61Hi06
	2325	10			-0.2	2					63Bo17
$^{102}\text{Ru}(\text{p,n})^{102}\text{Rh}$	-3115	15	-3105	6	0.6	2					83Do11
$^{102}\text{Rh}(\beta^-)^{102}\text{Pd}$	1150	6	1120	6	-5.0	B					61Hi06
$^{102}\text{Ag}(\beta^+)^{102}\text{Pd}$	5800	200	5656	8	-0.7	U					67Ch05
	5966	100			-3.1	B					67Ch05 *
	4910	140			5.3	C					70Be.A *
	5350	200			1.5	U					72We.A
	5880	110			-2.0	U					79Ve.A
$^{102}\text{Cd}(\beta^+)^{102}\text{Ag}$	2554	57	2587	8	0.6	U					72We.A
	2587	8				2			GSI		91Ke08 *
$^{102}\text{In}(\beta^+)^{102}\text{Cd}$	9250	380	8965	5	-0.8	U			Lvp		95Sz01 *
	8970	150			0.0	U			GSI		98Ka.A
	8910	170			0.3	U			GSI		03Gi06 *
$^{102}\text{Sn}(\beta^+)^{102}\text{In}$	5780	70	5760	100	-0.3	o			GSI		01St.A
	5760	100				2			GSI		02Fa13
* $^{102}\text{Y}-^{120}\text{Sn}_{,850}$	Associated with low-spin isomer										
* $^{102}\text{Ag}-u$	$M-A=-82260(28)$ keV for mixture gs+m at 9.40 keV										
* $^{102}\text{Pd}(\text{p,t})^{100}\text{Pd}$	Original error 12; added systematic error 21 keV										
* $^{102}\text{In}(\text{e}\text{p})^{101}\text{Ag}$	Estimated using proton spectrum from 1450 to 3200 keV										
* $^{102}\text{Ag}(\beta^+)^{102}\text{Pd}$	$E_{\beta^+}=3340(100), 3070(130)$ from $^{102}\text{Ag}^m$ at 9.40 to 1534.48, 2017.8 levels										
* $^{102}\text{Ag}(\beta^+)^{102}\text{Pd}$	$Q_{\beta^+}=4920(100)$ from $^{102}\text{Ag}^m$ at 9.40 keV										
* $^{102}\text{Cd}(\beta^+)^{102}\text{Ag}$	$E_{\beta^+}=1075(8)$ to $1^+$ level at 490.44 keV										
* $^{102}\text{In}(\beta^+)^{102}\text{Cd}$	From 9900 keV upper and 8600 keV lower limits										
* $^{102}\text{In}(\beta^+)^{102}\text{Cd}$	Good agreement with authors' earlier measurement, average=8950(120) keV										
											11Ha48 **
$^{103}\text{Y}-u$	-63060	183	-62757	12	1.1	o			GT1	1.5	04Ma.A
	-62803	106			0.2	U			GT2	2.5	08Kn.A
$^{103}\text{Y}-^{120}\text{Sn}_{,858}$	21154	12				2			JY1	1.0	11Ha48

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{103}\text{Zr}-u$	-72765	64	-72803	10	-0.6	U			JY0	1.0	04Ri12
$\text{C}_8 \text{H}_7-^{103}\text{Rh}$	149263.5	3.3	149281.2	2.5	2.1	U			M16	2.5	63Da10
	149261	19			0.7	U			R12	1.5	83De51
$\text{C}_7 \text{N H}_5-^{103}\text{Rh}$	136681	18	136705.1	2.5	0.9	U			R12	1.5	83De51
$^{103}\text{Pd}-u$	-93894.8	14.5	-93889.2	1.0	0.4	U			GS4	1.0	14Ya.A
$^{103}\text{Ag}-u$	-91091	52	-91039	4	1.0	U			GS2	1.0	05Li24 *
$^{103}\text{Ag}-^{85}\text{Rb}_{1,212}$	15875	14	15871	4	-0.3	U			SH1	1.0	07Ma92
	15871.4	4.4				2			MA8	1.0	11He10
$^{103}\text{Cd}-^{85}\text{Rb}_{1,212}$	20328	11	20327.8	1.9	0.0	U			SH1	1.0	07Ma92
	20328.2	2.1			-0.2	1	86	86 $^{103}\text{Cd}$	MA8	1.0	09Br09
$^{103}\text{In}-^{85}\text{Rb}_{1,212}$	26785	11	26789	10	0.4	1	88	88 $^{103}\text{In}$	SH1	1.0	07Ma92
$^{103}\text{Cd}-^{96}\text{Mo}_{1,073}$	15699.2	5.2	15700.9	1.9	0.3	1	14	14 $^{103}\text{Cd}$	JY1	1.0	09Ei08
$^{103}\text{Zr}-^{97}\text{Zr}_{1,062}$	21760.5	9.9				2			JY1	1.0	06Ha03
$^{103}\text{Mo}-^{97}\text{Zr}_{1,062}$	7648.4	9.9				2			JY1	1.0	06Ha03
$^{103}\text{Nb}-^{102}\text{Ru}_{1,010}$	16069.7	4.2				2			JY1	1.0	07Ha32
$^{103}\text{Cd}-^{102}\text{Cd}$	-1534	154	-1064.9	2.6	2.0	U			CR2	1.5	92Sh.A *
$^{103}\text{Rh}(p,u)^{101}\text{Rh}$	-8275	17	-8280	6	-0.3	1	13	12 $^{101}\text{Rh}$	Pri		64Th05
$^{102}\text{Ru}(n,\gamma)^{103}\text{Ru}$	6232.2	0.3	6232.05	0.15	-0.5	-					82Ba69 Z
	6232.00	0.17			0.3	-			Bdn		06Fi.A
$^{102}\text{Ru}(d,p)^{103}\text{Ru}$	4005	15	4007.49	0.15	0.2	U			ANL		71Fo01
$^{102}\text{Ru}(n,\gamma)^{103}\text{Ru}$	ave.	6232.05	6232.05	0.15	0.0	1	100	99 $^{103}\text{Ru}$			average
$^{103}\text{Rh}(\gamma,n)^{102}\text{Rh}$	-9307	32	-9320	7	-0.4	U			Phi		60Ge01
$^{103}\text{Rh}(p,d)^{102}\text{Rh}$	-7144	16	-7095	7	3.1	B			Pri		64Th05
$^{102}\text{Pd}(n,\gamma)^{103}\text{Pd}$	7624.6	1.5	7625.3	0.8	0.5	2					70Bo29
	7625.6	0.9			-0.3	2			Bdn		06Fi.A
$^{103}\text{Zr}(\beta^-)^{103}\text{Nb}$	6945	85	7213	10	3.2	B			Bwg		87Gr18
$^{103}\text{Nb}(\beta^-)^{103}\text{Mo}$	5535	35	5932	10	11.3	C			Bwg		87Gr.A
	5530	30			13.4	B			Bwg		87Gr18
$^{103}\text{Mo}(\beta^-)^{103}\text{Tc}$	3750	60	3643	13	-1.8	U			Bwg		87Gr18
$^{103}\text{Tc}(\beta^-)^{103}\text{Ru}$	2615	45	2663	10	1.1	U			Bwg		87Gr.A
$^{103}\text{Ru}(\beta^-)^{103}\text{Rh}$	764	4	764.5	2.3	0.1	-					58Ro09 *
	760	6			0.8	-					65Mu09 *
	762	5			0.5	-					70Pe04 *
	769	4			-1.1	-					82Oh04
	ave.	764.6	2.3			0.0	1	98	98 $^{103}\text{Rh}$		average
$^{103}\text{Pd}(\epsilon)^{103}\text{Rh}$	564	20	574.5	2.4	0.5	U					54Ri09 *
	543.0	0.8			39.4	B					86Be53
$^{103}\text{Ag}(\beta^+)^{103}\text{Pd}$	2580	100	2654	4	0.7	U					62Pa05 *
	2700	100			-0.5	U					66Ja12
	2320	50			6.7	B					69Ba02
	2622	34			1.0	U			Dlf		88Bo28 *
$^{103}\text{Cd}(\beta^+)^{103}\text{Ag}$	4200	130	4151	4	-0.4	U					70Be.A
	4250	200			-0.5	U					72We.A
	4131	23			0.9	U			Dlf		88Bo28 *
$^{103}\text{In}(\beta^+)^{103}\text{Cd}$	5380	200	6019	10	3.2	B			Brk		83Wo04
	6050	28			-1.1	1	12	12 $^{103}\text{In}$	Dlf		88Bo28 *
	6040	60			-0.3	U					98Ka42
$^{103}\text{Sn}(\beta^+)^{103}\text{In}$	7500	600	7660	70	0.3	o			GSI		04Mu32
	7660	70				2			GSI		05Ka34 *
* $^{103}\text{Ag}-u$	$M - A = -84784(29)$ keV for mixture gs+m at 134.45 keV										
* $^{103}\text{Cd}-^{102}\text{Cd}$	From $^{102}\text{Cd}/^{103}\text{Cd} = 0.99029800(150)$										
* $^{103}\text{Ru}(\beta^-)^{103}\text{Rh}$	$E_{\beta^-} = 227(4)$ to $5/2^+$ level at 536.840 keV, and other $E_{\beta^-}$										
* $^{103}\text{Ru}(\beta^-)^{103}\text{Rh}$	$E_{\beta^-} = 112(6)$ to $5/2^+$ level at 650.064 keV, and other $E_{\beta^-}$										
* $^{103}\text{Ru}(\beta^-)^{103}\text{Rh}$	$E_{\beta^-} = 225(5)$ to $5/2^+$ level at 536.840 keV, and other $E_{\beta^-}$										
* $^{103}\text{Pd}(\epsilon)^{103}\text{Rh}$	IBE=500(20) to $^{103}\text{Rh}^m$ $7/2^+$ at 39.753 keV										
* $^{103}\text{Ag}(\beta^+)^{103}\text{Pd}$	$E_{\beta^+} = 1290(100)$ to $5/2^+$ level at 266.861 keV, and other $E_{\beta^+}$										
* $^{103}\text{Ag}(\beta^+)^{103}\text{Pd}$	$E_{\beta^+} = 1601(27)$ , plus systematic error 20 keV estimated by evaluator										
* $^{103}\text{Cd}(\beta^+)^{103}\text{Ag}$	$E_{\beta^+} = 3109(11)$ , plus systematic error 20 keV estimated by evaluator										
* $^{103}\text{In}(\beta^+)^{103}\text{Cd}$	$E_{\beta^+} = 4841(20)$ , plus systematic error 20 keV estimated by evaluator										
* $^{103}\text{Sn}(\beta^+)^{103}\text{In}$	Original 7640(70) recalibrated										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{104}\text{Zr}-u$	-70661	54	-70558	10	1.9	U			JY0	1.0	04Ri12
$^{104}\text{Nb}-u$	-77470	140	-77100.9	2.9	1.1	U			GT2	2.5	08Kn.A *
$\text{C}_8 \text{H}_8-^{104}\text{Ru}$	157171.5	3.4	157174.9	2.7	0.4	1	10	$^{104}\text{Ru}$	M16	2.5	63Da10
$\text{C}_8 \text{H}_8-^{104}\text{Pd}$	158612	10	158569.9	1.4	-1.7	U			M16	2.5	63Da10
	158599	12			-1.6	U			R12	1.5	83De51
$\text{C}_7 \text{N H}_6-^{104}\text{Pd}$	146013	8	145993.8	1.4	-1.6	U			R12	1.5	83De51
$\text{C}_6 \text{ }^{13}\text{C N H}_5-^{104}\text{Pd}$	141552	20	141523.6	1.4	-0.9	U			R12	1.5	83De51
$^{104}\text{Pd}-u$	-95938	30	-95969.6	1.4	-1.1	U			GS2	1.0	05Li24
$^{104}\text{Ag}-u$	-91410	30	-91376	5	1.1	U			GS2	1.0	05Li24 *
$^{104}\text{Cd}-u$	-90147	30	-90143.8	1.8	0.1	U			GS2	1.0	05Li24
$^{104}\text{Cd}-^{85}\text{Rb}_{1.224}$	17813.7	5.5	17825.6	1.8	2.2	U			SH1	1.0	07Ma92
	17825.5	1.9			0.0	1	89	$^{104}\text{Cd}$	MA8	1.0	09Br09
$^{104}\text{In}-^{85}\text{Rb}_{1.224}$	26183.9	6.2				2			SH1	1.0	07Ma92
	26140.3	29.6	26184	6	1.5	U			JY1	1.0	09E108 *
$^{104}\text{Sn}-^{87}\text{Rb}_{1.195}$	31636.9	6.4	31634	6	-0.4	1	93	$^{104}\text{Sn}$	JY1	1.0	09E107
$^{104}\text{Cd}-^{96}\text{Mo}_{1.083}$	13094.2	5.5	13093.4	1.8	-0.1	1	11	$^{104}\text{Cd}$	JY1	1.0	09E108
$^{104}\text{Zr}-^{97}\text{Zr}_{1.072}$	24896	10				2			JY1	1.0	06Ha03
$^{104}\text{Nb}-^{97}\text{Zr}_{1.072}$	18352.8	2.9				2			JY1	1.0	07Ha32 *
$^{104}\text{Mo}-^{97}\text{Zr}_{1.072}$	9194.0	9.7	9194	10	0.0	1	97	$^{104}\text{Mo}$	JY1	1.0	06Ha03
$^{104}\text{In}-^{103}\text{In}$	-1241	231	-1664	12	-1.2	U			CR2	1.5	91Sh19 *
$^{104}\text{Pd}-^{102}\text{Pd}$	-1617	30	-1601.7	1.5	0.3	U			R12	1.5	83De51
$^{104}\text{Ru}(d,\alpha)^{102}\text{Tc}$	7180	10	7188	9	0.8	1	80	$^{102}\text{Tc}$	Pri		82De03
$^{102}\text{Ru}(t,p)^{104}\text{Ru}$	6648	30	6650.1	2.5	0.1	U			LAL		72Ca10
$^{104}\text{Ru}(d,^3\text{He})^{103}\text{Tc}$	-5289	10	-5287	9	0.2	2			VUn		83De20
$^{104}\text{Ru}(t,\alpha)^{103}\text{Tc}$	9048	30	9033	9	-0.5	2			LAL		81FI02
$^{104}\text{Ru}(d,t)^{103}\text{Ru}-^{148}\text{Gd}()^{147}\text{Gd}$	85	3	83.8	2.4	-0.4	1	65	$^{104}\text{Ru}$	Jul		86Ru04 *
$^{103}\text{Rh}(n,\gamma)^{104}\text{Rh}$	6998.96	0.10	6998.96	0.08	0.0	2			MMn		81Ke03 Z
	6998.95	0.14			0.0	2			Bdn		06Fi.A
$^{103}\text{Rh}(d,p)^{104}\text{Rh}$	4786	10	4774.39	0.08	-1.2	U			MIT		64Sp12
$^{104}\text{Pd}(d,t)^{103}\text{Pd}$	-3762	25	-3752.0	1.6	0.4	U			Pit		64Co11
$^{104}\text{Sb}(p)^{103}\text{Sn}$	510	100				3					94Pa12 *
$^{104}\text{Nb}^m(\text{IT})^{104}\text{Nb}$	215	120				3			Bwg		87Gr18 *
$^{104}\text{Nb}(\beta^-)^{104}\text{Mo}$	8105	90	8531	9	4.7	B			Bwg		87Gr18
$^{104}\text{Nb}^m(\beta^-)^{104}\text{Mo}$	8320	80	8750	120	5.3	B			Bwg		87Gr18 *
$^{104}\text{Mo}(\beta^-)^{104}\text{Tc}$	4800	200	2153	24	-13.2	B					64Te02
	2155	40			0.0	-			Bwg		87Gr18
	2160	40			-0.2	-			Jyv		94Jo.A
	ave.	2158	28		-0.1	1	73	$^{104}\text{Tc}$			average
$^{104}\text{Tc}(\beta^-)^{104}\text{Ru}$	5620	70	5592	25	-0.4	-					78Su03
	5590	60			0.0	-			Bwg		87Gr18
	ave.	5600	50		-0.2	1	30	$^{104}\text{Tc}$			average
$^{104}\text{Rh}(\beta^-)^{104}\text{Pd}$	2440	30	2435.8	2.7	-0.1	U					55Bu.A
$^{104}\text{Ag}(\beta^+)^{104}\text{Pd}$	4276	15	4279	4	0.2	U					60Nu02 *
	4350	200			-0.4	U					72We.A
	4306	31			-0.9	U			Dlf		88Bo28 *
$^{104}\text{Pd}(p,n)^{104}\text{Ag}$	-5061	4				3					79De44
$^{104}\text{Cd}(\beta^+)^{104}\text{Ag}$	1587	60	1148	5	-7.3	B					70Mu17 *
	1403	26			-9.8	B			GSI		79Pi06 *
$^{104}\text{In}(\beta^+)^{104}\text{Cd}$	7100	200	7786	6	3.4	B					78Hu06 *
	7260	250			2.1	U			Brk		83Wo04 *
	7800	250			-0.1	U			Dlf		88Bo28
	7890	120			-0.9	U			Dlf		90Re08
	7880	100			-0.9	U			GSI		98Ka.A
$^{104}\text{Sn}(\beta^+)^{104}\text{In}$	4550	300	4556	8	0.0	U					88Ba10 *
	4515	60			0.7	U			GSI		91Ke11
* $^{104}\text{Nb}-u$	$D_M=-77350(100) \mu\text{u}$ for mixture gs+m at 220(120); $M-A=-72051(93) \text{ keV}$										
* $^{104}\text{Ag}-u$	$M-A=-85144(28) \text{ keV}$ for mixture gs+m at 6.90 keV										
* $^{104}\text{In}-^{85}\text{Rb}_{1.224}$	$D_M=26190.5(5.5) \mu\text{u}$ for mixture gs+m at 93.48 keV; $M-A=-76176.5(5.1) \text{ keV}$										
* $^{104}\text{Nb}-^{97}\text{Zr}_{1.072}$	Only long-lived state is measured										
* $^{104}\text{In}-^{103}\text{In}$	From $^{103}\text{In}/^{104}\text{In}=0.99038900(222)$										
* $^{104}\text{Ru}(d,t)^{103}\text{Ru}-^{148}\text{Gd}()^{147}\text{Gd}$	$Q=82(3) \text{ to } 5/2^+$ level at 2.81 keV										
* $^{104}\text{Sb}(p)^{103}\text{Sn}$	Below 550 keV; value and error estimated by evaluator										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
* <sup>104</sup> Nb <sup>m</sup> (IT) <sup>104</sup> Nb	From difference in $Q_{\beta^-}$								GAu **	
* <sup>104</sup> Nb <sup>m</sup> ( $\beta^-$ ) <sup>104</sup> Mo	Better use the difference of the two $Q_{\beta^-}$ 's								GAu **	
* <sup>104</sup> Ag( $\beta^+$ ) <sup>104</sup> Pd	$E_{\beta^+}=2705(15)$ from <sup>104</sup> Ag <sup>m</sup> at 6.90 to 2 <sup>+</sup> level at 555.81 keV								Ens079 **	
* <sup>104</sup> Ag( $\beta^+$ ) <sup>104</sup> Pd	$E_{\beta^+}=2012(71)$ to 4 <sup>+</sup> level at 1323.59 keV								Ens079 **	
*	and $E_{\beta^+}=2729(24)$ from <sup>104</sup> Ag <sup>m</sup> at 6.90 to 2 <sup>+</sup> level at 555.81 keV								Ens079 **	
*	plus systematic error 20 keV estimated by evaluator								GAu **	
* <sup>104</sup> Cd( $\beta^+$ ) <sup>104</sup> Ag	$p^+=0.011(0.003)$ $0.0019(0.0005)$ respectively, to 1 <sup>+</sup> level at 90.6 keV; recalculated $E_{\beta^+}$								Ens079 **	
* <sup>104</sup> In( $\beta^+$ ) <sup>104</sup> Cd	$E_{\beta^+}=4600(200)$ $4750(250)$ respectively, to 4 <sup>+</sup> level at 1492.1 keV								Ens079 **	
* <sup>104</sup> Sn( $\beta^+$ ) <sup>104</sup> In	$p^+=0.71(0.07)$ to 1139.25 level								Ens079 **	
<sup>105</sup> Y-u	-55041	574					GT3	2.5	16Kn03	
<sup>105</sup> Rh-u	-94378	53	-94312.2	2.7	1.2	U	GS2	1.0	05Li24 *	
C <sub>8</sub> H <sub>9</sub> - <sup>105</sup> Pd	165357	14	165345.8	1.2	-0.3	U	M16	2.5	63Da10	
	165360	9			-1.1	U	R12	1.5	83De51	
C <sub>7</sub> N H <sub>7</sub> - <sup>105</sup> Pd	152773	18	152769.7	1.2	-0.1	U	R12	1.5	83De51	
C <sub>6</sub> <sup>13</sup> C N H <sub>6</sub> - <sup>105</sup> Pd	148309	26	148299.5	1.2	-0.2	U	R12	1.5	83De51	
C <sub>7</sub> O H <sub>5</sub> - <sup>105</sup> Pd	128970	18	128960.3	1.2	-0.4	U	R12	1.5	83De51	
<sup>105</sup> Ag-u	-93534	31	-93474	5	1.9	U	GS2	1.0	05Li24 *	
<sup>105</sup> Cd- <sup>96</sup> Mo <sub>1.094</sub>	13748.5	5.4	13749.7	1.5	0.2	U	JY1	1.0	09EI08	
<sup>105</sup> Cd- <sup>85</sup> Rb <sub>1.235</sub>	18403.4	1.5	18403.6	1.5	0.1	1	99 <sup>105</sup> Cd	MA8	1.0	09Br09
<sup>105</sup> In- <sup>85</sup> Rb <sub>1.235</sub>	23442	11				2				
<sup>105</sup> Sn- <sup>85</sup> Rb <sub>1.235</sub>	30204.1	7.1	30208	4	0.6	1	36 <sup>105</sup> Sn	SH1	1.0	07Ma92
<sup>105</sup> Sn- <sup>87</sup> Rb <sub>1.207</sub>	30890.0	5.6	30888	4	-0.4	1	58 <sup>105</sup> Sn	JY1	1.0	09EI07
<sup>105</sup> Zr- <sup>97</sup> Zr <sub>1.082</sub>	30359	13				2				
<sup>105</sup> Mo- <sup>97</sup> Zr <sub>1.082</sub>	13319.4	9.8	13319	10	0.0	1	98 <sup>105</sup> Mo	JY1	1.0	06Ha03
<sup>105</sup> Nb- <sup>102</sup> Ru <sub>1.029</sub>	23376.4	4.3				2				
<sup>105</sup> Pd- <sup>104</sup> Pd	1049	35	1049.1	0.8	0.0	U				
<sup>105</sup> In- <sup>104</sup> In	-3618	144	-3712	13	-0.4	U				
<sup>105</sup> Te( $\alpha$ ) <sup>101</sup> Sn	5079	50	5069	3	-0.2	U				
	5061.1	5.			1.6	o				
	5069.2	3.				3				
<sup>104</sup> Ru(n, $\gamma$ ) <sup>105</sup> Ru	5909.9	0.5	5910.10	0.11	0.4	-				
	5910.1	0.2			0.0	-				
	5910.11	0.14			-0.1	-				
<sup>104</sup> Ru(d,p) <sup>105</sup> Ru	3684	15	3685.53	0.11	0.1	U	Bdn			
	3684.5	1.0			1.0	U	ANL			
	ave.	5910.10	0.11	5910.10	0.11	0.0	1	100 <sup>105</sup> Ru	Mun	
<sup>104</sup> Ru(n, $\gamma$ ) <sup>105</sup> Ru	ave.	5910.10	0.11	5910.10	0.11	0.0	1	100 <sup>105</sup> Ru		average
<sup>104</sup> Pd(n, $\gamma$ ) <sup>105</sup> Pd	7094.1	0.7				2				70Bo29
<sup>104</sup> Pd(d,p) <sup>105</sup> Pd	4867	20	4869.5	0.7	0.1	U	Pit			64Co11
<sup>105</sup> Pd(d,t) <sup>104</sup> Pd	-851	30	-836.9	0.7	0.5	U	Pit			64Co11
<sup>105</sup> Sb(p) <sup>104</sup> Sn	482.6	15.	323	22	-10.7	F	Bkp			94Ti03 *
<sup>105</sup> Nb( $\beta^-$ ) <sup>105</sup> Mo	6485	70	7422	10	13.4	B	Bwg			87Gr18
<sup>105</sup> Mo( $\beta^-$ ) <sup>105</sup> Tc	4950	45	4950	40	0.1	1	61 <sup>105</sup> Tc	Bwg		87Gr18
<sup>105</sup> Tc( $\beta^-$ ) <sup>105</sup> Ru	3640	55	3640	40	0.1	1	41 <sup>105</sup> Tc	Bwg		87Gr18
<sup>105</sup> Ru( $\beta^-$ ) <sup>105</sup> Rh	1916	4	1916.8	2.9	0.2	1	51 <sup>105</sup> Rh			67Sc01
<sup>105</sup> Rh( $\beta^-$ ) <sup>105</sup> Pd	570	5	566.6	2.3	-0.7	-				51Du03
	560	5			1.3	-				56La24
	568	4			-0.3	-				64Ka23
	ave.	566.3	2.6		0.1	1	78 <sup>105</sup> Rh			average
<sup>105</sup> Ag( $\epsilon$ ) <sup>105</sup> Pd	1347	25	1347	5	0.0	U				67Pi03
	1310	25			1.5	U				67Sc26 *
<sup>105</sup> Cd( $\beta^+$ ) <sup>105</sup> Ag	2738	5	2737	4	-0.2	-				53Jo20 *
	2600	200			0.7	U				72We.A
	2742	11			-0.5	-				86Bo28 *
	ave.	2739	5		-0.4	1	92 <sup>105</sup> Ag			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{105}\text{In}(\beta^+)^{105}\text{Cd}$	5140	200	4693	10	-2.2	U			Brk		83Wo04
	4849	13			-12.0	B					86Bo28 *
$^{105}\text{Sn}(\beta^+)^{105}\text{In}$	6230	80	6303	11	0.9	U			GSI		06Ka74
* $^{105}\text{Rh}-u$	$M-A=-87847(32)$ keV for mixture gs+m at 129.782 keV										Nub16b **
* $^{105}\text{Ag}-u$	$M-A=-87113(28)$ keV for mixture gs+m at 25.479 keV										Nub16c **
* $^{105}\text{In}-^{104}\text{In}$	From $^{104}\text{In}/^{105}\text{In}=0.99050293(139)$										AHW **
* $^{105}\text{Te}(\alpha)^{101}\text{Sn}$	$E_\alpha=4720(50), 4703(5)$ to $7/2^+$ level at 171.7 keV (same group as next)										Ens07a **
* $^{105}\text{Te}(\alpha)^{101}\text{Sn}$	$E_\alpha=4711(3)$ to $7/2^+$ level at 171.7 keV; also $E_\alpha=4880(20)$ to ground state										Ens07a **
* $^{105}\text{Sb}(p)^{104}\text{Sn}$	F : expected 150 protons, no proton peak observed										05Li47 **
* $^{105}\text{Ag}(\epsilon)^{105}\text{Pd}$	$L/K=0.152(0.002) \rightarrow Q=222(12+\text{theor.error})$ to $3/2^-$ level at 1087.96 keV										Ens05b **
* $^{105}\text{Cd}(\beta^+)^{105}\text{Ag}$	$E_{\beta^+}=1691(5)$ to $^{105}\text{Ag}^m$ at 25.479 keV										Nub16b **
* $^{105}\text{Cd}(\beta^+)^{105}\text{Ag}$	$E_{\beta^+}=1695(11)$ to $^{105}\text{Ag}^m$ at 25.479 keV										Nub16b **
* $^{105}\text{In}(\beta^+)^{105}\text{Cd}$	$E_{\beta^+}=3696(13)$ to $7/2^+$ level at 131.11 keV										Ens05b **
$^{106}\text{Zr}-u$	-62674	322	-62860	470	-0.4	o			GT1	1.5	04Ma.A
	-62856	186				2			GT3	2.5	16Kn03
$^{106}\text{Nb}-u$	-70843	258	-71072	4	-0.6	U			GT1	1.5	04Ma.A
$^{106}\text{Mo}-^{97}\text{Zr}_{1.093}$	15589.8	9.8				2			JY1	1.0	06Ha03
$\text{C}_8 \text{H}_{10}-^{106}\text{Pd}$	174764.0	4.3	174770.0	1.2	0.6	U			M16	2.5	63Da10
	174751	32			0.4	U			R12	1.5	83De51
	174766	8			0.3	U			R12	1.5	83De51
$\text{C}_7 \text{ }^{13}\text{C} \text{H}_9-^{106}\text{Pd}$	170285	32	170299.8	1.2	0.3	U			R12	1.5	83De51
	170298	30			0.0	U			R12	1.5	83De51
$\text{C}_7 \text{ N} \text{H}_8-^{106}\text{Pd}$	162186	18	162194.0	1.2	0.3	U			R12	1.5	83De51
$\text{C}_7 \text{ O} \text{H}_6-^{106}\text{Pd}$	138378	20	138384.5	1.2	0.2	U			R12	1.5	83De51
$^{106}\text{Pd}-u$	-96495	30	-96519.7	1.2	-0.8	U			GS2	1.0	05Li24
	-96521.0	1.9			0.5	-			TG1	1.5	12Sm01
	-96524.9	4.7			0.7	-			TG1	1.5	12Sm01
	ave.	-96521.5	2.6		0.7	1	20	20 $^{106}\text{Pd}$			average
$^{106}\text{Ag}-u$	-93318	44	-93336	3	-0.4	U			GS2	1.0	05Li24 *
$\text{C}_8 \text{H}_{10}-^{106}\text{Cd}$	171789.3	2.7	171790.5	1.2	0.2	U			M16	2.5	63Da10
	171841	17			-2.0	U			R12	1.5	83De51
$\text{C}_7 \text{ N} \text{H}_8-^{106}\text{Cd}$	159210	15	159214.5	1.2	0.2	U			R12	1.5	83De51
$^{106}\text{Cd}-u$	-93540.6	1.7	-93540.2	1.2	0.2	-			TG1	1.5	12Sm01
	-93545.7	3.47			1.1	-			TG1	1.5	12Sm01
	ave.	-93541.6	2.3		0.6	1	27	27 $^{106}\text{Cd}$			average
$^{106}\text{Cd}-^{85}\text{Rb}_{1.247}$	16459.8	1.8	16458.0	1.2	-1.0	1	43	43 $^{106}\text{Cd}$	MA8	1.0	09Br09
$^{106}\text{In}-u$	-86516	32	-86536	13	-0.6	U			GS2	1.0	05Li24 *
$^{106}\text{Sn}-^{85}\text{Rb}_{1.247}$	26959.4	8.7	26956	5	-0.4	1	39	39 $^{106}\text{Sn}$	SH1	1.0	07Ma92
$^{106}\text{Sn}-^{87}\text{Rb}_{1.218}$	27578.0	7.6	27576	5	-0.3	1	52	52 $^{106}\text{Sn}$	JY1	1.0	09El07
$^{106}\text{Sb}-^{87}\text{Rb}_{1.218}$	39256.1	8.0				2			JY1	1.0	09El07
$^{106}\text{Nb}-^{102}\text{Ru}_{1.039}$	28318.2	4.4				2			JY1	1.0	07Ha32
$^{106}\text{Tc}-^{102}\text{Ru}_{1.039}$	13780.7	4.7	13747	13	-7.1	F			JY1	1.0	07Ha20 *
$^{106}\text{Ru}-^{105}\text{Ru}_{1.010}$	511.8	9.1	505	6	-0.7	1	42	37 $^{106}\text{Ru}$	JY1	1.0	07Ha20
$^{106}\text{Cd}-^{106}\text{Pd}$	2979.50	0.11	2979.50	0.11	0.0	1	100	70 $^{106}\text{Pd}$	SH1	1.0	11Go23
	2979.08	0.60			0.5	U			TG1	1.5	12Sm01
$^{106}\text{Pd}-^{105}\text{Pd}$	-1608	25	-1599.2	0.3	0.2	U			R12	1.5	83De51
$^{106}\text{Pd}-^{104}\text{Pd}$	-508	32	-550.1	0.8	-0.9	U			R12	1.5	83De51
$^{106}\text{Te}(\alpha)^{102}\text{Sn}$	4323.5	30.	4290	9	-1.1	U					81Sc17
	4290.2	9.				3					94Pa11
	4323.5	30.			-1.1	U					02Ma19
	4261.0	62.4			0.5	U					16Ca33
$^{106}\text{Cd}(\text{}^3\text{He}, \text{}^6\text{He})^{103}\text{Cd}$	-9173	17	-9141.4	2.1	1.9	U			MSU		78Pa11 *
$^{104}\text{Ru}(t,p)^{106}\text{Ru}$	5892	20	5888	5	-0.2	U			LAI		72Ca10
$^{104}\text{Pd}(\text{}^3\text{He}, p)^{106}\text{Ag}$	5153	6	5189.5	2.9	6.1	F			Bld		75An07 *

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{106}\text{Cd}(\text{p,t})^{104}\text{Cd}$	-10802	15	-10824.6	2.0	-1.5	U			MSU		82Cr01
	-10829	12			0.4	U			Pri		83De03
	-10819	12			-0.5	U			Ors		84Ro.A
$^{105}\text{Pd}(\text{n},\gamma)^{106}\text{Pd}$	9562.8	1.1	9560.96	0.28	-1.7	U					70Bo29
	9560.5	0.4			1.1	-			BNn		87Fo20 *
	9561.4	0.4			-1.1	-			Bdn		06Fi.A
$^{105}\text{Pd}(\text{d,p})^{106}\text{Pd}$	7349	30	7336.39	0.28	-0.4	U			Pit		64Co11
$^{106}\text{Pd}(\text{d,t})^{105}\text{Pd}$	-3311	25	-3303.73	0.28	0.3	U			Pit		64Co11
$^{105}\text{Pd}(\text{n},\gamma)^{106}\text{Pd}$	ave.	9560.95	0.28	9560.96	0.28	0.0	1	100	96 $^{105}\text{Pd}$		average
$^{105}\text{Pd}(\text{}^3\text{He,d})^{106}\text{Ag}$	322	8	320.0	2.8	-0.3	1	12	12 $^{106}\text{Ag}$	Bld		75An07
$^{106}\text{Cd}(\text{d,t})^{105}\text{Cd}$	-4661	50	-4612.4	1.8	1.0	U					73De16
$^{106}\text{Cd}(\text{}^3\text{He},\alpha)^{105}\text{Cd}$	9728	25	9708.0	1.8	-0.8	U			Man		75Ch21
$^{106}\text{Mo}(\beta^-)^{106}\text{Tc}$	3510	45	3642	15	2.9	U			Bwg		87Gr18
	3520	17			7.2	C			Bwg		92Gr.A
	3520	17			7.2	C			Jyv		94Jo.A
$^{106}\text{Tc}(\beta^-)^{106}\text{Ru}$	6545	45	6547	11	0.0	o			Bwg		87Gr18
	6547	11				2			Bwg		92Gr.A
$^{106}\text{Ru}(\beta^-)^{106}\text{Rh}$	39.2	0.3	39.40	0.21	0.7	-					50Ag01
	39.6	0.3			-0.7	-					58Gr07
	ave.	39.40	0.21		0.0	1	100	63 $^{106}\text{Ru}$			average
$^{106}\text{Rh}(\beta^-)^{106}\text{Pd}$	3530	10	3545	5	1.5	-					52Al06
	3550	10			-0.5	-					58Gr07
	3550	20			-0.3	-					60Se05
ave.	3541	7			0.6	1	64	63 $^{106}\text{Rh}$			average
$^{106}\text{Rh}^m(\beta^-)^{106}\text{Pd}$	3677	10				2					66De11 *
$^{106}\text{Ag}(\beta^+)^{106}\text{Pd}$	2980	20	2965.1	2.8	-0.7	U					53Be42
	3011	72			-0.6	U					86Bo28
$^{106}\text{Ag}(\epsilon)^{106}\text{Pd}$	2961	4			1.0	-					78Ge01 *
$^{106}\text{Pd}(\text{p,n})^{106}\text{Ag}$	-3754	13	-3747.5	2.8	0.5	U			Oak		64Jo11
	-3756	5			1.7	-					79De44
	ave.	2966	3	2965.1	2.8	-0.3	1	81	81 $^{106}\text{Ag}$		average
$^{106}\text{In}(\beta^+)^{106}\text{Cd}$	6516	30	6524	12	0.3	2					66Ca09 *
	6507	29			0.6	2					86Bo28 *
$^{106}\text{Cd}(\text{p,n})^{106}\text{In}$	-7312.9	15.	-7306	12	0.4	2			ANL		84Fi05 *
$^{106}\text{Sn}(\beta^+)^{106}\text{In}$	3195	60	3254	13	1.0	U			GSI		79PI06 *
	3200	100			0.5	U					88Ba10
* $^{106}\text{Ag}-\text{u}$	$M-A=-86880(32)$ keV for mixture gs+m at 89.66 keV										Nub16b **
* $^{106}\text{In}-\text{u}$	$M-A=-80575(29)$ keV for mixture gs+m at 28.6 keV										Nub16b **
* $^{106}\text{Tc}-^{102}\text{Ru}_{1.039}$	F : unidentified impurities in the trap										07Ha20 **
* $^{106}\text{Cd}(\text{}^3\text{He},\text{}^6\text{He})^{103}\text{Cd}$	First excited state at 187.89 keV yields $Q=-9146$ keV										GAu **
* $^{104}\text{Pd}(\text{}^3\text{He,p})^{106}\text{Ag}$	F : withdrawn by authors										AHW **
* $^{105}\text{Pd}(\text{n},\gamma)^{106}\text{Pd}$	Calculated from 13 $\gamma$ energies in 2 keV n-capture to levels in $^{106}\text{Pd}$ ; corrected for recoil										Ens086 **
*	$E_{\beta^-}=920(10)$ to $5^+$ level at 2757.06 keV										Ens086 **
* $^{106}\text{Rh}^m(\beta^-)^{106}\text{Pd}$	L/K=0.203(0.003) gives $Q_{\beta^+}=99(4)$ , recalculated $Q$										AHW **
* $^{106}\text{Ag}(\epsilon)^{106}\text{Pd}$	from $^{106}\text{Ag}^m 6^+$ at 89.66 keV to $5^+$ level at 2951.84 keV										Ens086 **
*	$E_{\beta^+}=4890(30)$ from $^{106}\text{In}^m 2^+$ at 28.6 to $2^+$ level at 632.64 keV										Ens086 **
* $^{106}\text{In}(\beta^+)^{106}\text{Cd}$	$E_{\beta^+}=2965(30)$ to $6^+$ level at 2491.66 keV and 4908(29) from										Ens086 **
* $^{106}\text{In}(\beta^+)^{106}\text{Cd}$	$^{106}\text{In}^m (2)^+$ at 28.6 to $2^+$ level at 632.64 keV										Ens086 **
*	T=7535(15) to $2^+$ level at 151.1 keV										Ens086 **
* $^{106}\text{Cd}(\text{p,n})^{106}\text{In}$	$p^+=0.253(0.021)$ to $1^+$ level at 893.0 keV										Ens086 **
* $^{106}\text{Sn}(\beta^+)^{106}\text{In}$											Ens086 **
$^{107}\text{Zr}-\text{u}$	-58379	482				2			GT3	2.5	16Kn03
$^{107}\text{Nb}-\text{u}$	-68326	279	-68410	9	-0.2	U			GT1	1.5	04Ma.A
$^{107}\text{Mo}-^{97}\text{Zr}_{1.103}$	20326.7	9.9				2			JY1	1.0	06Ha03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{107}\text{Pd-u}$	-95013	95	-94871.9	1.3	1.5	U			GS2	1.0	05Li24 *
$\text{C}_8 \text{H}_{11}-^{107}\text{Ag}$	180986.4	3.1	180983.8	2.6	-0.3	1	11	$^{11}^{107}\text{Ag}$	M16	2.5	63Da10
	180994	17			-0.4	U			R12	1.5	83De51
$\text{C}_7 \text{N H}_9-^{107}\text{Ag}$	168415	8	168407.8	2.6	-0.6	U			R12	1.5	83De51
$\text{C}_7 \text{O H}_7-^{107}\text{Ag}$	144595	18	144598.3	2.6	0.1	U			R12	1.5	83De51
$\text{C}_6 \text{ }^{13}\text{C O H}_6-^{107}\text{Ag}$	140131	16	140128.1	2.6	-0.1	U			R12	1.5	83De51
$\text{C}_6 \text{N O H}_5-^{107}\text{Ag}$	132025	16	132022.3	2.6	-0.1	U			R12	1.5	83De51
$^{107}\text{Cd-u}$	-93410	30	-93387.9	1.8	0.7	U			GS2	1.0	05Li24
$^{107}\text{Cd}-^{85}\text{Rb}_{1.259}$	17668.7	1.9	17668.8	1.8	0.1	1	88	$^{88}^{107}\text{Cd}$	MA8	1.0	09Br09
$^{107}\text{In-u}$	-89710	30	-89710	12	0.0	U			GS2	1.0	05Li24
$^{107}\text{Sn}-^{87}\text{Rb}_{1.230}$	27421.6	5.7				2			JY1	1.0	09El07
$^{107}\text{Sb}-^{87}\text{Rb}_{1.230}$	35866.2	5.8	35859	4	-1.3	1	59	$^{59}^{107}\text{Sb}$	JY1	1.0	09El07
$^{107}\text{Sb}-^{133}\text{Cs}_{.805}$	251.8	9.7	262	4	1.0	1	21	$^{21}^{107}\text{Sb}$	SH1	1.0	07Ma92
$^{107}\text{Nb}-^{102}\text{Ru}_{1.049}$	31936.7	8.6				2			JY1	1.0	07Ha32
$^{107}\text{Tc}-^{105}\text{Ru}_{1.019}$	9465.8	8.9				2			JY1	1.0	07Ha20
$^{107}\text{Ru}-^{105}\text{Ru}_{1.019}$	3977.2	8.9				2			JY1	1.0	07Ha20
$^{107}\text{Sn}-^{106}\text{Sn}$	-1148	86	-1244	8	-0.7	U			CR2	1.5	92Sh.A *
$^{107}\text{Te}(\alpha)^{103}\text{Sn}$	3982.2	15.	4008	5	1.7	3					79Se22
	4011.3	5.			-0.6	3					91He21
$^{107}\text{Ag}(\text{p,t})^{105}\text{Ag}$	-9015	15	-8997	5	1.2	1	11	$^9^{105}\text{Ag}$	Min		75Ku14 *
$^{106}\text{Pd}(\text{n},\gamma)^{107}\text{Pd}$	6536.4	0.5	6536.4	0.5	0.1	1	99	$^{94}^{107}\text{Pd}$	Bdn		06Fi.A
$^{107}\text{Ag}(\gamma,\text{n})^{106}\text{Ag}$	-9353	34	-9536	4	-5.4	B			Phi		60Ge01
$^{107}\text{Ag}(\text{p,d})^{106}\text{Ag}$	-7305	11	-7311	4	-0.5	1	10	$^7^{106}\text{Ag}$	Bld		75An07
$^{107}\text{Mo}(\beta^-)^{107}\text{Tc}$	6160	60	6198	13	0.6	U			Bwg		89Gr23
$^{107}\text{Tc}(\beta^-)^{107}\text{Ru}$	4820	85	5113	12	3.4	B			Bwg		89Gr23
$^{107}\text{Ru}(\beta^-)^{107}\text{Rh}$	3140	300	3001	15	-0.5	U					62Pi02 *
	2900	135			0.7	U			Bwg		89Gr23
$^{107}\text{Rh}(\beta^-)^{107}\text{Pd}$	1510	40	1509	12	0.0	U					62Pi02 *
$^{107}\text{Pd}(\beta^-)^{107}\text{Ag}$	33	3	34.0	2.3	0.3	1	60	$^{53}^{107}\text{Ag}$	49Pa.B		62La10 *
$^{107}\text{Cd}(\beta^+)^{107}\text{Ag}$	1417	4	1416.4	2.6	-0.1	1	41	$^{30}^{107}\text{Ag}$			62La10 *
$^{107}\text{In}(\beta^+)^{107}\text{Cd}$	3426	11				2					86Bo28 *
* $^{107}\text{Pd-u}$	$M - A = -88397(62)$ keV for mixture gs+n at 214.6 keV										Nub16b **
* $^{107}\text{Sn}-^{106}\text{Sn}$	From $^{107}\text{Sn}/^{106}\text{Sn} = 1.00943053(81)$										AHW **
* $^{107}\text{Ag}(\text{p,t})^{105}\text{Ag}$	Recalibrated with (p,t) results on $^{104}\text{Pd}$ , $^{105}\text{Pd}$ , $^{106}\text{Pd}$ and $^{108}\text{Pd}$										AHW **
* $^{107}\text{Ru}(\beta^-)^{107}\text{Rh}$	$E_{\beta^-} = 2100(300)$ to $(5/2^+, 7/2^+)$ level at 1041.950 keV										Ens085 **
* $^{107}\text{Rh}(\beta^-)^{107}\text{Pd}$	$E_{\beta^-} = 840(40)$ to $5/2^+$ level at 670.06 keV										Ens085 **
* $^{107}\text{Cd}(\beta^+)^{107}\text{Ag}$	$E_{\beta^+} = 302(4)$ to $^{107}\text{Ag}^m$ at 93.125 keV										Nub16b **
* $^{107}\text{In}(\beta^+)^{107}\text{Cd}$	$E_{\beta^+} = 2199(11)$ to $5/2^+$ level at 204.98 keV										Ens085 **
$^{108}\text{Nb-u}$	-64413	440	-63925	9	0.7	o			GT1	1.5	04Ma.A
	-63945	112			0.1	U			GT2	2.5	08Kn.A
$^{108}\text{Nb}-^{120}\text{Sn}_{.900}$	24093.3	8.8				2			JY1	1.0	11Ha48
$^{108}\text{Mo}-^{97}\text{Zr}_{1.113}$	23144.8	9.9				2			JY1	1.0	06Ha03
$^{108}\text{Mo-u}$	-76039	215	-75960	10	0.2	U			GT1	1.5	04Ma.A
$^{108}\text{Rh}-^{120}\text{Sn}_{.900}$	-3267	15				2			JY1	1.0	07Ha20
$^{108}\text{Rh}^m-^{120}\text{Sn}_{.900}$	-3144	13				2			JY1	1.0	07Ha20
$\text{C}_8 \text{H}_{12}-^{108}\text{Pd}$	190014	6	190008.6	1.2	-0.4	U			M16	2.5	63Da10
	190005	19			0.1	U			R12	1.5	83De51
$\text{C}_7 \text{ }^{13}\text{C H}_{11}-^{108}\text{Pd}$	185532	30	185538.4	1.2	0.1	U			R12	1.5	83De51
$\text{C}_7 \text{N H}_{10}-^{108}\text{Pd}$	177422	17	177432.5	1.2	0.4	U			R12	1.5	83De51
$\text{C}_6 \text{ }^{13}\text{C N H}_9-^{108}\text{Pd}$	172943	18	172962.3	1.2	0.7	U			R12	1.5	83De51
$\text{C}_7 \text{O H}_8-^{108}\text{Pd}$	153611	17	153623.1	1.2	0.5	U			R12	1.5	83De51
$\text{C}_6 \text{N O H}_6-^{108}\text{Pd}$	141031	16	141047.0	1.2	0.7	U			R12	1.5	83De51
$^{108}\text{Pd-u}$	-96109.6	1.3	-96108.2	1.2	0.7	-			TG1	1.5	12Sm01
	-96108.1	4.7			0.0	-			TG1	1.5	12Sm01
ave.	-96109.5	1.9			0.7	1	40	$^{40}^{108}\text{Pd}$			average



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{108}\text{Ag}-u$	-93973	50	-94049.7	2.6	-1.5	U			GS2	1.0	05Li24 *
$^{108}\text{Ag}-^{133}\text{Cs}_{.812}$	-17218	84	-17276.7	2.6	-0.7	U			MA8	1.0	08Br.A *
$\text{C}_8 \text{H}_{12}-^{108}\text{Cd}$	189715.6	2.9	189716.8	1.2	0.2	U			M16	2.5	63Da10
	189695	16			0.9	U			R12	1.5	83De51
$\text{C}_7 \text{N H}_{10}-^{108}\text{Cd}$	177140	30	177140.7	1.2	0.0	U			R12	1.5	83De51
$\text{C}_6 \text{ }^{13}\text{C N H}_9-^{108}\text{Cd}$	172653	15	172670.5	1.2	0.8	U			R12	1.5	83De51
$\text{C}_6 \text{ N O H}_6-^{108}\text{Cd}$	140746	15	140755.2	1.2	0.4	U			R12	1.5	83De51
$^{108}\text{Cd}-^{85}\text{Rb}_{1.271}$	16298.5	2.3	16298.8	1.2	0.1	1	27	27 $^{108}\text{Cd}$	MA8	1.0	09Br09
$^{108}\text{Cd}-u$	-95818.3	1.7	-95816.4	1.2	0.7	-			TG1	1.5	12Sm01
	-95817.1	4.87			0.1	-			TG1	1.5	12Sm01
	ave.	-95818.2	2.4		0.7	1	25	25 $^{108}\text{Cd}$			average
$^{108}\text{In}-u$	-90277	31	-90306	9	-0.9	U			GS2	1.0	05Li24 *
$^{108}\text{Sn}-u$	-88102	32	-88106	6	-0.1	U			GS2	1.0	05Li24
$^{108}\text{Sn}-^{87}\text{Rb}_{1.241}$	24599.8	5.9	24601	6	0.2	1	96	96 $^{108}\text{Sn}$	JY1	1.0	09EI07
$^{108}\text{Sb}-^{87}\text{Rb}_{1.241}$	34933.7	5.9				2			JY1	1.0	09EI07
$^{108}\text{Te}-^{87}\text{Rb}_{1.241}$	42085.3	6.0	42087	6	0.4	1	94	94 $^{108}\text{Te}$	JY1	1.0	09EI07
$^{108}\text{Tc}-^{105}\text{Ru}_{1.029}$	13423.4	9.0				2			JY1	1.0	07Ha20
$^{108}\text{Ru}-^{105}\text{Ru}_{1.029}$	5115.7	8.9				2			JY1	1.0	07Ha20
$^{108}\text{Pd}-^{108}\text{Cd}$	-292.05	0.59	-291.8	0.8	0.3	1	87	46 $^{108}\text{Cd}$	TG1	1.5	12Sm01
$^{108}\text{Sn}-^{107}\text{Sn}$	-3650	76	-3819	8	-1.5	U			CR2	1.5	92Sh.A *
$^{108}\text{Pd}-^{106}\text{Pd}$	425	40	411.5	1.7	-0.2	U			R12	1.5	83De51
$^{108}\text{Cd}-^{106}\text{Cd}$	-2232	32	-2276.2	1.7	-0.9	U			R12	1.5	83De51
$^{108}\text{Te}(\alpha)^{104}\text{Sn}$	3406.4	30.	3420	8	0.5	U					65Ma12
	3448.0	20.			-1.3	1	13	7 $^{104}\text{Sn}$			81Sc17
	3444.9	4.			-6.1	B					91He21
	3406.4	25.			0.6	U					91Pa05
$^{108}\text{I}(\alpha)^{104}\text{Sb}$	4034.7	25.	4100	50	1.3	F					91Pa05 *
	4099.1	5.				4					94Pa12
$^{108}\text{Pd}(d,^3\text{He})^{107}\text{Rh}$	-4456	12				2			Grn		86Ka43
$^{108}\text{Pd}(d,t)^{107}\text{Pd}$	-2977	30	-2965.7	1.6	0.4	U			Pit		64Co11
$^{107}\text{Ag}(n,\gamma)^{108}\text{Ag}$	7269.6	0.6	7271.41	0.17	3.0	B			ILn		85Ma54 Z
	7271.41	0.17				2			Bdn		06Fi.A
$^{107}\text{Ag}(d,p)^{108}\text{Ag}$	5051	7	5046.84	0.17	-0.6	U			MIT		67Sp09
$^{108}\text{Mo}(\beta^-)^{108}\text{Tc}$	5135	60	5167	13	0.5	U			Bwg		92Gr.A
	5120	40			1.2	o					94Jo.A
	5100	60			1.1	U					95Jo02
$^{108}\text{Tc}(\beta^-)^{108}\text{Ru}$	7720	50	7739	12	0.4	U			Bwg		89Gr23
$^{108}\text{Ru}(\beta^-)^{108}\text{Rh}$	1315	100	1370	16	0.6	U					62Pi02 *
	1420	185			-0.3	U			Bwg		89Gr23
	1380	80			-0.1	o			Jyv		92Jo05
	1350	60			0.3	U			Jyv		94Jo.A
$^{108}\text{Rh}(\beta^-)^{108}\text{Pd}$	4500	600	4492	14	0.0	U					62Pi02
	4505	105			-0.1	U			Bwg		89Gr23
$^{108}\text{Rh}^m(\beta^-)^{108}\text{Pd}$	4434	50	4607	12	3.5	B					69Pi08 *
	4510	100			1.0	U					84Bh02 *
$^{108}\text{Ag}(\beta^+)^{108}\text{Pd}$	1902	25	1917.4	2.6	0.6	U					62Fr07
$^{108}\text{Pd}(p,n)^{108}\text{Ag}$	-2675	100	-2699.8	2.6	-0.2	U			Oak		64Jo11
$^{108}\text{Ag}(\beta^-)^{108}\text{Cd}$	1650	40	1645.7	2.6	-0.1	U					60Wa10
$^{108}\text{In}(\beta^+)^{108}\text{Cd}$	5124	50	5133	9	0.2	U					62Ka23 *
	5125	14			0.5	-					86Bo28 *
$^{108}\text{Cd}(p,n)^{108}\text{In}$	-5927	12	-5915	9	1.0	-			ANL		84Fi05 *
$^{108}\text{In}(\beta^+)^{108}\text{Cd}$	ave.	5136	9	5133	9	-0.4	1	89	89 $^{108}\text{In}$		average
$^{108}\text{Sn}(\beta^+)^{108}\text{In}$	2078	25	2050	10	-1.1	1	15	11 $^{108}\text{In}$	GSI		79Pi06 *
* $^{108}\text{Ag}-u$	$M - A = -87480(34)$ keV for mixture gs+m at 109.466 keV										
* $^{108}\text{Ag}-^{133}\text{Cs}_{.812}$	$D_M = -17159(77)$ $\mu\text{u}$ for mixture gs+m at 109.466 keV; $M - A = -87497(72)$ keV										
* $^{108}\text{In}-u$	$M - A = -84078(28)$ keV for mixture gs+m at 29.75 keV										
* $^{108}\text{Sn}-^{107}\text{Sn}$	From $^{107}\text{Sn}/^{108}\text{Sn} = 0.99076701(70)$										
* $^{108}\text{I}(\alpha)^{104}\text{Sb}$	F : Same authors say: Consistent with new value, if recalibrated										
* $^{108}\text{Ru}(\beta^-)^{108}\text{Rh}$	$E_{\beta^-} = 1150(100)$ to $1^+$ level at 164.98 keV										
* $^{108}\text{Rh}^m(\beta^-)^{108}\text{Pd}$	$E_{\beta^-} = 1570(50)$ to $(4^+, 5^+, 6^+)$ level at 2863.70 keV										
* $^{108}\text{Rh}^m(\beta^-)^{108}\text{Pd}$	$E_{\beta^-} = 1970(100)$ to $4^+$ level at 2540.2 keV										
* $^{108}\text{In}(\beta^+)^{108}\text{Cd}$	$E_{\beta^+} = 1290(80)$ to $6^+$ level at 2807.81 keV, and $E_{\beta^+} = 3500(50)$ from $^{108}\text{In}^m$										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* $^{108}\text{In}(\beta^+)^{108}\text{Cd}$	at 29.75 to 2 <sup>+</sup> level at 632.986 keV								Nub16b **
* $^{108}\text{Cd}(\text{p,n})^{108}\text{In}$	$E_{\beta^+}=1887(28)$ to 4 <sup>+</sup> level at 2239.35 keV, and $E_{\beta^+}=3494(14)$ from $^{108}\text{In}^m$ at 29.75 to 2 <sup>+</sup> level at 632.986 keV								Ens08a **
* $^{108}\text{Sn}(\beta^+)^{108}\text{In}$	$Q(\text{not T, PrvCom Fi})=-6191(8), -6244(9)$ , errors statistical only, to 3 <sup>+</sup> levels at respectively, 198.36 and 266.06 keV								AHW **
	$p^+ =35(6)\times 10^{-4}$ to 1 <sup>+</sup> level at 698.85 keV								Ens08a **
$^{109}\text{Nb}-u$	-60784 376	-60860 280	-0.1	o			GT1	1.5	04Ma.A
	-60859 185			2			GT1	1.5	16Kn03
$^{109}\text{Mo}-^{97}\text{Zr}_{1.124}$	28515 12			2			JY1	1.0	06Ha03
$^{109}\text{Mo}-u$	-71552 247	-71569 12	0.0	U			GT1	1.5	04Ma.A
$^{109}\text{Rh}-^{120}\text{Sn}_{908}$	-2463.6 7.2	-2450 4	1.9	1	37	36 $^{109}\text{Rh}$	JY1	1.0	07Ha20
$\text{C}_8 \text{H}_{13}-^{109}\text{Ag}$	196972.1 3.8	196969.6 1.4	-0.3	U			M16	2.5	63Da10
	196972 6		-0.3	U			R12	1.5	83De51
$\text{C}_6 \text{ }^{13}\text{C O H}_8-^{109}\text{Ag}$	156110 16	156113.9 1.4	0.2	U			R12	1.5	83De51
$\text{C}_6 \text{ N O H}_7-^{109}\text{Ag}$	148006 16	148008.1 1.4	0.1	U			R12	1.5	83De51
$^{109}\text{Ag}-^{133}\text{Cs}_{820}$	-17766 83	-17714.8 1.4	0.6	U			MA8	1.0	08Br.A *
$^{109}\text{Cd}-^{85}\text{Rb}_{1.282}$	18072.9 1.9	18072.3 1.6	-0.3	1	75	75 $^{109}\text{Cd}$	MA8	1.0	09Br09
$^{109}\text{Sn}-u$	-88747 30	-88707 9	1.3	U			GS2	1.0	05Li24
$^{109}\text{Sb}-^{87}\text{Rb}_{1.253}$	31937.9 5.9	31938 6	0.0	1	92	92 $^{109}\text{Sb}$	JY1	1.0	09El07
$^{109}\text{Sb}-^{133}\text{Cs}_{820}$	-4360 19	-4329 6	1.6	U			SH1	1.0	07Ma92
$^{109}\text{Te}-^{87}\text{Rb}_{1.253}$	41101.6 6.4	41101 5	0.0	1	54	54 $^{109}\text{Te}$	JY1	1.0	09El07
$^{109}\text{Te}-^{133}\text{Cs}_{820}$	4835.6 8.3	4834 5	-0.2	1	32	32 $^{109}\text{Te}$	SH1	1.0	07Ma92
$^{109}\text{Tc}-^{105}\text{Ru}_{1.038}$	16014.3 10.0			2			JY1	1.0	07Ha20
$^{109}\text{Ru}-^{105}\text{Ru}_{1.038}$	9083.9 9.2			2			JY1	1.0	07Ha20
$^{109}\text{Ag}-^{107}\text{Ag}$	-335 28	-335.8 2.9	0.0	U			R12	1.5	83De51
$^{109}\text{Te}(\alpha)^{105}\text{Sn}$	3197.6 30.	3198 6	0.0	U					65Ma12
	3197.6 15.		0.0	1	13	7 $^{109}\text{Te}$			79Sc22
	3225.6 4.		-7.0	C					91He21
$^{109}\text{I}(\alpha)^{105}\text{Sb}$	3918.1 20.8			3			ORa		07Ma35
$^{109}\text{Xe}(\alpha)^{105}\text{Te}$	4217.0 7.3			4					06Li41 *
$^{109}\text{Ag}(\text{p,t})^{107}\text{Ag}$	-7995 15	-7973.6 2.7	1.4	U			Min		75Ku14 *
$^{108}\text{Pd}(\text{n},\gamma)^{109}\text{Pd}$	6153.8 0.3	6153.59 0.15	-0.7	-			ILn		80Ca02 Z
	6153.54 0.17		0.3	-			Bdn		06Fi.A
$^{108}\text{Pd}(\text{d,p})^{109}\text{Pd}$	3936 30	3929.02 0.15	-0.2	U			Pit		64Co11
$^{108}\text{Pd}(\text{n},\gamma)^{109}\text{Pd}$	ave. 6153.60 0.15	6153.59 0.15	-0.1	1	100	81 $^{109}\text{Pd}$			average
$^{109}\text{Ag}(\gamma,\text{n})^{108}\text{Ag}$	-9196 26	-9184.0 2.7	0.5	U			Phi		60Ge01
$^{109}\text{Ag}(\text{d,t})^{108}\text{Ag}$	-2947 30	-2926.8 2.7	0.7	U			Pit		64Co11
$^{108}\text{Cd}(\text{}^3\text{He,d})^{109}\text{In}-^{110}\text{Cd}(\text{}^{111}\text{In}$	-806.5 2.6	-807.0 2.5	-0.2	1	91	70 $^{109}\text{In}$			80Ta07
$^{109}\text{Te}(\epsilon\text{p})^{108}\text{Sn}$	7140 60	7066 7	-1.2	U					73Bo20
$^{109}\text{I}(\text{p})^{108}\text{Te}$	819 5	820 4	0.2	o					84Fa04
	819.6 2.0		0.3	o					92He.A
	820.2 4.0			2					95Ho26
$^{109}\text{Tc}(\beta^-)^{109}\text{Ru}$	6315 70	6456 13	2.0	U			Bwg		89Gr23
$^{109}\text{Ru}(\beta^-)^{109}\text{Rh}$	4160 65	4261 10	1.6	U			Bwg		89Gr23
$^{109}\text{Rh}(\beta^-)^{109}\text{Pd}$	2577 50	2607 4	0.6	U					78Ka10 *
$^{109}\text{Pd}(\beta^-)^{109}\text{Ag}$	1116 2	1112.9 1.4	-1.5	1	49	30 $^{109}\text{Ag}$			62Br15 *
$^{109}\text{Cd}(\epsilon)^{109}\text{Ag}$	182 3	215.1 1.8	11.0	C					68Go.A *
	214 3		0.4	1	35	22 $^{109}\text{Cd}$			Averag *
$^{109}\text{In}(\beta^+)^{109}\text{Cd}$	2015 8	2015 4	0.0	-					62No06 *
	2030 15		-1.0	-					71Ba08 *
	ave. 2018 7		-0.5	1	33	30 $^{109}\text{In}$			average
$^{109}\text{Sn}(\beta^+)^{109}\text{In}$	4120 50	3859 9	-5.2	B					70Sh05
$^{109}\text{Sb}(\beta^+)^{109}\text{Sn}$	6380 16	6379 9	0.0	1	30	22 $^{109}\text{Sn}$			82Jo03 *
* $^{109}\text{Ag}-^{133}\text{Cs}_{820}$	$D_M=-17719(78) \mu\text{u}$ for mixture gs+m at 88.0341 keV; $M-A=-88723(73)$ keV								Nub16b **
* $^{109}\text{Xe}(\alpha)^{105}\text{Te}$	Also $E_\alpha=3918(9)$ keV to 150(13) level								06Li41 **
* $^{109}\text{Ag}(\text{p,t})^{107}\text{Ag}$	Recalibrated with (p,t) results on $^{104}\text{Pd}$ , $^{105}\text{Pd}$ , $^{106}\text{Pd}$ and $^{108}\text{Pd}$								AHW **
* $^{109}\text{Rh}(\beta^-)^{109}\text{Pd}$	$E_{\beta^-}=2250(50)$ to 5/2 <sup>+</sup> level at 326.869 keV								Ens062 **
* $^{109}\text{Pd}(\beta^-)^{109}\text{Ag}$	$E_{\beta^-}=1028(2)$ to $^{109}\text{Ag}^m$ at 88.0341 keV								Nub16b **
* $^{109}\text{Cd}(\epsilon)^{109}\text{Ag}$	IBE=68(3) gives 94(3) to $^{109}\text{Ag}^m$ at 88.0341 keV								Nub16b **
* $^{109}\text{Cd}(\epsilon)^{109}\text{Ag}$	From average LM/K=0.2265(0.0026) $\rightarrow Q_{\beta^+}=126(3)$ ; recalculated $Q$								AHW **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
*	to $^{109}\text{Ag}^m$ at 88.0341 keV										Nub16b **
*	LMN/K=0.228(0.003)										65Le06 **
*	L/K=0.195(0.005) $\rightarrow$ LMN/K=0.258(0.006) $\rightarrow Q_{\beta^+}=109(5)$ not used										65Le06 **
*	LMN/K=0.226(0.003)										70Go39 **
* $^{109}\text{In}(\beta^+)^{109}\text{Cd}$	$E_{\beta^+}=790(8)$ 805(15) respectively, to $7/2^+$ level at 203.30 keV										Ens062 **
* $^{109}\text{Sb}(\beta^+)^{109}\text{Sn}$	$E_{\beta^+}=4416(21)$ to $3/2^+$ level at 925.6 keV, and other $E_{\beta^+}$										Ens062 **
$^{110}\text{Nb}-u$	-56157	360				2			GT3	2.5	16Kn03
$^{110}\text{Mo}-^{97}\text{Zr}_{1.134}$	31685	26				2			JY1	1.0	06Ha03
$^{110}\text{Mo}-u$	-69544	268	-69289	26	0.6	U			GT1	1.5	04Ma.A
$^{110}\text{Ru}-u$	-85897	78	-85961	10	-0.8	U			JY0	1.0	03Ko.A
$^{110}\text{Rh}-u$	-88835	130	-88920	19	-0.7	U			JY0	1.0	03Ko.A *
$^{110}\text{Rh}-^{120}\text{Sn}_{917}$	815	110	761	19	-0.5	U			JY1	1.0	07Ha20 *
$\text{C}_8 \text{H}_{14}-^{110}\text{Pd}$	204389	9	204377.6	0.7	-0.5	U			M16	2.5	63Da10
	204380	20			-0.1	U			R12	1.5	83De51
$\text{C}_7 \text{ }^{13}\text{C} \text{H}_{13}-^{110}\text{Pd}$	199913	20	199907.4	0.7	-0.2	U			R12	1.5	83De51
$\text{C}_6 \text{N O H}_8-^{110}\text{Pd}$	155418	17	155416.0	0.7	-0.1	U			R12	1.5	83De51
$\text{C}_5 \text{ }^{13}\text{C} \text{N O H}_7-^{110}\text{Pd}$	150946	17	150945.8	0.7	0.0	U			R12	1.5	83De51
$\text{C}_6 \text{O}_2 \text{H}_6-^{110}\text{Pd}$	131609	18	131606.6	0.7	-0.1	U			R12	1.5	83De51
$^{110}\text{Pd}-u$	-94829.7	1.5	-94827.1	0.7	1.7	-			MA8	1.0	12Fi01
	-94829.5	1.7			0.9	-			TG1	1.5	12Sm01
	-94830.9	3.0			0.8	-			TG1	1.5	12Sm01
ave.	-94829.7	1.2			2.1	1	28	28 $^{110}\text{Pd}$			average
$\text{C}_8 \text{H}_{14}-^{110}\text{Cd}$	206548.4	4.6	206543.0	0.4	-0.5	U			M16	2.5	63Da10
	206550	45			-0.1	U			R12	1.5	83De51
	206569	13			-1.3	U			R12	1.5	83De51
$\text{C}_7 \text{ }^{13}\text{C} \text{H}_{13}-^{110}\text{Cd}$	202093	14	202072.8	0.4	-1.0	U			R12	1.5	83De51
	202053	28			0.5	U			R12	1.5	83De51
$\text{C}_7 \text{O H}_{10}-^{110}\text{Cd}$	170156	16	170157.5	0.4	0.1	U			R12	1.5	83De51
$\text{C}_6 \text{N O H}_8-^{110}\text{Cd}$	157614	17	157581.4	0.4	-1.3	U			R12	1.5	83De51
$\text{C}_5 \text{ }^{13}\text{C} \text{N O H}_7-^{110}\text{Cd}$	153131	17	153111.2	0.4	-0.8	U			R12	1.5	83De51
$\text{C}_6 \text{O}_2 \text{H}_6-^{110}\text{Cd}$	133801	18	133772.0	0.4	-1.1	U			R12	1.5	83De51
$\text{C}_9 \text{H}_2-^{110}\text{Cd}$	112661	19	112642.6	0.4	-0.6	U			R12	1.5	83De51
$^{110}\text{Cd}-u$	-96993.6	1.5	-96992.5	0.4	0.7	-			MA8	1.0	12Fi01
	-96997.0	1.5			2.0	-			TG1	1.5	12Sm01
	-96992.2	2.4			-0.1	-			TG1	1.5	12Sm01
ave.	-96994.4	1.2			1.6	1	12	12 $^{110}\text{Cd}$			average
$^{110}\text{In}-u$	-92898	36	-92829	12	1.9	U			GS2	1.0	05Li24 *
$^{110}\text{Sn}-u$	-92189	30	-92155	15	1.1	2			GS2	1.0	05Li24
$^{110}\text{Sb}-^{87}\text{Rb}_{1.264}$	31650.1	6.4				2			JY1	1.0	09El07
$^{110}\text{Te}-^{133}\text{Cs}_{827}$	643.8	7.7	649	7	0.7	1	84	84 $^{110}\text{Te}$	SH1	1.0	07Ma92
$^{110}\text{Tc}-^{105}\text{Ru}_{1.048}$	20424.0	9.8				2			JY1	1.0	07Ha20
$^{110}\text{Ru}-^{105}\text{Ru}_{1.048}$	10719.5	9.3	10721	9	0.2	1	97	97 $^{110}\text{Ru}$	JY1	1.0	07Ha20
$^{110}\text{Cd }^{35}\text{Cl}-^{108}\text{Cd }^{37}\text{Cl}$	1764	5	1774.0	1.3	0.5	U			H11	4.0	63Bi12
$^{110}\text{Pd}-^{110}\text{Cd}$	2166.24	0.69	2165.4	0.6	-1.2	-			MA8	1.0	12Fi01
	2166.2	1.3			-0.4	-			TG1	1.5	12Sm01
ave.	2166.2	0.7			-1.3	1	80	71 $^{110}\text{Pd}$			average
$^{110}\text{Pd}-^{108}\text{Pd}$	1288	35	1281.1	1.3	-0.1	U			R12	1.5	83De51
$^{110}\text{Cd}-^{108}\text{Cd}$	-1219	34	-1176.1	1.3	0.8	U			R12	1.5	83De51
$^{110}\text{Te}(\alpha)^{106}\text{Sn}$	2723.2	15.6	2699	8	-1.6	1	25	16 $^{110}\text{Te}$			81Sc17
$^{110}\text{I}(\alpha)^{106}\text{Sb}$	3574.2	10.	3580	50	0.2	3					81Sc17
	3586.7	5.			-0.1	3					91He21
$^{110}\text{Xe}(\alpha)^{106}\text{Te}$	3878.3	30.	3872	9	-0.2	4					81Sc17
	3886.6	15.			-0.9	4					92He.A
	3871.0	30.			0.0	4					02Ma19
	3857.5	19.7			0.7	4			Jya		07Sa36
	3860.7	20.8			0.6	4					16Ca33

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{110}\text{Pd}(p,t)^{108}\text{Pd}$	-6495	15	-6467.5	1.3	1.8	U			Min		75Ku14 *
$^{110}\text{Pd}(d,^3\text{He})^{109}\text{Rh}$	-5134	5	-5127	4	1.4	1	65	64 $^{109}\text{Rh}$	VUn		87Ka29
$^{110}\text{Pd}(t,\alpha)^{109}\text{Rh}$	9206	25	9193	4	-0.5	U			LAl		82Fl09
$^{109}\text{Ag}(n,\gamma)^{110}\text{Ag}$	6809.2	0.1	6809.19	0.10	-0.1	1	100	57 $^{109}\text{Ag}$			81Bo.B
	6808.20	0.16			6.2	C			Bdn		06Fi.A
$^{109}\text{Ag}(d,p)^{110}\text{Ag}$	4590	5	4584.63	0.10	-1.1	U			MIT		64Sp12
$^{110}\text{Cd}(d,t)^{109}\text{Cd}$	-3664	30	-3657.7	1.6	0.2	U			Pit		64Ro17
$^{110}\text{Tc}(\beta^-)^{110}\text{Ru}$	6680	120	9038	13	19.7	C			Jyv		89Jo.A
	9021	55			0.3	U			Jyv		00Kr.A
$^{110}\text{Ru}(\beta^-)^{110}\text{Rh}$	2810	50	2756	19	-1.1	1	15	12 $^{110}\text{Rh}$	Jyv		91Jo11 *
$^{110}\text{Rh}(\beta^-)^{110}\text{Pd}$	5500	500	5502	18	0.0	U					63Ka21
	5400	100			1.0	U					70Pi01 *
	5510	19			-0.4	1	88	88 $^{110}\text{Rh}$	Bwg		00Kr.A
$^{110}\text{Ag}(\beta^-)^{110}\text{Cd}$	2891.4	3.0	2890.7	1.3	-0.2	-					63Da03 *
	2892.9	2.0			-1.1	-					67Mo12 *
ave.	2892.4	1.7			-1.1	1	59	57 $^{110}\text{Ag}$			average
$^{110}\text{In}(\beta^+)^{110}\text{Cd}$	3928	20	3878	12	-2.5	2					51Mc11 *
	3868	20			0.5	2					53Bl44 *
	3838	20			2.0	2					62Ka08 *
$^{110}\text{Sb}(\beta^+)^{110}\text{Sn}$	8750	200	8392	15	-1.8	U					72Mi26 *
	9085	100			-6.9	B					72Si28 *
* $^{110}\text{Rh}-u$	$M-A=-82639(73)$ keV for mixture gs+m at 220#(150#) keV										Nub16b **
* $^{110}\text{Rh}-^{120}\text{Sn}_{917}$	$D_M=933.3(7.2)$ $\mu\text{u}$ for mixture gs+m at 220#(150#) keV; $M-A=-82668.1(6.8)$ keV										Nub16b **
* $^{110}\text{In}-u$	$M-A=-86503(28)$ keV for mixture gs+m at 62.08 keV										Nub16b **
* $^{110}\text{Pd}(p,t)^{108}\text{Pd}$	Recalibrated with (p,t) results on $^{104}\text{Pd}$ , $^{105}\text{Pd}$ , $^{106}\text{Pd}$ and $^{108}\text{Pd}$										AHW **
* $^{110}\text{Ru}(\beta^-)^{110}\text{Rh}$	$E_{\beta^-}=2700(50)$ to $1^+$ level at 112.19 keV										Ens126 **
* $^{110}\text{Rh}(\beta^-)^{110}\text{Pd}$	$E_{\beta^-}=2600(100)$ to $(4^+)$ levels at 2790.64 and 2805.03 keV										Ens126 **
* $^{110}\text{Ag}(\beta^-)^{110}\text{Cd}$	$E_{\beta^-}=529(3)$ from $^{110}\text{Ag}^n$ at 117.59 to $6^+$ level at 2479.9339 keV										Ens126 **
* $^{110}\text{Ag}(\beta^-)^{110}\text{Cd}$	$E_{\beta^-}=2891(4)$ ; and 531(2) from $^{110}\text{Ag}^n$ at 117.59 to $6^+$ level at 2479.9339 keV										Ens126 **
* $^{110}\text{In}(\beta^+)^{110}\text{Cd}$	$E_{\beta^+}=2310(20)$ 2250(20) 2220(20) respectively, from $^{110}\text{In}^m$ at 62.08(0.04) keV										Nub16b **
*	to $2^+$ level at 657.7623 keV										Ens126 **
* $^{110}\text{Sb}(\beta^+)^{110}\text{Sn}$	$E_{\beta^+}=6500(200)$ 6850(100) respectively, to $2^+$ level at 1211.72; and other $E_{\beta^+}$										Ens126 **
$^{111}\text{Mo}-u$	-64348	279	-64348	14	0.0	U			GT1	1.5	04Ma.A
$^{111}\text{Ru}-u$	-82302	79	-82432	10	-1.7	U			JY0	1.0	03Ko.A
$^{111}\text{Rh}-u$	-88282	79	-88357	7	-1.0	U			JY0	1.0	03Ko.A
$^{111}\text{Rh}-^{120}\text{Sn}_{925}$	2105.8	7.3				2			JY1	1.0	07Ha20
$^{111}\text{Ag}-u$	-94741	51	-94703.2	1.6	0.7	U			GS2	1.0	05Li24 *
$\text{C}_8 \text{H}_{15}-^{111}\text{Cd}$	213184.4	3.9	213191.7	0.4	0.8	U			M16	2.5	63Da10
	213197	40			-0.1	U			R12	1.5	83De51
$\text{C}_7 \text{ } ^{13}\text{C} \text{H}_{14}-^{111}\text{Cd}$	208719	19	208721.5	0.4	0.1	U			R12	1.5	83De51
$\text{C}_7 \text{ O} \text{H}_{11}-^{111}\text{Cd}$	176814	16	176806.2	0.4	-0.3	U			R12	1.5	83De51
$\text{C}_9 \text{ H}_3-^{111}\text{Cd}$	119317	18	119291.3	0.4	-1.0	U			R12	1.5	83De51
$\text{C}_8 \text{ N} \text{H}-^{111}\text{Cd}$	106723	17	106715.3	0.4	-0.3	U			R12	1.5	83De51
$^{111}\text{Cd}-u$	-95774	30	-95816.2	0.4	-1.4	U			GS2	1.0	05Li24 *
$^{111}\text{Sb}-u$	-86837	30	-86782	10	1.8	U			GS2	1.0	05Li24
$^{111}\text{Sb}-^{133}\text{Cs}_{835}$	-7834.2	9.5				2			SH1	1.0	07Ma92
$^{111}\text{Te}-^{133}\text{Cs}_{835}$	-51.8	6.9				2			SH1	1.0	07Ma92
$^{111}\text{I}-^{87}\text{Rb}_{1,276}$	46150.4	6.1	46155	5	0.7	1	70	70 $^{111}\text{I}$	JY1	1.0	09El07
$^{111}\text{I}-^{133}\text{Cs}_{835}$	9197	19	9217	5	1.0	U			SH1	1.0	07Ma92
$^{111}\text{Tc}-^{105}\text{Ru}_{1,057}$	23412	11				2			JY1	1.0	07Ha20
$^{111}\text{Ru}-^{105}\text{Ru}_{1,057}$	15080.6	10.0				2			JY1	1.0	07Ha20
$^{110}\text{Cd} \text{H}-^{111}\text{Cd}$	6638	18	6648.73	0.18	0.4	U			R12	1.5	83De51
$^{111}\text{Mo}-^{111}\text{Tc}$	9753.0	7.3				3			JY1	1.0	11Ha48 *
$^{111}\text{Cd}-^{110}\text{Cd}$	1180	11	1176.31	0.18	-0.1	U			M16	2.5	63Da10
	1208	34			-0.6	U			R12	1.5	83De51

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{111}\text{Cd H-}^{110}\text{Cd}$	8994	35	9001.34	0.18	0.1	U			R12	1.5	83De51
$^{111}\text{I}(\alpha)^{107}\text{Sb}$	3270.1	10.	3275	5	0.4	-					79Sc22
	3293.0	10.			-1.8	-					92He.A
ave.	3281	7			-0.9	1	50	30 $^{111}\text{I}$			average
$^{111}\text{Xe}(\alpha)^{107}\text{Te}$	3693.3	25.	3720	50	0.5	4					79Sc22
	3714.1	30.			0.1	4					81Sc17
	3723.5	10.			-0.1	4					91He21
$^{110}\text{Pd}(\text{n},\gamma)^{111}\text{Pd}$	5726.3	0.4				2			Bdn		06Fi.A
$^{110}\text{Cd}(\text{n},\gamma)^{111}\text{Cd}$	6980	100	6975.60	0.17	0.0	U					61Ja21
	6975.5	0.5			0.2	-					86Ba72
	6975.9	0.2			-1.5	-					90Ne.B
	6975.1	0.4			1.2	-			Bdn		06Fi.A
$^{111}\text{Cd}(\gamma,\text{n})^{110}\text{Cd}$	-6975	3	-6975.60	0.17	-0.2	U			McM		79Ba06
$^{110}\text{Cd}(\text{d},\text{p})^{111}\text{Cd}$	4740	30	4751.03	0.17	0.4	U			Pit		64Ro17
	4750.68	0.88			0.4	U			Rez		90Pi05 *
$^{111}\text{Cd}(\text{d},\text{t})^{110}\text{Cd}$	-745	30	-718.37	0.17	0.9	U			Pit		64Co11
ave.	6975.71	0.17	6975.60	0.17	-0.7	1	97	77 $^{110}\text{Cd}$			average
$^{110}\text{Cd}(\text{n},\gamma)^{111}\text{Cd}$	5070	70	4966	15	-1.5	U					68Ba53
$^{111}\text{Te}(\epsilon\text{p})^{110}\text{Sn}$	7480	80	7761	14	3.5	B			Jyv		96Kl.A
$^{111}\text{Tc}(\beta^-)^{111}\text{Ru}$	7449	80			3.9	B			Jyv		00Kr.A
$^{111}\text{Ru}(\beta^-)^{111}\text{Rh}$	5039	50	5519	12	9.6	C			Jyv		00Kr.A
$^{111}\text{Rh}(\beta^-)^{111}\text{Pd}$	3640	50	3681	7	0.8	U			Jyv		00Kr.A
	3650	33			1.0	U			Bwg		00Kr.A
$^{111}\text{Pd}(\beta^-)^{111}\text{Ag}$	2210	100	2229.6	1.6	0.2	U					52Mc34 *
	2190	50			0.8	U					57Kn.A *
	2160	100			0.7	U					60Pr07 *
$^{111}\text{Ag}(\beta^-)^{111}\text{Cd}$	1028	3	1036.8	1.4	2.9	B					67Le06
	1035	2			0.9	2					71Na02
	1038.6	2.			-0.9	2					77Re12
$^{111}\text{Cd}(\text{p},\text{n})^{111}\text{In}$	-1635	20	-1643	3	-0.4	U			Oak		74Ki02
$^{111}\text{Sn}(\beta^+)^{111}\text{In}$	2530	30	2453	6	-2.6	U					51Mc11
$^{111}\text{Sb}(\beta^+)^{111}\text{Sn}$	4470	50	5102	10	12.6	B					72Si28 *
* $^{111}\text{Ag-u}$	$M-A=-88221(44)$ keV for mixture gs+m at 59.82 keV										
* $^{111}\text{Cd-u}$	$M-A=-88817(28)$ keV for $^{111}\text{Cd}^m$ $11/2^-$ at 396.214 keV										
* $^{111}\text{Mo}-^{111}\text{Tc}$	Taken as low-spin isomer (see also $^{102}\text{Y}$ and $^{114}\text{Tc}$ doublets in same work)										
* $^{110}\text{Cd}(\text{d},\text{p})^{111}\text{Cd}$	Estimated systematic error 0.5 added to statistical error 0.73 keV										
* $^{111}\text{Pd}(\beta^-)^{111}\text{Ag}$	$Q_{\beta^-}=2150(100)$ 2130(50) 2100(100) respectively, to $^{111}\text{Ag}^m$ at 59.82 keV										
* $^{111}\text{Sb}(\beta^+)^{111}\text{Sn}$	$E_{\beta^+}=3290(50)$ to $5/2^+$ level at 154.48 keV										
$^{112}\text{Tc}-^{102}\text{Ru}_{1,098}$	34976.0	5.9				2			JY1	1.0	07Ha20
$^{112}\text{Ru-u}$	-81033	78	-81193	10	-2.1	U			JY0	1.0	03Ko.A
$^{112}\text{Rh-u}$	-85506	119	-85600	50	-0.8	1	16	16 $^{112}\text{Rh}$	JY0	1.0	03Ko.A *
$^{112}\text{Ag}-^{133}\text{Cs}_{,842}$	-13342.0	2.6				2			MA8	1.0	10Br02
$\text{C}_8 \text{H}_{16}-^{112}\text{Cd}$	222445.3	3.9	222436.63	0.27	-0.9	U			M16	2.5	63Da10
$\text{C}_7 \text{O H}_{12}-^{112}\text{Cd}$	186063	16	186051.12	0.27	-0.5	U			R12	1.5	83De51
$\text{C}_9 \text{H}_4-^{112}\text{Cd}$	128541	19	128536.25	0.27	-0.2	U			R12	1.5	83De51
	128550	10			-0.9	U			R12	1.5	83De51
$\text{C}_8 \text{N H}_2-^{112}\text{Cd}$	115979	14	115960.19	0.27	-0.9	U			R12	1.5	83De51
$^{112}\text{In-u}$	-94366	58	-94461	5	-1.6	U			GS2	1.0	05Li24 *
$\text{C}_8 \text{H}_{16}-^{112}\text{Sn}$	220384	9	220375.6	0.3	-0.4	U			M16	2.5	63Da10
	220385	8			-0.8	U			R13	1.5	83De51
$^{112}\text{Sn}-^{86}\text{Kr}_{1,302}$	21210.3	2.5	21209.8	0.3	-0.2	U			JY1	1.0	11Ha48
$^{112}\text{Sb-u}$	-87597	30	-87600	19	-0.1	2			GS2	1.0	05Li24
$^{112}\text{Te}-^{133}\text{Cs}_{,842}$	-3662.7	9.0				2			SH1	1.0	07Ma92
$^{112}\text{I}-^{133}\text{Cs}_{,842}$	7614	11				2			SH1	1.0	07Ma92

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{112}\text{Rh}-^{120}\text{Sn}_{933}$	5640	110	5650	50	0.1	1	19	19 $^{112}\text{Rh}$	JY1	1.0	07Ha20 *
$^{112}\text{Pd}-^{120}\text{Sn}_{933}$	-1423.7	7.4	-1424	7	-0.1	1	89	89 $^{112}\text{Pd}$	JY1	1.0	07Ha20
$^{112}\text{Sn}-^{120}\text{Sn}_{933}$	-3930.2	1.9	-3929.5	0.9	0.4	1	24	22 $^{120}\text{Sn}$	JY1	1.0	11Ha48
$^{112}\text{Ru}-^{105}\text{Ru}_{1,067}$	17242.5	9.9				2			JY1	1.0	07Ha20
$^{112}\text{Cd } ^{35}\text{Cl}-^{110}\text{Cd } ^{37}\text{Cl}$	2701	2	2706.5	0.3	0.7	U			H11	4.0	63Bi12
$^{111}\text{Cd H}-^{112}\text{Cd}$	9255	20	9244.9	0.3	-0.3	U			R12	1.5	83De51
$^{112}\text{Sn}-^{112}\text{Cd}$	2061.01	0.17	2060.99	0.17	-0.1	1	99	97 $^{112}\text{Sn}$	JY1	1.0	09Ra11
$^{112}\text{Cd}-^{110}\text{Cd H}$	-8060	40	-8068.6	0.3	-0.1	U			R12	1.5	83De51
$^{112}\text{Cd}-^{111}\text{Cd}$	-1419	11	-1419.9	0.3	0.0	U			M16	2.5	63Da10
	-1410	42			-0.2	U			R12	1.5	83De51
$^{112}\text{Cd}-^{110}\text{Cd}$	-238	39	-243.6	0.3	-0.1	U			R12	1.5	83De51
$^{112}\text{Cd H}-^{111}\text{Cd}$	6402	35	6405.2	0.3	0.1	U			R12	1.5	83De51
$^{112}\text{I}(\alpha)^{108}\text{Sb}$	2986.9	31.1	2957	12	-0.6	U					81Sc17
$^{112}\text{Xe}(\alpha)^{108}\text{Te}$	3329.1	20.	3330	6	0.1	2					81Sc17
	3308.4	15.6			1.4	2					92He.A
	3335.4	7.			-0.7	2					94Pa11
$^{112}\text{Sn}(^3\text{He},^6\text{He})^{109}\text{Sn}$	-8686	9	-8686	8	0.0	1	78	78 $^{109}\text{Sn}$	MSU		78Pa11
$^{110}\text{Pd}(\text{t,p})^{112}\text{Pd}$	5659	20	5651	7	-0.4	1	11	11 $^{112}\text{Pd}$	LAI		72Ca10
$^{112}\text{Cd}(^{14}\text{C},^{16}\text{O})^{110}\text{Pd}$	5543	29	5512.9	0.6	-1.0	U			LAI		84Co19
$^{110}\text{Cd}(\text{t,p})^{112}\text{Cd}$	7888	20	7887.7	0.3	0.0	U			Ald		67Hi01
$^{112}\text{Cd}(\text{p,t})^{110}\text{Cd}$	-7891	5	-7887.7	0.3	0.7	U			Min		73Oo01
$^{112}\text{Sn}(\text{p,t})^{110}\text{Sn}$	-10485	15	-10474	14	0.7	R			Roc		70FI08
$^{111}\text{Cd}(\text{n},\gamma)^{112}\text{Cd}$	9460	50	9393.93	0.28	-1.3	U					61Ja21
	9394.3	0.3			-1.2	1	89	81 $^{111}\text{Cd}$	ILn		93Dr.A
$^{112}\text{Cd}(\gamma,\text{n})^{111}\text{Cd}$	-9403	5	-9393.93	0.28	1.8	U			McM		79Ba06
$^{111}\text{Cd}(\text{d,p})^{112}\text{Cd}$	7183	30	7169.36	0.28	-0.5	U			Pit		64Co11
	7170	10			-0.1	U			Yal		67Ba15
	7171	5			-0.3	U			MIT		67Sp09
$^{112}\text{Cd}(\text{d,t})^{111}\text{Cd}$	-3129	30	-3136.70	0.28	-0.3	U			Pit		64Ro17
$^{112}\text{Sn}(\text{d},^3\text{He})^{111}\text{In}$	-2050	50	-2059	3	-0.2	U			Sac		69Co03
$^{112}\text{Sn}(\text{p,d})^{111}\text{Sn}$	-8574	15	-8563	5	0.7	2			Har		70Ca01
$^{112}\text{Sn}(\text{d,t})^{111}\text{Sn}$	-4529.0	5.7	-4531	5	-0.3	2			SPa		75Be09
$^{112}\text{Cs}(\text{p})^{111}\text{Xe}$	814.3	7.	816	4	0.3	5					94Pa12
	817.3	5.			-0.2	5					12Wa10
$^{112}\text{Tc}(\beta^-)^{112}\text{Ru}$	6060	130	10372	11	33.2	C			Jyv		89Jo.A
	9484	100			8.9	C			Jyv		00Kr.A
$^{112}\text{Ru}(\beta^-)^{112}\text{Rh}$	4520	80	4100	50	-5.2	B			Jyv		91Jo11 *
$^{112}\text{Rh}(\beta^-)^{112}\text{Pd}$	6200	500	6590	40	0.8	U			Jyv		88Ay02
	6573	54			0.3	1	66	66 $^{112}\text{Rh}$	Bwg		00Kr.A
$^{112}\text{Rh}^m(\beta^-)^{112}\text{Pd}$	6929	56				2			Bwg		00Kr.A
$^{112}\text{Pd}(\beta^-)^{112}\text{Ag}$	299	20	262	7	-1.8	U					55Nu11 *
$^{112}\text{Ag}(\beta^-)^{112}\text{Cd}$	4057	20	3991.1	2.4	-3.3	C					57Je.A *
	3940	40			1.3	U					62In01 *
$^{112}\text{In}(\beta^+)^{112}\text{Cd}$	2582	20	2585	4	0.1	U					62Ru05
$^{112}\text{Cd}(\text{p,n})^{112}\text{In}$	-3399.3	20.	-3367	4	1.6	U			Oak		64Jo11
	-3376	6			1.5	1	50	50 $^{112}\text{In}$	Tky		80Ad04
$^{112}\text{In}(\beta^-)^{112}\text{Sn}$	656	6	665	4	1.5	1	50	50 $^{112}\text{In}$			53B144
$^{112}\text{Sb}(\beta^+)^{112}\text{Sn}$	7530	100	7056	18	-4.7	B					72Mi27 *
	7029	50			0.5	R					72Si28 *
	7062	26			-0.2	R					82Jo03 *
$^{112}\text{Sn}(\text{p,n})^{112}\text{Sb}$	-7995	55	-7838	18	2.8	U			VUn		76Ka19
* $^{112}\text{Rh}-\text{u}$	Average of 2 values; $M-A=-79479(36)$ keV for mixture gs+m at 340(70) keV										Nub16b **
* $^{112}\text{In}-\text{u}$	$M-A=-87823(30)$ keV for mixture gs+m at 156.592 keV										Nub16b **
* $^{112}\text{Rh}-^{120}\text{Sn}_{933}$	$D_M=5822.6(7.4)$ $\mu\text{u}$ for mixture gs+m at 340(70) keV; $M-A=-79571.2(7.0)$ keV										Nub16b **
* $^{112}\text{Ru}(\beta^-)^{112}\text{Rh}$	$E_{\beta^-}=4190(80)$ to $1^+$ level at 327.03 keV										Ens14c **
* $^{112}\text{Pd}(\beta^-)^{112}\text{Ag}$	$E_{\beta^-}=280(20)$ to $(1^+)$ level at 18.5 keV										Ens14c **
* $^{112}\text{Ag}(\beta^-)^{112}\text{Cd}$	$E_{\beta^-}=3440(20)$ to $2^+$ level at 617.518 keV										Ens14c **
* $^{112}\text{Ag}(\beta^-)^{112}\text{Cd}$	$E_{\beta^-}=3350(20)$ to $2^+$ level at 617.518 keV; error increased by evaluator										Ens14c **
* $^{112}\text{Sb}(\beta^+)^{112}\text{Sn}$	$E_{\beta^+}=5200(100)$ 4750(50) 4783(26) respectively, to $2^+$ level at 1256.69 keV										Ens14c **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{113}\text{Mo}-u$	-57317	337	-56350#	320#	1.1	D			GT3	2.5	16Kn03 *
$^{113}\text{Tc}-u$	-67633	268	-67431	4	0.5	o			GT1	1.5	04Ma.A
	-67502	106			0.3	U			GT2	2.5	08Kn.A
$^{113}\text{Tc}-^{129}\text{Xe}_{.876}$	15981.0	3.6				2			JY1	1.0	11Ha48
$^{113}\text{Ru}-u$	-77031	93	-77150	40	-1.3	-			JY0	1.0	03Ko.A *
	-77240	110			0.3	o			GT2	2.5	08Kn.A
	-77295	140			0.4	-			GT2	2.5	08Su19
	ave.	-77050	90		-1.2	1	19	19 $^{113}\text{Ru}$			average
$^{113}\text{Rh}-u$	-84464	83	-84560	8	-1.2	U			JY0	1.0	03Ko.A
$\text{C}_9 \text{H}_5-^{113}\text{Cd}$	134721.1	3.9	134717.06	0.26	-0.4	U			M16	2.5	63Da10
	134727	19			-0.3	U			R12	1.5	83De51
	134728	5			-1.5	U			R12	1.5	83De51
$\text{C}_7 \text{O H}_{13}-^{113}\text{Cd}$	192250	16	192231.94	0.26	-0.8	U			R12	1.5	83De51
$\text{C}_6 \text{ }^{13}\text{C O H}_{12}-^{113}\text{Cd}$	187772	17	187761.74	0.26	-0.4	U			R12	1.5	83De51
$\text{C}_8 \text{N H}_3-^{113}\text{Cd}$	122161	19	122141.00	0.26	-0.7	U			R12	1.5	83De51
$^{113}\text{Cd}-u$	-95506	93	-95591.90	0.26	-0.9	U			GS2	1.0	05Li24 *
$\text{C}_9 \text{H}_5-^{113}\text{In}$	135015	9	135064.71	0.20	2.2	U			M16	2.5	63Da10
	135087	6			-2.5	U			R12	1.5	83De51
$\text{C}_8 \text{N H}_3-^{113}\text{In}$	122506	14	122488.65	0.20	-0.8	U			R12	1.5	83De51
$^{113}\text{In}-u$	-95969	126	-95939.55	0.20	0.2	U			GS2	1.0	05Li24 *
$^{113}\text{Sn}-u$	-94796	39	-94824.2	1.7	-0.7	U			GS2	1.0	05Li24 *
$^{113}\text{Sb}-u$	-90635	30	-90625	18	0.3	R			GS2	1.0	05Li24
$^{113}\text{Te}-u$	-84109	30				2			GS2	1.0	05Li24
$^{113}\text{I}-^{133}\text{Cs}_{.850}$	4015.9	8.6				2			SH1	1.0	07Ma92
$^{113}\text{Xe}-^{133}\text{Cs}_{.850}$	13585.5	8.1	13588	7	0.2	1	82	82 $^{113}\text{Xe}$	SH1	1.0	07Ma92
$^{113}\text{Ru}-^{105}\text{Ru}_{1.076}$	22087	44	22110	40	0.6	1	81	81 $^{113}\text{Ru}$	JY1	1.0	07Ha20 *
$^{113}\text{Rh}-^{120}\text{Sn}_{.942}$	7565.4	7.6				2			JY1	1.0	07Ha20
$^{113}\text{Pd}-^{120}\text{Sn}_{.942}$	2387.1	7.4				2			JY1	1.0	07Ha20
$^{113}\text{Cd }^{35}\text{Cl}-^{111}\text{Cd }^{37}\text{Cl}$	3174	2	3174.4	0.4	0.1	U			H11	4.0	63Bi12
$^{112}\text{Cd H}-^{113}\text{Cd}$	6164	20	6180.82	0.23	0.6	U			R12	1.5	83De51
$^{113}\text{Cd}-^{112}\text{Cd}_{1.009}$	2519.36	0.29	2519.33	0.23	-0.1	1	65	35 $^{112}\text{Cd}$	MS1	1.0	16Ga24
$^{113}\text{In}-^{112}\text{Cd}_{1.009}$	2171.26	0.32	2171.68	0.26	1.3	1	65	48 $^{112}\text{Cd}$	MS1	1.0	16Ga24
$^{113}\text{Cd}-^{111}\text{Cd H}$	-7623	42	-7600.7	0.4	0.4	U			R12	1.5	83De51
$^{113}\text{Cd}-^{112}\text{Cd}$	1642	11	1644.21	0.23	0.1	U			M16	2.5	63Da10
	1620	40			0.4	U			R12	1.5	83De51
$^{113}\text{In}-^{112}\text{Cd}$	1297	45	1296.56	0.26	0.0	U			R12	1.5	83De51
$^{113}\text{Cd}-^{110}\text{Cd H}$	-6412	32	-6424.4	0.4	-0.3	U			R12	1.5	83De51
$^{113}\text{Cd}-^{111}\text{Cd}$	242	35	224.3	0.4	-0.3	U			R12	1.5	83De51
$^{113}\text{Cd H}-^{112}\text{Cd}$	9467	35	9469.25	0.23	0.0	U			R12	1.5	83De51
$^{113}\text{I}(\alpha)^{109}\text{Sb}$	2706.0	41.5	2707	10	0.0	U					81Sc17
$^{113}\text{Xe}(\alpha)^{109}\text{Te}$	3094.8	15.	3087	8	-0.5	1	24	18 $^{113}\text{Xe}$			79Sc22
$^{111}\text{Cd}(t,p)^{113}\text{Cd}$	7456	20	7451.9	0.3	-0.2	U			Ald		67Hi01
$^{113}\text{Cd}(p,t)^{111}\text{Cd}$	-7456	5	-7451.9	0.3	0.8	U			Min		73Oo01
$^{113}\text{In}(p,t)^{111}\text{In}-^{115}\text{In}^{113}\text{In}$	-810	10	-806	3	0.4	1	12	12 $^{111}\text{In}$	Roc		74Ma09
$^{113}\text{In}(p,t)^{111}\text{In}-^{112}\text{Cd}^{110}\text{Cd}$	-746.3	4.1	-748	3	-0.5	1	69	69 $^{111}\text{In}$	SPa		80Ta07
$^{112}\text{Cd}(n,\gamma)^{113}\text{Cd}$	6550	100	6539.74	0.22	-0.1	U					61Ja21
	6542.0	0.2			-11.3	C					90Ne.A
$^{112}\text{Cd}(d,p)^{113}\text{Cd}$	4318	30	4315.18	0.22	-0.1	U			Pit		64Ro17
	4315.56	0.64			-0.6	1	11	6 $^{112}\text{Cd}$	Rez		90Pi05 *
$^{113}\text{Cd}(d,t)^{112}\text{Cd}$	-254	30	-282.51	0.22	-1.0	U			Pit		64Co11
$^{113}\text{In}(d,t)^{112}\text{In}$	-3180	25	-3191	4	-0.4	U			Pit		67Hj03
$^{112}\text{Sn}(n,\gamma)^{113}\text{Sn}$	7741.9	2.3	7744.4	1.6	1.1	-			ORn		75Sl.A
$^{112}\text{Sn}(d,p)^{113}\text{Sn}$	5504	25	5519.8	1.6	0.6	U			Pit		64Co11
	5518.2	3.2			0.5	-			SPa		75Be09
$^{112}\text{Sn}(n,\gamma)^{113}\text{Sn}$	ave.	7742.2	7744.4	1.6	1.2	1	70	69 $^{113}\text{Sn}$			average
$^{112}\text{Sn}(^3\text{He},d)^{113}\text{Sb}$	-2400	40	-2443	17	-1.1	R			Sac		68Co22

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{113}\text{Xe}(\text{ep})^{112}\text{Te}$	7920	150	8075	11	1.0	o					82Pl05
	8300	150			-1.5	U					05Ja10
$^{113}\text{Cs}(\text{p})^{112}\text{Xe}$	967	4	972.8	2.2	1.5	o					84Fa04 *
	982.7	4.			-2.5	3					92He.A
	967.6	6.			0.9	3					94Pa12
	968.6	3.			1.4	3					95Ho26
$^{113}\text{Ru}(\beta^-)^{113}\text{Rh}$	6480	50	6900	40	8.4	C			Jyv		00Kr.A
$^{113}\text{Rh}(\beta^-)^{113}\text{Pd}$	5008	50	4824	10	-3.7	C			Jyv		00Kr.A
$^{113}\text{Pd}(\beta^-)^{113}\text{Ag}$	3360	150	3436	18	0.5	U					75Br.A
	3340	35			2.7	B			Stu		90Fo07
$^{113}\text{Ag}(\beta^-)^{113}\text{Cd}$	2010	20	2016	17	0.3	2					57Je.A
	2070	150			-0.4	U					70Ma47 *
	2031	30			-0.5	2			Stu		90Fo07 *
$^{113}\text{Cd}(\beta^-)^{113}\text{In}$	326.4	15.	323.83	0.27	-0.2	U					51Ca43 *
	316.4	30.			0.2	U					54De13 *
	320	10			0.4	U			CIT		88Mi13
	344.9	21.0			-1.0	U					07Be61
	322.2	0.9			1.8	U					09Da03
$^{113}\text{Sn}(\beta^+)^{113}\text{In}$	1034.6	5.0	1039.0	1.6	0.9	-					93Li10 *
$^{113}\text{In}(\text{p,n})^{113}\text{Sn}$	-1809	6	-1821.3	1.6	-2.1	-			Oak		73Ra13
$^{113}\text{Sn}(\beta^+)^{113}\text{In}$	ave.	1031	1039.0	1.6	2.0	1	17	17 $^{113}\text{Sn}$			average
$^{113}\text{Sb}(\beta^+)^{113}\text{Sn}$	3934	30	3911	17	-0.8	2					61Se08 *
	3945	50			-0.7	2					69Ki16 *
$^{113}\text{Te}(\beta^+)^{113}\text{Sb}$	5520	300	6070	30	1.8	U					74Bu21
	5720	200			1.7	U					74Ch17
* $^{113}\text{Mo}-\text{u}$	Trends from Mass Surface TMS suggest $^{113}\text{Mo}$ 900 less bound										GAu **
* $^{113}\text{Ru}-\text{u}$	$M-A=-71689(77)$ keV for mixture gs+m at 130(18) keV										Nub16b **
* $^{113}\text{Ru}-\text{u}$	$M-A=-71882(93)$ keV for mixture gs+m at 130(18) keV										Nub16b **
* $^{113}\text{Ru}-\text{u}$	$M-A=-71931(120)$ keV for mixture gs+m at 130(18) keV										Nub16b **
* $^{113}\text{Cd}-\text{u}$	$M-A=-88832(41)$ keV for mixture gs+m at 263.54 keV										Nub16b **
* $^{113}\text{In}-\text{u}$	$M-A=-89199(30)$ keV for mixture gs+m at 391.699 keV										Nub16b **
* $^{113}\text{Sn}-\text{u}$	$M-A=-88263(29)$ keV for mixture gs+m at 77.389 keV										Nub16b **
* $^{113}\text{Ru}-^{105}\text{Ru}_{1,076}$	$D_M=22157(12)$ $\mu\text{u}$ for mixture gs+m at 130(18) keV; $M-A=-71822(12)$ keV										Nub16b **
* $^{112}\text{Cd}(\text{d,p})^{113}\text{Cd}$	Estimated systematic error 0.5 added to statistical error 0.40										AHW **
* $^{113}\text{Cs}(\text{p})^{112}\text{Xe}$	$E_p$ from another GSI work; superseded by 95Ho26										87Gi07 **
* $^{113}\text{Ag}(\beta^-)^{113}\text{Cd}$	$E_{\beta^-}=1530(150)$ from $^{113}\text{Ag}^m$ at 43.50 to $5/2^+$ level at 583.962 keV										Ens106 **
* $^{113}\text{Ag}(\beta^-)^{113}\text{Cd}$	$Q_{\beta^-}=2075(30)$ from $^{113}\text{Ag}^m$ at 43.50 keV										Nub16b **
* $^{113}\text{Cd}(\beta^-)^{113}\text{In}$	$Q_{\beta^-}=590(15)$ 580(30) respectively, from $^{113}\text{Cd}^m$ at 263.54 keV										Nub16b **
* $^{113}\text{Sn}(\beta^+)^{113}\text{In}$	$Q_{\beta^+}=642.9(5.0)$ to $1/2^-$ level at 391.699 keV										Ens106 **
* $^{113}\text{Sb}(\beta^+)^{113}\text{Sn}$	$E_{\beta^+}=2420(20)$ 2430(50) respectively, to $3/2^+$ level at 498.06 keV,										Ens106 **
*	plus 6% to $5/2^+$ at 409.83 keV										Ens106 **
$^{114}\text{Tc}-\text{u}$	-62459	365	-62910	470	-0.8	o			GT1	1.5	04Ma.A
	-62910	186				2			GT3	2.5	16Kn03
$^{114}\text{Ru}-^{105}\text{Ru}_{1,086}$	24805	13	24802	5	-0.2	U			JY1	1.0	07Ha20
$^{114}\text{Ru}-\text{u}$	-75642	236	-75386	4	0.7	U			GT1	1.5	04Ma.A
$^{114}\text{Rh}-\text{u}$	-81193	120	-81280	80	-0.7	1	41	41 $^{114}\text{Rh}$	JY0	1.0	03Ko.A *
$^{114}\text{Ag}-^{133}\text{Cs}_{,857}$	-10149.3	4.9				2			MA8	1.0	10Br02
$\text{C}_8 \text{H}_{18}-^{114}\text{Cd}$	237487.6	4.	237485.59	0.30	-0.2	U			M16	2.5	63Da10
$\text{C}_6 \text{O}_2 \text{H}_{10}-^{114}\text{Cd}$	164713	15	164714.57	0.30	0.1	U			R12	1.5	83De51
	164711	15			0.2	U			R12	1.5	83De51
$\text{C}_9 \text{H}_6-^{114}\text{Cd}$	143591	5	143585.20	0.30	-0.8	U			R12	1.5	83De51
	143586	8			-0.1	U			R12	1.5	83De51
$\text{C}_8 \text{ } ^{13}\text{C} \text{H}_5-^{114}\text{Cd}$	139117	17	139115.01	0.30	-0.1	U			R12	1.5	83De51
$\text{C}_8 \text{ N} \text{H}_4-^{114}\text{Cd}$	131017	12	131009.14	0.30	-0.4	U			R12	1.5	83De51
	131009	20			0.0	U			R12	1.5	83De51



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{114}\text{Cd}-^{133}\text{Cs}_{.857}$	-15611.4	4.2	-15607.34	0.30	1.0	U			MA8	1.0	10Br02
$^{114}\text{In}-u$	-94986	68	-95083.6	0.3	-1.4	U			GS2	1.0	05Li24 *
$\text{C}_8 \text{H}_{18}-^{114}\text{Sn}$	238092	10	238070.45	0.03	-0.9	U			M16	2.5	63Da10
	238066	8			0.4	U			R13	1.5	83De51
$^{114}\text{Sb}-u$	-90731	30	-90711	23	0.7	1	61	61 $^{114}\text{Sb}$	GS2	1.0	05Li24
$^{114}\text{Te}-u$	-87911	30				2			GS2	1.0	05Li24
$^{114}\text{Xe}-^{133}\text{Cs}_{.857}$	9008	12				2			MA6	1.0	04Di18
$^{114}\text{Ru}-^{120}\text{Sn}_{.950}$	17522.0	3.7				2			JY1	1.0	11Ha48
$^{114}\text{Rh}-^{120}\text{Sn}_{.950}$	11570	100	11630	80	0.6	1	59	59 $^{114}\text{Rh}$	JY1	1.0	07Ha20 *
$^{114}\text{Pd}-^{120}\text{Sn}_{.950}$	3277.0	7.4				2			JY1	1.0	07Ha20
$^{114}\text{Cd}-^{35}\text{Cl}-^{112}\text{Cd}-^{37}\text{Cl}$	3546	3	3551.22	0.28	0.4	U			H11	4.0	63Bi12
	3547	3			0.6	U			H20	2.5	66Ma05
	3548.5	1.0			1.1	U			H26	2.5	73Me28
$^{113}\text{Cd H}-^{114}\text{Cd}$	8859	18	8868.14	0.15	0.3	U			R12	1.5	83De51
$^{114}\text{Tc}^m-^{114}\text{Ru}$	12651	13				3			JY1	1.0	11Ha48 *
$^{114}\text{Cd}-^{112}\text{Cd H}$	-7225	33	-7223.93	0.27	0.0	U			R12	1.5	83De51
$^{114}\text{Cd}-^{113}\text{Cd}$	-1040	11	-1043.11	0.15	-0.1	U			M16	2.5	63Da10
	-1032	33			-0.2	U			R12	1.5	83De51
$^{114}\text{Cd}-^{113}\text{In}$	-679	45	-695.5	0.3	-0.2	U			R12	1.5	83De51
$^{114}\text{Cd}-^{111}\text{Cd H}$	-8651	35	-8643.8	0.4	0.1	U			R12	1.5	83De51
$^{114}\text{Cd}-^{112}\text{Cd}$	587	33	601.11	0.27	0.3	U			R12	1.5	83De51
$^{114}\text{Cd H}-^{113}\text{Cd}$	6821	35	6781.92	0.15	-0.7	U			R12	1.5	83De51
$^{114}\text{Ba}(\gamma,^{12}\text{C})^{102}\text{Sn}$	18110	780	19029	23	1.2	U					95Gu01
$^{114}\text{Cs}(\alpha)^{110}\text{I}$	3343.5	30.	3360	50	0.3	o			GSa		80Ro04
	3357.0	30.				4			GSa		81Sc17
$^{114}\text{Ba}(\alpha)^{110}\text{Xe}$	3534.2	40.	3592	19	1.4	5					02Ma19
	3606.8	20.			-0.7	5					16Ca33
$^{112}\text{Cd}(\text{t,p})^{114}\text{Cd}$	7105	20	7100.91	0.25	-0.2	U			Ald		67Hi01
$^{114}\text{Cd}(\text{p,t})^{112}\text{Cd}$	-7106	5	-7100.91	0.25	1.0	U			Min		73Oo01
$^{112}\text{Sn}(\text{t,p})^{114}\text{Sn}$	9579	15	9565.51	0.30	-0.9	U			Ald		69Bj01
$^{114}\text{Sn}(\text{p,t})^{112}\text{Sn}$	-9582	15	-9565.51	0.30	1.1	U			Roc		70Fl08
$^{113}\text{Cd}(\text{n},\gamma)^{114}\text{Cd}$	9042.76	0.20	9042.97	0.14	1.0	-			ILn		79Br25 Z
	9043.18	0.19			-1.1	-			Bdn		06Fi.A
$^{114}\text{Cd}(\gamma,\text{n})^{113}\text{Cd}$	-9050	4	-9042.97	0.14	1.8	U			McM		79Ba06
$^{113}\text{Cd}(\text{d,p})^{114}\text{Cd}$	6817	30	6818.40	0.14	0.0	U			Pit		64Co11
	6822	8			-0.5	U			MIT		67Co15
$^{114}\text{Cd}(\text{d,t})^{113}\text{Cd}$	-2801	30	-2785.74	0.14	0.5	U			Pit		64Ro17
$^{113}\text{Cd}(\text{n},\gamma)^{114}\text{Cd}$	ave. 9042.98	0.14	9042.97	0.14	-0.1	1	98	93 $^{114}\text{Cd}$			average
$^{113}\text{In}(\text{n},\gamma)^{114}\text{In}$	7274.0	1.2	7274.00	0.25	0.0	U					75Ra07 Z
	7273.83	0.27			0.6	1	88	82 $^{114}\text{In}$	Bdn		06Fi.A
$^{113}\text{In}(\text{d,p})^{114}\text{In}$	5082	25	5049.44	0.25	-1.3	U			Pit		67Hj03
$^{114}\text{Sn}(\text{d},^3\text{He})^{113}\text{In}$	-2980	50	-2988.10	0.19	-0.2	U			Sac		69Co03
$^{114}\text{Sn}(\text{p,d})^{113}\text{Sn}$	-8101	15	-8078.3	1.6	1.5	U			Har		70Ca01
$^{114}\text{Sn}(\text{d,t})^{113}\text{Sn}$	-4052	20	-4045.7	1.6	0.3	U			Pit		64Co11
	-4043.7	4.2			-0.5	1	14	14 $^{113}\text{Sn}$	SPa		75Be09
$^{114}\text{Cs}(\varepsilon\text{p})^{113}\text{I}$	8730	150	9150	70	2.8	B					82Pi05
$^{114}\text{Ru}(\beta^-)^{114}\text{Rh}$	6100	200	5490	70	-3.1	B			Jyv		92Jo05 *
	6120	200			-3.2	C			Jyv		94Jo.A
$^{114}\text{Rh}(\beta^-)^{114}\text{Pd}$	6500	500	7780	70	2.6	U			Jyv		88Ay02
	7392	53			7.3	C			Jyv		00Kr.A
$^{114}\text{Pd}(\beta^-)^{114}\text{Ag}$	1450	100	1440	8	-0.1	U					75Br.A
	1450	100			-0.1	o			Jyv		89Ay.A
	1450	100			-0.1	o			Jyv		89Ko22
	1414	30			0.9	U			Stu		90Fo07
	1451	25			-0.4	U			Jyv		94Jo.A
$^{114}\text{Ag}(\beta^-)^{114}\text{Cd}$	4850	150	5084	5	1.6	U					71Ro19
	4900	260			0.7	U					72Wa06
	5160	110			-0.7	o			Stu		84Lu02
	5018	35			1.9	U			Stu		90Fo07

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{114}\text{In}(\beta^+)^{114}\text{Cd}$	1422	25	1445.1	0.4	0.9	U					56Gr35	
	1417	20			1.4	U					57Dz64	
$^{114}\text{In}(\beta^-)^{114}\text{Sn}$	1987	2	1989.9	0.3	1.5	U					61Da01	
	1989	1			0.9	-					61Ni02	
	1980	2			5.0	B					64An12	
	1988.5	1.0			1.4	-					68Ze04	
ave.	1988.8	0.7			1.7	1	18	18	$^{114}\text{In}$		average	
$^{114}\text{Sb}(\beta^+)^{114}\text{Sn}$	5690	100	6063	22	3.7	C					69Bu.A *	
	6370	100			-3.1	B					72Mi27 *	
$^{114}\text{Sn}(p,n)^{114}\text{Sb}$	-6875	35	-6845	22	0.8	1	39	39	$^{114}\text{Sb}$	VUn	76Ka19	
* $^{114}\text{Rh}-u$	Average of 2 values; $M - A = -75531(60)$ keV for mixture gs+m at 200#150 keV										Nub16b **	
* $^{114}\text{In}-u$	$M - A = -88384(31)$ keV for mixture gs+m at 190.2682 keV										Nub16b **	
* $^{114}\text{Rh}-^{120}\text{Sn}_{950}$	$D_M = 11678.0(7.8)$ $\mu\text{u}$ for mixture gs+m at 200#150 keV; $M - A = -75665.6(7.3)$ keV										Nub16b **	
* $^{114}\text{Tc}^m - ^{114}\text{Ru}$	Mixture of two isomeric states with stronger component of low-spin state										11Ri01 **	
*	however, estimates from TMS suggest this is $^{114}\text{Tc}^m$										GAu **	
* $^{114}\text{Ru}(\beta^-)^{114}\text{Rh}$	$E_{\beta^-} = 5910(120)$ doublet to $(2)^+$ level at 127.0, $1^+$ at 255.2 keV										Ens123 **	
* $^{114}\text{Sb}(\beta^+)^{114}\text{Sn}$	$E_{\beta^+} = 3365(50)$ to $2^+$ at 1299.907, original error doubled see $^{114}\text{Sn}(p,n)$										Ens123 **	
* $^{114}\text{Sb}(\beta^+)^{114}\text{Sn}$	$E_{\beta^+} = 4050(100)$ to $2^+$ at 1299.907 level, see $^{112}\text{Sb}(\beta^+)$										Ens123 **	
$^{115}\text{Tc}-u$	-60462	339				2			GT3	2.5	16Kn03	
$^{115}\text{Rh}-u$	-79664	85	-79689	8	-0.3	U			JY0	1.0	03Ko.A	
$^{115}\text{Ag}-^{133}\text{Cs}_{865}$	-9439	24	-9449	20	-0.4	1	67	67	$^{115}\text{Ag}$	MA8	1.0	10Br02 *
$\text{C}_9 \text{H}_7 - ^{115}\text{In}$	150910	8	150896.452	0.012	-0.7	U			M16	2.5	63Da10	
	150932	16			-1.5	U			R12	1.5	83De51	
$\text{C}_6 \text{O}_2 \text{H}_{11} - ^{115}\text{In}$	172055	16	172025.820	0.012	-1.2	U			R12	1.5	83De51	
$\text{C}_8 \text{N H}_5 - ^{115}\text{In}$	138355	13	138320.392	0.012	-1.8	U			R12	1.5	83De51	
$^{115}\text{In}-u$	-96095	30	-96121.226	0.013	-0.9	U			GS2	1.0	05Li24	
$\text{C}_9 \text{H}_7 - ^{115}\text{Sn}$	151411	8	151430.528	0.016	1.0	U			M16	2.5	63Da10	
	151440	8			-0.8	U			R13	1.5	83De51	
$^{115}\text{Sb}-u$	-93402	30	-93402	17	0.0	2			GS2	1.0	05Li24	
$^{115}\text{Te}-u$	-88098	30				2			GS2	1.0	05Li24 *	
$^{115}\text{Tl}-u$	-81952	31				2			GS2	1.0	05Li24	
$^{115}\text{Xe}-^{133}\text{Cs}_{865}$	8078	13				2			MA6	1.0	04Di18	
$^{115}\text{Ru}-^{120}\text{Sn}_{958}$	22633	95				2			JY1	1.0	07Ha20 *	
$^{115}\text{Rh}-^{120}\text{Sn}_{958}$	14001.6	7.8				2			JY1	1.0	07Ha20	
$^{115}\text{Pd}-^{120}\text{Sn}_{958}$	7347	15	7349	15	0.2	1	94	94	$^{115}\text{Pd}$	JY1	1.0	07Ha20 *
$^{115}\text{Sn}-^{120}\text{Sn}_{958}$	-2963.9	2.0	-2964.7	0.9	-0.4	1	21	21	$^{120}\text{Sn}$	JY1	1.0	11Ha48
$^{113}\text{Cd}-^{115}\text{In}_{983}$	-1104.76	0.34	-1104.74	0.26	0.1	1	59	59	$^{113}\text{Cd}$	MS1	1.0	16Ga24
$^{113}\text{In}-^{115}\text{In}_{983}$	-1452.08	0.23	-1452.39	0.20	-1.3	1	77	77	$^{113}\text{In}$	MS1	1.0	16Ga24
$^{115}\text{In}-^{115}\text{Sn}$	534.0768	0.0104	534.077	0.010	0.0	1	100	100	$^{115}\text{Sn}$	FS1	1.0	09Mo23
	534.28	0.18			-1.1	U			JY1	1.0	09Wi10	
$^{115}\text{In}-^{114}\text{Cd}$	483	45	513.78	0.30	0.5	U			R12	1.5	83De51	
$^{115}\text{Sn}-^{114}\text{Sn}$	573	11	564.565	0.027	-0.3	U			M16	2.5	63Da10	
$^{115}\text{In}-^{113}\text{In}$	-200	28	-181.67	0.20	0.4	U			R12	1.5	83De51	
$^{115}\text{In}-^{129}\text{Xe}$	-902.0845	0.0111	-902.085	0.011	0.0	1	100	100	$^{115}\text{In}$	FS1	1.0	09Mo23
$^{115}\text{Sn}-^{129}\text{Xe}$	-1436.1613	0.0130	-1436.162	0.015	0.0	o			FS1	1.0	09Mo23 *	
$^{113}\text{Cd}(t,p)^{115}\text{Cd}$	6702	20	6702.0	0.6	0.0	U			Ald		67Hi01	
$^{114}\text{Cd}(n,\gamma)^{115}\text{Cd}$	6160	100	6140.9	0.6	-0.2	U					61Ja21	
$^{114}\text{Cd}(d,p)^{115}\text{Cd}$	3923	30	3916.3	0.6	-0.2	U					64Ro17	
	3929	20			-0.6	U			Oak		64Si18	
	3916.30	0.59			0.0	1	100	100	$^{115}\text{Cd}$	Rez		90Pi05 *
$^{114}\text{Cd}(^3\text{He,d})^{115}\text{In}$	1320	15	1316.91	0.28	-0.2	U					70Th.A	
$^{115}\text{In}(\gamma,n)^{114}\text{In}$	-9025	29	-9037.9	0.3	-0.4	U					60Ge01	
	-9039	5			0.2	U			McM		79Ba06	
$^{115}\text{In}(d,t)^{114}\text{In}$	-2789	30	-2780.6	0.3	0.3	U					64Co11	
	-2766	25			-0.6	U					67Hj03	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{114}\text{Sn}(n,\gamma)^{115}\text{Sn}$	7545.5	2.0	7545.429	0.025	0.0	U			ORn		78Ra16 Z
	7545.427	0.025			0.1	1	100	100 $^{114}\text{Sn}$			16Ur03
$^{114}\text{Sn}(d,p)^{115}\text{Sn}$	5329	25	5320.863	0.025	-0.3	U			Pit		64Co11
	5320.6	3.4			0.1	U			SPa		75Be09
$^{115}\text{Sn}(d,t)^{114}\text{Sn}$	-1304	30	-1288.200	0.025	0.5	U			Pit		64Co11
$^{115}\text{Xe}(\text{ep})^{114}\text{Te}$	6200	130	5940	30	-2.0	U					72Ho18
$^{115}\text{Ru}(\beta^-)^{115}\text{Rh}$	7780	100	8040	90	2.6	C			Jyv		00Kr.A
$^{115}\text{Rh}(\beta^-)^{115}\text{Pd}$	6000	500	6197	15	0.4	U			Jyv		88Ay01
	6566	50			-7.4	C			Jyv		00Kr.A
$^{115}\text{Pd}(\beta^-)^{115}\text{Ag}$	4584	50	4556	22	-0.6	1	19	12 $^{115}\text{Ag}$	Stu		90Fo07
$^{115}\text{Ag}(\beta^-)^{115}\text{Cd}$	3180	100	3102	18	-0.8	U					64Ba36 *
	3118	100			-0.2	U					78Ma18 *
	3091	40			0.3	1	21	21 $^{115}\text{Ag}$			90Fo07 *
$^{115}\text{Cd}(\beta^-)^{115}\text{In}$	1460	4	1451.9	0.7	-2.0	U					74Bo26 *
	1431	5			4.2	B					75Bo29 *
	1440	2			5.9	B					76Ra16 *
	494	20	497.489	0.010	0.2	U					49Be53 *
$^{115}\text{In}(\beta^-)^{115}\text{Sn}$	630	30			-4.4	B					50Ma76
	625	70			-1.8	U					61Be15
	494	30			0.1	U					62Se03 *
	480	30			0.6	U					62Wa15
	495	20			0.1	U					72Mu02
	482	15			1.0	U					78Pf01 *
	3030	20	3030	16	0.0	R					61Se08 *
$^{115}\text{Sb}(\beta^+)^{115}\text{Sn}$											61Se08 *
$^{115}\text{Ag}-^{133}\text{Cs}_{.865}$	$D_M=-9416.7(9.2) \mu\text{u}$ for ground state or $^{115}\text{Ag}^m$ at 41.16 keV; $M-A=-84952.9(8.6) \text{ keV}$										Nub16b **
$^{115}\text{Te}-u$	$M-A=-82058(28) \text{ keV}$ for mixture gs+m at 10(7) keV										Nub16b **
$^{115}\text{Ru}-^{120}\text{Sn}_{.958}$	$D_M=22767.3(7.3) \mu\text{u}$ for mixture gs+m at 250#(100#); $M-A=-66064.8(6.9) \text{ keV}$										Nub16b **
$^{115}\text{Pd}-^{120}\text{Sn}_{.958}$	$D_M=7348(15), 7442(15) \mu\text{u}$ for ground state, 89.21 isomer										Nub16b **
$^{115}\text{Sn}-^{129}\text{Xe}$	Used are the equations for the $^{115}\text{In}-^{129}\text{Xe}$ and $^{115}\text{In}-^{115}\text{Sn}$ doublets										GAU **
$^{114}\text{Cd}(d,p)^{115}\text{Cd}$	Estimated systematic error 0.5 added to statistical error 0.32 keV										AHW **
$^{115}\text{Ag}(\beta^-)^{115}\text{Cd}$	$E_{\beta^-}=2950(100)$ to $(3/2)^+$ level at 229.1 keV, and other $E_{\beta^-}$										Ens12a **
$^{115}\text{Ag}(\beta^-)^{115}\text{Cd}$	$E_{\beta^-}=721(100)$ to $(23/2^-)$ level at 2397.2 keV, and other $E_{\beta^-}$										Ens12a **
$^{115}\text{Ag}(\beta^-)^{115}\text{Cd}$	$Q_{\beta^-}=3132(40)$ from $^{115}\text{Ag}^m$ at 41.16 keV										Nub16b **
$^{115}\text{Cd}(\beta^-)^{115}\text{In}$	$E_{\beta^-}=593(2), 636(2)$ to $1/2^+$ level at 864.139, $3/2^+$ at 828.588 keV										Ens12a **
$^{115}\text{Cd}(\beta^-)^{115}\text{In}$	$E_{\beta^-}=320(5), 679(6)$ from $^{115}\text{Cd}^m$ 181.0 to $13/2^+$ 1290.592, $7/2^+$ 933.780										Ens12a **
$^{115}\text{Cd}(\beta^-)^{115}\text{In}$	$Q_{\beta^-}=1621(2)$ from $^{115}\text{Cd}^m$ at 181.0 keV										Nub16b **
$^{115}\text{In}(\beta^-)^{115}\text{Sn}$	$Q_{\beta^-}=830(20)$ from $^{115}\text{In}^m$ at 336.244 keV										Nub16b **
$^{115}\text{In}(\beta^-)^{115}\text{Sn}$	$Q_{\beta^-}=830(30)$ from $^{115}\text{In}^m$ at 336.244 keV										Nub16b **
$^{115}\text{In}(\beta^-)^{115}\text{Sn}$	$Q_{\beta^-}$ is larger than first excitation energy 497.334(0.023) in $^{115}\text{Sn}$										05Ca03 **
$^{115}\text{Sb}(\beta^+)^{115}\text{Sn}$	$E_{\beta^+}=1510(20)$ to $3/2^+$ level at 497.334 keV										Ens12a **
$^{116}\text{Ru}-^{129}\text{Xe}_{.899}$	16821.2	4.0				2			JY1	1.0	11Ha48
	-75936	140	-75940	80	0.0	-			JY0	1.0	03Ko.A *
$^{116}\text{Rh}-u$	-75960	140			0.1	-			GT2	2.5	08Kn.A *
	ave. -75940	130			0.0	1	37	37 $^{116}\text{Rh}$			average
$^{116}\text{Ag}-^{133}\text{Cs}_{.872}$	-6167.3	3.5				2			MA8	1.0	10Br02
$\text{C}_9 \text{H}_8-^{116}\text{Cd}$	157837.4	2.9	157837.03	0.17	-0.1	U			M16	2.5	63Da10
	157851	5			-1.9	U			R12	1.5	83De51
	157846	22			-0.3	U			R12	1.5	83De51
$\text{C}_6 \text{O}_2 \text{H}_{12}-^{116}\text{Cd}$	178982	15	178966.40	0.17	-0.7	U			R12	1.5	83De51
	153376	8	153366.83	0.17	-0.8	U			R12	1.5	83De51
$\text{C}_8 \text{ }^{13}\text{C} \text{H}_7-^{116}\text{Cd}$	153382	22			-0.5	U			R12	1.5	83De51
	145262	17	145260.97	0.17	0.0	U			R12	1.5	83De51
$\text{C}_9 \text{H}_8-^{116}\text{Sn}$	160861	8	160857.43	0.10	-0.2	U			M16	2.5	63Da10
	160857	8			0.0	U			R13	1.5	83De51

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{116}\text{Sb}-u$	-93128	129	-93207	6	-0.6	U			GS2	1.0	05Li24 *
$^{116}\text{Te}-u$	-91540	30				2			GS2	1.0	05Li24
$^{116}\text{Xe}-^{133}\text{Cs}_{.872}$	4027	14				2			MA6	1.0	04Di18
$^{116}\text{Rh}-^{120}\text{Sn}_{.967}$	18633	100	18630	80	0.0	1	63	63 $^{116}\text{Rh}$	JY1	1.0	07Ha20 *
$^{116}\text{Pd}-^{120}\text{Sn}_{.967}$	8868.0	7.6				2			JY1	1.0	07Ha20
$^{116}\text{Cd } ^{35}\text{Cl}-^{114}\text{Cd } ^{37}\text{Cl}$	4353	3	4348.4	0.3	-0.4	U			H11	4.0	63Bi12
	4344	2			0.9	U			H20	2.5	66Ma05
	4348.7	1.2			-0.1	o			H26	2.5	73Me28
	4347.46	0.44			0.8	1	10	7 $^{114}\text{Cd}$	H49	2.5	10Mc04
$^{116}\text{Cd}-^{116}\text{Sn}$	3020.42	0.14	3020.41	0.14	-0.1	1	99	98 $^{116}\text{Cd}$	JY1	1.0	11Ra24
$^{116}\text{Cd}-^{114}\text{Cd } H$	-6452	32	-6426.8	0.3	0.5	U			R12	1.5	83De51
$^{116}\text{Sn}-^{115}\text{Sn}$	-1602	11	-1601.87	0.10	0.0	U			M16	2.5	63Da10
$^{116}\text{Cd}-^{113}\text{Cd } H$	-7458	32	-7469.9	0.3	-0.2	U			R12	1.5	83De51
$^{116}\text{Cd}-^{114}\text{Cd}$	1370	32	1398.2	0.3	0.6	U			R12	1.5	83De51
$^{116}\text{Cs}(\epsilon\alpha)^{112}\text{Te}$	12300	400	13100#	100#	2.0	D					77Bo28
	12400	900			0.8	D					76Jo.A *
$^{116}\text{Cd}(^{14}\text{C},^{16}\text{O})^{114}\text{Pd}$	2497	29	2535	7	1.3	U			LAI		84Co19
$^{114}\text{Cd}(t,p)^{116}\text{Cd}$	6362	20	6358.4	0.3	-0.2	U			Ald		67Hi01
$^{116}\text{Cd}(p,t)^{114}\text{Cd}$	-6363	5	-6358.4	0.3	0.9	U			Min		73Oo01
$^{116}\text{Sn}(p,t)^{114}\text{Sn}$	-8619	15	-8627.09	0.10	-0.5	U			Roc		70FI08
$^{116}\text{Cd}(\gamma,n)^{115}\text{Cd}$	-8702	4	-8699.3	0.7	0.7	U			McM		79Ba06
$^{116}\text{Cd}(d,t)^{115}\text{Cd}$	-2458	30	-2442.1	0.7	0.5	U			Pit		64Ro17
$^{115}\text{In}(n,\gamma)^{116}\text{In}$	6783.8	1.2	6784.72	0.22	0.8	U					72Ra39 Z
	6784.4	1.1			0.3	U					74Co35
	6784.72	0.22				2			Bdn		06Fi.A
$^{115}\text{In}(d,p)^{116}\text{In}$	4494	25	4560.15	0.22	2.6	U			Pit		64Co11
	4580	30			-0.7	U			Pit		67Hj03
$^{116}\text{Sn}(d,^3\text{He})^{115}\text{In}$	-3740	50	-3785.12	0.10	-0.9	U			Sac		69Co03
$^{115}\text{Sn}(n,\gamma)^{116}\text{Sn}$	9562.2	1.5	9563.45	0.09	0.8	U					72Mc08
	9563.5	0.5			-0.1	U					84Ga.B
	9563.41	0.11			0.4	-			ORn		91Ra01 Z
	9563.55	0.19			-0.5	-			Bdn		06Fi.A
$^{115}\text{Sn}(d,p)^{116}\text{Sn}$	7358	30	7338.89	0.09	-0.6	U			Pit		64Co11
$^{116}\text{Sn}(p,d)^{115}\text{Sn}$	-7344	15	-7338.89	0.09	0.3	U			Har		70Ca01
$^{116}\text{Sn}(d,t)^{115}\text{Sn}$	-3309	20	-3306.22	0.09	0.1	U			Pit		64Co11
	-3305.0	2.5			-0.5	U			SPa		75Be09
$^{115}\text{Sn}(n,\gamma)^{116}\text{Sn}$	ave. 9563.45	0.10	9563.45	0.09	0.1	1	99	99 $^{116}\text{Sn}$			average
$^{115}\text{Sn}(^3\text{He},d)^{116}\text{Sb}-^{120}\text{Sn}()^{121}\text{Sb}$	-1722	10	-1714	5	0.8	1	30	25 $^{116}\text{Sb}$	VUn		78Ka12
$^{116}\text{Cs}(\epsilon p)^{115}\text{I}$	6350	300	7010#	100#	2.2	U					78Da07 *
$^{116}\text{Rh}(\beta^-)^{116}\text{Pd}$	8000	500	9100	70	2.2	U			Jyv		88Ay02
$^{116}\text{Pd}(\beta^-)^{116}\text{Ag}$	2615	100	2711	8	1.0	U					75Br.A
	2620	100			0.9	o			Jyv		89Ay.A
	2607	30			3.5	B			Stu		90Fo07
	2620	100			0.9	U			Jyv		94Jo.A
	6062	130	6170	3	0.8	o			Stu		82A129 *
$^{116}\text{Ag}(\beta^-)^{116}\text{Cd}$	5800	200			1.8	U					82Br10
	6194	50			-0.5	U			Stu		90Fo07 *
	3290	60	3276.22	0.24	-0.2	U					54Bo39
$^{116}\text{Sb}(\beta^+)^{116}\text{Sn}$	4586	100	4704	5	1.2	U					61Fi05 *
$^{116}\text{Sn}(p,n)^{116}\text{Sb}$	4606	50			2.0	U					68Ki07 *
	-5515	40	-5486	5	0.7	U			VUn		76Ka19
$^{116}\text{Sb}(\beta^+)^{116}\text{Sn}$	-5483.2	6.			-0.5	1	75	75 $^{116}\text{Sb}$	Oak		77Jo03
	5090	40				2					60Je03 *
$^{116}\text{Te}(\beta^+)^{116}\text{Sb}$	1554	100	1553	28	0.0	U					61Fi05 *
$^{116}\text{I}(\beta^+)^{116}\text{Te}$	7760	130	7780	100	0.1	R					70Be.A
$^{116}\text{Xe}(\beta^+)^{116}\text{I}$	7710	200			0.3	R					76Go02
	4340	200	4450	100	0.5	3					76Go02
* $^{116}\text{Rh}-u$	$M - A = -70634(96)$ keV for mixture gs+m at 200#150 keV										Nub16b **
* $^{116}\text{Rh}-u$	$M - A = -70662(93)$ keV for mixture gs+m at 200#150 keV										Nub16b **
* $^{116}\text{Sb}-u$	$M - A = -86553(34)$ keV for mixture gs+n at 390(40) keV										Nub16b **
* $^{116}\text{Rh}-^{120}\text{Sn}_{.967}$	$D_M = 18740.7(8.4)$ $\mu\text{u}$ for mixture gs+m at 200#150 keV ; $M - A = -70635.5(7.9)$ keV										Nub16b **
* $^{116}\text{Cs}(\epsilon\alpha)^{112}\text{Te}$	$Q = 12500(900)$ from $^{116}\text{Cs}^m$ estimated at 100#60 keV										Nub16b **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>116</sup> Cs( $\epsilon\alpha$ ) <sup>112</sup> Te	Trends from Mass Surface TMS suggest <sup>116</sup> Cs 780 less bound										GAu **
* <sup>116</sup> Cs( $\epsilon p$ ) <sup>115</sup> I	$Q=6450(300)$ from <sup>116</sup> Cs <sup>m</sup> at estimated 100#60 keV										Nub16b **
* <sup>116</sup> Ag( $\beta^-$ ) <sup>116</sup> Cd	$Q_{\beta^-}=6110(130)$ from <sup>116</sup> Ag <sup>m</sup> at 47.90 keV										Nub16b **
* <sup>116</sup> Ag( $\beta^-$ ) <sup>116</sup> Cd	$Q_{\beta^-}=6199(100)$ ; and 6241(50) from <sup>116</sup> Ag <sup>m</sup> at 47.90 keV										Nub16b **
* <sup>116</sup> Sb( $\beta^+$ ) <sup>116</sup> Sn	$E_{\beta^+}=2270(100)$ 2290(50) respectively, to 2 <sup>+</sup> level at 1293.560 keV										Ens104 **
* <sup>116</sup> Sb <sup>n</sup> ( $\beta^+$ ) <sup>116</sup> Sn	$E_{\beta^+}=1160(40)$ to 7 <sup>-</sup> level at 2908.85 keV										Ens104 **
* <sup>116</sup> Te( $\beta^+$ ) <sup>116</sup> Sb	$E_{\beta^+}=440(100)$ to 1 <sup>+</sup> level at 93.99 keV										Ens104 **
<sup>117</sup> Ru-u	-63897	419	-63870	470	0.1	o			GT1	1.5	04Ma.A
	-63865	186				2			GT3	2.5	16Kn03
<sup>117</sup> Rh-u	-73903	408	-73964	10	-0.1	U			GT1	1.5	04Ma.A
<sup>117</sup> Ag- <sup>133</sup> Cs. <sub>880</sub>	-5029	16	-5024	15	0.3	1	83	83 <sup>117</sup> Ag	MA8	1.0	10Br02 *
C <sub>9</sub> H <sub>9</sub> - <sup>117</sup> Sn	167486	12	167471.3	0.5	-0.5	U			M16	2.5	63Da10
	167471	8			0.0	U			R13	1.5	83De51
C <sup>35</sup> Cl <sub>3</sub> - <sup>117</sup> Sn	3596	2	3604.1	0.5	1.0	U			H14	4.0	62Ba24
<sup>117</sup> Te-u	-91318	30	-91354	14	-1.2	-			GS2	1.0	05Li24
	-91359	30			0.2	-			GS2	1.0	05Li24 *
	ave.	-91339	21		-0.7	1	46	46 <sup>117</sup> Te			average
<sup>117</sup> I-u	-86350	30	-86352	28	-0.1	1	88	88 <sup>117</sup> I	GS2	1.0	05Li24
<sup>117</sup> Xe-u	-79647	30	-79641	11	0.2	R			GS2	1.0	05Li24
<sup>117</sup> Xe- <sup>133</sup> Cs. <sub>880</sub>	3562	12	3561	11	-0.1	2			MA6	1.0	04Di18
<sup>117</sup> Cs- <sup>133</sup> Cs. <sub>880</sub>	11819	67				2			MA4	1.0	99Am05 *
<sup>117</sup> Rh- <sup>120</sup> Sn. <sub>975</sub>	21388.8	9.5				2			JY1	1.0	07Ha20
<sup>117</sup> Pd- <sup>120</sup> Sn. <sub>975</sub>	13309.4	7.9	13308	8	-0.2	1	96	96 <sup>117</sup> Pd	JY1	1.0	07Ha20
<sup>117</sup> Sn- <sup>116</sup> Sn	1219	11	1211.2	0.5	-0.3	U			M16	2.5	63Da10
<sup>116</sup> Cd(d,p) <sup>117</sup> Cd	3538	30	3552.7	1.0	0.5	U			Pit		64Ro17
	3550	20			0.1	U			Oak		64Si18
	3552.66	1.0				2			Rez		90Pi05 *
<sup>116</sup> Sn(n, $\gamma$ ) <sup>117</sup> Sn	6943.5	2.0	6943.1	0.5	-0.2	U					75Bh01 Z
	6943.3	1.5			-0.1	U					78Ra16 Z
	6942.9	0.5			0.4	-			Bdn		06Fi.A
<sup>116</sup> Sn(d,p) <sup>117</sup> Sn	4721.0	1.8	4718.5	0.5	-1.4	-			SPa		75Be09
<sup>116</sup> Sn(n, $\gamma$ ) <sup>117</sup> Sn	ave.	6943.1	0.5	6943.1	0.5	0.0	1	97	97 <sup>117</sup> Sn		average
<sup>116</sup> Sn( <sup>3</sup> He,d) <sup>117</sup> Sb	-1091	10	-1091	8	0.0	1	71	71 <sup>117</sup> Sb	VUn		78Ka12 *
<sup>117</sup> Xe( $\epsilon p$ ) <sup>116</sup> Te	4100	200	3795	30	-1.5	U					72Ho18
<sup>117</sup> Ba( $\epsilon p$ ) <sup>116</sup> Xe	7900	300	8300	250	1.3	F					78Bo20 *
	8300	250				3			GSI		05Ja06
<sup>117</sup> La(p) <sup>116</sup> Ba	789.8	6.	820	3	5.0	B					01So02 *
	813.0	5.			1.4	o			Arp		01Ma69
	820.1	3.				3			Arp		11Li28
<sup>117</sup> Pd( $\beta^-$ ) <sup>117</sup> Ag	5735	32	5758	15	0.7	1	21	17 <sup>117</sup> Ag	Jyv		00Kr.A
<sup>117</sup> Ag( $\beta^-$ ) <sup>117</sup> Cd	4160	50	4236	14	1.5	U			Stu		82Al29 *
<sup>117</sup> Cd( $\beta^-$ ) <sup>117</sup> In	2535	20	2525	5	-0.5	U					75Ta06 *
<sup>117</sup> In( $\beta^-$ ) <sup>117</sup> Sn	1456.6	5.	1455	5	-0.4	1	94	94 <sup>117</sup> In			55Mc17 *
<sup>117</sup> Sb( $\beta^+$ ) <sup>117</sup> Sn	1751	40	1758	8	0.2	U					64Ba46 *
<sup>117</sup> Sn(p,n) <sup>117</sup> Sb	-2525	20	-2541	8	-0.8	1	18	18 <sup>117</sup> Sb	Oak		71Ke21
<sup>117</sup> Te( $\beta^+$ ) <sup>117</sup> Sb	3552	20	3544	13	-0.4	-					62Kh05 *
	3492	30			1.7	-					67Be46 *
	ave.	3534	17		0.6	1	62	51 <sup>117</sup> Te			average
<sup>117</sup> I( $\beta^+$ ) <sup>117</sup> Te	4680	100	4659	29	-0.2	-					69La33 *
	4610	110			0.4	-					70Be.A *
	ave.	4650	70		0.1	1	15	12 <sup>117</sup> I			average
<sup>117</sup> Xe( $\beta^+$ ) <sup>117</sup> I	6270	300	6251	28	-0.1	U					85Le10 *
<sup>117</sup> Cs <sup>x</sup> (IT) <sup>117</sup> Cs	50	50				3					AHW
* <sup>117</sup> Ag- <sup>133</sup> Cs. <sub>880</sub>	$D_M=-5013.3(4.0)$ $\mu$ u for ground state or <sup>117</sup> Ag <sup>m</sup> at 28.6 keV; $M-A=-82172.3(3.7)$ keV										Nub16b **
* <sup>117</sup> Te-u	$M-A=-84804(28)$ keV for <sup>117</sup> Te <sup>m</sup> 11/2 <sup>-</sup> at 296.1 keV										Nub16b **
* <sup>117</sup> Cs- <sup>133</sup> Cs. <sub>880</sub>	$D_M=11900(21)$ $\mu$ u for mixture gs+m at 150#80 keV; $M-A=-66418(19)$ keV										Nub16b **
* <sup>116</sup> Cd(d,p) <sup>117</sup> Cd	Estimated systematic error 0.5 added to statistical error 0.85 keV										AHW **
* <sup>116</sup> Sn( <sup>3</sup> He,d) <sup>117</sup> Sb	$Q-Q(^{120}\text{Sn}(^3\text{He,d}))=1373(10,\text{Ka})$ , $Q(120)=282.1(2.0)$ keV										AHW **
* <sup>117</sup> Ba( $\epsilon p$ ) <sup>116</sup> Xe	$F$ : disagrees with next and with TMS=8480#										GAu **
* <sup>117</sup> La(p) <sup>116</sup> Ba	Reports also an isomer <sup>117</sup> La <sup>m</sup> $E_p=933(10)$ $Q_p=941.1$ keV T=10(5) ms,										01So02 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
*	not observed in reference using similar set-up and far greater statistics										11Li28	**
* <sup>117</sup> Ag( $\beta^-$ ) <sup>117</sup> Cd	$Q_{\beta^-}$ =4260(110); and 4170(50) from <sup>117</sup> Ag <sup>m</sup> at 28.6 keV										Nub16b	**
* <sup>117</sup> Cd( $\beta^-$ ) <sup>117</sup> In	$Q_{\beta^-}$ =2220(20) to <sup>117</sup> In <sup>m</sup> at 315.303 keV										Nub16b	**
* <sup>117</sup> In( $\beta^-$ ) <sup>117</sup> Sn	$E_{\beta^-}$ =740(10) to 7/2 <sup>+</sup> level at 711.54; and 1772(5), 1616(5) from <sup>117</sup> In <sup>m</sup> at 315.303 to 1/2 <sup>+</sup> ground state, 3/2 <sup>+</sup> level at 158.56 keV										Ens111	**
*											Nub16b	**
* <sup>117</sup> Sb( $\beta^+$ ) <sup>117</sup> Sn	$E_{\beta^+}$ =570(40) to 3/2 <sup>+</sup> level at 158.562 keV										Ens111	**
* <sup>117</sup> Te( $\beta^+$ ) <sup>117</sup> Sb	$E_{\beta^+}$ =1810(20) 1750(30) respectively, to 1/2 <sup>+</sup> level at 719.7 keV										Ens111	**
* <sup>117</sup> I( $\beta^+$ ) <sup>117</sup> Te	$E_{\beta^+}$ =3500(50), 3250(50) to 5/2 <sup>+</sup> level at 274.4, (3/2) <sup>+</sup> at 325.9 keV										Ens111	**
* <sup>117</sup> I( $\beta^+$ ) <sup>117</sup> Te	$Q_{\beta^+}$ =4310(100) assumed to 5/2 <sup>+</sup> level at 274.4, (3/2) <sup>+</sup> at 325.9 keV										Ens111	**
* <sup>117</sup> Xe( $\beta^+$ ) <sup>117</sup> I	May be lower limit										AHW	**
<sup>118</sup> Ru-u	-61879	196	-61470#	220#	0.8	D			GT3	2.5	16Kn03	*
<sup>118</sup> Rh-u	-69598	290	-69660	26	-0.1	U			GT1	1.5	04Ma.A	
<sup>118</sup> Pd- <sup>129</sup> Xe. <sub>915</sub>	6193.6	4.3	6192.4	2.7	-0.3	1	39	39 <sup>118</sup> Pd	JY1	1.0	11Ha48	
<sup>118</sup> Ag- <sup>133</sup> Cs. <sub>887</sub>	-1540.4	2.7				2			MA8	1.0	10Br02	*
C <sub>9</sub> H <sub>10</sub> - <sup>118</sup> Sn	176645	7	176643.7	0.5	-0.1	U			M16	2.5	63Da10	
	176637	8			0.6	U			R13	1.5	83De51	
<sup>118</sup> Te-u	-94162	30	-94146	20	0.5	R			GS2	1.0	05Li24	
<sup>118</sup> I-u	-86932	30	-86926	21	0.2	2			GS2	1.0	05Li24	
	-86920	30			-0.2	2			GS2	1.0	05Li24	*
<sup>118</sup> Xe-u	-83785	30	-83821	11	-1.2	R			GS2	1.0	05Li24	
<sup>118</sup> Xe- <sup>133</sup> Cs. <sub>887</sub>	37	12	43	11	0.5	2			MA6	1.0	04Di18	
<sup>118</sup> Cs <sup>x</sup> - <sup>133</sup> Cs. <sub>887</sub>	10424	16	10429	13	0.3	o			MA1	1.0	90St25	
	10429	13				2			MA1	1.0	99Am05	
<sup>118</sup> Rh- <sup>120</sup> Sn. <sub>983</sub>	26476	26				2			JY1	1.0	07Ha20	
<sup>118</sup> Pd- <sup>120</sup> Sn. <sub>983</sub>	15199.7	7.9	15202.4	2.6	0.3	-			JY1	1.0	07Ha20	
	15202.1	3.6			0.1	-			JY1	1.0	11Ha48	
ave.	15202	3			0.2	1	64	61 <sup>118</sup> Pd			average	
<sup>118</sup> Sn <sup>35</sup> Cl- <sup>116</sup> Sn <sup>37</sup> Cl	2814	2	2813.9	0.5	0.0	U			H15	4.0	62Ba23	
<sup>118</sup> Sn- <sup>117</sup> Sn	-1338	11	-1347.41	0.14	-0.3	U			M16	2.5	63Da10	
<sup>117</sup> Cs <sup>x</sup> - <sup>118</sup> Cs <sup>x</sup> . <sub>496</sub> <sup>116</sup> Cs. <sub>504</sub>	-1160	400	-1250#	100#	-0.1	U			P32	2.5	86Au02	
<sup>118</sup> Cs( $\epsilon\alpha$ ) <sup>114</sup> Te	10600	200	11050	30	2.3	U					77Bo28	
	10750	200			1.5	U					78Da07	*
<sup>116</sup> Cd(t,p) <sup>118</sup> Cd	5650	20				2			Ald		67Hi01	
<sup>116</sup> Sn(t,p) <sup>118</sup> Sn	7769	15	7787.7	0.5	1.2	U			Ald		68Bj02	
<sup>118</sup> Sn(p,t) <sup>116</sup> Sn	-7790	10	-7787.7	0.5	0.2	U			Roc		70Fi08	
<sup>118</sup> Sn(d, <sup>3</sup> He) <sup>117</sup> In	-4440	40	-4505	5	-1.6	U			Sac		69Co03	
	-4481	15			-1.6	U			MSU		71We01	
<sup>117</sup> Sn(n, $\gamma$ ) <sup>118</sup> Sn	9326.5	2.	9326.42	0.13	0.0	U					70Or.A	
	9324.8	2.1			0.8	U			ORn		75Sl.A	
	9326.42	0.13			0.0	1	100	97 <sup>118</sup> Sn			02Bo11	
	9327.9	1.1			-1.3	U			Bdn		06Fi.A	
<sup>117</sup> Sn(d,p) <sup>118</sup> Sn	7090	12	7101.85	0.13	1.0	U			Tal		64No06	
<sup>118</sup> Sn(p,d) <sup>117</sup> Sn	-7097	15	-7101.85	0.13	-0.3	U			Har		70Ca01	
<sup>118</sup> Cs( $\epsilon p$ ) <sup>117</sup> I	4700	300	4738	29	0.1	U					78Da07	
<sup>118</sup> Pd( $\beta^-$ ) <sup>118</sup> Ag	4100	200	4165	4	0.3	U			Jyv		89Ko22	*
<sup>118</sup> Ag( $\beta^-$ ) <sup>118</sup> Cd	7122	100	7148	20	0.3	U			Stu		82Al29	*
	7110	470			0.1	U			Stu		82Al29	*
	7155	76			-0.1	U					95Ap.A	
<sup>118</sup> In( $\beta^-$ ) <sup>118</sup> Sn	4200	400	4425	8	0.6	U					61Gl02	
	4200	300			0.7	U					64Ka10	
	4310	100			1.1	U					87Ga.A	
<sup>118</sup> In <sup>m</sup> ( $\beta^-$ ) <sup>118</sup> Sn	4270	100	4530#	50#	2.5	D					64Ka10	*
<sup>118</sup> Sb( $\beta^+$ ) <sup>118</sup> Sn	3610	50	3656.6	3.0	0.9	U					61Fi05	
<sup>118</sup> Sn(p,n) <sup>118</sup> Sb	-4477.7	5.7	-4439.0	3.0	6.8	F			Tkm		63Ok01	*
	-4439.0	3.				2			Oak		77Jo03	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{118}\text{Sb}^n(\beta^+)^{118}\text{Sn}$	3907	5				2					61Bo13 *
$^{118}\text{I}(\beta^+)^{118}\text{Te}$	7080	150	6726	27	-2.4	U					68La18 *
	7068	100			-3.4	C					70Be.A *
$^{118}\text{Xe}(\beta^+)^{118}\text{I}$	3720	110	2892	22	-7.5	F					70Be.A *
$^{118}\text{Cs}(\beta^+)^{118}\text{Xe}$	9300	1000	9670	16	0.4	U					76Da.C *
$^{118}\text{Cs}^x(\text{IT})^{118}\text{Cs}$	5	4				3					82Au01 *
* $^{118}\text{Ru}-\text{u}$	Trends from Mass Surface TMS suggest $^{118}\text{Ru}$ 380 less bound										GAu **
* $^{118}\text{Ag}-^{133}\text{Cs}_{.887}$	$D_M=-1403.5(2.7) \mu\text{u}$ for $^{118}\text{Ag}^n$ at 127.63(0.10) keV; $M-A=-79426.3(2.5)$ keV										Nub16b **
* $^{118}\text{I}-\text{u}$	$M-A=-80775(28)$ keV for $^{118}\text{I}^m 7^-$ at 188.8 keV										Nub16b **
* $^{118}\text{Cs}(\varepsilon\alpha)^{114}\text{Te}$	As read from Fig. 2 (p.401)										GAu **
* $^{118}\text{Pd}(\beta^-)^{118}\text{Ag}$	Original value 4000(200) corrected for new branching ratios										93Ja03 **
* $^{118}\text{Ag}(\beta^-)^{118}\text{Cd}$	$E_{\beta^-}=4330(240), 3960(170), 3810(150)$ reinterpreted as feeding										95Ap.A **
*	(1) level at 2788.72, (1) at 3224.32, (2,3,4) at 3265.77 keV										Ens959 **
* $^{118}\text{Ag}(\beta^-)^{118}\text{Cd}$	$E_{\beta^-}=3990(720), 3910(630)$ reinterpreted as $^{118}\text{Ag}^n$ at 127.63(0.10) keV										95Ap.A **
*	to (2,3,4) level at 3181.73, 3381.8 keV										Ens959 **
* $^{118}\text{In}^m(\beta^-)^{118}\text{Sn}$	$E_{\beta^-}=2000(100)$ to $4^+$ level at 2280.342 level, and other $E_{\beta^-}$										Ens959 **
* $^{118}\text{In}^m(\beta^-)^{118}\text{Sn}$	Trends from Mass Surface TMS suggest $^{118}\text{In}^m$ 255 less bound										GAu **
* $^{118}\text{Sn}(p,n)^{118}\text{Sb}$	F : see note added in proof to reference										77Jo03 **
* $^{118}\text{Sb}^n(\beta^+)^{118}\text{Sn}$	$p^+ = 16(1) \times 10^{-4}$ to $7^-$ level at 2574.91 keV, recalculated $Q$										Ens959 **
* $^{118}\text{I}(\beta^+)^{118}\text{Te}$	$E_{\beta^+}=5450(150)$ to $2^+$ level at 605.70 keV										Ens959 **
* $^{118}\text{I}(\beta^+)^{118}\text{Te}$	$E_{\beta^+}=5440(100)$ to $2^+$ level at 605.70 keV										Ens959 **
* $^{118}\text{Xe}(\beta^+)^{118}\text{I}$	F : probably contaminated by isobars										GAu **
* $^{118}\text{Cs}^x(\text{IT})^{118}\text{Cs}$	Original 24(19) corrected for new estimated IT=100(60)#										Nub16b **
$^{119}\text{Rh}-\text{u}$	-67698	268	-67443	10	0.6	U			GT1	1.5	04Ma.A
$^{119}\text{Rh}-^{129}\text{Xe}_{.922}$	20349	10				2			JY1	1.0	11Ha48
$^{119}\text{Pd}-\text{u}$	-76844	208	-76660	9	0.6	U			GT1	1.5	04Ma.A
$^{119}\text{Ag}-^{133}\text{Cs}_{.895}$	188	16	191	16	0.2	1	97	97 $^{119}\text{Ag}$	MA8	1.0	10Br02 *
$\text{C}_9 \text{H}_{11}-^{119}\text{Sn}$	182778	7	182764.1	0.8	-0.8	U			M16	2.5	63Da10
	182762	8			0.2	U			R13	1.5	83De51
$^{119}\text{I}-\text{u}$	-89926	30				2			GS2	1.0	05Li24
$^{119}\text{Xe}-\text{u}$	-84601	30	-84589	11	0.4	R			GS2	1.0	05Li24
$^{119}\text{Xe}-^{133}\text{Cs}_{.895}$	33	12	31	11	-0.1	2			MA6	1.0	04Di18
$^{119}\text{Cs}-\text{u}$	-77532	57	-77623	15	-1.6	U			GS2	1.0	05Li24 *
$^{119}\text{Cs}^x-^{133}\text{Cs}_{.895}$	7011	16	7015	9	0.3	o			MA1	1.0	90St25
	7018	13			-0.2	2			MA1	1.0	99Am05
	7012	13			0.2	2			MA4	1.0	99Am05
$^{119}\text{Sn}^{35}\text{Cl}-^{117}\text{Sn}^{37}\text{Cl}$	3306	2	3307.3	0.6	0.2	U			H15	4.0	62Ba23
$^{119}\text{Pd}-^{120}\text{Sn}_{.992}$	20356.2	8.8				2			JY1	1.0	07Ha20
$^{119}\text{Sn}-^{118}\text{Sn}$	1709	12	1704.6	0.6	-0.1	U			M16	2.5	63Da10
$^{119}\text{I}-^{118}\text{I}$	-2747	155	-3000	40	-1.1	U			CR2	1.5	92Sh.A *
$^{119}\text{I}-^{117}\text{I}$	-3570	155	-3570	40	0.0	U			CR2	1.5	92Sh.A *
$^{118}\text{Cs}^x-^{119}\text{Cs}_{.661}^{116}\text{Cs}_{.339}$	530	80	410#	40#	-0.6	U			P32	2.5	86Au02
$^{118}\text{Cs}^x-^{119}\text{Cs}_{.496}^{117}\text{Cs}_{.504}$	870	50	940	40	0.5	U			P22	2.5	82Au01
	980	40			-0.4	U			P32	2.5	86Au02
$^{119}\text{Sn}(t,\alpha)^{118}\text{In}-^{118}\text{Sn}(0)^{117}\text{In}$	-127	6	-127	6	0.0	1	100	100 $^{118}\text{In}$	McM		85Pi03
$^{118}\text{Sn}(n,\gamma)^{119}\text{Sn}$	6484.6	1.5	6483.5	0.5	-0.7	-			ORn		78Ra16
	6483.3	0.6			0.3	-			Bdn		06Fi.A
$^{118}\text{Sn}(d,p)^{119}\text{Sn}$	4238	12	4258.9	0.5	1.7	U			MIT		67Sp09
$^{118}\text{Sn}(n,\gamma)^{119}\text{Sn}$	ave.	6483.5	0.6	6483.5	0.5	0.0	1	96	93 $^{119}\text{Sn}$		average
$^{118}\text{Sn}(^3\text{He},d)^{119}\text{Sb}$	-388	10	-383	8	0.5	1	59	59 $^{119}\text{Sb}$	VUn		78Ka12 *
$^{119}\text{Ba}(\text{ep})^{118}\text{Xe}$	6200	200				3					78Bo20 *
$^{119}\text{Ag}(\beta^-)^{119}\text{Cd}$	5350	40	5330	40	-0.5	1	81	78 $^{119}\text{Cd}$	Stu		82A129
$^{119}\text{Cd}(\beta^-)^{119}\text{In}$	3797	80	3720	40	-0.9	1	23	22 $^{119}\text{Cd}$	Stu		82A129 *
$^{119}\text{In}(\beta^-)^{119}\text{Sn}$	2387	100	2366	7	-0.2	U					60Yu01 *
	2413	200			-0.2	U					61Gl06 *
$^{119}\text{Sb}(\varepsilon)^{119}\text{Sn}$	579	20	591	8	0.6	-					57Ol05 *
$^{119}\text{Sn}(p,n)^{119}\text{Sb}$	-1369	15	-1373	8	-0.3	-			Oak		71Ke21
$^{119}\text{Sb}(\varepsilon)^{119}\text{Sn}$	ave.	584	12	591	8	0.6	1	41	41 $^{119}\text{Sb}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{119}\text{Te}(\beta^+)^{119}\text{Sb}$	2293	2				2					60Ko12 *
$^{119}\text{I}(\beta^+)^{119}\text{Te}$	3630	100	3416	29	-2.1	U					69La33 *
	3370	100			0.5	U					70Be.A
$^{119}\text{Xe}(\beta^+)^{119}\text{I}$	4990	120	4971	30	-0.2	U					70Be.A
$^{119}\text{Cs}(\beta^+)^{119}\text{Xe}$	6260	290	6489	17	0.8	U					83Pa.A *
$^{119}\text{Cs}^x(\text{IT})^{119}\text{Cs}$	16	11				3					82Au01 *
* $^{119}\text{Ag}-^{133}\text{Cs}_{.895}$	$D_M=198.4(5.7) \mu\text{u}$ for ground state or $^{119}\text{Ag}^m$ at 20#20 keV; $M-A=-78638.7(5.3) \text{keV}$										Nub16b **
* $^{119}\text{Cs}-\text{u}$	$M-A=-72195(48) \text{keV}$ for mixture gs+m at 50#30 keV										Nub16b **
* $^{119}\text{I}-^{118}\text{I}$	From $^{118}\text{I}^{119}\text{I}=0.99161584(117)$ , originally $D_M=-3039(139) \mu\text{u}$ , revised by										GAu **
*	authors: $-2849(139) \mu\text{u}$ for $^{118}\text{I}$ gs+m mixture at 188.8 keV										Nub16b **
* $^{119}\text{I}-^{117}\text{I}$	From $^{117}\text{I}^{119}\text{I}=0.98321059(130)$										GAu **
* $^{118}\text{Sn}(\beta^+\text{He,d})^{119}\text{Sb}$	$Q-Q(^{120}\text{Sn}(\beta^+\text{He,d})^{121}\text{Sb})=-673(10)$ , $Q(120)=285.1(2.1) \text{keV}$										AHW **
* $^{119}\text{Ba}(\epsilon\text{p})^{118}\text{Xe}$	Trends from Mass Surface TMS suggest $^{119}\text{Ba}$ 180 less bound										GAu **
* $^{119}\text{Cd}(\beta^-)^{119}\text{In}$	$Q_{\beta^-}=3800(90)$ ; and $3940(80)$ from $^{119}\text{Cd}^m$ at 146.54 keV										Nub16b **
* $^{119}\text{In}(\beta^-)^{119}\text{Sn}$	$E_{\beta^-}=1600(100)$ to $7/2^+$ level at 787.01 keV										Ens09a **
* $^{119}\text{In}(\beta^-)^{119}\text{Sn}$	$E_{\beta^-}=2700(200)$ from $^{119}\text{In}^m$ at 311.37(0.03) to $3/2^+$ level at 23.871 keV										Ens09a **
* $^{119}\text{Sb}(\epsilon)^{119}\text{Sn}$	IBE=526(20) to $3/2^+$ level at 23.871 keV										Ens09a **
* $^{119}\text{Te}(\beta^+)^{119}\text{Sb}$	$E_{\beta^+}=627(2)$ to $1/2^+$ level at 644.03 keV										Ens09a **
* $^{119}\text{I}(\beta^+)^{119}\text{Te}$	$E_{\beta^+}=2350(100)$ to $3/2^+$ level at 257.484 keV										Ens09a **
* $^{119}\text{Cs}(\beta^+)^{119}\text{Xe}$	$E_{\beta^+}=4980(290)$ to $9/2^+$ level at 257.84 keV										Ens09a **
* $^{119}\text{Cs}^x(\text{IT})^{119}\text{Cs}$	Original 33(22) corrected for new estimated IT=50(30)#										GAu **
$^{120}\text{Pd}-^{129}\text{Xe}_{.930}$	13107.0	4.4	13105.1	2.5	-0.4	1	31	31 $^{120}\text{Pd}$	JY1	1.0	11Ha48
$^{120}\text{Ag}-^{133}\text{Cs}_{.902}$	4067.1	4.8				2			MA8	1.0	10Br02
	4086	12	4067	5	-1.6	o			MA8	1.0	10Br02
$^{120}\text{Cd}-^{133}\text{Cs}_{.902}$	-4849.6	4.0				2			MA8	1.0	10Br02
$\text{C}_9 \text{H}_{12}-^{120}\text{Sn}$	191709	11	191698.5	1.0	-0.4	U			M16	2.5	63Da10
	191705	8			-0.5	U			R13	1.5	83De51
$^{13}\text{C} \ ^{35}\text{Cl}_2 \ ^{37}\text{Cl}-^{120}\text{Sn}$	4758	3	4760.9	1.0	0.2	U			H14	4.0	62Ba24
$^{120}\text{Sb}-\text{u}$	-94796	76	-94920	8	-1.6	U			GS2	1.0	05Li24 *
$\text{C}_9 \text{H}_{12}-^{120}\text{Te}$	189879	9	189841	3	-1.7	U			M16	2.5	63Da10
	189868	8			-2.3	U			R13	1.5	83De51
$^{120}\text{I}-\text{u}$	-90222	104	-89913	16	3.0	C			GS2	1.0	05Li24 *
$^{120}\text{Xe}-\text{u}$	-88231	30	-88216	13	0.5	R			GS2	1.0	05Li24
$^{120}\text{Xe}-^{133}\text{Cs}_{.902}$	-2930	14	-2933	13	-0.2	2			MA6	1.0	04Di18
$^{120}\text{Cs}-\text{u}$	-79342	54	-79323	11	0.4	U			GS2	1.0	05Li24 *
$^{120}\text{Cs}^x-^{133}\text{Cs}_{.902}$	5947	16	5965	10	1.1	o			MA1	1.0	90St25
	5956	12			0.7	2			MA1	1.0	99Am05
	5983	17			-1.1	2			MA4	1.0	99Am05
$^{120}\text{Sn} \ ^{35}\text{Cl}-^{118}\text{Sn} \ ^{37}\text{Cl}$	3546	2	3545.4	1.1	-0.1	U			H15	4.0	62Ba23
$^{120}\text{Pd}-^{120}\text{Sn}$	22317.1	9.7	22349.4	2.4	3.3	B			JY1	1.0	07Ha20
	22348.6	2.8			0.3	1	72	69 $^{120}\text{Pd}$	JY1	1.0	11Ha48
$^{120}\text{Te}-^{120}\text{Sn}$	1842.2	1.7	1858	3	9.1	B			CP1	1.0	09Sc19
	1839.7	1.7			10.6	B			CP1	1.0	09Sc19
$^{120}\text{Sn}-^{119}\text{Sn}$	-1113	11	-1109.3	1.2	0.1	U			M16	2.5	63Da10
$^{118}\text{Cs}^x-^{120}\text{Cs}^x \ ^{117}\text{Cs}^x_{.328} \ ^{117}\text{Cs}^x_{.672}$	460	120	480	60	0.1	U			P22	2.5	82Au01
$^{119}\text{Cs}^x-^{120}\text{Cs}^x \ ^{117}\text{Cs}^x_{.661} \ ^{117}\text{Cs}^x_{.339}$	-940	50	-928	29	0.1	U			P22	2.5	82Au01
$^{119}\text{Cs}^x-^{120}\text{Cs}^x \ ^{118}\text{Cs}^x_{.496} \ ^{118}\text{Cs}^x_{.504}$	-1220	30	-1167	11	0.7	U			P22	2.5	82Au01
	-1180	60			0.1	U			P32	2.5	86Au02
	-1200	30			0.4	U			P32	2.5	86Au02
	-1270	50			0.8	F			P32	2.5	86Au02 *
$^{120}\text{Cs}(\epsilon\alpha)^{116}\text{Te}$	9200	300	8955	30	-0.8	U					76Jo.A
$^{118}\text{Sn}(\text{t,p})^{120}\text{Sn}$	7107	15	7106.4	1.0	0.0	U			Ald		68Bj02
$^{120}\text{Sn}(\text{p,t})^{118}\text{Sn}$	-7109	10	-7106.4	1.0	0.3	U			Roc		70Fl08
$^{120}\text{Te}(\text{p,t})^{118}\text{Te}$	-9343	24	-9332	18	0.4	2			Win		74De31 *



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{120}\text{Sn}(d, ^3\text{He})^{119}\text{In}$	-5160	40	-5195	7	-0.9	U			Sac		69Co03
	-5169	20			-1.3	1	13	13 $^{119}\text{In}$	MSU		71We01
$^{120}\text{Sn}(t, \alpha)^{119}\text{In} - ^{118}\text{Sn}(^0)^{117}\text{In}$	-692	6	-689	6	0.5	1	92	86 $^{119}\text{In}$	McM		85Pi03
$^{119}\text{Sn}(d, p)^{120}\text{Sn}$	6890	12	6880.1	1.1	-0.8	U			Tal		64No06
$^{120}\text{Sn}(p, d)^{119}\text{Sn}$	-6889	15	-6880.1	1.1	0.6	U			Har		70Ca01
$^{120}\text{Sn}(d, t)^{119}\text{Sn}$	-2847.0	2.5	-2847.4	1.1	-0.2	1	19	12 $^{120}\text{Sn}$	SPa		75Be09
$^{120}\text{Pd}(\beta^-)^{120}\text{Ag}$	5500	100	5371	5	-1.3	U			Jyv		94Jo.A
$^{120}\text{Ag}(\beta^-)^{120}\text{Cd}$	8200	100	8306	6	1.1	U			Stu		82Al29
	8450	100			-1.4	U					95Ap.A
$^{120}\text{In}(\beta^-)^{120}\text{Sn}$	5300	170	5370	40	0.4	U			Stu		78Al18
	5370	40				2					87Ga.A
$^{120}\text{In}^m(\beta^-)^{120}\text{Sn}$	5280	200	5420#	50#	0.7	D					64Ka10 *
	5340	170			0.5	D			Stu		78Al18 *
$^{120}\text{Sb}(\beta^+)^{120}\text{Sn}$	2720	20	2681	7	-2.0	U					50B192
	2770	30			-3.0	B					69Ki15
$^{120}\text{Sn}(p, n)^{120}\text{Sb}$	-3462.9	7.1				2			Tkm		63Ok01
$^{120}\text{I}(\beta^+)^{120}\text{Te}$	5615	15				2					70Ga32 *
	5608	150	5615	15	0.0	U					68La18 *
$^{120}\text{Xe}(\beta^+)^{120}\text{I}$	1960	40	1581	19	-9.5	B					74Mu10 *
$^{120}\text{Cs}(\beta^+)^{120}\text{Xe}$	7300	500	8284	15	2.0	U					76Ba.A *
	7800	1000			0.5	U					76Da.C *
	7380	230			3.9	C					83Pa.A *
	8210	200			0.4	U			IRS		93Al03
$^{120}\text{Cs}^x(\text{IT})^{120}\text{Cs}$	5	4				3					82Au01 *
$^{120}\text{Ba}(\beta^+)^{120}\text{Cs}$	5000	300				4					92Xu04
* $^{120}\text{Sb}-u$	$M-A=-88302(50)$ keV for mixture gs+m at 0#100 keV										Nub16b **
* $^{120}\text{I}-u$	$M-A=-83881(28)$ keV for mixture gs+n at 320(15) keV										Nub16b **
* $^{120}\text{Cs}-u$	$M-A=-73856(29)$ keV for mixture gs+m at 100#60 keV										Nub16b **
* $^{119}\text{Cs}^x - ^{120}\text{Cs}^x$ $^{118}\text{C}$	F : rejection based on line-shape analysis										86Au02 **
* $^{120}\text{Te}(p, t)^{118}\text{Te}$	Original error 12; added systematic error 21 keV										GAu **
* $^{120}\text{In}^m(\beta^-)^{120}\text{Sn}$	$E_{\beta^-}=3100(200), 2200(200)$ to $4^+$ levels at 2194.299, 3057.946 keV										Ens029 **
* $^{120}\text{In}^m(\beta^-)^{120}\text{Sn}$	$E_{\beta^-}=3100(170)$ to $4^+$ level at 2194.299 keV, and other $E_{\beta^-}$										Ens029 **
* $^{120}\text{In}^m(\beta^-)^{120}\text{Sn}$	Trends from Mass Surface TMS suggest $^{120}\text{In}^m$ 105 less bound										GAu **
* $^{120}\text{I}(\beta^+)^{120}\text{Te}$	$E_{\beta^+}=4595(15), 4030(20)$ to ground state, $2^+$ level at 560.438 keV										Ens029 **
* $^{120}\text{I}(\beta^+)^{120}\text{Te}$	$E_{\beta^+}=3130(150)$ from $^{120}\text{I}^m$ at 320(15) to $6^+$ level at 1776.23 keV										Ens029 **
* $^{120}\text{Xe}(\beta^+)^{120}\text{I}$	$p^+ = 0.07(0.01)$ to $1^+$ level at 25.1 keV, recalculated $Q$										Ens029 **
* $^{120}\text{Cs}(\beta^+)^{120}\text{Xe}$	$E_{\beta^+}=6000(500) 6500(1000) 6040(230)$ , respectively, to $2^+$ level at 322.61 keV										Ens029 **
* $^{120}\text{Cs}^x(\text{IT})^{120}\text{Cs}$	Original 24(19) corrected for new estimated IT=100(60)#										Nub16b **
$^{121}\text{Rh}-u$	-60387	266				2			GT3	2.5	16Kn03
$^{121}\text{Pd}-u$	-71820	311	-71050	4	1.7	U			GT1	1.5	04Ma.A
$^{121}\text{Ag}-^{133}\text{Cs}_{910}$	6164	13				2			MA8	1.0	10Br02 *
	6170	17	6164	13	-0.4	o			MA8	1.0	10Br02 *
$^{121}\text{Cd}-^{130}\text{Xe}_{931}$	2796.2	3.0	2796.5	2.1	0.1	2			JY1	1.0	12Ha25
	2796.7	2.9			-0.1	2			JY1	1.0	13Ka08 *
$\text{C}_9 \text{H}_{13} - ^{121}\text{Sb}$	197910.5	3.7	197915.3	2.8	0.5	U			M16	2.5	63Da10
	197910	8			0.4	U			R13	1.5	83De51
$^{121}\text{Sb}-\text{C } ^{35}\text{Cl } ^{37}\text{Cl}_2$	3162	3	3152.2	2.8	-0.8	U			H14	4.0	62Ba24
$^{121}\text{Sb}-u$	-96180	30	-96189.9	2.8	-0.3	U			GS2	1.0	05Li24
$^{121}\text{I}-u$	-92609	30	-92595	6	0.5	U			GS2	1.0	05Li24
$^{121}\text{Xe}-u$	-88562	30	-88547	11	0.5	R			GS2	1.0	05Li24
$^{121}\text{Xe}-^{133}\text{Cs}_{910}$	-2495	13	-2508	11	-1.0	-			MA6	1.0	04Di18
	ave.	-2499	12		-0.7	1	85	85 $^{121}\text{Xe}$			average
$^{121}\text{Cs}-^{133}\text{Cs}_{910}$	3247	25	3266	15	0.8	o			MA1	1.0	90St25 *
	3248	25			0.7	1	38	38 $^{121}\text{Cs}$	MA1	1.0	99Am05 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{121}\text{Cs}-u$	-82821	38	-82773	15	1.3	1	16	$^{16}\text{ }^{121}\text{Cs}$	GS2	1.0	05Li24 *
$^{121}\text{Pd}-^{129}\text{Xe}_{938}$	18265.9	3.6				2			JY1	1.0	11Ha48 *
$^{121}\text{Sb }^{35}\text{Cl}-^{119}\text{Sn }^{37}\text{Cl}$	3452	2	3449.0	2.9	-0.4	U			H14	4.0	62Ba24
$^{119}\text{Cs}^x-^{121}\text{Cs}^x_{328}$ $^{118}\text{Cs}^x_{672}$	-1080	30	-1047	13	0.4	U			P22	2.5	82Au01
$^{120}\text{Cs}^x-^{121}\text{Cs}^x_{661}$ $^{118}\text{Cs}^x_{339}$	280	30	240	15	-0.5	U			P22	2.5	82Au01
$^{120}\text{Cs}^x-^{121}\text{Cs}^x_{496}$ $^{119}\text{Cs}^x_{504}$	790	30	770	13	-0.3	U			P22	2.5	82Au01
	860	50			-0.7	U			P32	2.5	86Au02
	813	14			-1.2	U			P32	2.5	86Au02
$^{120}\text{Sn}(n,\gamma)^{121}\text{Sn}$	6170.3	2.	6170.2	0.3	0.0	U					76Ca24
	6170.5	0.7			-0.4	-					81Ba53
	6170.1	0.4			0.3	-			Bdn		06Fi.A
$^{120}\text{Sn}(d,p)^{121}\text{Sn}$	3946.2	1.7	3945.6	0.3	-0.3	-			SPa		75Be09
$^{120}\text{Sn}(n,\gamma)^{121}\text{Sn}$	ave. 6170.2	0.3	6170.2	0.3	0.0	1	99	$^{99}\text{ }^{121}\text{Sn}$			average
$^{121}\text{Sb}(\gamma,n)^{120}\text{Sb}$	-9310	60	-9254	8	0.9	U			Phi		60Ge01
	-9240	25			-0.6	U			MeM		79Ba06
$^{120}\text{Te}(^3\text{He},d)^{121}\text{I}$	-1320.5	4.4	-1321	4	-0.1	1	99	$^{99}\text{ }^{121}\text{I}$	Hei		78Sz09
$^{121}\text{Ba}(\epsilon p)^{120}\text{Xe}$	4200	300	4140	140	-0.2	R					78Bo20
$^{121}\text{Pr}(p)^{120}\text{Ce}$	837	50	890	10	1.1	F					90Bo39 *
	889.6	10.				3			Arp		05Ro19 *
$^{121}\text{Ag}(\beta^-)^{121}\text{Cd}$	6400	120	6671	12	2.3	U			Stu		82Al29
$^{121}\text{Cd}(\beta^-)^{121}\text{In}$	4780	80	4762	27	-0.2	U			Stu		82Al29 *
$^{121}\text{In}(\beta^-)^{121}\text{Sn}$	3426	200	3361	27	-0.3	U					60Yu01 *
	3406	50			-0.9	R			Stu		78Al18 *
$^{121}\text{Sn}(\beta^-)^{121}\text{Sb}$	383	5	403.1	2.7	4.0	B					49Du15
	383.4	3.			6.6	B					68Sn01 *
$^{121}\text{Te}(\beta^+)^{121}\text{Sb}$	1080	30	1055	26	-0.8	1	74	$^{74}\text{ }^{121}\text{Te}$			75Me23 *
$^{121}\text{I}(\beta^+)^{121}\text{Te}$	2364	50	2294	26	-1.4	1	27	$^{27}\text{ }^{121}\text{Te}$			53Fi.A *
	2384	100			-0.9	U					65Bu03 *
$^{121}\text{Xe}(\beta^+)^{121}\text{I}$	3790	100	3770	12	-0.2	U					60Mo.A
	4160	140			-2.8	U					70Be.A
$^{121}\text{Cs}(\beta^+)^{121}\text{Xe}$	5650	490	5379	14	-0.6	U					75We23
	5400	20			-1.1	-					81So06
	5210	220			0.8	U					83Pa.A *
	5300	100			0.8	U			IRS		93Al03 *
	5400	40			-0.5	-			JAE		96Os04 *
	ave. 5400	18			-1.2	1	61	$^{46}\text{ }^{121}\text{Cs}$			average
$^{121}\text{Cs}^x(\text{IT})^{121}\text{Cs}$	46	8				2					GAu
$^{121}\text{Ba}(\beta^+)^{121}\text{Cs}$	6340	160	6360	140	0.1	2			JAE		96Os04
* $^{121}\text{Ag}-^{133}\text{Cs}_{910}$	$D_M=6175.1(5.0)\mu\text{u}$ for ground state or $^{121}\text{Ag}^m$ at 20#20 keV; $M-A=-74392.5(4.7)\text{keV}$										
* $^{121}\text{Ag}-^{133}\text{Cs}_{910}$	$D_M=6180(12)\mu\text{u}$ for ground state or $^{121}\text{Ag}^m$ at 20#20 keV; $M-A=-74388(11)\text{keV}$										
* $^{121}\text{Cd}-^{130}\text{Xe}_{931}$	$D_M=3027.4(2.9)\mu\text{u}$ for $^{121}\text{Cd}^m$ at 214.86(0.15) keV; $M-A=-80858.7(2.7)\text{keV}$										
* $^{121}\text{Cs}-^{133}\text{Cs}_{910}$	$D_M=3284(16)\mu\text{u}$ for mixture gs+m at 68.5 keV										
* $^{121}\text{Cs}-^{133}\text{Cs}_{910}$	$D_M=3285(13)\mu\text{u}$ for mixture gs+m at 68.5 keV; $M-A=-77084(12)\text{keV}$										
* $^{121}\text{Cs}-u$	$M-A=-77113(29)\text{keV}$ for mixture gs+m at 68.5 keV										
* $^{121}\text{Pd}-^{129}\text{Xe}_{938}$	Taken as low-spin isomer (see also $^{102}\text{Y}$ and $^{114}\text{Tc}$ doublets in same paper)										
* $^{121}\text{Pr}(p)^{120}\text{Ce}$	F : misassigned according to reference										
* $^{121}\text{Pr}(p)^{120}\text{Ce}$	$E_p=882(10)$ ; in publication $Q_p=900(10)\text{keV}$										
* $^{121}\text{Cd}(\beta^-)^{121}\text{In}$	$Q_{\beta^-}=4890(150)$ ; and $4960(80)$ from $^{121}\text{Cd}^m$ at 214.86 keV										
* $^{121}\text{In}(\beta^-)^{121}\text{Sn}$	$E_{\beta^-}=3700(200)$ from $^{121}\text{In}^m$ at 313.68(0.07) to ground state and $1/2^+$ level at 60.34 keV										
* $^{121}\text{In}(\beta^-)^{121}\text{Sn}$	$E_{\beta^-}=2480(50)$ to $7/2^+$ level at 925.59 keV										
* $^{121}\text{Sn}(\beta^-)^{121}\text{Sb}$	$E_{\beta^-}=383(3)$ ; and $354(5)$ from $^{121}\text{Sn}^m$ at 6.31 to $7/2^+$ level at 37.1298 keV										
* $^{121}\text{Te}(\beta^+)^{121}\text{Sb}$	$p^+=0.024(0.011)$ gives $Q_{\beta^+}=315(30)$ , recalculated $Q_{\beta^+}$										
*	from $^{121}\text{Te}^m$ at 293.974 to $7/2^+$ level at 37.1298 keV										
* $^{121}\text{I}(\beta^+)^{121}\text{Te}$	$E_{\beta^+}=1130(50)$ $1150(100)$ respectively, to $3/2^+$ level at 212.191 keV										
* $^{121}\text{Cs}(\beta^+)^{121}\text{Xe}$	$E_{\beta^+}=3730(220)$ to $7/2^+$ level at 459.59 keV										
* $^{121}\text{Cs}(\beta^+)^{121}\text{Xe}$	$Q_{\beta^+}=5370(100)$ $5470(40)$ respectively, from $^{121}\text{Cs}^m$ at 68.5 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{122}\text{Pd}-u$	-69308	397	-69368	21	-0.1	U			GT1	1.5	04Ma.A
$^{122}\text{Ag}-u$	-76280	110	-76340	40	-0.2	o			GT2	2.5	08Kn.A *
	-76340	130			0.0	U			GT2	2.5	08Su19 *
$^{122}\text{Ag}-^{133}\text{Cs}_{.917}$	10365	41				2			MA8	1.0	10Br02 *
$^{122}\text{Cd}-^{133}\text{Cs}_{.917}$	155.1	4.7	159.6	2.5	1.0	1	28	28 $^{122}\text{Cd}$	MA8	1.0	10Br02
$\text{C}_8 \text{H}_{12} \text{N}-^{122}\text{Sn}$	193541	8	193530.4	2.6	-0.5	U			M16	2.5	63Da10
	193558	8			-2.3	U			R13	1.5	83De51
$\text{C}_8 \text{H}_{12} \text{N}-^{122}\text{Te}$	193925	9	193931.0	1.6	0.3	U			M16	2.5	63Da10
	193926	8			0.4	U			R13	1.5	83De51
$^{122}\text{Xe}-u$	-91637	30	-91632	12	0.2	R			GS2	1.0	05Li24
$^{122}\text{Xe}-^{133}\text{Cs}_{.917}$	-4931	13	-4932	12	-0.1	2			MA6	1.0	04Di18
$^{122}\text{Cs}-^{133}\text{Cs}_{.917}$	2812	48	2810	40	-0.1	o			MA1	1.0	90St25 *
	2805	48			0.1	1	57	57 $^{122}\text{Cs}$	MA1	1.0	99Am05 *
$^{122}\text{Cs}-u$	-83887	55	-83890	40	-0.1	1	43	43 $^{122}\text{Cs}$	GS2	1.0	05Li24 *
$^{122}\text{Cs}^n-^{133}\text{Cs}_{.917}$	2952	16	2959	10	0.4	o			MA1	1.0	90St25
	2961	12			-0.2	2			MA1	1.0	99Am05
	2955	17			0.2	2			MA4	1.0	99Am05
$^{122}\text{Ba}-u$	-80096	30				2			GS2	1.0	05Li24
$^{122}\text{Cd}-^{130}\text{Xe}_{.938}$	3969.0	2.9	3967.3	2.5	-0.6	1	72	72 $^{122}\text{Cd}$	JY1	1.0	12Ha25
$^{122}\text{Pd}-^{129}\text{Xe}_{.946}$	20709	21				2			JY1	1.0	11Ha48
$^{122}\text{Sn} \text{ } ^{35}\text{Cl}-^{120}\text{Sn} \text{ } ^{37}\text{Cl}$	4196	2	4192.2	2.5	-0.5	U			H15	4.0	62Ba23
$^{119}\text{Cs}^x-^{122}\text{Cs}^x_{.244} \text{ } ^{118}\text{Cs}^x_{.756}$	-1600	80	-1511	15	0.4	U			P32	2.5	86Au02
$^{120}\text{Cs}^x-^{122}\text{Cs}^x_{.492} \text{ } ^{118}\text{Cs}^x_{.508}$	-724	27	-694	20	0.4	U			P32	2.5	86Au02
$^{120}\text{Cs}^x-^{122}\text{Cs}^x_{.328} \text{ } ^{119}\text{Cs}^x_{.672}$	350	50	321	16	-0.2	U			P22	2.5	82Au01
	360	17			-0.9	U			P32	2.5	86Au02
$^{121}\text{Cs}^x-^{122}\text{Cs}^x_{.496} \text{ } ^{120}\text{Cs}^x_{.504}$	-1100	40	-1066	24	0.3	U			P32	2.5	86Au02
	-1169	15			2.7	U			P32	2.5	86Au02
$^{122}\text{Sn}(p,t)^{120}\text{Sn}$	-6504	15	-6503.8	2.3	0.0	U			Roc		70FI08
$^{122}\text{Te}(p,t)^{120}\text{Te}$	-8560	24	-8607.3	2.7	-2.0	U			Win		74De31 *
$^{122}\text{Te}(p,t)^{120}\text{Te}-^{132}\text{Ba}()^{130}\text{Ba}$	227.0	0.2	227.00	0.20	0.0	1	100	80 $^{120}\text{Te}$			08Su14
$^{122}\text{Te}(p,t)^{120}\text{Te}-^{144}\text{Sm}()^{142}\text{Sm}$	2032.6	0.4	2032.6	0.4	0.0	1	100	79 $^{142}\text{Sm}$			09Bu.A
$^{122}\text{Sn}(d,^3\text{He})^{121}\text{In}$	-5910	50	-5901	27	0.2	2			Sac		69Co03
	-5861	43			-0.9	2			MSU		71We01
$^{122}\text{Sn}(p,d)^{121}\text{Sn}$	-6587	15	-6590.8	2.3	-0.3	U			Har		70Ca01
$^{122}\text{Sn}(d,t)^{121}\text{Sn}$	-2558.8	3.0	-2558.2	2.3	0.2	1	60	57 $^{122}\text{Sn}$	SPa		75Be09
$^{121}\text{Sb}(n,\gamma)^{122}\text{Sb}$	6806.4	0.3	6806.37	0.13	-0.1	-					72Sh.A Z
	6806.36	0.15			0.0	-			Bdn		06Fi.A
ave.	6806.37	0.13			0.0	1	100	95 $^{121}\text{Sb}$			average
$^{122}\text{In}(\beta^-)^{122}\text{Sn}$	6440	200	6370	50	-0.4	U					71Ta07 *
	6510	230			-0.6	U			Stu		78A118
$^{122}\text{Sn}(t,^3\text{He})^{122}\text{In}$	-6350	50				2			LAl		78Aj01
$^{122}\text{In}^n(\beta^-)^{122}\text{Sn}$	6736	200	6660	130	-0.4	2					71Ta07 *
	6590	180			0.4	2			Stu		78A118
$^{122}\text{Sb}(\beta^+)^{122}\text{Sn}$	1587	25	1606	3	0.8	U					58Pe17
$^{122}\text{Sb}(\beta^-)^{122}\text{Te}$	1970	5	1979.1	2.1	1.8	-					55Fa33
	1980	3			-0.3	-					68Hs02
ave.	1977.4	2.6			0.7	1	68	67 $^{122}\text{Sb}$			average
$^{122}\text{I}(\beta^+)^{122}\text{Te}$	4140	40	4234	5	2.4	U					54Ma75
	4140	40			2.4	U					60Mo.A
	4234	5				2					77Re.A
$^{122}\text{Cs}(\beta^+)^{122}\text{Xe}$	7150	700	7210	40	0.1	U					75We23 *
	7050	180			0.9	U					83Pa.A *
	7000	150			1.4	U			IRS		93Al03
	7080	50			2.6	B			JAE		96Os04
$^{122}\text{Cs}^n(\beta^+)^{122}\text{Xe}$	6950	250	7350	14	1.6	U					83Pa.A *
	7300	150			0.3	U			IRS		93Al03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{122}\text{Cs}^x(\text{IT})^{122}\text{Cs}$	14	7				2					82Au01 *	
* $^{122}\text{Ag}-\text{u}$	$M-A=-71014(94)$ keV for mixture gs+m at 80#(50#) keV										Nub16b **	
* $^{122}\text{Ag}-\text{u}$	$M-A=-71065(120)$ keV for mixture gs+m at 80#(50#) keV										Nub16b **	
* $^{122}\text{Ag}-^{133}\text{Cs}_{.917}$	$D_M=10408(18)$ $\mu\text{u}$ for mixture gs+m at 80#(50#) keV; $M-A=-71066(17)$ keV										Nub16b **	
* $^{122}\text{Cs}-^{133}\text{Cs}_{.917}$	$D_M=2887(12)$ $\mu\text{u}$ for mixture gs+n at 140(30) keV										Nub16b **	
* $^{122}\text{Cs}-^{133}\text{Cs}_{.917}$	$D_M=2880(12)$ $\mu\text{u}$ for mixture gs+n at 140(30) keV; $M-A=-78078(12)$ keV										Nub16b **	
* $^{122}\text{Cs}-\text{u}$	$M-A=-78070(28)$ keV for mixture gs+m at 140(30) keV										Nub16b **	
* $^{122}\text{Te}(\text{p,t})^{120}\text{Te}$	Original error 12; added systematic error 21 keV										GAu **	
* $^{122}\text{In}(\beta^-)^{122}\text{Sn}$	$E_{\beta^-}=5300(200)$ to $2^+$ level at 1140.51 keV										Ens074 **	
* $^{122}\text{In}^n(\beta^-)^{122}\text{Sn}$	$E_{\beta^-}=4400(200)$ to $4^+$ level at 2331.09 keV										Ens074 **	
* $^{122}\text{Cs}(\beta^+)^{122}\text{Xe}$	$E_{\beta^+}=5800(700)$ 5690(180) respectively, to $2^+$ level at 331.28 keV										Ens074 **	
* $^{122}\text{Cs}^n(\beta^+)^{122}\text{Xe}$	$E_{\beta^+}=3710(250)$ to $8^+$ level at 2217.69 keV										Ens074 **	
* $^{122}\text{Cs}^x(\text{IT})^{122}\text{Cs}$	Original was 45(33); revised using $^{122}\text{Cs}^n=140(30)$ keV										Nub16b **	
$^{123}\text{Pd}-\text{u}$	-64423	290	-64870	850	-1.0	o			GT1	1.5	04Ma.A	
	-64874	339				2			GT3	2.5	16Kn03	
$^{123}\text{Ag}-\text{u}$	-74729	215	-74660	30	0.2	o			GT1	1.5	04Ma.A	
	-74479	130			-0.6	U			GT2	2.5	08Su19 *	
$^{123}\text{Ag}-^{133}\text{Cs}_{.925}$	12700	120	12790	30	0.8	U			MA8	1.0	10Br02	
	12794	33				2			MA8	1.0	10Br02 *	
$^{123}\text{Cd}-^{133}\text{Cs}_{.925}$	4491	52	4349.4	2.9	-2.7	U			MA8	1.0	10Br02 *	
$\text{C}_8 \text{H}_{13} \text{N}-^{123}\text{Sb}$	200580.0	3.3	200585.4	1.6	0.7	U			M16	2.5	63Da10	
	200615	8			-2.5	U			R13	1.5	83De51	
$\text{C}_8 \text{H}_{13} \text{N}-^{123}\text{Te}$	200538	16	200529.7	1.6	-0.2	U			M16	2.5	63Da10	
	200515	8			1.2	U			R13	1.5	83De51	
$^{123}\text{Te}-\text{u}$	-95615	83	-95730.3	1.6	-1.4	U			GS2	1.0	05Li24 *	
$^{123}\text{I}-\text{u}$	-94444	30	-94411	4	1.1	U			GS2	1.0	05Li24	
$^{123}\text{Xe}-^{133}\text{Cs}_{.925}$	-4048	13	-4061	10	-1.0	1	62	62	$^{123}\text{Xe}$	MA6	1.0	04Di18
$^{123}\text{Cs}-\text{u}$	-87007	57	-87004	13	0.1	U			GS2	1.0	05Li24 *	
$^{123}\text{Cs}-^{133}\text{Cs}_{.925}$	447	17	453	13	0.4	o			MA1	1.0	90St25	
	453	13				2			MA1	1.0	99Am05	
$^{123}\text{Ba}-^{133}\text{Cs}_{.925}$	6238	13				2			MA5	1.0	00Be42	
$^{123}\text{Ba}-\text{u}$	-81327	30	-81219	13	3.6	C			GS2	1.0	05Li24	
$^{123}\text{Cd}-^{130}\text{Xe}_{.946}$	8172.5	2.9	8172.6	2.9	0.0	1	100	100	$^{123}\text{Cd}$	JY1	1.0	12Ha25
$^{123}\text{Cd}^m-^{130}\text{Xe}_{.946}$	8326.5	3.3				2			JY1	1.0	13Ka08	
$^{123}\text{Sb } ^{35}\text{Cl}-^{121}\text{Sb } ^{37}\text{Cl}$	3343	2	3354.0	2.3	1.4	U			H14	4.0	62Ba24	
$^{123}\text{Te}-^{123}\text{Sb}$	55.729	0.071	55.73	0.07	0.0	1	100	96	$^{123}\text{Sb}$	SH2	1.0	16Fi07
$^{119}\text{Cs}^x-^{123}\text{Cs}_{.193}^x$ $^{118}\text{Cs}_{.807}^x$	-1480	60	-1447	13	0.2	U			P32	2.5	86Au02	
$^{123}\text{Te}(\text{n},\alpha)^{120}\text{Sn}$	7564	30	7572.6	1.7	0.3	U			ILL		75Em04	
$^{121}\text{Sb}(\text{t,p})^{123}\text{Sb}$	7295	20	7284.6	2.1	-0.5	U			Ald		67Hi01	
$^{122}\text{Sn}(\text{n},\gamma)^{123}\text{Sn}$	5948	3	5946.2	1.2	-0.6	-					75Bh01	
	5945.8	1.5			0.2	-					77Ca09	
$^{122}\text{Sn}(\text{d,p})^{123}\text{Sn}$	3726	12	3721.6	1.2	-0.4	U			Tal		64Ne10	
	3716	11			0.5	U					72Ca33	
	3721.8	2.6			-0.1	-			SPa		75Be09	
$^{122}\text{Sn}(\text{n},\gamma)^{123}\text{Sn}$	ave.	5946.3	5946.2	1.2	-0.1	1	94	51	$^{123}\text{Sn}$		average	
$^{123}\text{Sb}(\gamma,\text{n})^{122}\text{Sb}$	-8980	50	-8960.0	2.1	0.4	U			Phi		60Ge01	
	-8966	4			1.5	1	28	28	$^{122}\text{Sb}$	McM	79Ba06	
$^{122}\text{Te}(\text{n},\gamma)^{123}\text{Te}$	6937	5	6929.01	0.08	-1.6	U					68Ch.A	
	6929.1	0.5			-0.2	U					91Ho08	
	6928.97	0.09			0.5	-					00Bo24	
	6929.16	0.17			-0.9	-			Bdn		06Fi.A	
$^{122}\text{Te}(\text{d,p})^{123}\text{Te}$	4706	6	4704.45	0.08	-0.3	U			MIT		75Li22	
$^{122}\text{Te}(\text{n},\gamma)^{123}\text{Te}$	ave.	6929.01	6929.01	0.08	0.0	1	100	98	$^{122}\text{Te}$		average	
$^{122}\text{Te}(\text{He},\text{d})^{123}\text{I}$	-574.2	3.5	-575	3	-0.3	1	97	96	$^{123}\text{I}$	Hei	78Sz04	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{123}\text{Cd}(\beta^-)^{123}\text{In}$	6033	35	6016	20	-0.5	1	32	32 $^{123}\text{In}$	Stu		87Sp09 *
$^{123}\text{In}(\beta^-)^{123}\text{Sn}$	4400	30	4386	20	-0.5	1	44	43 $^{123}\text{In}$	Stu		87Sp09 *
$^{123}\text{Sn}(\beta^-)^{123}\text{Sb}$	1395	10	1407.9	2.7	1.3	-					49Du15 *
		10			-1.2	-					50Ke11
		20			0.4	U					66Au04 *
ave.	1408	7			0.1	1	14	11 $^{123}\text{Sn}$			average
$^{123}\text{I}(\beta^+)^{123}\text{Te}$	1260	7	1228	3	-4.5	C					86Ag.A
$^{123}\text{Xe}(\beta^+)^{123}\text{I}$	2720	100	2695	10	-0.2	U					54Ma75
	2676	15			1.3	1	42	38 $^{123}\text{Xe}$			60Mo.A *
$^{123}\text{Cs}(\beta^+)^{123}\text{Xe}$	4110	310	4205	15	0.3	U					75We23 *
	4000	140			1.5	U					81So06 *
	4050	180			0.9	U					83Pa.A *
	4200	100			0.1	U			IRS		93Al03
	4110	30			3.2	B			JAE		96Os04
$^{123}\text{Cs}^x(\text{IT})^{123}\text{Cs}$	7	4				3					82Au01 *
$^{123}\text{Ba}(\beta^+)^{123}\text{Cs}$	5330	100	5389	17	0.6	U			JAE		96Os04
$^{123}\text{Ag}-u$	Isomer should be expected, but $^{123}\text{Ag}^m$ at 20#(20#) keV, correction negligible										GAu **
$^{123}\text{Ag}-^{133}\text{Cs}_{925}$	$D_M=12805(30) \mu\text{u}$ for mixture gs+m at 20#(20#) keV; $M-A=-69538(28)$										Nub16b **
$^{123}\text{Cd}-^{133}\text{Cs}_{925}$	$D_M=4568(26) \mu\text{u}$ for mixture gs+m at 143(4) keV; $M-A=-77210(25)$										Nub16b **
$^{123}\text{Te}-u$	$M-A=-88941(30)$ keV for mixture gs+m at 247.47 keV										Nub16b **
$^{123}\text{Cs}-u$	$M-A=-80968(28)$ keV for mixture gs+m at 156.27 keV										Nub16b **
$^{123}\text{Cd}(\beta^-)^{123}\text{In}$	$Q=3590(51) 3464(41) 3547(36)$ from ground state to 2393 2529 2541 levels										89Hu03 **
$^{123}\text{In}(\beta^-)^{123}\text{Sn}$	$Q_{\beta^-}=4410(31)$ ; and 4645(72) from $^{123}\text{In}^m$ at 327.21 keV										Nub16b **
$^{123}\text{Sn}(\beta^-)^{123}\text{Sb}$	$E_{\beta^-}=1260(10)$ from $^{123}\text{Sn}^m$ at 24.6 to $5/2^+$ level at 160.33 keV										Ens04a **
$^{123}\text{Sb}(\beta^-)^{123}\text{Sb}$	$E_{\beta^-}=310(20)$ to $9/2^+$ level at 1088.64 keV										Ens04a **
$^{123}\text{Xe}(\beta^+)^{123}\text{I}$	$E_{\beta^+}=1505(15)$ to $1/2^+$ level at 148.92 keV										Ens04a **
$^{123}\text{Cs}(\beta^+)^{123}\text{Xe}$	$E_{\beta^+}=2990(310)$ to $3/2^+$ level at 97.30 keV										Ens04a **
$^{123}\text{Cs}(\beta^+)^{123}\text{Xe}$	$E_{\beta^+}=2370(140)$ to $(1/2,3/2)^+$ level at 596.65 keV, and other $E_{\beta^+}$										Ens04a **
$^{123}\text{Cs}(\beta^+)^{123}\text{Xe}$	$E_{\beta^+}=2930(180)$ to $3/2^+$ level at 97.30 keV										Ens04a **
$^{123}\text{Cs}^x(\text{IT})^{123}\text{Cs}$	Based on $^{123}\text{Cs}^m(\text{IT})=156.27$ and isomeric ratio $R<0.1$										Nub16b **
$^{124}\text{Pd}-u$	-64617	399	-62680#	320#	1.9	D			GT3	2.5	16Kn03 *
$^{124}\text{Ag}-^{133}\text{Cs}_{932}$	17050	270				2			MA8	1.0	10Br02 *
$^{124}\text{Cd}-^{133}\text{Cs}_{932}$	5781	10	5776	3	-0.5	1	10	10 $^{124}\text{Cd}$	MA8	1.0	10Br02
$\text{C}_7 \text{ }^{13}\text{C} \text{ H}_{13} \text{ N}-^{124}\text{Sn}$	202886	8	202877.6	1.1	-0.4	U			M16	2.5	63Da10
	202891	8			-1.1	U			R13	1.5	83De51
$^{124}\text{Sn}-^{13}\text{C} \text{ }^{37}\text{Cl}_3$	4210.47	0.71	4214.1	1.1	2.0	1	38	37 $^{124}\text{Sn}$	H39	2.5	84Ha20
$^{124}\text{Sn}-^{133}\text{Cs}_{932}$	-6598	21	-6604.5	1.1	-0.3	U			MA8	1.0	05Si34
$^{124}\text{Te}-^{13}\text{C} \text{ }^{37}\text{Cl}_3$	1754.63	1.26	1754.5	1.6	0.0	1	26	26 $^{124}\text{Te}$	H39	2.5	84Ha20
$^{124}\text{Te}-^{54}\text{Fe} \text{ }^{35}\text{Cl}_2$	25501.65	2.56	25503.4	1.7	0.3	U			H39	2.5	84Ha20
$\text{C}_7 \text{ }^{13}\text{C} \text{ H}_{13} \text{ N}-^{124}\text{Te}$	205336	13	205337.2	1.6	0.0	U			M16	2.5	63Da10
	205325	8			1.0	U			R13	1.5	83De51
$^{124}\text{I}-u$	-93786	30	-93791.0	2.6	-0.2	U			GS2	1.0	05Li24
$^{124}\text{Xe}-^{13}\text{C} \text{ }^{37}\text{Cl}_3$	4831.15	1.58	4829.0	1.9	-0.5	1	24	24 $^{124}\text{Xe}$	H39	2.5	84Ha20
$^{124}\text{Xe}-^{54}\text{Fe} \text{ }^{35}\text{Cl}_2$	28575.78	0.99	28577.9	1.9	0.9	1	60	59 $^{124}\text{Xe}$	H39	2.5	84Ha20
$^{124}\text{Xe}-^{133}\text{Cs}_{932}$	-5986	13	-5989.6	1.9	-0.3	U			MA6	1.0	04Di18
$^{124}\text{Cs}-^{133}\text{Cs}_{932}$	361	16	377	9	1.0	o			MA1	1.0	90St25
	370	13			0.5	R			MA1	1.0	99Am05
	361	15			1.0	R			MA8	1.0	05Gu37
$^{124}\text{Cs}-u$	-87696	30	-87742	9	-1.5	2			GS2	1.0	05Li24
	-87693	30			-1.6	2			GS2	1.0	05Li24 *
$^{124}\text{Ba}-^{133}\text{Cs}_{932}$	3212	15	3212	13	0.0	2			MA1	1.0	99Am05
$^{124}\text{Ba}-u$	-84905	30	-84906	13	0.0	R			GS2	1.0	05Li24
$^{124}\text{La}-u$	-75464	71	-75430	60	0.5	2			GS2	1.0	05Li24 *
$^{124}\text{Cd}-^{130}\text{Xe}_{954}$	9708.9	3.4	9709	3	0.2	1	89	89 $^{124}\text{Cd}$	JY1	1.0	12Ha25

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{124}\text{Sn}-^{129}\text{Xe}_{.961}$	-3214.3	2.1	-3217.7	1.1	-1.6	1	27	27 $^{124}\text{Sn}$	JY1	1.0	11Ha48
$^{124}\text{Sn}-^{120}\text{Sn}_{1.033}$	6305.1	2.1	6302.2	1.3	-1.4	1	36	20 $^{124}\text{Sn}$	JY1	1.0	11Ha48
$^{124}\text{Sn } ^{35}\text{Cl}-^{122}\text{Sn } ^{37}\text{Cl}$	4784	2	4782.8	2.6	-0.1	U			H15	4.0	62Ba23
$^{124}\text{Te } ^{35}\text{Cl}-^{122}\text{Te } ^{37}\text{Cl}$	2728	2	2723.74	0.15	-0.5	U			H16	4.0	63Ba47
$^{124}\text{Sn}-^{124}\text{Te}$	2458.51	0.89	2459.6	1.6	0.5	1	53	41 $^{124}\text{Te}$	H39	2.5	84Ha20
$^{124}\text{Xe}-^{124}\text{Te}$	3076.00	1.78	3074.5	2.3	-0.3	1	27	16 $^{124}\text{Xe}$	H39	2.5	84Ha20
$^{124}\text{Sn}-^{122}\text{Sn}$	1838	22	1832.7	2.6	-0.1	U			M16	2.5	63Da10
$^{120}\text{Cs}^x-^{124}\text{Cs}^x_{194} \quad ^{119}\text{Cs}^x_{807}$	310	30	304	12	-0.1	U			P22	2.5	82Au01
$^{121}\text{Cs}^x-^{124}\text{Cs}^x_{244} \quad ^{120}\text{Cs}^x_{756}$	-1360	30	-1265	19	1.3	U			P22	2.5	82Au01
$^{123}\text{Cs}^x-^{124}\text{Cs}^x_{744} \quad ^{120}\text{Cs}^x_{256}$	-1390	30	-1337	21	0.7	U			P22	2.5	82Au01
$^{124}\text{Sn}(d, ^6\text{Li})^{120}\text{Cd}$	-5216	24	-5228	4	-0.5	U					79Ja21
$^{124}\text{Sn}(^3\text{He}, ^7\text{Be})^{120}\text{Cd}$	-5098	30	-5115	4	-0.6	U			MSU		76St11
$^{124}\text{Sn}(^{18}\text{O}, ^{20}\text{Ne})^{122}\text{Cd}$	-1266	39	-1362.7	2.5	-2.5	U					97Gu32 *
$^{122}\text{Sn}(t,p)^{124}\text{Sn}$	5931	15	5953.7	2.4	1.5	U			Roc		70FI05
$^{124}\text{Sn}(p,t)^{122}\text{Sn}$	-5956	10	-5953.7	2.4	0.2	U					64Al29
$^{124}\text{Sn}(d, ^3\text{He})^{123}\text{In}$	-6610	50	-6599	20	0.2	-			Sac		69Co03
	-6572	66			-0.4	-			MSU		71We01
ave.	-6600	40			-0.1	1	25	25 $^{123}\text{In}$			average
$^{124}\text{Sn}(p,d)^{123}\text{Sn}$	-6279	15	-6264.8	2.4	0.9	U			Har		70Ca01
$^{124}\text{Sn}(d,t)^{123}\text{Sn}$	-2260	35	-2232.1	2.4	0.8	U			Pit		64Co11
	-2233.4	3.7			0.4	1	42	38 $^{123}\text{Sn}$	SPa		75Be09
$^{123}\text{Sb}(n,\gamma)^{124}\text{Sb}$	6467.55	0.10	6467.50	0.06	-0.5	2					73Sh.A Z
	6467.40	0.10			1.0	2					81Su.A Z
	6467.58	0.14			-0.6	2			Bdn		06Fi.A
$^{123}\text{Te}(n,\gamma)^{124}\text{Te}$	9425	2	9424.48	0.09	-0.3	U					69Bu05
	9423.7	1.5			0.5	U					70Or.A
	9424.05	0.30			1.4	-			Ltn		95Ge06 Z
	9423.89	0.20			3.0	C			Bdn		06Fi.A
	9424.53	0.10			-0.5	-					06Vo09
ave.	9424.48	0.09			0.0	1	100	94 $^{123}\text{Te}$			average
$^{124}\text{Cd}(\beta^-)^{124}\text{In}$	4166	39	4170	30	0.1	1	61	61 $^{124}\text{In}$	Stu		87Sp09
$^{124}\text{In}(\beta^-)^{124}\text{Sn}$	7180	50	7360	30	3.7	B			Stu		78Al18
	7360	49			0.1	1	39	39 $^{124}\text{In}$	Stu		87Sp09
$^{124}\text{Sn}(t, ^3\text{He})^{124}\text{In}$	-7590	50	-7350	30	4.9	B			LAI		78Aj01
$^{124}\text{In}^m(\beta^-)^{124}\text{Sn}$	7370	210	7340	50	-0.1	o			Stu		78Al18
	7341	51				2			Stu		87Sp09
$^{124}\text{Sb}(\beta^-)^{124}\text{Te}$	2907.7	5.	2905.07	0.13	-0.5	U					65Hs02 *
	2903.7	4.			0.3	U					66Ca10 *
	2904.7	2.			0.2	U					69Na05 *
$^{124}\text{I}(\beta^+)^{124}\text{Te}$	3157	4	3159.6	1.9	0.6	2					71Bo01 *
	3160.3	2.1			-0.3	2					92Wo03
$^{124}\text{Cs}(\beta^+)^{124}\text{Xe}$	5920	460	5930	8	0.0	U					75We23
	5900	90			0.3	U			IRS		93Al03
	5910	30			0.7	U			JAE		96Os04
$^{124}\text{Cs}^x(\text{IT})^{124}\text{Cs}$	30	20				3					AHW *
$^{124}\text{La}(\beta^+)^{124}\text{Ba}$	8930	110	8830	60	-0.9	R			JAE		98Ko66
* $^{124}\text{Pd}-u$	Trends from Mass Surface TMS suggest $^{124}\text{Pd}$ 1800 less bound										WgM168**
* $^{124}\text{Ag}-^{133}\text{Cs}_{.932}$	$D_M=17050(270) \mu u$ for mixture gs+m at 0#(100#); $M-A=-66200(250) \text{ keV}$										Nub16b **
* $^{124}\text{Cs}-u$	$M-A=-81223(28) \text{ keV}$ for $^{124}\text{Cs}^m$ at 462.63 keV										Nub16b **
* $^{124}\text{La}-u$	$M-A=-70244(32) \text{ keV}$ for mixture gs+m at 100#100 keV										Nub16b **
* $^{124}\text{Sn}(^{18}\text{O}, ^{20}\text{Ne})^{122}\text{Cd}$	Original $Q=-1250(39)$ calibrated with $^{120}\text{Cd}=-83973(19) \text{ keV}$										GAu **
* $^{124}\text{Sb}(\beta^-)^{124}\text{Te}$	$E_{\beta^-}=2305(5) \quad 2301(4) \quad 2302(2)$ respectively, to $2^+$ level at 602.7271 keV										Ens087 **
* $^{124}\text{I}(\beta^+)^{124}\text{Te}$	Original error increased, see $^{84}\text{Rb}(\beta^+)$										AHW **
* $^{124}\text{Cs}^x(\text{IT})^{124}\text{Cs}$	Based on $^{124}\text{Cs}^m(\text{IT})=462.63 \text{ keV}$										Nub16b **
* $^{124}\text{Cs}^x(\text{IT})^{124}\text{Cs}$	Isomeric ratio assumed $<0.1$ as for $^{118}\text{Cs}, ^{120}\text{Cs}, ^{122}\text{Cs}$										AHW **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{125}\text{Ag}-u$	-68954	429	-69270	470	-0.5	o			GT1	1.5	04Ma.A
	-69265	186				2			GT3	2.5	16Kn03
$^{125}\text{Cd}-u$	-78770	120	-78742	3	0.1	o			GT2	2.5	08Kn.A *
	-78780	140			0.1	U			GT2	2.5	08Su19 *
$\text{C}_7 \text{H}_6 \text{ } ^{35}\text{Cl}-^{125}\text{Te}$	111363	6	111373.0	1.6	0.7	U			M16	2.5	63Da10
	111368	8			0.4	U			R13	1.5	83De51
$^{125}\text{I}-u$	-95374	30	-95370.7	1.6	0.1	U			GS2	1.0	05Li24
$^{125}\text{Cs}-u$	-90280	30	-90272	8	0.3	U			GS2	1.0	05Li24
$^{125}\text{Ba}-u$	-85569	30	-85528	12	1.4	R			GS2	1.0	05Li24
$^{125}\text{La}-u$	-79191	30	-79184	28	0.2	1	87	87 $^{125}\text{La}$	GS2	1.0	05Li24
$^{125}\text{Cd}-^{133}\text{Cs}_{,940}$	10133	13	10133	3	0.0	U			TT1	1.0	16La.A
$^{125}\text{Cd}^m-^{133}\text{Cs}_{,940}$	10334	13	10333	3	-0.1	U			TT1	1.0	16La.A
$^{125}\text{Cs}-^{133}\text{Cs}_{,940}$	-1392	17	-1397	8	-0.3	o			MA1	1.0	90St25
	-1382	14			-1.1	-			MA1	1.0	99Am05
	-1386	14			-0.8	-			MA4	1.0	99Am05
ave.	-1384	10			-1.3	1	71	71 $^{125}\text{Cs}$			average
$^{125}\text{Ba}-^{133}\text{Cs}_{,940}$	3356	13	3347	12	-0.7	-			MA5	1.0	00Be42
	ave.	3348	12		-0.1	1	98	98 $^{125}\text{Ba}$			average
$^{125}\text{Cd}-^{130}\text{Xe}_{,962}$	14081.6	3.1	14082	3	0.0	1	100	100 $^{125}\text{Cd}$	JY1	1.0	12Ha25
$^{125}\text{Cd}^m-^{130}\text{Xe}_{,962}$	14281.6	3.4				2			JY1	1.0	13Ka08
$^{125}\text{Te } ^{35}\text{Cl}-^{123}\text{Te } ^{37}\text{Cl}$	3090	2	3110.26	0.13	2.5	U			H16	4.0	63Ba47
$^{122}\text{Cs}^x-^{125}\text{Cs}_{,244} \text{ } ^{121}\text{Cs}^x_{,756}$	715	23	640	40	-1.3	U			P32	2.5	86Au02
$^{123}\text{Sb}(t,p)^{125}\text{Sb}$	6696	20	6693.0	2.1	-0.1	U			Ald		67Hi01
$^{124}\text{Sn}(n,\gamma)^{125}\text{Sn}$	5733.1	1.5	5733.50	0.20	0.3	U					77Ca09 Z
	5733.1	0.6			0.7	U					81Ba53
	5733.5	0.2			0.0	1	100	100 $^{125}\text{Sn}$			11To04
$^{124}\text{Sn}(d,p)^{125}\text{Sn}$	3530	30	3508.93	0.20	-0.7	U			Pit		64Co11
	3506	12			0.2	U			Tal		64Ne10
	3515	11			-0.6	U					72Ca33
	3509.4	3.6			-0.1	U			SPa		75Be09
$^{124}\text{Te}(n,\gamma)^{125}\text{Te}$	6569.0	1.0	6568.970	0.030	0.0	U					71Gr.A
	6568.97	0.03			0.0	1	100	83 $^{125}\text{Te}$	Pm		99Ho01
	6569.39	0.19			-2.2	U			Bdn		06Fi.A
$^{125}\text{Te}(\gamma,n)^{124}\text{Te}$	-6560	60	-6568.970	0.030	-0.1	U			Phi		60Ge01
$^{124}\text{Te}(d,p)^{125}\text{Te}$	4344	8	4344.40	0.03	0.1	U			MIT		69Gr24
$^{124}\text{Te}(^3\text{He},d)^{125}\text{I}$	115.1	3.0	107.38	0.07	-2.6	U			Hei		78Sz04
$^{124}\text{Te}(\alpha,t)^{125}\text{I}$	-14203	7	-14213.01	0.07	-1.4	U			Hei		78Sz04
$^{124}\text{Xe}(n,\gamma)^{125}\text{Xe}$	7603.3	0.4	7603.3	0.4	-0.1	1	100	99 $^{125}\text{Xe}$			82Ka.A
$^{125}\text{Cd}(\beta^-)^{125}\text{In}$	7122	62	7129	27	0.1	1	19	19 $^{125}\text{In}$	Stu		87Sp09 *
$^{125}\text{Cd}^m(\beta^-)^{125}\text{In}$	7172	35	7315	27	4.1	B			Stu		87Sp09 *
$^{125}\text{In}(\beta^-)^{125}\text{Sn}$	5418	30	5420	27	0.1	1	81	81 $^{125}\text{In}$	Stu		87Sp09 *
	2330	10	2359.9	2.6	3.0	B					50Ha58
$^{125}\text{Sn}(\beta^-)^{125}\text{Sb}$	2370	20			-0.5	U					50Ke11
	2335	40			0.6	U					64De02 *
	767.7	3.	766.7	2.1	-0.3	2					64Ma30 *
$^{125}\text{Sb}(\beta^-)^{125}\text{Te}$	765.7	3.			0.3	2					66Ma49 *
	184	7	185.77	0.06	0.3	U					64Le05 *
$^{125}\text{I}(\epsilon)^{125}\text{Te}$	185	8			0.1	U					66Sm05 *
	177.2	2.			4.3	C					68Go.A *
	186.1	0.3			-1.1	U					86Bo46
	179.3	2.0			3.2	B					90Li14 *
	185.77	0.06				2					94Hi04
	1735	40	1643.8	2.2	-2.3	U					69Lu09 *
$^{125}\text{Xe}(\epsilon)^{125}\text{I}$	3072	20	3105	8	1.7	-					54Ma54
	3082	20			1.2	-					75We23
	3100	100			0.1	U			IRS		93AI03
	ave.	3077	14		2.0	1	31	29 $^{125}\text{Cs}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{125}\text{Ba}(\beta^+)^{125}\text{Cs}$	4560	250	4419	13	-0.6	U					68Da09 *
	4380	50			0.8	U			JAE		96Os04
$^{125}\text{La}(\beta^+)^{125}\text{Ba}$	5950	70	5909	28	-0.6	1	16	13 $^{125}\text{La}$	JAE		98Ko66
* $^{125}\text{Cd}-\text{u}$	$M-A=-73274(93)$ keV for mixture gs+m at 186(4) keV										
* $^{125}\text{Cd}-\text{u}$	$M-A=-73287(120)$ keV for mixture gs+m at 186(4) keV										
* $^{125}\text{Cd}(\beta^-)^{125}\text{In}$	$E_{\beta^-}=4625(62)$ to $(1/2^+, 3/2^+)$ level at 2497.43 keV										
* $^{125}\text{Cd}^m(\beta^-)^{125}\text{In}$	$E_{\beta^-}=5009(109), 4581(126), 4533(39)$ to 2101.40, 2640.29, 2641.26 levels										
* $^{125}\text{In}(\beta^-)^{125}\text{Sn}$	$Q_{\beta^-}=5443(31);$ and 5730(43) from $^{125}\text{In}^m$ at 360.12 keV										
* $^{125}\text{Sn}(\beta^-)^{125}\text{Sb}$	$E_{\beta^-}=2030(40)$ from $^{125}\text{Sn}^m$ at 27.50(0.14) to $5/2^+$ level at 332.06 keV										
* $^{125}\text{Sb}(\beta^-)^{125}\text{Te}$	$E_{\beta^-}=623(3)$ 621(3) respectively, to $1/2^-$ level at 144.775 keV										
* $^{125}\text{I}(\epsilon)^{125}\text{Te}$	LMK=0.254(0.003) 0.253(0.005) IBE=110(2) 150.6(0.3) respectively, all to $3/2^+$ level at 35.4925 keV. $Q(\text{LMK})$ recalculated, error mainly theory										
* $^{125}\text{I}(\epsilon)^{125}\text{Te}$	IBE=112.0(2.0)(1s)+31.8 to $3/2^+$ level at 35.4925 keV										
* $^{125}\text{Xe}(\epsilon)^{125}\text{I}$	$E_{\beta^+}=470(40)$ to $3/2^+$ at 188.416 and $1/2^+$ at 243.382 keV, ratio 1:2										
* $^{125}\text{Ba}(\beta^+)^{125}\text{Cs}$	$E_{\beta^+}=3450(250)$ to $(5/2^+)$ level at 84.82 level										
$^{126}\text{Ag}-\text{u}$	-65926	329	-65140#	220#	1.0	D			GT3	2.5	16Kn03 *
$\text{C}_{10}\text{H}_6-^{126}\text{Te}$	143623	9	143639.3	1.6	0.7	U			M16	2.5	63Da10
	143640	8			-0.1	U			R13	1.5	83De51
$^{126}\text{Xe}-\text{u}$	-95647	30	-95703	4	-1.9	U			GS2	1.0	05Li24
$^{126}\text{Ba}-\text{u}$	-88745	30	-88750	13	-0.2	R			GS2	1.0	05Li24
$^{126}\text{La}-\text{u}$	-80503	232	-80490	100	0.1	2			GS2	1.0	05Li24 *
$^{126}\text{Ce}-\text{u}$	-76029	30				2			GS2	1.0	05Li24
$^{126}\text{Xe}-^{134}\text{Xe}_{940}$	-6772.8	2.9	-6773	4	0.0	o			MA8	1.0	05He.A
	-6773.2	3.8			0.1	1	98	98 $^{126}\text{Xe}$	MA8	1.0	06He29
$^{126}\text{Cd}-^{133}\text{Cs}_{947}$	11966.5	4.5	11966.1	2.7	-0.1	1	35	35 $^{126}\text{Cd}$	MA8	1.0	10Br02
	11956	15			0.7	U			MA8	1.0	10Br02
	11958	13			0.6	U			TT1	1.0	16La.A
$^{126}\text{Cs}-^{133}\text{Cs}_{947}$	-1027	17	-1017	11	0.6	o			MA1	1.0	90St25
	-1011	13			-0.5	1	74	74 $^{126}\text{Cs}$	MA1	1.0	99Am05
$^{126}\text{Ba}-^{133}\text{Cs}_{947}$	786	15	787	13	0.1	2			MA1	1.0	99Am05
$^{126}\text{Cd}-^{130}\text{Xe}_{969}$	15928.6	3.3	15928.6	2.7	0.0	1	65	65 $^{126}\text{Cd}$	JY1	1.0	12Ha25
$^{126}\text{Te}-^{35}\text{Cl}-^{124}\text{Te}-^{37}\text{Cl}$	3432	2	3443.91	0.11	1.5	U			H16	4.0	63Ba47
	3441.28	1.54			1.1	U			H43	1.5	90Dy04
$^{123}\text{Cs}^x-^{126}\text{Cs}_{390}$	$^{121}\text{Cs}^x_{610}$	-1160	30	-1135	17	0.3	U		P22	2.5	82Au01
$^{124}\text{Cs}^x-^{126}\text{Cs}_{590}$	$^{121}\text{Cs}^x_{410}$	-340	30	-341	23	0.0	U		P22	2.5	82Au01
$^{124}\text{Cs}^x-^{126}\text{Cs}_{492}$	$^{122}\text{Cs}^x_{508}$	-570	30	-510	28	0.8	U		P22	2.5	82Au01
$^{124}\text{Cs}^x-^{126}\text{Cs}_{328}$	$^{123}\text{Cs}^x_{672}$	390	30	422	24	0.4	U		P22	2.5	82Au01
$^{125}\text{Cs}-^{126}\text{Cs}_{496}$	$^{124}\text{Cs}^x_{504}$	-1130	30	-1072	14	0.8	U		P22	2.5	82Au01
$^{124}\text{Sn}(\text{t,p})^{126}\text{Sn}$	5445	15	5442	10	-0.2	-			Ald		69Bj01
	5444	15			-0.1	-			Roc		70Fl05
	ave.	11			-0.2	1	96	96 $^{126}\text{Sn}$			average
$^{125}\text{Te}(\text{n},\gamma)^{126}\text{Te}$	9113.7	0.4	9113.69	0.08	0.0	U					77Ko.A
	9113.69	0.08			0.0	1	100	83 $^{126}\text{Te}$			03Vo03
$^{126}\text{Te}(\gamma,\text{n})^{125}\text{Te}$	-8840	120	-9113.69	0.08	-2.3	U			Phi		60Ge01
$^{125}\text{Te}(\text{d,p})^{126}\text{Te}$	6892	6	6889.13	0.08	-0.5	U			MIT		71Gr01
$^{126}\text{Cd}(\beta^-)^{126}\text{In}$	5486	36	5516	27	0.8	1	56	56 $^{126}\text{In}$	Stu		87Sp09
$^{126}\text{In}(\beta^-)^{126}\text{Sn}$	8207	39	8242	27	0.9	1	48	44 $^{126}\text{In}$	Stu		87Sp09
$^{126}\text{In}^m(\beta^-)^{126}\text{Sn}$	8309	51				2			Stu		87Sp09
$^{126}\text{Sn}(\beta^-)^{126}\text{Sb}$	378	30				2					71Or04 *
$^{126}\text{Sb}(\beta^-)^{126}\text{Te}$	3667	150	3670	30	0.0	U					71Or04 *
$^{126}\text{I}(\beta^+)^{126}\text{Te}$	2151	5	2154	4	0.6	1	54	52 $^{126}\text{I}$			59Ha27
$^{126}\text{I}(\beta^-)^{126}\text{Xe}$	1258	5	1236	5	-4.5	B					55Ko14 *
$^{126}\text{Cs}(\beta^+)^{126}\text{Xe}$	4670	140	4796	11	0.9	U					75We23 *
	4810	100			-0.1	U					76Pa11 *
	4830	40			-0.8	U			JAE		92Os07
	4730	100			0.7	U			IRS		93Al03
	4780	20			0.8	1	28	26 $^{126}\text{Cs}$	JAE		96Os04



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{126}\text{La}(\beta^+)^{126}\text{Ba}$	7700	100	7700	90	0.0	R			JAE		98Ko66
$^{126}\text{La}^m(\beta^+)^{126}\text{Ba}$	7910	400				3			JAE		98Ko66
* $^{126}\text{Ag}-u$	Trends from Mass Surface TMS suggest $^{126}\text{Ag}$ 730 less bound										
* $^{126}\text{La}-u$	$M-A=-74883(28)$ keV for mixture gs+m at 210(410) keV										
* $^{126}\text{Sn}(\beta^-)^{126}\text{Sb}$	$E_{\beta^-}=250(30)$ to $2^+$ level at 127.9 keV										
* $^{126}\text{Sb}(\beta^-)^{126}\text{Te}$	$E_{\beta^-}=1900(150)$ from mixture ground state and $^{126}\text{Sb}^m$ at 17.7 to $6^+$ level at 1776.19 keV										
* $^{126}\text{I}(\beta^-)^{126}\text{Xe}$	$E_{\beta^-}=865(5)$ to $2^+$ level at 388.631 keV, and other $E_{\beta^-}$										
* $^{126}\text{Cs}(\beta^+)^{126}\text{Xe}$	$E_{\beta^+}=3260(140)$ 3400(100) respectively, to $2^+$ level at 388.631 keV										
$\text{C}_{10} \text{H}_7-^{127}\text{I}$	150297	6	150303	4	0.4	U			M16	2.5	63Da10
	150305.3	3.4			-0.2	1	21	21 $^{127}\text{I}$	M16	2.5	63Da10
	150322	8			-1.6	U			R13	1.5	83De51
$^{127}\text{Cs}-u$	-92571	30	-92583	6	-0.4	U			GS2	1.0	05Li24
$^{127}\text{Ba}-u$	-88923	39	-88909	12	0.4	R			GS2	1.0	05Li24 *
$^{127}\text{La}-u$	-83640	30	-83625	28	0.5	1	87	87 $^{127}\text{La}$	GS2	1.0	05Li24 *
$^{127}\text{Ce}-u$	-77273	31				2			GS2	1.0	05Li24 *
$^{127}\text{Cd}-^{133}\text{Cs}_{955}$	16490	13				2			TT1	1.0	16La.A
$^{127}\text{Cd}^m-^{133}\text{Cs}_{955}$	16799	11	16786	9	-1.2	1	61	61 $^{127}\text{Cd}^m$	TT1	1.0	16La.A
$^{127}\text{Sn}^{34}\text{S}-^{133}\text{Cs}_{1,211}$	-7237	12	-7245	11	-0.7	1	81	81 $^{127}\text{Sn}$	MA8	1.0	08Dw01 *
$^{127}\text{Cs}-^{133}\text{Cs}_{955}$	-2303	17	-2289	6	0.8	o			MA1	1.0	90St25
	-2287	13			-0.2	-			MA1	1.0	99Am05
	-2293.3	7.7			0.5	-			MA8	1.0	05Gu37
ave.	-2292	7			0.4	1	82	82 $^{127}\text{Cs}$			average
$^{127}\text{Ba}-^{133}\text{Cs}_{955}$	1389	13	1385	12	-0.3	-			MA5	1.0	00Be42
ave.	1387	12			-0.2	1	98	98 $^{127}\text{Ba}$			average
$^{127}\text{Cd}^m-^{130}\text{Xe}_{977}$	20741	14	20764	9	1.6	1	38	38 $^{127}\text{Cd}^m$	JY1	1.0	12Ha25 *
$^{125}\text{Cs}-^{127}\text{Cs}_{591}$ $^{122}\text{Cs}_{410}^x$	-1098	18	-1086	16	0.3	U			P32	2.5	86Au02
$^{126}\text{Te}(n,\gamma)^{127}\text{Te}$	6289	3	6287.65	0.18	-0.5	U					72Mu.A
	6287.8	0.4			-0.4	-			Bdn		06Fi.A
	6287.6	0.2			0.2	-			Prn		05Ho15
$^{126}\text{Te}(d,p)^{127}\text{Te}$	4044	8	4063.08	0.18	2.4	U			MIT		68Gr16
$^{126}\text{Te}(n,\gamma)^{127}\text{Te}$	ave. 6287.64	0.18	6287.65	0.18	0.0	1	100	98 $^{127}\text{Te}$			average
$^{127}\text{I}(\gamma,n)^{126}\text{I}$	-9135	22	-9143.9	2.7	-0.4	U			Phi		60Ge01
	-9145	3			0.4	1	83	48 $^{126}\text{I}$	MMn		86Ts04
$^{127}\text{Cd}^m(\beta^-)^{127}\text{In}$	8468	63	8425	22	-0.7	1	13	11 $^{127}\text{In}$	Stu		87Sp09 *
$^{127}\text{In}(\beta^-)^{127}\text{Sn}$	6514	31	6575	19	2.0	o			Stu		87Sp09 *
	6579	20			-0.2	1	91	89 $^{127}\text{In}$	Stu		04Ga24 *
$^{127}\text{In}^n(\beta^-)^{127}\text{Sn}$	8442	56				2			Stu		04Ga24
$^{127}\text{Sn}(\beta^-)^{127}\text{Sb}$	3201	24	3229	11	1.2	1	21	17 $^{127}\text{Sn}$	Stu		77Lu06 *
$^{127}\text{Sb}(\beta^-)^{127}\text{Te}$	1581	5	1582	5	0.2	1	97	96 $^{127}\text{Sb}$			67Ra13 *
$^{127}\text{Te}(\beta^-)^{127}\text{I}$	683	10	702	4	1.9	-					55Da37
	695	10			0.7	-					56Kn20
ave.	689	7			1.9	1	26	24 $^{127}\text{I}$			average
$^{127}\text{Xe}(\epsilon)^{127}\text{I}$	663.3	2.2	662.3	2.0	-0.4	-					68Sc14
$^{127}\text{I}(\beta^-\text{He,t})^{127}\text{Xe}$	-676	6	-680.9	2.0	-0.8	-			Pri		89Ch01
$^{127}\text{Xe}(\epsilon)^{127}\text{I}$	ave. 662.6	2.1	662.3	2.0	-0.1	1	98	91 $^{127}\text{Xe}$			average
$^{127}\text{Cs}(\beta^+)^{127}\text{Xe}$	2115	25	2081	6	-1.3	-					54Ma54 *
	2076	20			0.3	-					67Sp08 *
	2089	20			-0.4	-					75We23 *
ave.	2090	12			-0.7	1	27	18 $^{127}\text{Cs}$			average
$^{127}\text{Ba}(\beta^+)^{127}\text{Cs}$	3450	100	3422	13	-0.3	U					76Be11 *
$^{127}\text{La}(\beta^+)^{127}\text{Ba}$	5010	70	4922	28	-1.3	1	16	13 $^{127}\text{La}$	JAE		98Ko66
* $^{127}\text{Ba}-u$	$M-A=-82791(28)$ keV for mixture gs+m at 80.32 keV										
* $^{127}\text{La}-u$	$M-A=-77903(28)$ keV for mixture gs+m at 14.2 keV										
* $^{127}\text{Ce}-u$	$M-A=-71976(29)$ keV for mixture gs+m at 7.3 keV										
* $^{127}\text{Sn}^{34}\text{S}-^{133}\text{Cs}_{1,211}$	$D_M=-7234.3(11.6)$ $\mu\text{u}$ for mixture gs+m at 5.07(0.06) keV										
* $^{127}\text{Cd}^m-^{130}\text{Xe}_{977}$	Re-assigned to $^{127}\text{Cd}^m$ by the evaluator										
* $^{127}\text{Cd}^m(\beta^-)^{127}\text{In}$	Also $E_{\beta^-}=7910(200)$ to $^{127}\text{In}^m$ at 408.9 keV										
* $^{127}\text{Cd}^m(\beta^-)^{127}\text{In}$	Re-assigned to $^{127}\text{Cd}^m$ by the evaluator										
* $^{127}\text{In}(\beta^-)^{127}\text{Sn}$	Also $E_{\beta^-}=6976(64)$ from $^{127}\text{In}^m$ at 408.9 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference		
* <sup>127</sup> In( $\beta^-$ ) <sup>127</sup> Sn	Also $E_{\beta^-}$ =6999(63) from <sup>127</sup> In <sup>m</sup> at 408.9 keV								Nub16b **		
* <sup>127</sup> Sn( $\beta^-$ ) <sup>127</sup> Sb	$Q_{\beta^-}$ =3206(24) from <sup>127</sup> Sn <sup>m</sup> at 5.07 keV								Nub16b **		
* <sup>127</sup> Sb( $\beta^-$ ) <sup>127</sup> Te	$E_{\beta^-}$ =1493(5) to 11/2 <sup>-</sup> level at 88.23 keV, and other $E_{\beta^-}$								Ens118 **		
* <sup>127</sup> Cs( $\beta^+$ ) <sup>127</sup> Xe	$E_{\beta^+}$ =1063(10), 685(25) to mixture gs+124.751 and 1/2 <sup>+</sup> level at 411.965 keV								Ens118 **		
* <sup>127</sup> Cs( $\beta^+$ ) <sup>127</sup> Xe	$E_{\beta^+}$ =1068(20), 910(30), 650(30) to ground state, 3/2 <sup>+</sup> at 124.751, 1/2 <sup>+</sup> at 411.965								Ens118 **		
* <sup>127</sup> Cs( $\beta^+$ ) <sup>127</sup> Xe	$E_{\beta^+}$ =1040(20), 650(20) to mixture gs+124.751 and 1/2 <sup>+</sup> level at 411.965 keV								Ens118 **		
* <sup>127</sup> Ba( $\beta^+$ ) <sup>127</sup> Cs	$E_{\beta^+}$ =2230(100) to 3/2 <sup>+</sup> level at 180.92 keV, and other $E_{\beta^+}$								Ens118 **		
<sup>128</sup> Sn-u	-89512	25	-89493	19	0.8	1	58	58 <sup>128</sup> Sn	GS3	1.0	12Ch19
<sup>C10</sup> H <sub>8</sub> - <sup>128</sup> Te	158112	9	158138.9	0.9	1.2	U			M16	2.5	63Da10
	158141.2	7.			-0.1	U			C3	2.5	70Ke05
	158151	8			-1.0	U			R13	1.5	83De51
<sup>C10</sup> H <sub>8</sub> - <sup>128</sup> Xe	159068.2	4.2	159069.3	1.1	0.1	U			M16	2.5	63Da10
	159069.7	0.7			-0.3	1	42	42 <sup>128</sup> Xe	C3	2.5	70Ke05
<sup>128</sup> Cs-u	-92181	30	-92251	6	-2.3	U			GS2	1.0	05Li24
<sup>128</sup> Ba-u	-91663	30	-91658	6	0.2	R			GS2	1.0	05Li24
<sup>128</sup> La-u	-84436	69	-84410	60	0.4	2			GS2	1.0	05Li24 *
<sup>128</sup> Ce-u	-81089	30				2			GS2	1.0	05Li24
<sup>128</sup> Pr-u	-71209	32				2			GS2	1.0	05Li24
<sup>128</sup> Cd- <sup>133</sup> Cs <sub>962</sub>	18759	11	18768	8	0.8	1	50	50 <sup>128</sup> Cd	MA8	1.0	10Br02
<sup>128</sup> Sn <sup>34</sup> S- <sup>133</sup> Cs <sub>1,218</sub>	-6396	14	-6466	19	-5.0	F			MA8	1.0	08Dw01 *
<sup>128</sup> Cs- <sup>133</sup> Cs <sub>962</sub>	-1306	17	-1296	6	0.6	o			MA1	1.0	90St25
<sup>128</sup> Ba- <sup>133</sup> Cs <sub>962</sub>	-1293	13			-0.2	1	20	20 <sup>128</sup> Cs	MA1	1.0	99Am05
	-720	13	-702	6	1.4	-			MA1	1.0	99Am05
ave.	-718	12			1.3	1	22	22 <sup>128</sup> Ba	average		
<sup>128</sup> Cd- <sup>130</sup> Xe <sub>985</sub>	22865	11	22856	8	-0.8	1	50	50 <sup>128</sup> Cd	JY1	1.0	12Ha25
<sup>128</sup> Te <sup>35</sup> Cl- <sup>126</sup> Te <sup>37</sup> Cl	4106	2	4100.6	1.8	-0.7	U			H16	4.0	63Ba47
	4102.3	1.8			-0.4	1	16	12 <sup>126</sup> Te	C3	2.5	70Ke05
<sup>128</sup> Te- <sup>128</sup> Xe	931.26	1.20	930.3	1.0	-0.5	-			H43	1.5	90Dy04
	929.6	1.4			0.5	-			CP1	1.0	09Sc19
ave.	930.2	1.1			0.1	1	77	56 <sup>128</sup> Xe	average		
<sup>128</sup> Xe- <sup>126</sup> Xe	-774	45	-766	4	0.1	U			M16	2.5	63Da10
<sup>126</sup> Cs- <sup>128</sup> Cs <sub>656</sub> <sup>122</sup> Cs <sub>344</sub>	-1130	30	-1103	16	0.4	U			P22	2.5	82Au01
<sup>124</sup> Cs <sup>x</sup> - <sup>128</sup> Cs <sub>323</sub> <sup>122</sup> Cs <sub>678</sub>	-1070	30	-970	30	1.3	U			P22	2.5	82Au01
<sup>126</sup> Cs- <sup>128</sup> Cs <sub>591</sub> <sup>123</sup> Cs <sub>410</sub>	-350	30	-340	12	0.1	U			P22	2.5	82Au01
<sup>124</sup> Cs <sup>x</sup> - <sup>128</sup> Cs <sub>194</sub> <sup>123</sup> Cs <sub>807</sub>	370	50	366	24	0.0	U			P22	2.5	82Au01
<sup>125</sup> Cs- <sup>128</sup> Cs <sub>244</sub> <sup>124</sup> Cs <sub>756</sub>	-1440	30	-1354	18	1.1	U			P22	2.5	82Au01
<sup>126</sup> Cs- <sup>128</sup> Cs <sub>492</sub> <sup>124</sup> Cs <sub>508</sub>	-610	30	-568	15	0.6	U			P22	2.5	82Au01
<sup>127</sup> Cs- <sup>128</sup> Cs <sub>661</sub> <sup>125</sup> Cs <sub>339</sub>	-965	16	-934	7	0.8	U			P32	2.5	86Au02
<sup>127</sup> Cs- <sup>128</sup> Cs <sub>496</sub> <sup>126</sup> Cs <sub>504</sub>	-1160	30	-1105	8	0.7	U			P22	2.5	82Au01
<sup>128</sup> Te( $\gamma,n$ ) <sup>127</sup> Te	-8410	120	-8783.4	1.7	-3.1	B			Phi		60Ge01
<sup>127</sup> I(n, $\gamma$ ) <sup>128</sup> I	6825.7	0.5	6826.13	0.05	0.9	U					71Sc07
	6826.12	0.05			0.2	-			MMn		90Is03 Z
	6826.22	0.14			-0.6	-			Bdn		06Fi.A
	ave.	6826.13	0.05			0.0	1	100	87 <sup>128</sup> I	average	
<sup>128</sup> Cd( $\beta^-$ ) <sup>128</sup> In	7070	290	6900	150	-0.6	1	28	28 <sup>128</sup> In	Stu		87Sp09
<sup>128</sup> In( $\beta^-$ ) <sup>128</sup> Sn	9280	180	9220	150	-0.4	1	72	72 <sup>128</sup> In	Stu		78Al18 *
	8984	37			6.3	F			Stu		87Sp09 *
	8950	103			2.6	F			Gsn		90St13 *
<sup>128</sup> In <sup>m</sup> ( $\beta^-$ ) <sup>128</sup> Sn	9390	220	9298	28	-0.4	o			Stu		78Al18 *
	9306	30			-0.3	2			Stu		87Sp09 *
	9230	90			0.8	2			Gsn		90St13 *
<sup>128</sup> Sn( $\beta^-$ ) <sup>128</sup> Sb <sup>m</sup>	1265	30	1258	12	-0.2	-					76Nu01 *
	1290	40			-0.8	-			Stu		77Lu06 *
	1260	15			-0.1	-			Gsn		90St13 *
	ave.	1264	13			-0.4	1	87	45 <sup>128</sup> Sb <sup>m</sup>	average	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{128}\text{Sb}^m(\text{IT})^{128}\text{Sb}$	10	7				2					AHW *
$^{128}\text{Sb}(\beta^-)^{128}\text{Te}$	4640	100	4363	19	-2.8	U					71Ki15 *
$^{128}\text{Sb}^m(\beta^-)^{128}\text{Te}$	4391	40	4373	18	-0.4	-			Stu		77Lu06 *
	4395	30			-0.7	-			Gsn		90St13 *
ave.	4394	24			-0.8	1	55	55 $^{128}\text{Sb}^m$			average
$^{128}\text{I}(\beta^+)^{128}\text{Te}$	1277	13	1255	4	-1.7	U					61La16
$^{128}\text{I}(\beta^-)^{128}\text{Xe}$	2116	10	2122	4	0.6	1	14	13 $^{128}\text{I}$			56Be18 *
$^{128}\text{Cs}(\beta^+)^{128}\text{Xe}$	3855	90	3929	5	0.8	U					75We23 *
	3928	6			0.1	1	80	80 $^{128}\text{Cs}$			76Cr.B
	3907	40			0.5	o			IRS		83A106
	3930	100			0.0	U			IRS		93A103
$^{128}\text{La}(\beta^+)^{128}\text{Ba}$	6650	400	6750	50	0.3	U					66Li04 *
	6820	100			-0.7	R			JAE		98Ko66 *
* $^{128}\text{La-u}$	$M-A=-78601(28)$ keV for mixture gs+m at 100#100 keV										
* $^{128}\text{Sn }^{34}\text{S}-^{133}\text{Cs}_{1,218}$	F : authors say "possible contamination, measurement abandoned"										
* $^{128}\text{In}(\beta^-)^{128}\text{Sn}$	$E_{\beta^-}=4980(180)$ to $(2)^+$ level at 4297.70 keV										
* $^{128}\text{In}(\beta^-)^{128}\text{Sn}$	$E_{\beta^-}=5464(37)$ to $(2)^+$ at 3519.86; others 6986(170), 7857(109) to 2104.07, 1168.82; different equipment/method than in previous; low $E_{\beta^-}$ not seen										
* $^{128}\text{In}(\beta^-)^{128}\text{Sn}$	$E_{\beta^-}=4650(120), 5440(200)$ to 4297, 3520 levels and others										
* $^{128}\text{In}(\beta^-)^{128}\text{Sn}$	F : above 2 items conflict with 1st one and with trends in Ex of $8^-$ isomer										
* $^{128}\text{In}^m(\beta^-)^{128}\text{Sn}$	$E_{\beta^-}=5430(220)$ to 3958 level										
* $^{128}\text{In}^m(\beta^-)^{128}\text{Sn}$	$E_{\beta^-}=5239(40), 5350(44)$ to 4066, 3958 level										
* $^{128}\text{In}^m(\beta^-)^{128}\text{Sn}$	$E_{\beta^-}=5160(170), 5250(130)$ to 4066, 3958 levels										
* $^{128}\text{Sn}(\beta^-)^{128}\text{Sb}^m$	$E_{\beta^-}=630(30) 655(40) 625(15)$ , to $1^+$ level 635.2 above $^{128}\text{Sb}^m$ at 10(7) keV										
* $^{128}\text{Sb}^m(\text{IT})^{128}\text{Sb}$	From 3.6% IT for M3 transition										
* $^{128}\text{Sb}(\beta^-)^{128}\text{Te}$	$E_{\beta^-}=2300(100)$ to $(7)^-$ level at 2337.68 keV										
* $^{128}\text{Sb}^m(\beta^-)^{128}\text{Te}$	$E_{\beta^-}=2580(40) 2585(30)$ respectively, to $6^+$ level at 1811.13 keV										
* $^{128}\text{I}(\beta^-)^{128}\text{Xe}$	$E_{\beta^-}=2120(10)$ and $E_{\beta^-}=1665(15)$ to ground state and $2^+$ level at 442.911 keV										
* $^{128}\text{Cs}(\beta^+)^{128}\text{Xe}$	$E_{\beta^+}=2390(90)$ to $2^+$ level at 442.911 keV										
* $^{128}\text{La}(\beta^+)^{128}\text{Ba}$	$E_{\beta^+}=3200(400) 3370(100)$ respectively, to $(4^-, 5^+)$ level at 2425.45 keV										
$^{129}\text{Cd-u}$	-67789	186	-67696	18	0.2	U			GT3	2.5	16Kn02
$^{129}\text{Sn-u}$	-86521	31	-86518	19	0.1	1	36	36 $^{129}\text{Sn}$	MA8	1.0	05Si34 *
$^{129}\text{Xe}-^{120}\text{Sn}_{1,075}$	9913.3	2.4	9913.9	1.0	0.2	1	19	19 $^{120}\text{Sn}$	JY1	1.0	11Ha48
$\text{C}_{10} \text{H}_9-^{129}\text{Xe}$	165643.6	3.6	165644.431	0.006	0.1	U			M16	2.5	63Da10
$^{129}\text{Xe-u}$	-95228.7	5.4	-95219.141	0.006	0.7	U			ACC	2.5	90Me08
$^{129}\text{Xe}-\text{C}_2 \text{ } ^{35}\text{Cl}_3$	-1777.98	0.68	-1777.23	0.11	0.7	U			H47	1.5	94Hy01
$^{129}\text{Xe}_2-^{86}\text{Kr}_3$	77729.8547	0.0250	77729.839	0.014	-0.6	1	30	15 $^{86}\text{Kr}$	FS1	1.0	05Sh38 *
$^{129}\text{La-u}$	-87300	30	-87306	23	-0.2	1	58	58 $^{129}\text{La}$	GS2	1.0	05Li24
$^{129}\text{Ce-u}$	-81898	30				2			GS2	1.0	05Li24
$^{129}\text{Pr-u}$	-74905	32				2			GS2	1.0	05Li24 *
$^{129}\text{Xe}-^{134}\text{Xe}_{.963}$	-4114.7	3.8	-4112.633	0.009	0.5	o			MA8	1.0	05He.A
	-4119.3	5.1			1.3	U			MA8	1.0	06He29
$^{129}\text{Cd}-^{133}\text{Cs}_{.970}$	24016	18				2			MA8	1.0	15At03
$^{129}\text{Cs}-^{133}\text{Cs}_{.970}$	-2234	18	-2223	5	0.6	o			MA1	1.0	90St25
	-2216	14			-0.5	1	12	12 $^{129}\text{Cs}$	MA1	1.0	99Am05
$^{129}\text{In}-^{130}\text{Xe}_{.992}$	17523.9	2.9	17524.2	2.9	0.1	1	99	99 $^{129}\text{In}$	JY1	1.0	12Ha25
$^{129}\text{In}^m-^{130}\text{Xe}_{.992}$	18016.5	3.5	18016	3	-0.1	1	99	99 $^{129}\text{In}^m$	JY1	1.0	13Ka08
$\text{C}_{10} \text{H}_{10}-^{129}\text{Xe}$	173469.4660	0.0147	173469.463	0.006	-0.2	1	15	15 $^{129}\text{Xe}$	FS1	1.0	09Re03
$^{129}\text{Xe}-^{128}\text{Xe}$	1247	12	1249.9	1.1	0.1	U			M16	2.5	63Da10
$\text{C}_3 \text{O}_6-^{129}\text{Xe}$	64706.8420	0.0255	64706.859	0.006	0.7	o			FS1	1.0	05Sh38 *
	64706.8516	0.0181			0.4	1	10	10 $^{129}\text{Xe}$	FS1	1.0	09Re03
$^{129}\text{Xe}_2-^{84}\text{Kr}_3$	75068.5115	0.0405	75068.532	0.014	0.5	1	12	7 $^{84}\text{Kr}$	FS1	1.0	05Sh38 *
$^{128}\text{Cs}-^{129}\text{Cs}_{.661}$ $^{126}\text{Cs}_{.339}$	510	30	500	7	-0.1	U			P22	2.5	82Au01
$^{128}\text{Te}(n,\gamma)^{129}\text{Te}$	6085	3	6082.41	0.08	-0.9	U					72Mu.A
	6082.42	0.09			-0.1	-			Prn		03Wi02
	6082.36	0.19			0.3	-			Bdn		06Fi.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{128}\text{Te}(\text{d,p})^{129}\text{Te}$		3857	10	3857.84	0.08	0.1	U		MIT		67Mo22
$^{128}\text{Te}(\text{n},\gamma)^{129}\text{Te}$	ave.	6082.41	0.08	6082.41	0.08	0.0	1	100	98 $^{129}\text{Te}$		average
$^{129}\text{Nd}(\text{ep})^{128}\text{Ce}$		5300	300	5930#	200#	2.1	D				78Bo.A *
$^{129}\text{In}(\beta^-)^{129}\text{Sn}$		7655	32	7753	17	3.1	B		Stu		87Sp09 *
		7780	26			-1.0	1	44	44 $^{129}\text{Sn}$		04Ga24 *
$^{129}\text{In}^m(\beta^-)^{129}\text{Sn}$		8033	66	8211	17	2.7	B		Stu		87Sp09 *
		8149	38			1.6	1	21	21 $^{129}\text{Sn}$		04Ga24 *
$^{129}\text{In}^p(\beta^-)^{129}\text{Sn}$		9410	50				2		Stu		04Ga24 *
$^{129}\text{Sn}(\beta^-)^{129}\text{Sb}$		3996	120	4038	27	0.4	U		Stu		77Lu06 *
$^{129}\text{Sb}(\beta^-)^{129}\text{Te}$		2345	30	2375	21	1.0	2				70Oh05 *
$^{129}\text{Te}(\beta^-)^{129}\text{I}$		1453	28	1502	3	1.8	U				56Gr10 *
		1485	10			1.7	U				64De10 *
		1503	4			-0.2	1	62	60 $^{129}\text{I}$		68Go34 *
$^{129}\text{I}(\beta^-)^{129}\text{Xe}$		190	5	189	3	-0.2	1	40	40 $^{129}\text{I}$		54De17 *
$^{129}\text{Cs}(\beta^+)^{129}\text{Xe}$		1197	5	1197	5	0.0	1	83	83 $^{129}\text{Cs}$		76Ma35 *
$^{129}\text{Ba}(\beta^+)^{129}\text{Cs}$		2446	15	2436	11	-0.7	1	50	45 $^{129}\text{Ba}$		61Ar05 *
$^{129}\text{La}(\beta^+)^{129}\text{Ba}$		3720	50	3739	22	0.4	-				79Br05 *
		3740	40			0.0	-		JAE		98Ko66 *
	ave.	3730	30			0.2	1	48	42 $^{129}\text{La}$		average
$^{129}\text{Ce}(\beta^+)^{129}\text{La}$		5600	200	5040	40	-2.8	U		IRS		93Al03 *
* $^{129}\text{Sn}-\text{u}$	$M - A = -80576(27)$ keV for mixture gs+m at 35.15 keV										
* $^{129}\text{Xe}_{2-86}\text{Kr}_3$	Corrected in reference of same group										
* $^{129}\text{Pr}-\text{u}$	Isomer at 382.57 with estimated $T=1\#$ ms not considered										
* $\text{C}_3 \text{O}_6-^{129}\text{Xe}$	Corrected in reference of same group										
* $^{129}\text{Xe}_{2-84}\text{Kr}_3$	Corrected in reference of same group										
* $^{129}\text{Nd}(\text{ep})^{128}\text{Ce}$	Trends from Mass Surface TMS suggest $^{129}\text{Nd}$ 630 less bound										
* $^{129}\text{In}(\beta^-)^{129}\text{Sn}$	$E_{\beta^-} = 7780(26), 9410(50)$ from ground state, 1688.0(0.5) levels										
* $^{129}\text{Sn}(\beta^-)^{129}\text{Sb}$	$E_{\beta^-} = 3350(120)$ to $(5/2^+)$ level at 645.14 keV										
* $^{129}\text{Sb}(\beta^-)^{129}\text{Te}$	$E_{\beta^-} = 1800(30)$ to $5/2^+$ level at 544.585 keV, and other $E_{\beta^-}$										
* $^{129}\text{Te}(\beta^-)^{129}\text{I}$	$E_{\beta^-} = 1453(5)$ to $5/2^+$ level at 27.793 keV and 1530(5) from $^{129}\text{Te}^m$										
*	at 105.51 to ground state (Birge=8.0: arithmetic average used)										
* $^{129}\text{Te}(\beta^-)^{129}\text{I}$	$E_{\beta^-} = 1452(10)$ to $5/2^+$ level at 27.793 keV and 1595(10) from $^{129}\text{Te}^m$ at 105.51										
* $^{129}\text{Te}(\beta^-)^{129}\text{I}$	$E_{\beta^-} = 1476(4)$ to $5/2^+$ level at 27.793 keV and 1607(7) from $^{129}\text{Te}^m$ at 105.51										
* $^{129}\text{I}(\beta^-)^{129}\text{Xe}$	$E_{\beta^-} = 150(5)$ to $3/2^+$ level at 39.5774 keV										
* $^{129}\text{Ba}(\beta^+)^{129}\text{Cs}$	$E_{\beta^+} = 1425(15);$ and 1243(35), 975(60) from $^{129}\text{Ba}^m$ at 8.42(0.06) keV										
*	to $7/2^+$ level at 188.91, $(9/2)^+$ at 426.47 keV										
* $^{129}\text{La}(\beta^+)^{129}\text{Ba}$	$E_{\beta^+} = 2420(50)$ to $1/2^+$ level at 278.57 keV, and other $E_{\beta^+}$										
$^{130}\text{Cd}-\text{u}$	-66700	441	-65612	24	1.0	U			GT3	2.5	16Kn02
$\text{C}_9 \text{H}_8 \text{N}-^{130}\text{Te}$	159446	10	159451.515	0.012	0.2	U			M16	2.5	63Da10
$^{13}\text{C} \text{C}_8 \text{N} \text{H}_7-^{130}\text{Te}$	154990.6	7.	154981.318	0.012	-0.5	U			C3	2.5	70Ke05
$\text{C}_9 \text{H}_8 \text{N}-^{130}\text{Te}$	159449	8	159451.515	0.012	0.2	U			R13	1.5	83De51
$\text{C}_{10} \text{H}_{10}-^{130}\text{Xe}$	174743.6	4.2	174740.973	0.010	-0.3	U			M16	2.5	63Da10
$^{13}\text{C} \text{C}_8 \text{N} \text{H}_7-^{130}\text{Xe}$	157695.4	0.7	157694.716	0.010	-0.4	U			C3	2.5	70Ke05
$^{130}\text{Xe}-\text{C}^{13}\text{C}^{35}\text{Cl}_3$	-6407.63	1.21	-6403.57	0.11	2.2	U			H47	1.5	94Hy01
$^{130}\text{Cs}-\text{u}$	-93181	60	-93291	9	-1.8	U			GS2	1.0	05Li24 *
$\text{C}_{10} \text{H}_{10}-^{130}\text{Ba}$	171926	68	171929.4	2.7	0.0	U			R07	1.5	68De17
$^{130}\text{Ba}-^{85}\text{Rb}_{1,529}$	41195.8	3.4	41194.4	2.7	-0.4	1	65	65 $^{130}\text{Ba}$	MA8	1.0	05Gu37
$^{130}\text{La}-\text{u}$	-87635	30	-87631	28	0.1	2			GS2	1.0	05Li24
$^{130}\text{Ce}-\text{u}$	-85264	30				2			GS2	1.0	05Li24
$^{130}\text{Pr}-\text{u}$	-76410	69				2			GS2	1.0	05Li24 *
$^{130}\text{Nd}-\text{u}$	-71494	30				2			GS2	1.0	05Li24
$^{130}\text{Xe}-^{134}\text{Xe}_{,970}$	-4726.6	5.6	-4721.891	0.012	0.8	o			MA8	1.0	05He.A
	-4724.8	7.0			0.4	U			MA8	1.0	06He29
$^{130}\text{Cd}-^{133}\text{Cs}_{,977}$	26761	24				2			MA8	1.0	15At03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{130}\text{In}-^{133}\text{Cs}_{977}$	17384	130	17350	40	-0.3	U			CP1	1.0	13Va12 *
$^{130}\text{Sn}-^{133}\text{Cs}_{977}$	6346	17	6348.0	2.0	0.1	U			MA8	1.0	05Si34
	6344	12			0.3	U			MA8	1.0	05Si34 *
	6349.5	3.9			-0.4	1	27	27 $^{130}\text{Sn}$	CP1	1.0	13Va12
$^{130}\text{Xe}-^{133}\text{Cs}_{977}$	-4114	13	-4117.217	0.011	-0.2	U			MA6	1.0	04Di18
$^{130}\text{Cs}-^{133}\text{Cs}_{977}$	-928	17	-917	9	0.6	o			MA1	1.0	90St25
	-916	13			-0.1	1	48	48 $^{130}\text{Cs}$	MA1	1.0	99Am05
$^{130}\text{Nd } ^{19}\text{F}-^{133}\text{Cs}_{1,120}$	32902	130	32800	30	-0.8	U			MA5	1.0	00Be42 *
$^{130}\text{Te } ^{35}\text{Cl}-^{128}\text{Te } ^{37}\text{Cl}$	4715	2	4711.5	0.9	-0.4	o			H16	4.0	63Ba47
	4711.7	1.8			0.0	U			C3	2.5	70Ke05
	4711.57	0.72			0.0	1	74	74 $^{128}\text{Te}$	H43	1.5	90Dy04
$^{130}\text{Xe}-^{129}\text{Xe}_{1,008}$	-509.78	0.34	-509.755	0.009	0.1	U			CP1	1.0	09Sc19
	-509.96	0.26			0.8	U			SH2	1.0	13El01 *
	-509.02	0.94			-0.8	U			SH1	1.0	13El01 *
$^{130}\text{Sn}-^{130}\text{Xe}$	10463.9	3.6	10465.2	2.0	0.4	-			JY1	1.0	12Ha25
	10465.4	3.1			-0.1	-			JY1	1.0	13Ka08 *
	ave.	10464.8	2.3		0.2	1	73	73 $^{130}\text{Sn}$			average
$^{130}\text{Te}-^{130}\text{Xe}$	2706.2	7.	2713.398	0.012	0.4	U			C3	2.5	70Ke05
	2712.98	3.02			0.1	U			H43	1.5	90Dy04
	2713.416	0.034			-0.5	-			FS1	1.0	09Re07
	2713.402	0.026			-0.2	-			FS1	1.0	09Re07 *
	2713.402	0.014			-0.3	o			FS1	1.0	09Re07 *
	2712.86	0.34			1.6	U			CP1	1.0	09Sc19
	2712.82	0.25			2.3	U			JY1	1.0	11Ra24
	ave.	2713.407	0.021		-0.4	1	35	22 $^{130}\text{Te}$			average
$^{130}\text{Te}-^{129}\text{Xe}$	1441.885	0.012	1441.888	0.011	0.3	1	78	78 $^{130}\text{Te}$	FS1	1.0	09Re07
$^{130}\text{Xe}-^{129}\text{Xe}$	-1277	12	-1271.510	0.009	0.2	U			M16	2.5	63Da10
	-1271.517	0.012			0.6	1	52	50 $^{130}\text{Xe}$	FS1	1.0	09Re07
$^{129}\text{Cs}-^{130}\text{Cs}_{794}^{125}\text{Cs}_{206}$	-1270	40	-1200	14	0.7	U			P22	2.5	82Au01
$^{130}\text{Ba}(p,t)^{128}\text{Ba}$	-9482	32	-9543.9	2.8	-1.9	U			Win		74De31 *
$^{130}\text{Ba}(p,t)^{128}\text{Ba}-^{144}\text{Sm}()^{142}\text{Sm}$	1095.9	1.0	1096.0	1.0	0.1	1	99	78 $^{128}\text{Ba}$			09Pa25
$^{130}\text{Te}(d,^3\text{He})^{129}\text{Sb}$	-4550	30	-4519	21	1.0	R			Oak		68Au04
$^{129}\text{I}(n,\gamma)^{130}\text{I}$	6500.33	0.04				2			ILn		89Sa11
$^{129}\text{Xe}(n,\gamma)^{130}\text{Xe}$	9255.3	1.0	9255.721	0.008	0.4	U					71Gr28 Z
	9256.1	0.8			-0.5	U					74Ge05 Z
	9255.57	0.30			0.5	U			Bdn		06Fi.A
$^{129}\text{Xe}(^3\text{He},d)^{130}\text{Cs}$	5	20	-1	8	-0.3	1	17	17 $^{130}\text{Cs}$	ChR		81Ha08
$^{130}\text{Ba}(d,t)^{129}\text{Ba}$	-4001	15	-4012	11	-0.8	1	50	48 $^{129}\text{Ba}$	Tal		74Gr22
$^{130}\text{Eu}(p)^{129}\text{Sm}$	1028.0	15.0				3			Arp		04Da04
$^{130}\text{Cd}(\beta^-)^{130}\text{In}$	8350	160	8770	40	2.6	B			Bwg		03Di06 *
$^{130}\text{In}(\beta^-)^{130}\text{Sn}$	10249	38				2			Stu		87Sp09
	9880	90	10250	40	4.1	B			Gsn		90St13
$^{130}\text{In}^m(\beta^-)^{130}\text{Sn}$	10300	37				2			Stu		87Sp09
	10170	170	10300	40	0.8	U			Gsn		90St13
$^{130}\text{In}^n(\beta^-)^{130}\text{Sn}$	10650	49				2			Stu		87Sp09
	9880	200	10650	50	3.8	B			Gsn		90St13
$^{130}\text{Sn}(\beta^-)^{130}\text{Sb}$	2195	35	2153	14	-1.2	-			Stu		77Lu06 *
	2080	40			1.8	-					77Nu01 *
	2149	18			0.2	-			Gsn		90St13 *
	ave.	2148	15		0.4	1	90	90 $^{130}\text{Sb}$			average
$^{130}\text{Sb}(\beta^-)^{130}\text{Te}$	5046	100	5067	14	0.2	U					71Ki15 *
	5015	100			0.5	U			Stu		77Lu06 *
	4990	70			1.1	U			Gsn		90St13 *
	5015	45			1.2	1	10	10 $^{130}\text{Sb}$	Stu		95Me16 *
$^{130}\text{I}(\beta^-)^{130}\text{Xe}$	2983	10	2944	3	-3.9	B					65Da01 *
	2964	50			-0.4	U					70Qa03 *
$^{130}\text{Cs}(\beta^+)^{130}\text{Xe}$	2992	20	2981	8	-0.6	-					52Sm41
	2972	20			0.4	-					75We23
	ave.	2982	14		-0.1	1	35	35 $^{130}\text{Cs}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{130}\text{Cs}^x(\text{IT})^{130}\text{Cs}$	27	15							AHW *
$^{130}\text{Cs}(\beta^-)^{130}\text{Ba}$	442	50	362	9	-1.6	U			52Sm41 *
$^{130}\text{La}(\beta^+)^{130}\text{Ba}$	5660	70	5634	26	-0.4	R	JAE		98Ko66
* $^{130}\text{Cs}-u$	$M - A = -86716(30)$ keV for mixture gs+m at 163.25 keV								
* $^{130}\text{Pr}-u$	$M - A = -71125(29)$ keV for mixture gs+m at 100#100 keV								
* $^{130}\text{In}-^{133}\text{Cs}_{977}$	$M - A = -69652(20)$ keV for mixture gs+n at 400(60) keV								
*	assuming no $8^-$ isomer								
* $^{130}\text{Sn}-^{133}\text{Cs}_{977}$	$D_M = 8434(12)$ $\mu\text{u}$ for $^{130}\text{Sn}^m$ at 1946.88 keV; $M - A = -78189(11)$ keV								
* $^{130}\text{Nd } ^{19}\text{F}-^{133}\text{Cs}_{1,120}$	Tentative result, low statistics								
* $^{130}\text{Xe}-^{129}\text{Xe}_{1,008}$	Respectively for PI-ICR and Ramsey ToF-ICR techniques								
* $^{130}\text{Sn}-^{130}\text{Xe}$	$D_M = 12555.5(3.1)$ $\mu\text{u}$ for $^{130}\text{Sn}^m$ at 1946.88 keV; $M - A = -78185.1(2.9)$ keV								
* $^{130}\text{Te}-^{130}\text{Xe}$	First item 1 ion; second item 2 ions - considered independent								
* $^{130}\text{Te}-^{130}\text{Xe}$	Combination of $^{130}\text{Xe}-^{129}\text{Xe}$ and $^{130}\text{Te}-^{129}\text{Xe}$								
* $^{130}\text{Ba}(p,t)^{128}\text{Ba}$	Not resolved peak. Original uncertainty 16 increased to 24 keV and added systematic error 21 keV								
*									
* $^{130}\text{Cd}(\beta^-)^{130}\text{In}$	$E_{\beta^-} = 6224(+165-157)$ to $1^+$ level at 2120.2 keV								
* $^{130}\text{Sn}(\beta^-)^{130}\text{Sb}$	$E_{\beta^-} = 1490(90), 1150(35)$ to $1^+$ levels at 702.32, 1047.67 keV								
* $^{130}\text{Sn}(\beta^-)^{130}\text{Sb}$	$E_{\beta^-} = 1280(80), 1060(40)$ to $1^+$ levels at 702.32, 1047.67 keV								
* $^{130}\text{Sn}(\beta^-)^{130}\text{Sb}$	$E_{\beta^-} = 1415(30), 1112(18)$ to $1^+$ levels at 702.32, 1047.67 keV								
*	and a $3\sigma$ conflicting 3955(50) from $^{130}\text{Sn}^m$ at 1946.88 keV								
* $^{130}\text{Sb}(\beta^-)^{130}\text{Te}$	$E_{\beta^-} = 2900(100)$ to $^{130}\text{Te}^m$ at 2146.41 keV								
* $^{130}\text{Sb}(\beta^-)^{130}\text{Te}$	$Q = 5020(100)$ from $^{130}\text{Sb}^m$ at 4.80 keV								
* $^{130}\text{Sb}(\beta^-)^{130}\text{Te}$	Also 4960(25) from $^{130}\text{Sb}^m$ at 4.80, in disagreement								
* $^{130}\text{Sb}(\beta^-)^{130}\text{Te}$	Derived from given average=5008(38) with 90St13=4990(70) keV								
* $^{130}\text{I}(\beta^-)^{130}\text{Xe}$	$E_{\beta^-} = 1702(10) 1042(10) 618(10)$ to $4^+ 1204.614, 6^+ 1944.140, 5^+ 2362.073$								
* $^{130}\text{I}(\beta^-)^{130}\text{Xe}$	$E_{\beta^-} = 2480(50), 1850(80)$ from $^{130}\text{I}^m$ at 39.9525 to $2^+$ levels at 536.068,								
*	and 1122.112 keV								
* $^{130}\text{Cs}^x(\text{IT})^{130}\text{Cs}$	Combining isomer ratio of reference								
*	with $^{130}\text{Cs}^m(\text{IT}) = 163.25$ keV								
* $^{130}\text{Cs}(\beta^-)^{130}\text{Ba}$	Value given without associated error								
$^{131}\text{Cd}-u$	-59671	1023	-59280	110	0.2	U			GT3 2.5 16Kn02
	-59280	110				2			MR1 1.0 15At03
$^{131}\text{Sn}-u$	-82958	26	-82947	4	0.4	U			MA8 1.0 05Si34 *
	-82950	130			0.0	U			GT2 2.5 08Su19 *
$^{131}\text{Sb}-u$	-88170	530	-88010.7	2.2	0.1	U			GT2 2.5 08Kn.A *
$\text{C}_{10} \text{H}_{11} - ^{131}\text{Xe}$	180991.6	3.0	180991.218	0.009	-0.1	U			M16 2.5 63Da10
$^{131}\text{Xe}-u$	-94925.5	5.7	-94915.864	0.009	0.7	U			ACC 2.5 90Me08
$^{131}\text{Xe}-\text{C}_2 \text{ } ^{35}\text{Cl}_2 \text{ } ^{37}\text{Cl}$	1472.65	0.80	1476.16	0.09	2.9	B			H47 1.5 94Hy01
$^{131}\text{Ba}-u$	-92955	66	-93058.8	2.8	-1.6	U			GS2 1.0 05Li24 *
$^{131}\text{La}-u$	-89930	30				2			GS2 1.0 05Li24
$^{131}\text{Ce}-u$	-85579	36	-85570	40	0.2	1	96	96 $^{131}\text{Ce}$	GS2 1.0 05Li24 *
$^{131}\text{Pr}-u$	-79741	56	-79770	50	-0.4	1	81	81 $^{131}\text{Pr}$	GS2 1.0 05Li24 *
$^{131}\text{Nd}-u$	-72753	30	-72752	30	0.0	1	97	97 $^{131}\text{Nd}$	GS2 1.0 05Li24
$^{131}\text{In}-^{133}\text{Cs}_{985}$	20262	37	20101.9	2.9	-4.3	C			CP1 1.0 13Va12 *
$^{131}\text{Sn } ^{34}\text{S}-^{133}\text{Cs}_{1,241}$	2253	11	2254	4	0.1	-			MA8 1.0 08Dw01 *
$^{131}\text{Sn}-^{133}\text{Cs}_{985}$	10188.2	4.7	10183	4	-1.1	-			CP1 1.0 13Va12
$^{131}\text{Sn } ^{34}\text{S}-^{133}\text{Cs}_{1,241}$	ave. 2259	4	2254	4	-1.0	1	81	81 $^{131}\text{Sn}$	average
$^{131}\text{Sb}-^{133}\text{Cs}_{985}$	5114	11	5119.2	2.2	0.5	U			CP1 1.0 13Va12
$^{131}\text{Xe}-^{129}\text{Xe}_{1,016}$	1826.7855	0.0096	1826.787	0.008	0.1	1	66	62 $^{131}\text{Xe}$	FS1 1.0 13Ho22
$^{131}\text{Cs}-^{133}\text{Cs}_{985}$	-1429	18	-1405	5	1.3	o			MA1 1.0 90St25
	-1419	14			1.0	1	15	15 $^{131}\text{Cs}$	MA1 1.0 99Am05
$^{131}\text{Ba}-^{133}\text{Cs}_{985}$	72	14	71.0	2.8	-0.1	U			MA5 1.0 00Be42
$^{131}\text{In}-^{130}\text{Xe}_{1,008}$	24234.7	2.9				2			JY1 1.0 12Ha25
$^{131}\text{In}^m - ^{130}\text{Xe}_{1,008}$	24626.8	7.7				2			JY1 1.0 13Ka08

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{131}\text{Sn}-^{132}\text{Xe}_{.992}$	12134	21	12131	4	-0.1	U			JY1	1.0	12Ha25 *
$^{131}\text{Sb}-^{130}\text{Xe}_{1.008}$	9250.7	2.3	9251.9	2.2	0.5	1	95	95 $^{131}\text{Sb}$	JY1	1.0	12Ha25
$^{131}\text{Xe}-^{132}\text{Xe}_{.992}$	162.283	0.030	162.288	0.008	0.2	-			SH1	1.0	14Ne15
	162.292	0.014			-0.3	-			FS1	1.0	13Ho22
	ave. 162.290	0.013			-0.2	1	40	38 $^{131}\text{Xe}$			average
$^{131}\text{Xe}-^{130}\text{Xe}$	1574	11	1574.787	0.011	0.0	U			M16	2.5	63Da10
$^{128}\text{Cs}-^{131}\text{Cs}_{.391}$ $^{126}\text{Cs}_{.610}$	-100	30	-47	9	0.7	U			P22	2.5	82Au01
$^{128}\text{Cs}-^{131}\text{Cs}_{.244}$ $^{127}\text{Cs}_{.756}$	783	21	752	7	-0.6	F			P33	2.5	86Au02 *
$^{129}\text{Cs}-^{131}\text{Cs}_{.328}$ $^{128}\text{Cs}_{.672}$	-1030	30	-870	6	2.1	U			P22	2.5	82Au01
$^{130}\text{Te}(n,\gamma)^{131}\text{Te}$	5929.7	0.5	5929.38	0.06	-0.6	U					77Ko.A
	5929.5	0.4			-0.3	U					80Ho29 Z
	5929.38	0.06				2			Prn		03To08
	5930.16	0.19			-4.1	C			Bdn		06Fi.A
$^{130}\text{Te}(d,p)^{131}\text{Te}$	3703	6	3704.81	0.06	0.3	U			MIT		67Gr21
$^{130}\text{Ba}(n,\gamma)^{131}\text{Ba}$	7493.5	0.3	7493.50	0.30	0.0	1	100	95 $^{131}\text{Ba}$			82Ka.A
$^{130}\text{Ba}(d,p)^{131}\text{Ba}$	5269	15	5268.94	0.30	0.0	U			ANL		70Vo04
$^{131}\text{Nd}(\epsilon p)^{130}\text{Ce}$	4600	400	4370	40	-0.6	U					78Bo.A
$^{131}\text{Eu}(p)^{130}\text{Sm}$	957.4	8.	947	5	-1.3	3					98Da03
	939.2	7.			1.1	3					99So17
$^{131}\text{In}(\beta^-)^{131}\text{Sn}$	8820	200	9240	5	2.1	U					80De35
	8930	150			2.1	o			Stu		84Fo19
	9184	33			1.7	o			Stu		88Fo05
	9165	30			2.5	o			Stu		95Me16
	9174	22			3.0	B			Stu		99Fo01
	9222	18			1.0	U			Stu		04Fo06
$^{131}\text{In}^m(\beta^-)^{131}\text{Sn}$	9230	220	9605	8	1.7	o			Stu		84Fo19
	9547	46			1.3	o			Stu		88Fo05
	9480	70			1.8	o			Stu		95Me16
	9524	26			3.1	B			Stu		04Fo06
$^{131}\text{In}^n(\beta^-)^{131}\text{Sn}$	13000	500	12990	90	0.0	o			Stu		84Fo19
	13450	163			-2.8	B			Stu		88Fo05
	13230	80			-3.1	B			Stu		95Me16
	12986	86				2			Stu		04Fo06
$^{131}\text{Sn}(\beta^-)^{131}\text{Sb}$	4582	120	4717	4	1.1	o			Bwg		79Ke02 *
	4640	20			3.8	B			Stu		84Fo19 *
	4610	110			1.0	U			Bwg		87Gr.A *
	4689	14			2.0	o			Stu		95Me16
	4688	14			2.1	o			Stu		99Fo01
	4701	8			2.0	1	25	19 $^{131}\text{Sn}$	Stu		04Fo06 *
$^{131}\text{Sb}(\beta^-)^{131}\text{Te}$	3190	70	3229.6	2.1	0.6	o			Stu		77Lu06
	3217	20			0.6	o			Stu		95Me16
	3200	26			1.1	U			Stu		99Fo01
	2275	10	2231.7	0.6	-4.3	B					61Be20 *
$^{131}\text{I}(\beta^-)^{131}\text{Xe}$	2278	15			-3.1	B					65De22 *
	971.0	0.7	970.8	0.6	-0.2	2					51Ve05 *
$^{131}\text{Cs}(\epsilon)^{131}\text{Xe}$	970.4	1.2			0.4	2					52Ro16 *
	355	10	355	5	0.0	-					54Sa22
	355	10			0.0	-					56Ho66
	360	15			-0.3	-					57Mi63
$^{131}\text{Ba}(\beta^+)^{131}\text{Cs}$	ave. 356	6			-0.2	1	60	60 $^{131}\text{Cs}$			average
	1370	16	1375	5	0.3	-					76Ge14 *
	1371	12			0.3	-					78Va04 *
$^{131}\text{La}(\beta^+)^{131}\text{Ba}$	ave. 1371	10			0.5	1	30	25 $^{131}\text{Cs}$			average
	2960	100	2914	28	-0.5	U					60Cr01
$^{131}\text{Ce}(\beta^+)^{131}\text{La}$	4020	400	4060	40	0.1	U					66No05 *
$^{131}\text{Pr}(\beta^+)^{131}\text{Ce}$	5250	150	5410	60	1.1	1	14	9 $^{131}\text{Pr}$	IRS		93Al03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{131}\text{Nd}(\beta^+)^{131}\text{Pr}$	6560	150	6530	50	-0.2	1	13	9	$^{131}\text{Pr}$	IRS	93Al03	
* $^{131}\text{Sn}-u$	$M-A=-77242(15)$ keV for mixture gs+m at 65.1 keV											
*	next $^{131}\text{Sn } ^{34}\text{S}-^{133}\text{Cs}_{1,241}$ also from 1.4 GeV p on UC target											
* $^{131}\text{Sn}-u$	$M-A=-77238(120)$ keV for mixture gs+m at 65.1 keV											
* $^{131}\text{Sb}-u$	$M-A=-81291(96)$ keV for mixture gs+m at 1676.06 keV											
* $^{131}\text{Ba}-u$	$M-A=-86494(30)$ keV for mixture gs+m at 187.995 keV											
* $^{131}\text{Ce}-u$	$M-A=-79685(28)$ keV for mixture gs+m at 63.09 keV											
* $^{131}\text{Pr}-u$	$M-A=-74202(28)$ keV for mixture gs+m at 152.4 keV											
* $^{131}\text{In}-^{133}\text{Cs}_{985}$	Freq. ratio mistyped, derived from mass excess in col.3 Table 1 of paper											
* $^{131}\text{Sn } ^{34}\text{S}-^{133}\text{Cs}_{1,241}$	$D_M=2300.3(3.6)$ $\mu\text{u}$ for mixture $^{131}\text{Sn}$ gs+m at 65.1 keV with $R=0.65(0.15)$											
* $^{131}\text{Sn}-^{132}\text{Xe}_{.992}$	$D_M=12168.7(3.4)$ $\mu\text{u}$ for mixture gs+m at 65.1 keV; $M-A=-77229.6(3.2)$ keV											
*	identical to result given in reference											
* $^{128}\text{Cs}-^{131}\text{Cs}_{.244}$ $^{127}\text{Cs}$ .	$F$ : rejection based on line-shape analysis											
* $^{131}\text{Sn}(\beta^-)^{131}\text{Sb}$	$E_{\beta^-}=3870(120), 2620(180)$ from $^{131}\text{Sn}^m$ at 65.1 keV											
*	to $(5/2^+)$ level at 798.494, $(13/2^-)$ at 1980.39 keV											
* $^{131}\text{Sn}(\beta^-)^{131}\text{Sb}$	$Q_{\beta^-}=4638(20)$ ; and 4796(80) from $^{131}\text{Sn}^m$ at 65.1 keV											
* $^{131}\text{Sn}(\beta^-)^{131}\text{Sb}$	$Q_{\beta^-}=4600(110)$ ; and 4680(120) from $^{131}\text{Sn}^m$ at 65.1 keV											
* $^{131}\text{Sn}(\beta^-)^{131}\text{Sb}$	$Q_{\beta^-}=4698(11)$ ; and 4767(8) from $^{131}\text{Sn}^m$ at 65.1 keV											
* $^{131}\text{Te}(\beta^-)^{131}\text{I}$	$Q_{\beta^-}=2457(10)$ 2460(15) respectively, from $^{131}\text{Te}^m$ at 182.258 keV											
* $^{131}\text{I}(\beta^-)^{131}\text{Xe}$	$E_{\beta^-}=606.5(0.7)$ 605.9(1.2) respectively, to $5/2^+$ level at 364.490 keV											
* $^{131}\text{Ba}(\beta^+)^{131}\text{Cs}$	$p^+=22(11)\times 10^{-6}$ to $3/2^+$ level at 216.086, recalculated $p^+$ and $Q$											
* $^{131}\text{Ba}(\beta^+)^{131}\text{Cs}$	$L/K=0.165(0.001)$ to $3/2^+$ level at 1047.682 keV											
* $^{131}\text{Ce}(\beta^+)^{131}\text{La}$	$E_{\beta^+}=2800(400)$ to $7/2^+$ level at 195.68 keV											
$^{132}\text{Sb}-u$	-85850	150	-85492.0	2.6	1.0	U			GT2	2.5	08Su19 *	
$\text{C}_{10} \text{H}_{12}-^{132}\text{Xe}$	189740.8	3.3	189745.300	0.006	0.5	U			M16	2.5	63Da10	
$^{132}\text{Xe}-u$	-95856.2	4.0	-95844.913	0.006	1.1	U			ACC	2.5	90Me08	
$^{132}\text{Xe}-\text{C } ^{13}\text{C } ^{35}\text{Cl}_2$ $^{37}\text{Cl}$	-2803.73	1.40	-2807.72	0.09	-1.9	U			H47	1.5	94Hy01	
$^{132}\text{Xe}-\text{C}_3 \text{O}_6$	-65332.6117	0.0248	-65332.631	0.006	-0.8	o			FS1	1.0	05Sh38 *	
	-65332.6238	0.0140			-0.5	1	16	15	$^{132}\text{Xe}$	FS1	1.0	09Re03
$\text{C}_{10} \text{H}_{12}-^{132}\text{Ba}$	188863	70	188839.3	1.1	-0.2	U			R07	1.5	68De17	
	188821	88			0.1	U			R07	1.5	68De17	
$^{132}\text{La}-u$	-89874	67	-89880	40	-0.1	1	34	34	$^{132}\text{La}$	GS2	1.0	05Li24 *
$^{132}\text{Ce}-u$	-88542	30	-88536	22	0.2	1	54	54	$^{132}\text{Ce}$	GS2	1.0	05Li24
$^{132}\text{Ce}-\text{O}-^{142}\text{Sm}_{1.042}$	-5258	32	-5265	22	-0.2	1	47	46	$^{132}\text{Ce}$	MA7	1.0	01Bo59 *
$^{132}\text{Pr}-u$	-80760	31				2			GS2	1.0	05Li24 *	
$^{132}\text{Nd}-u$	-76690	30	-76679	26	0.4	2			GS2	1.0	05Li24	
$^{132}\text{Xe}-^{129}\text{Xe}_{1.023}$	1564.20	0.32	1564.268	0.004	0.2	U			CP1	1.0	09Sc19	
	1565.4	1.0			-1.1	U			CP1	1.0	09Sc19	
$^{132}\text{Sb}-^{130}\text{Xe}_{1.015}$	12445.7	2.9	12446.0	2.6	0.1	1	83	83	$^{132}\text{Sb}$	JY1	1.0	12Ha25
$^{132}\text{Te}-^{130}\text{Xe}_{1.015}$	6482.9	4.3	6485	4	0.4	1	76	76	$^{132}\text{Te}$	JY1	1.0	12Ha25
$^{132}\text{Sn}-^{133}\text{Cs}_{.992}$	11621	19	11615.6	2.1	-0.3	U			MA8	1.0	05Si34	
	11613.1	2.9			0.8	-			CP1	1.0	13Va12	
$^{132}\text{Sn } ^{34}\text{S}-^{133}\text{Cs}_{1,248}$	3686.3	7.7	3686.9	2.1	0.1	-			MA8	1.0	08Dw01	
$^{132}\text{Sn}-^{133}\text{Cs}_{.992}$	ave. 11613.3	2.7	11615.6	2.1	0.8	1	61	61	$^{132}\text{Sn}$			average
$^{132}\text{Sb}-^{133}\text{Cs}_{.992}$	8301.3	6.5	8299.7	2.6	-0.3	1	17	17	$^{132}\text{Sb}$	CP1	1.0	13Va12
$^{132}\text{Cs}-^{133}\text{Cs}_{.992}$	223	18	229.4	1.1	0.4	o			MA1	1.0	90St25	
	232	14			-0.2	U			MA1	1.0	99Am05	
	246.9	5.9			-3.0	C			MA8	1.0	09Bo.A	
	230.8	1.3			-1.1	1	73	73	$^{132}\text{Cs}$	MA8	1.0	15At.A
$^{132}\text{Nd}-^{133}\text{Cs}_{.992}$	17147	52	17113	26	-0.7	R			MA5	1.0	00Be42	
$^{132}\text{Sn}-^{132}\text{Xe}$	13672.3	3.4	13668.8	2.1	-1.0	1	39	39	$^{132}\text{Sn}$	JY1	1.0	12Ha25
$^{132}\text{Xe}-^{131}\text{Xe}$	-930	11	-929.049	0.008	0.0	U			M16	2.5	63Da10	
$^{132}\text{Xe}-\text{C}_{10} \text{H}_{10}$	-174095.2367	0.0095	-174095.235	0.006	0.1	1	34	33	$^{132}\text{Xe}$	FS1	1.0	09Re03
$^{132}\text{Xe}-^{130}\text{Xe}$	645.724	0.014	645.738	0.009	1.0	1	40	38	$^{130}\text{Xe}$	FS1	1.0	09Re07



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{132}\text{Ba}-^{130}\text{Ba}$	-1241	4	-1259.8	2.9	-1.9	U			M17	2.5	66Be10
	-1253	68			-0.1	U			R07	1.5	68De17
$^{132}\text{Xe}-^{129}\text{Xe}$	-625.7755	0.0156	-625.772	0.004	0.2	o			FS1	1.0	05Sh38 *
	-625.7703	0.0119			-0.1	-			FS1	1.0	09Re03 *
	-625.7732	0.0125			0.1	-			FS1	1.0	09Re03 *
	-625.771	0.013			-0.1	-			FS1	1.0	09Re07
	-625.7771	0.0083			0.6	-			FS1	1.0	10Mo30
	ave.	-625.774	0.005		0.4	1	48	28 $^{129}\text{Xe}$			average
$^{132}\text{Xe}_2-^{84}\text{Kr}_3$	73816.9775	0.0594	73816.988	0.014	0.2	U			FS1	1.0	05Sh38 *
$^{132}\text{Xe}_2-^{86}\text{Kr}_3$	76478.3099	0.0412	76478.295	0.014	-0.4	1	11	6 $^{86}\text{Kr}$	FS1	1.0	05Sh38 *
$^{14}\text{N}_{10}-^{132}\text{Xe}$	126584.9632	0.0168	126584.958	0.006	-0.3	1	11	10 $^{132}\text{Xe}$	FS1	1.0	09Re03
$^{131}\text{Cs}-^{132}\text{Cs}_{.794}$ $^{127}\text{Cs}_{.206}$	-1118	16	-1094	5	0.6	F			P33	2.5	86Au02 *
$^{131}\text{Cs}-^{132}\text{Cs}_{.744}$ $^{128}\text{Cs}_{.256}$	-1200	30	-1219	5	-0.2	U			P22	2.5	82Au01
$^{130}\text{Cs}^x-^{132}\text{Cs}_{.492}$ $^{128}\text{Cs}_{.508}$	-210	40	-340	17	-1.3	U			P22	2.5	82Au01
$^{131}\text{Xe}(n,\gamma)^{132}\text{Xe}$	8936.3	1.0	8936.721	0.007	0.4	U					71Ge05
	8935	2			0.9	U					71Gr28
	8936.65	0.22			0.3	U			Bdn		06Fi.A
$^{132}\text{In}(\beta^-)^{132}\text{Sn}$	13600	400	14140	60	1.3	U					86Bj01
	14135	60				2			Stu		95Me16
$^{132}\text{Sn}(\beta^-)^{132}\text{Sb}$	3080	40	3089	3	0.2	o			Stu		77Al09
	3103	10			-1.4	o			Stu		95Me16
	3115	10			-2.6	U			Stu		99Fo01
$^{132}\text{Sb}(\beta^-)^{132}\text{Te}$	5530	70	5553	4	0.3	o			Stu		77Al09
	5486	24			2.8	U			Stu		95Me16
	5491	20			3.1	B			Stu		99Fo01
$^{132}\text{Te}(\beta^-)^{132}\text{I}$	493	4	515	3	5.6	B					65Iv01 *
	517	4			-0.4	1	76	52 $^{132}\text{I}$	Stu		99Fo01 *
$^{132}\text{I}(\beta^-)^{132}\text{Xe}$	3596	15	3575	4	-1.4	-					61De17 *
	3558	15			1.2	-					65Jo13 *
	3580	7			-0.6	-			Stu		99Fo01
	ave.	3579	6		-0.6	1	48	48 $^{132}\text{I}$			average
$^{132}\text{I}^m(\beta^-)^{132}\text{Xe}$	3685	10				2					74Di03 *
$^{132}\text{Cs}(\beta^+)^{132}\text{Xe}$	2090	25	2126.3	1.0	1.5	U					63Ta05 *
	2127.7	6.			-0.2	U					87De33 *
$^{132}\text{La}(\beta^+)^{132}\text{Ba}$	4820	100	4710	40	-1.1	-					60Wa03
	4680	50			0.6	-					67Fr02
	ave.	4710	40		0.1	1	66	66 $^{132}\text{La}$			average
* $^{132}\text{Sb}-u$	$M-A=-79870(124)$ keV for mixture gs+m at 200(30) keV										
* $^{132}\text{Xe}-\text{C}_3 \text{O}_6$	Corrected in reference of same group										
* $^{132}\text{La}-u$	$M-A=-83623(30)$ keV for mixture gs+m at 188.20 keV										
* $^{132}\text{Ce O}-^{142}\text{Sm}_{1.042}$	Original error (22 keV) increased by 23 for BaF contamination in trap										
* $^{132}\text{Pr}-u$	$M-A=-75213(28)$ keV for mixture gs+m at 30#30 keV										
* $^{132}\text{Xe}-^{129}\text{Xe}$	Corrected in reference of same group										
* $^{132}\text{Xe}-^{129}\text{Xe}$	First item $5^+$ ions; second item $3^+$ ions - considered to be independent										
* $^{132}\text{Xe}_2-^{84}\text{Kr}_3$	Corrected in reference of same group										
* $^{132}\text{Xe}_2-^{86}\text{Kr}_3$	Corrected in reference of same group										
* $^{131}\text{Cs}-^{132}\text{Cs}_{.794}$ $^{127}\text{Cs}_{.206}$	F : Rejection based on line-shape analysis										
* $^{132}\text{Te}(\beta^-)^{132}\text{I}$	$E_{\beta^-}=215(4)$ $239(4)$ respectively, to $1^+$ level at 277.86 keV										
* $^{132}\text{I}(\beta^-)^{132}\text{Xe}$	$E_{\beta^-}=2156(15)$ $2118(15)$ respectively, to $4^+$ level at 1440.323 keV										
* $^{132}\text{I}^m(\beta^-)^{132}\text{Xe}$	$E_{\beta^-}=1465(10)$ to $7^-$ level at 2214.01 level, and other $E_{\beta^-}$										
* $^{132}\text{Cs}(\beta^+)^{132}\text{Xe}$	$E_{\beta^+}=400(25)$ to $2^+$ level at 667.715 keV										
* $^{132}\text{Cs}(\beta^+)^{132}\text{Xe}$	$p^+=0.0042(0.0001)$ gives $E_{\beta^+}=438(6)$ recalculated $Q$										
*	to $2^+$ level at 667.715 keV										
											Ens054 **
											Ens054 **
											Ens054 **
											Ens054 **
											AHW **
											Ens054 **
$^{133}\text{Sb}-u$	-84766	100	-84728	3	0.2	o			GT2	2.5	08Kn.A
	-84795	129			0.2	U			GT2	2.5	08Su19
	-84702	25			-1.0	U			GS3	1.0	12Ch19

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_{10} H_{13} - {}^{133}Cs$	196266	64	196273.458	0.009	0.1	U			R07	1.5	68De17
	196279	25			-0.1	U			R07	1.5	68De17
$C_9 {}^{13}C H_{12} - {}^{133}Cs$	191796	34	191803.261	0.009	0.1	U			R07	1.5	68De17
$C_8 O N H_7 - {}^{133}Cs$	147321	25	147311.889	0.009	-0.2	U			R07	1.5	68De17
$C_7 {}^{13}C N O H_6 - {}^{133}Cs$	142835	31	142841.692	0.009	0.1	U			R07	1.5	68De17
${}^{133}Cs - {}^{85}Rb_{1.565}$	43500	13	43501.027	0.011	0.1	U			MA5	1.0	00Be42
	43499.3	1.6			1.1	U			MA8	1.0	07Ke09
	43500.9	6.7			0.0	U			MA8	1.0	02Ke.A
	43500.1	6.7			0.1	U			MA8	1.0	09Na.A
	43470	47			0.7	U			MA9	1.0	09Na.A
	43501.2	1.7			-0.1	U			MA8	1.0	11He10
${}^{133}Cs - u$	-94548.41	0.41	-94548.039	0.009	0.9	U			ST2	1.0	99Ca46 *
${}^{133}La - u$	-91810	120	-91780	30	0.2	U			GS1	1.0	00Ra23
	-91782	30				2			GS2	1.0	05Li24
${}^{133}Ce - u$	-88471	32	-88480	18	-0.3	2			GS2	1.0	05Li24 *
${}^{133}Ce O - {}^{142}Sm_{1.049}$	-4618	21	-4615	18	0.2	R			MA7	1.0	01Bo59 *
${}^{133}Pr - u$	-83663	30	-83669	13	-0.2	R			GS2	1.0	05Li24
${}^{133}Nd - u$	-77652	50				2			GS2	1.0	05Li24 *
${}^{133}Pm - u$	-70218	54				2			GS2	1.0	05Li24 *
${}^{133}Sb - {}^{136}Xe_{0.978}$	6022	10	6016	3	-0.6	1	11	11 ${}^{133}Sb$	CP1	1.0	12Va02
${}^{133}Sb - {}^{130}Xe_{1.023}$	13984.7	4.0	13982	3	-0.7	1	70	70 ${}^{133}Sb$	JY1	1.0	12Ha25
${}^{133}Te - {}^{130}Xe_{1.023}$	9672.1	2.3	9673.3	2.2	0.5	1	93	93 ${}^{133}Te$	JY1	1.0	12Ha25
${}^{133}Sn - {}^{134}Xe_{0.993}$	17856.5	2.4	17858.5	2.0	0.8	1	73	73 ${}^{133}Sn$	JY1	1.0	12Ha25
${}^{133}Sn {}^{34}S - {}^{133}Cs_{1.256}$	10562	25	10533.1	2.0	-1.2	U			MA8	1.0	08Dw01
${}^{133}Sn - {}^{133}Cs$	18467.0	3.9	18461.8	2.0	-1.3	1	27	27 ${}^{133}Sn$	CP1	1.0	13Va12
${}^{133}Sb - {}^{133}Cs$	9818	13	9820	3	0.2	U			CP1	1.0	13Va12
${}^{133}Te - {}^{133}Cs_{1.000}$	5551.4	6.9	5511.4	2.2	-5.8	B			CP1	1.0	13Va12
${}^{133}I - {}^{133}Cs$	2375.4	6.9				2			CP1	1.0	13Va12
${}^{133}Pr - {}^{133}Cs$	10877	15	10879	13	0.1	2			MA5	1.0	00Be42
${}^{133}Cs - C_3 O_6$	-64035.786	0.026	-64035.757	0.009	1.1	1	11	11 ${}^{133}Cs$	MI2	1.0	99Br47
${}^{133}Cs - C_{10} H_{12}$	-188448.445	0.057	-188448.426	0.009	0.3	U			MI2	1.0	99Br47
${}^{133}Cs - {}^{132}Xe$	1296.8803	0.0103	1296.874	0.007	-0.6	1	50	45 ${}^{133}Cs$	FS1	1.0	10Mo30
${}^{133}Cs - {}^{129}Xe$	671.1007	0.0103	671.102	0.007	0.1	1	50	44 ${}^{133}Cs$	FS1	1.0	10Mo30
${}^{133}Cs(\gamma, n) {}^{132}Cs$	-8988	33	-8989.6	1.0	0.0	U			Phi		60Ge01
	-8986	2			-1.8	1	27	27 ${}^{132}Cs$	MMn		85Ts02
${}^{132}Ba(n, \gamma) {}^{133}Ba$	7189.91	0.36	7189.9	0.4	0.0	1	100	98 ${}^{132}Ba$	MMn		90Is07 Z
${}^{132}Ba(d, p) {}^{133}Ba$	4977	15	4965.3	0.4	-0.8	U			ANL		70Vo04
${}^{133}Sn(\beta^-) {}^{133}Sb$	7830	70	8050	4	3.1	B			Stu		83B116 *
	8013	50			0.7	o			Stu		92Sp.A *
	7990	25			2.4	U			Stu		95Me16
	3966	50	4014	4	1.0	o			Stu		70Ru.A *
${}^{133}Sb(\beta^-) {}^{133}Te$	4003	10			1.1	o			Stu		95Me16
	4002	7			1.7	1	25	18 ${}^{133}Sb$	Stu		99Fo01
	2960	100	2921	7	-0.4	U					68Mc09
	2876	100			0.5	U					68Pa03 *
${}^{133}Te(\beta^-) {}^{133}I$	3392	100			-4.7	C			Stu		70Ru.A *
	2890	15			2.1	o			Stu		95Me16
	2942	24			-0.9	U			Stu		99Fo01
	1800	50	1785	7	-0.3	U					59Ho97 *
	1760	30			0.8	U					66Ei01 *
${}^{133}I(\beta^-) {}^{133}Xe$	1757	4			7.1	B			Stu		99Fo01
	${}^{133}Xe(\beta^-) {}^{133}Cs$	428.0	4.	427.4	2.4	-0.2	2				52Be55 *
	427.0	3.			0.1	2					61Er04 *
${}^{133}Ba(\epsilon) {}^{133}Cs$	424	11			0.3	U			Stu		99Fo01
	517.3	1.0	517.3	1.0	0.0	1	98	98 ${}^{133}Ba$			67Sc10 *
	498	5			3.9	F					68Mc06 *
	486	2			15.7	F					69Bo49 *
	521	5			-0.7	U					69To14 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{133}\text{La}(\beta^+)^{133}\text{Ba}$	2230	200	2059	28	-0.9	U					50Na09 *
* $^{133}\text{Cs}-u$	As revised in reference. Original: $-94548.20(0.28) \mu\text{u}$										
* $^{133}\text{Ce}-u$	$M-A=-82392(28)$ keV for mixture gs+m at 37.2 keV										
* $^{133}\text{Ce O}-^{142}\text{Sm}_{1.049}$	$D_M=-4599(16) \mu\text{u}$ for mixture gs+m at 37.2 keV; $M-A=-87150(16)$ keV										
* $^{133}\text{Nd}-u$	$M-A=-72268(28)$ keV for mixture gs+m at 127.97 keV										
* $^{133}\text{Pm}-u$	$M-A=-65342(33)$ keV for mixture gs+m at 129.7 keV										
* $^{133}\text{Sn}(\beta^-)^{133}\text{Sb}$	$E_{\beta^-}=6870(70)$ to $5/2^+$ level at 962.30 keV										
* $^{133}\text{Sn}(\beta^-)^{133}\text{Sb}$	Private communication to reference										
* $^{133}\text{Sb}(\beta^-)^{133}\text{Te}$	$E_{\beta^-}=1210(50)$ to $(5/2^+)$ level at 2755.51 keV; re-evaluated										
* $^{133}\text{Te}(\beta^-)^{133}\text{I}$	$Q_{\beta^-}=3210(100)$ from $^{133}\text{Te}^m$ at 334.26 keV										
*	reported as belonging to ground state, reinterpreted										
* $^{133}\text{Te}(\beta^-)^{133}\text{I}$	$E_{\beta^-}=850(100)$ to $(3/2^+, 5/2^+)$ level at 2541.74 keV										
* $^{133}\text{I}(\beta^-)^{133}\text{Xe}$	$E_{\beta^-}=1270(50) 1230(30)$ respectively, to $5/2^+$ level at 529.872 keV										
* $^{133}\text{Xe}(\beta^-)^{133}\text{Cs}$	$E_{\beta^-}=347(4) 346(3)$ respectively, to $5/2^+$ level at 80.9979 keV										
* $^{133}\text{Ba}(\epsilon)^{133}\text{Cs}$	From $L/K=0.371(0.007)$ to $1/2^+$ level at 437.0113; recalculated $Q$										
*	and $L/K=0.221(0.005)$ to $3/2^+$ level at 383.8491 keV; $Q=521(5)$ keV										
* $^{133}\text{Ba}(\epsilon)^{133}\text{Cs}$	$F$ : badly resolved L-peak										
* $^{133}\text{Ba}(\epsilon)^{133}\text{Cs}$	$L/K=0.67(0.15) LM/K=1.11(0.05) 0.45(0.04)$ respectively, to $1/2^+$ 437.0113 keV										
* $^{133}\text{La}(\beta^+)^{133}\text{Ba}$	$E_{\beta^+}=1200(200)$ to $3/2^+$ level at 12.327 keV										
$^{134}\text{Te}-u$	-88844	130	-88603.6	2.9	0.7	U			GT2	2.5	08Su19
$\text{C}_{10} \text{H}_{14}-^{134}\text{Xe}$	204155.5	3.2	204157.417	0.010	0.2	U			M16	2.5	63Da10
$^{134}\text{Xe}-u$	-94634.4	5.4	-94606.966	0.010	2.0	U			ACC	2.5	90Me08
$^{134}\text{Xe}-\text{C}^{13}\text{C}^{35}\text{Cl}^{37}\text{Cl}_2$	1381.76	0.60	1380.33	0.12	-1.6	U			H47	1.5	94Hy01
$\text{C}_{10} \text{H}_{14}-^{134}\text{Ba}$	205025	20	205042.1	0.3	0.3	U			M17	2.5	66Be10
	205010	46			0.5	U			R07	1.5	68De17
$\text{C}_{11} \text{H}_2-^{134}\text{Ba}$	111125	48	111141.7	0.3	0.2	U			R07	1.5	68De17
$\text{C}_8 \text{N O H}_8-^{134}\text{Ba}$	156063	78	156080.5	0.3	0.1	U			R07	1.5	68De17
$\text{C}_{12} \text{H}_6-^{134}\text{Ba O}$	147531	64	147527.2	0.3	0.0	U			R07	1.5	68De17
$^{134}\text{La}-u$	-91456	34	-91486	21	-0.9	2			GS2	1.0	05Li24
$^{134}\text{Ce}-u$	-91190	130	-91072	22	0.9	U			GS1	1.0	00Ra23
	-91056	30			-0.5	2			GS2	1.0	05Li24
$^{134}\text{Ce O}-^{142}\text{Sm}_{1.056}$	-6631	32	-6613	22	0.6	R			MA7	1.0	01Bo59 *
$^{134}\text{Pr}-u$	-84285	37	-84303	22	-0.5	2			GS2	1.0	05Li24 *
$^{134}\text{Nd}-u$	-81234	30	-81210	13	0.8	R			GS2	1.0	05Li24
$^{134}\text{Pm}-u$	-71647	62				2			GS2	1.0	05Li24 *
$^{134}\text{Sb}-^{130}\text{Xe}_{1.031}$	20016.8	2.2	20017.5	1.8	0.3	2			JY1	1.0	12Ha25
	20019.2	3.3			-0.5	2			JY1	1.0	13Ka08 *
$^{134}\text{Te}-^{130}\text{Xe}_{1.031}$	10877.1	3.5	10878.2	2.9	0.3	1	71	71	$^{134}\text{Te}$	1.0	12Ha25
$^{134}\text{Sb}-^{136}\text{Xe}_{.985}$	11906	36	11929.4	1.8	0.7	U			CP1	1.0	12Va02 *
$^{134}\text{Te}-^{136}\text{Xe}_{.985}$	2791.4	6.5	2790.1	2.9	-0.2	1	21	21	$^{134}\text{Te}$	1.0	12Va02
$^{134}\text{Xe}-^{132}\text{Xe}_{1.015}$	2675.6193	0.0084	2675.619	0.008	0.0	1	100	100	$^{134}\text{Xe}$	1.0	13Ho22
$^{134}\text{Sn}^{34}\text{S}-^{133}\text{Cs}_{1.263}$	16080	160	15962	3	-0.7	U			MA8	1.0	08Dw01
$^{134}\text{Sn}-^{133}\text{Cs}_{1.008}$	23974	17	23985	3	0.6	U			CP1	1.0	13Va12
$^{134}\text{Sb}-^{133}\text{Cs}_{1.008}$	15850	11	15840.1	1.8	-0.9	U			CP1	1.0	13Va12
$^{134}\text{I}-^{133}\text{Cs}_{1.008}$	5083.0	6.8	5080	5	-0.4	1	59	59	$^{134}\text{I}$	1.0	13Va12
$^{134}\text{Cs}-^{133}\text{Cs}_{1.008}$	2025	18	2022.929	0.015	-0.1	o			MA1	1.0	90St25
	2033	14			-0.7	U			MA1	1.0	99Am05
$^{134}\text{Pr}-^{133}\text{Cs}_{1.008}$	10992	27	11001	22	0.3	R			MA5	1.0	00Be42 *
$^{134}\text{Nd}-^{133}\text{Cs}_{1.008}$	14100	14	14095	13	-0.4	2			MA5	1.0	00Be42
$^{134}\text{Sn}-^{134}\text{Xe}$	23287.4	3.4				2			JY1	1.0	12Ha25
$^{134}\text{Ba}-\text{C}_{10} \text{H}_{13}$	-197229	20	-197217.0	0.3	0.2	U			M17	2.5	66Be10
$^{134}\text{Ba}-^{132}\text{Ba}$	-553	4	-552.7	1.2	0.0	U			M17	2.5	66Be10
	-550	121			0.0	U			R07	1.5	68De17
$^{131}\text{Cs}-^{134}\text{Cs}_{.244} \ ^{130}\text{Cs}_{.756}^x$	-1313	50	-1182	14	1.1	U			P22	2.5	82Au01 *
$^{133}\text{Cs}(n,\gamma)^{134}\text{Cs}$	6891.540	0.017	6891.540	0.014	0.0	-			MMn		84Ke11 Z
	6891.540	0.027			0.0	-			ILn		87Bo24 Z
	6891.39	0.14			1.1	U			Bdn		06Fi.A
ave.	6891.540	0.014			0.0	1	100	100	$^{134}\text{Cs}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{134}\text{Sn}(\beta^-)^{134}\text{Sb}$	7370	90	7587	4	2.4	U			Stu		95Me16
$^{134}\text{Sb}(\beta^-)^{134}\text{Te}$	8306	210	8513	3	1.0	o			Stu		72Ke21 *
	7960	240			2.3	o			Stu		77Lu06 *
	8310	80			2.5	U			Bwg		79Ke02 *
	8390	45			2.7	U			Stu		95Me16
$^{134}\text{Te}(\beta^-)^{134}\text{I}$	1560	90	1510	5	-0.6	U			Stu		77Lu06 *
	1550	30			-1.3	U			Stu		95Me16
	1513	7			-0.5	1	50	41 $^{134}\text{I}$	Stu		99Fo01
$^{134}\text{I}(\beta^-)^{134}\text{Xe}$	4170	60	4082	5	-1.5	U					61Jo08 *
	4175	15			-6.2	B			Stu		95Me16
	4052	8			3.8	B			Stu		99Fo01
$^{134}\text{Cs}(\beta^-)^{134}\text{Ba}$	2058.6	0.4	2058.7	0.3	0.2	1	58	58 $^{134}\text{Ba}$			68Hs01 *
$^{134}\text{La}(\beta^+)^{134}\text{Ba}$	3772	50	3731	20	-0.8	R					65Bi12
	3692	30			1.3	R					73Al20
$^{134}\text{Ce}(\epsilon)^{134}\text{La}$	500	200	386	29	-0.6	U					76Gr09 *
$^{134}\text{Pr}(\beta^+)^{134}\text{Ce}$	6190	90	6305	29	1.3	U			Dbn		95Ve08 *
$^{134}\text{Nd}(\beta^+)^{134}\text{Pr}$	2770	150	2882	24	0.7	U					77Ko.B
$^{134}\text{Pm}(\beta^+)^{134}\text{Nd}$	9170	200	8910	60	-1.3	U			Dbn		95Ve08 *
* $^{134}\text{Ce O}-^{142}\text{Sm}_{1.056}$	Original error (22 keV) increased by 23 for BaF contamination in trap										
* $^{134}\text{Pr-u}$	$M - A = -78477(28)$ keV for mixture gs+m at 68(1) keV										
* $^{134}\text{Pm-u}$	$M - A = -66739(30)$ keV for mixture gs+m at 0#100 keV										
* $^{134}\text{Sb}-^{130}\text{Xe}_{1.031}$	$D_M = 20318.7(3.1)$ $\mu\text{u}$ for $^{134}\text{Sb}^m$ at 279(1) keV; $M - A = -73740.0(2.9)$ keV										
* $^{134}\text{Sb}-^{136}\text{Xe}_{985}$	$D_M = 12206(36)$ $\mu\text{u}$ for $^{134}\text{Sb}^m$ at 279(1) keV; $M - A = -73762(33)$ keV										
*	assuming high spin is favored, as for $^{136}\text{P}^m$										
* $^{134}\text{Pr}-^{133}\text{Cs}_{1.008}$	Most certainly gs. Mixture with isomer not completely excluded										
* $^{134}\text{Pr}-^{133}\text{Cs}_{1.008}$	$D_M = 11029(16)$ $\mu\text{u}$ for mixture gs+m at 68(1) keV; $M - A = -78503(15)$ keV										
* $^{131}\text{Cs}-^{134}\text{Cs}_{244}$ $^{130}\text{Cs}^x$	$D_M = -1330(50)$ keV for mixture gs+m at 138.7441 keV										
* $^{134}\text{Sb}(\beta^-)^{134}\text{Te}$	$E_{\beta^-} = 8400(300)$ , and 6800(300) from $^{134}\text{Sb}^m$ at 279(1) to $^{134}\text{Te}^m$ at 1691.34										
* $^{134}\text{Sb}(\beta^-)^{134}\text{Te}$	$E_{\beta^-} = 5840(240)$ from $^{134}\text{Sb}^m$ at 279(1) to (6) $^+$ level at 2397.70 keV										
* $^{134}\text{Sb}(\beta^-)^{134}\text{Te}$	$E_{\beta^-} = 8420(120)$ , and 6710(210), 6980(210), 6070(150) from $^{134}\text{Sb}^m$ at 279(1)										
*	to 6 $^+$ levels at 1691.34, 1691.34, 2397.70 keV										
* $^{134}\text{Te}(\beta^-)^{134}\text{I}$	$E_{\beta^-} = 730(110)$ 610(160) to 1 $^+$ levels at 846.688 923.432 keV										
* $^{134}\text{I}(\beta^-)^{134}\text{Xe}$	$E_{\beta^-} = 2410(60)$ 1680(60) 1250(60) to 4 $^+$ levels at 1731.17 2588.46 2867.38 keV										
* $^{134}\text{Cs}(\beta^-)^{134}\text{Ba}$	$E_{\beta^-} = 658.0(0.4)$ to 4 $^+$ level at 1400.590 keV										
* $^{134}\text{Ce}(\epsilon)^{134}\text{La}$	LK=0.798(0.04); also $Q_{\beta^+} > 375$ keV										
* $^{134}\text{Pr}(\beta^+)^{134}\text{Ce}$	$E_{\beta^+} = 4120(90)$ to 4 $^+$ level at 1048.68 keV										
* $^{134}\text{Pm}(\beta^+)^{134}\text{Nd}$	$E_{\beta^+} = 7360(200)$ to 4 $^+$ level at 788.92 keV										
$^{135}\text{Sb-u}$	-74932	103	-74815.6	2.8	0.5	o			GT2	2.5	08Kn.A
	-74943	130			0.4	U			GT2	2.5	08Su19
$^{135}\text{Te-u}$	-83643	106	-83445.3	1.8	0.7	U			GT2	2.5	08Kn.A
	-83441	132			0.0	U			GT2	2.5	08Su19
$\text{C}_8 \text{ N O H}_9 - ^{135}\text{Ba}$	162731	48	162725.3	0.3	-0.1	U			R07	1.5	68De17
$\text{C}_{11} \text{ H}_3 - ^{135}\text{Ba}$	117822	77	117786.5	0.3	-0.3	U			R07	1.5	68De17
$\text{C}_{12} \text{ H}_7 - ^{135}\text{Ba O}$	154160	46	154172.0	0.3	0.2	U			R07	1.5	68De17
$^{135}\text{Ce-u}$	-90779	30	-90839	11	-2.0	1	13	13 $^{135}\text{Ce}$	GS2	1.0	05Li24 *
$^{135}\text{Pr-u}$	-86897	30	-86888	13	0.3	R			GS2	1.0	05Li24
$^{135}\text{Nd-u}$	-81800	130	-81819	21	-0.1	o			GS1	1.0	00Ra23
	-81811	36			-0.2	R			GS2	1.0	05Li24 *
$^{135}\text{Pm-u}$	-75204	81				2			GS2	1.0	05Li24 *
$^{135}\text{Sm-u}$	-67480	166				2			GS2	1.0	05Li24 *
$^{135}\text{Sn}-^{130}\text{Xe}_{1.038}$	35065.9	3.3				2			JY1	1.0	12Ha25
$^{135}\text{Sb}-^{130}\text{Xe}_{1.038}$	25342.4	3.1	25341.7	2.8	-0.2	1	84	84 $^{135}\text{Sb}$	JY1	1.0	12Ha25
$^{135}\text{Te}-^{130}\text{Xe}_{1.038}$	16713.0	2.9	16712.0	1.8	-0.3	1	41	41 $^{135}\text{Te}$	JY1	1.0	12Ha25
$^{135}\text{Sn}-^{133}\text{Cs}_{1.015}$	30927	37	30875	3	-1.4	U			CP1	1.0	13Va12

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{135}\text{Sb}-^{133}\text{Cs}_{1.015}$	21146.8	7.0	21150.6	2.8	0.5	1	16	16 $^{135}\text{Sb}$	CP1	1.0	13Va12
$^{135}\text{Te}-^{133}\text{Cs}_{1.015}$	12520.3	2.4	12521.0	1.8	0.3	1	59	59 $^{135}\text{Te}$	CP1	1.0	13Va12
$^{135}\text{I}-^{133}\text{Cs}_{1.015}$	6026.6	2.3	6025.6	2.2	-0.4	1	93	93 $^{135}\text{I}$	CP1	1.0	13Va12
$^{135}\text{Cs}-^{133}\text{Cs}_{1.015}$	1958	18	1943.5	1.1	-0.8	o			MA1	1.0	90St25
	1957	14			-1.0	U			MA1	1.0	99Am05
$^{135}\text{Pr}-^{133}\text{Cs}_{1.015}$	9080	14	9078	13	-0.1	2			MA5	1.0	00Be42
$^{135}\text{Nd}-^{133}\text{Cs}_{1.015}$	14144	25	14148	21	0.1	2			MA5	1.0	00Be42 *
$^{135}\text{Te}-^{136}\text{Xe}_{.993}$	8686	10	8690.7	1.8	0.5	U			CP1	1.0	12Va02
$^{135}\text{I}-^{136}\text{Xe}_{.993}$	2186.3	8.3	2195.4	2.2	1.1	U			CP1	1.0	12Va02
$^{135}\text{Ba}-\text{C}_{10}\text{H}_{14}$	-203860	20	-203861.8	0.3	0.0	U			M17	2.5	66Be10
$^{135}\text{Ba}-^{134}\text{Ba}$	1177	2	1180.21	0.11	0.6	U			M17	2.5	66Be10
	1161	70			0.2	U			R07	1.5	68De17
	1168	78			0.1	U			R07	1.5	68De17
$^{134}\text{Cs}(n,\gamma)^{135}\text{Cs}$	8762	1	8761.8	1.0	-0.2	1	98	98 $^{135}\text{Cs}$	ILn		92Ul.A
$^{134}\text{Ba}(n,\gamma)^{135}\text{Ba}$	6973.2	0.4	6971.96	0.10	-3.1	C			BNn		77Ko.A
	6972.17	0.18			-1.2	-			MMn		90Is07 Z
	6971.84	0.17			0.7	-			Ltn		93Bo01 Z
	6973.24	0.22			-5.8	B			BNn		93Ch21
	6971.87	0.18			0.5	-			Bdn		06Fi.A
$^{134}\text{Ba}(d,p)^{135}\text{Ba}$	4746	15	4747.40	0.10	0.1	U			ANL		70Vo04
$^{134}\text{Ba}(n,\gamma)^{135}\text{Ba}$	ave.	6971.96	0.10	6971.96	0.10	0.1	1	97	55 $^{135}\text{Ba}$		average
$^{135}\text{Tb}(p)^{134}\text{Gd}$	1188	7				3			Arp		04Wo07
$^{135}\text{Sb}(\beta^-)^{135}\text{Te}$	8120	50	8038	3	-1.6	U			Stu		89Ho08
$^{135}\text{Te}(\beta^-)^{135}\text{I}$	5950	240	6050.4	2.7	0.4	o			Stu		77Lu06
	5950	100			1.0	o			Bwg		79Ke02 *
	5970	200			0.4	U					85Sa15 *
	5960	100			0.9	U			Bwg		87Gr.A
	5888	13			12.5	B			Stu		07Fo02
$^{135}\text{I}(\beta^-)^{135}\text{Xe}$	2780	80	2634	4	-1.8	U					70Ma19
	2590	50			0.9	U			Stu		76Lu04 *
	2627	6			1.2	1	42	34 $^{135}\text{Xe}$	Stu		99Fo01
$^{135}\text{Xe}(\beta^-)^{135}\text{Cs}$	1155	10	1168	4	1.3	-					52Be55 *
	1167	5			0.3	-			Stu		99Fo01 *
	ave.	1165	4			0.9	1	68	66 $^{135}\text{Xe}$		average
$^{135}\text{Cs}(\beta^-)^{135}\text{Ba}$	205	5	268.9	1.0	12.8	B					53Li01
$^{135}\text{La}(\beta^+)^{135}\text{Ba}$	1200	10	1207	9	0.7	1	89	89 $^{135}\text{La}$			71Ba18 *
$^{135}\text{Ce}(\beta^+)^{135}\text{La}$	2027	5	2027	5	0.0	-					76Ga.A *
	2016	13			0.9	-					81Sa09 *
	ave.	2026	5			0.3	1	98	87 $^{135}\text{Ce}$		average
$^{135}\text{Pr}(\beta^+)^{135}\text{Ce}$	3720	150	3680	16	-0.3	U					54Ha68 *
$^{135}\text{Pm}^m(\beta^+)^{135}\text{Nd}$	6040	150	6390#	50#	2.3	D			Dbn		95Ve08 *
* $^{135}\text{Ce}-u$	$M-A=-84114(28)$ keV for $^{135}\text{Ce}^m$ at 445.81 keV										Nub16b **
* $^{135}\text{Nd}-u$	$M-A=-76174(28)$ keV for mixture gs+m at 64.95 keV										Nub16b **
* $^{135}\text{Pm}-u$	$M-A=-69952(28)$ keV for mixture gs+m at 200#80 keV										Nub16b **
* $^{135}\text{Sm}-u$	$M-A=-62857(38)$ keV for mixture gs+m at 0#300 keV										Nub16b **
* $^{135}\text{Nd}-^{133}\text{Cs}_{1.015}$	$D_M=14179(14)$ $\mu$ u for gs+m mixture at 64.95 keV; $M-A=-76185(13)$ keV										Nub16b **
* $^{135}\text{Te}(\beta^-)^{135}\text{I}$	$E_{\beta^-}=5120(120)$ to $5/2^-$ level at 870.52 keV and other $E_{\beta^-}$										Ens083 **
* $^{135}\text{Te}(\beta^-)^{135}\text{I}$	$E_{\beta^-}=5370(100)$ to $5/2^+$ level at 603.68 keV										Ens083 **
* $^{135}\text{I}(\beta^-)^{135}\text{Xe}$	$E_{\beta^-}=1320(50)$ to $5/2^+$ level at 1260.416 keV, and other $E_{\beta^-}$										Ens083 **
* $^{135}\text{Xe}(\beta^-)^{135}\text{Cs}$	$E_{\beta^-}=905(10)$ 917(5) respectively, to $5/2^+$ level at 249.767 keV										Ens083 **
* $^{135}\text{La}(\beta^+)^{135}\text{Ba}$	$p^+=7(1)\times 10^{-5}$ , from reanalysis										AHW **
* $^{135}\text{La}(\beta^+)^{135}\text{Ba}$	But 65Mo05 says $p^+ < 0.002\%$ ( $2\times 10^{-5}$ ) or $E_{\beta^+} < 125$ keV										65Mo05 **
* $^{135}\text{Ce}(\beta^+)^{135}\text{La}$	$E_{\beta^+}=705(5)$ 694(13) respectively, to $1/2^+$ level at 300.052 keV										Ens083 **
* $^{135}\text{Pr}(\beta^+)^{135}\text{Ce}$	$E_{\beta^+}=2500(100)$ to levels ( $5/2^+$ ) 296.11 and $3/2^+$ 82.67 keV, roughly equal										Ens083 **
* $^{135}\text{Pm}^m(\beta^+)^{135}\text{Nd}$	$E_{\beta^+}=4920(150)$ to mixture ground state and ( $11/2^-$ ) level at 198.5 keV										Ens083 **
* $^{135}\text{Pm}^m(\beta^+)^{135}\text{Nd}$	Trends from Mass Surface TMS suggest $^{135}\text{Pm}^m$ 350 less bound (see Nubase)										GAu **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{136}\text{Te}-u$	-79945	25	-79898.8	2.4	1.8	U			GS3	1.0	12Ch19
$^{136}\text{Xe}-^{120}\text{Sn}_{1,133}$	18019.8	3.1	18019.8	1.1	0.0	U			JY1	1.0	11Ha48
$\text{C}_{10}\text{H}_{16}-^{136}\text{Xe}$	217982.	3.9	217986.039	0.007	0.4	U			M16	2.5	63Da10
$^{136}\text{Xe}-u$	-92793.6	9.0	-92785.524	0.007	0.4	U			ACC	2.5	90Me08
$^{136}\text{Xe}-^{28}\text{Si}_4\text{D}_{12}$	-169712.9828	0.0162	-169713.001	0.007	-1.1	1	20	18	$^{136}\text{Xe}$	1.0	07Re03
$\text{C}_{11}\text{H}_4-^{136}\text{Ba}$	126737	56	126724.2	0.3	-0.2	U			R07	1.5	68De17
$\text{C}_8\text{N O H}_{10}-^{136}\text{Ba}$	171635	56	171663.0	0.3	0.3	U			R07	1.5	68De17
$\text{C}_{12}\text{H}_8-^{136}\text{Ba O}$	163094	40	163109.7	0.3	0.3	U			R07	1.5	68De17
$^{136}\text{La}-u$	-92394	88	-92370	60	0.3	2			GS2	1.0	05Li24 *
$\text{C}_{10}\text{H}_{16}-^{136}\text{Ce}$	218128	50	218071.1	0.4	-0.3	U			R05	4.0	65De13
$\text{C}_{12}\text{H}_8-^{136}\text{Ce O}$	160563	36	160556.2	0.4	0.0	U			R05	4.0	65De13
$^{136}\text{Nd}-u$	-85044	30	-85024	13	0.7	R			GS2	1.0	05Li24
$^{136}\text{Pm}-u$	-76378	79	-76400	70	-0.3	2			GS2	1.0	05Li24 *
$^{136}\text{Sm}-u$	-71768	30	-71724	13	1.5	R			GS2	1.0	05Li24
$^{136}\text{Sb}-^{130}\text{Xe}_{1,046}$	31675.1	6.8	31678	6	0.5	1	85	85	$^{136}\text{Sb}$	1.0	12Ha25
$^{136}\text{Te}-^{130}\text{Xe}_{1,046}$	21029.9	3.1	21030.4	2.4	0.2	1	62	62	$^{136}\text{Te}$	1.0	12Ha25
$^{136}\text{Sb}-^{133}\text{Cs}_{1,023}$	27489	16	27472	6	-1.1	1	15	15	$^{136}\text{Sb}$	1.0	13Va12
$^{136}\text{Te}-^{133}\text{Cs}_{1,023}$	16827.7	6.8	16823.8	2.4	-0.6	1	13	13	$^{136}\text{Te}$	1.0	13Va12
$^{136}\text{Xe}-^{133}\text{Cs}_{1,023}$	3936.5	1.9	3937.121	0.011	0.3	U			MA8	1.0	09Ne11
$^{136}\text{Cs}-^{133}\text{Cs}_{1,023}$	4007	18	4034.2	2.0	1.5	o			MA1	1.0	90St25
	4021	14			0.9	U			MA1	1.0	99Am05
$^{136}\text{Pr}-^{133}\text{Cs}_{1,023}$	9418	15	9400	12	-1.2	1	67	67	$^{136}\text{Pr}$	1.0	00Be42
$^{136}\text{Nd}-^{133}\text{Cs}_{1,023}$	11703	14	11699	13	-0.3	2			MA5	1.0	00Be42
$^{136}\text{Pm}^m-^{133}\text{Cs}_{1,023}$	20429	100				2			MA5	1.0	00Be42 *
$^{136}\text{Sm}-^{133}\text{Cs}_{1,023}$	25009	15	24998	13	-0.7	2			MA5	1.0	00Be42
$^{136}\text{Xe}-^{134}\text{Xe}_{1,015}$	3245.8	3.8	3240.546	0.012	-1.4	o			MA8	1.0	05He.A
	3244.3	4.0			-0.9	U			MA8	1.0	06He29
$^{136}\text{Te}-^{136}\text{Xe}$	12887.9	5.0	12886.7	2.4	-0.2	1	24	24	$^{136}\text{Te}$	1.0	12Va02
$^{136}\text{Pm}-^{136}\text{Xe}$	7611.2	4.9				2			CP1	1.0	12Va02 *
$^{136}\text{Xe}-^{136}\text{Ba}$	2639.6	0.6	2638.5	0.3	-0.7	-			H49	2.5	10Mc04
	2638.62	0.52			-0.2	-			JY1	1.0	11Ko03
ave.	2638.7	0.5			-0.4	1	45	45	$^{136}\text{Ba}$		average
$^{136}\text{Ce}-^{136}\text{Ba}$	2553.46	0.29	2553.48	0.29	0.1	1	100	100	$^{136}\text{Ce}$	1.0	11Ko03
$^{136}\text{Xe}-^{13}\text{C}_3\text{O}_6$	-72337.7553	0.0109	-72337.747	0.007	0.8	-			FS1	1.0	07Re03
	-72337.7464	0.0109			0.0	-			FS1	1.0	07Re03
ave.	-72337.751	0.008			0.5	1	82	82	$^{136}\text{Xe}$		average
$^{136}\text{Ba}-^{135}\text{Ba}$	-1115	3	-1112.65	0.04	0.3	U			M17	2.5	66Be10
	-1119	50			0.1	U			R07	1.5	68De17
	-1074	50			-0.5	U			R07	1.5	68De17
$^{136}\text{Ba}-^{134}\text{Ba}$	67	5	67.56	0.12	0.0	U			M17	2.5	66Be10
	69	128			0.0	U			R07	1.5	68De17
	72	78			0.0	U			R07	1.5	68De17
$\text{N}_{10}-^{136}\text{Xe}$	123525.5778	0.0235	123525.568	0.007	-0.4	U			FS1	1.0	07Re03
$^{136}\text{Te}(\beta^-n)^{135}\text{I}$	1285	50	1283	3	0.0	U					84Kr.B
$^{136}\text{Xe}(d,^3\text{He})^{135}\text{I}$	-4438	30	-4445.5	2.1	-0.3	U			Oak		71Wi04
$^{136}\text{Xe}(d,t)^{135}\text{Xe}$	-1723	40	-1830	4	-2.7	U			Oak		68Mo21
$^{135}\text{Ba}(n,\gamma)^{136}\text{Ba}$	9106.4	0.8	9107.74	0.04	1.7	U					69Ge07
	9107.74	0.04			0.0	-			MMn		90Is07 Z
	9107.73	0.19			0.1	-			Bdn		06Fi.A
$^{135}\text{Ba}(d,p)^{136}\text{Ba}$	6886	15	6883.17	0.04	-0.2	U			ANL		70Vo04
$^{135}\text{Ba}(n,\gamma)^{136}\text{Ba}$	9107.74	0.04	9107.74	0.04	0.0	1	100	55	$^{136}\text{Ba}$		average
$^{136}\text{Te}(\beta^-)^{136}\text{I}$	5100	150	5120	14	0.1	U					77Sc21
	5095	100			0.2	U			Bwg		87Gr.A
	5086	20			1.7	1	50	50	$^{136}\text{I}$		07Fo02
$^{136}\text{I}(\beta^-)^{136}\text{Xe}$	6960	100	6884	14	-0.8	U					59Jo37 *
	6690	150			1.3	U			Stu		76Lu04 *
	6925	70			-0.6	U			Bwg		87Gr.A
	6850	20			1.7	1	50	50	$^{136}\text{I}$		07Fo02

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{136}\text{I}^m(\beta^-)^{136}\text{Xe}$	7100	230	7090	5	0.0	o			Stu		76Lu04 *
	7705	120			-5.1	C			Bwg		87Gr.A
	7051	12			3.2	B			Stu		07Fo02
$^{136}\text{Cs}(\beta^-)^{136}\text{Ba}$	2548.1	2.0	2548.2	1.9	0.1	2					54OI05 *
	2549	5			-0.2	2					65Re07 *
$^{136}\text{La}(\beta^+)^{136}\text{Ba}$	2870	70	2850	50	-0.3	R					59Gi50
$^{136}\text{Pr}(\beta^+)^{136}\text{Ce}$	5084	50	5168	11	1.7	U					68Zh04 *
	5114	75			0.7	U					71Ke07 *
	5134	20			1.7	1	33	33 $^{136}\text{Pr}$	IRS		83Al.B *
$^{136}\text{Nd}(\beta^+)^{136}\text{Pr}$	2501	50	2141	16	-7.2	B					68Zh04 *
	2211	25			-2.8	B					75Br16 *
$^{136}\text{Pm}(\beta^+)^{136}\text{Nd}$	7850	200	8030	70	0.9	R			IRS		83Al06 *
* $^{136}\text{La-u}$	$M-A=-85935(32)$ keV for mixture gs+m at 259.3 keV										Nub16b **
* $^{136}\text{Pm-u}$	$M-A=-71091(28)$ keV for mixture gs+m at 110(120) keV										Nub16b **
* $^{136}\text{Pm}^m-^{133}\text{Cs}_{1.023}$	Slightly contaminated by ground state, original error (20) increased										00Be42 **
* $^{136}\text{I}^m-^{136}\text{Xe}$	High spin isomer is preferred (see also $^{134}\text{Sb}$ )										GAu **
* $^{136}\text{I}(\beta^-)^{136}\text{Xe}$	$E_{\beta^-}=7000(100), 5610(150), 4280(150)$ to ground state, $2^+$ levels 1313.027, 2634.16 keV										Ens026 **
* $^{136}\text{I}(\beta^-)^{136}\text{Xe}$	$E_{\beta^-}=5370(400), 4700(320), 3920(220)$ to $2^+$ levels 1313.027, 2289.53, 2634.16										Ens026 **
* $^{136}\text{I}^m(\beta^-)^{136}\text{Xe}$	$E_{\beta^-}=5170(400) 4670(270)$ to $^{136}\text{Xe}^m 6^+$ at 1891.703 and 2444.39 level										Ens026 **
* $^{136}\text{Cs}(\beta^-)^{136}\text{Ba}$	$E_{\beta^-}=341(2) 342(5)$ respectively, to $6^+$ level at 2207.077 keV										Ens026 **
* $^{136}\text{Pr}(\beta^+)^{136}\text{Ce}$	$E_{\beta^+}=2970(50) 3000(75) 3020(20)$ respectively, to $2^+$ level at 1092.09 keV										Ens026 **
* $^{136}\text{Nd}(\beta^+)^{136}\text{Pr}$	$E_{\beta^+}=1330(50)$ to $1^+$ level at 149.11 keV										Ens026 **
* $^{136}\text{Nd}(\beta^+)^{136}\text{Pr}$	$K/\beta^+=13.2(0.5)$ to $1^+$ level at 149.11 keV										Ens026 **
* $^{136}\text{Pm}(\beta^+)^{136}\text{Nd}$	$E_{\beta^-}=4732(70)$ probably from high spin isomer going to several										AHW **
*	high spin levels around 2100 keV										AHW **
$^{137}\text{Sb-u}$	-64445	215	-64480	60	-0.1	o			GT1	1.5	04Ma.A
	-65068	186			1.3	U			GT3	2.5	16Kn03
$^{137}\text{Te-u}$	-74528	101	-74400.6	2.3	0.5	o			GT2	2.5	08Kn.A
	-74386	129			0.0	U			GT2	2.5	08Su19
$^{137}\text{I-u}$	-82145	130	-81972	9	0.5	U			GT2	2.5	08Su19
$\text{C}_{11} \text{H}_5-^{137}\text{Ba}$	133366	24	133297.8	0.3	-1.9	U			R07	1.5	68De17
$\text{C}_7 \text{ } ^{13}\text{C} \text{ N O H}_{10}-^{137}\text{Ba}$	173792	73	173766.4	0.3	-0.2	U			R07	1.5	68De17
$\text{C}_{12} \text{H}_9-^{137}\text{Ba O}$	169692	39	169683.3	0.3	-0.1	U			R07	1.5	68De17
$^{137}\text{La-u}$	-93556	30	-93549.4	1.8	0.2	U			GS2	1.0	05Li24
$^{137}\text{Ce-u}$	-92101	85	-92237.4	0.5	-1.6	U			GS2	1.0	05Li24 *
$^{137}\text{Nd-u}$	-85438	30	-85438	13	0.0	1	18	18 $^{137}\text{Nd}$	GS2	1.0	05Li24 *
$^{137}\text{Pm-u}$	-79608	62	-79520	14	1.4	U			GS2	1.0	05Li24 *
$^{137}\text{Sm-u}$	-73025	69	-73030	50	-0.1	1	44	44 $^{137}\text{Sm}$	GS2	1.0	05Li24 *
$^{137}\text{Te}-^{130}\text{Xe}_{1.054}$	27300.0	2.7	27300.5	2.3	0.2	1	70	70 $^{137}\text{Te}$	JY1	1.0	12Ha25
$^{137}\text{Sb}-^{133}\text{Cs}_{1.030}$	32907	56				2			CP1	1.0	13Va12
$^{137}\text{Te}-^{133}\text{Cs}_{1.030}$	22985.0	4.1	22983.8	2.3	-0.3	1	30	30 $^{137}\text{Te}$	CP1	1.0	13Va12
$^{137}\text{Xe}-^{133}\text{Cs}_{1.030}$	8943.6	2.0	8942.25	0.11	-0.7	U			MA8	1.0	09Ne11
$^{137}\text{Cs}-^{133}\text{Cs}_{1.030}$	4452	19	4473.9	0.4	1.2	o			MA1	1.0	90St25
	4470	14			0.3	U			MA1	1.0	99Am05
$^{137}\text{Pr}-^{133}\text{Cs}_{1.030}$	8095	15	8064	9	-2.1	1	34	34 $^{137}\text{Pr}$	MA5	1.0	00Be42
$^{137}\text{Nd}-^{133}\text{Cs}_{1.030}$	11947	14	11947	13	0.0	1	81	81 $^{137}\text{Nd}$	MA5	1.0	00Be42
$^{137}\text{Pm}-^{133}\text{Cs}_{1.030}$	17864	14				2			MA5	1.0	00Be42
$^{137}\text{Sm}-^{133}\text{Cs}_{1.030}$	24350	78	24350	50	0.1	1	34	34 $^{137}\text{Sm}$	MA5	1.0	00Be42 *
$^{137}\text{Eu}-^{133}\text{Cs}_{1.030}$	32815.2	4.7				2			MA8	1.0	13Wo05
$^{137}\text{Ba } ^{35}\text{Cl}-^{135}\text{Ba } ^{37}\text{Cl}$	3089.1	0.6	3088.88	0.11	-0.1	U			H49	2.5	10Mc04
$^{137}\text{Te}-^{136}\text{Xe}_{1.007}$	19057	18	19034.4	2.3	-1.3	U			CP1	1.0	12Va02
$^{137}\text{I}-^{136}\text{Xe}_{1.007}$	11463.2	9.0				2			CP1	1.0	12Va02
$^{137}\text{Xe}-^{136}\text{Xe}_{1.007}$	5004	11	4992.79	0.11	-1.0	U			CP1	1.0	12Va02
$^{137}\text{Ba}-^{136}\text{Ba}$	1249	3	1251.42	0.07	0.3	U			M17	2.5	66Be10
	1222	50			0.4	U			R07	1.5	68De17
	1227	44			0.4	U			R07	1.5	68De17

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{137}\text{Ba}-^{135}\text{Ba}$	143	3	138.77	0.09	-0.6	U			M17	2.5	66Be10
	69	63			0.7	U			R07	1.5	68De17
	106	46			0.5	U			R07	1.5	68De17
$^{137}\text{I}(\beta^-n)^{136}\text{Xe}$	1850	30	2002	8	5.1	C					84Kr.B
$^{136}\text{Xe}(n,\gamma)^{137}\text{Xe}$	4025.5	0.5	4025.56	0.10	0.1	U					77Fo02 Z
	4025.8	0.3			-0.8	2					77Pr07 Z
	4025.53	0.11			0.3	2			Bdn		06Fi.A
$^{136}\text{Xe}(d,p)^{137}\text{Xe}$	1637	20	1801.00	0.10	8.2	F			Oak		68Mo21 *
$^{136}\text{Xe}(^3\text{He},d)^{137}\text{Cs}$	1918	12	1911.9	0.4	-0.5	U			ChR		81Ha08
$^{136}\text{Ba}(n,\gamma)^{137}\text{Ba}$	6891	5	6905.63	0.07	2.9	U					69Gr31
	6905.54	0.10			0.9	-			MMn		90Is07 Z
	6905.70	0.12			-0.6	-			Mtn		95Bo03 Z
	6905.74	0.16			-0.7	-			Bdn		06Fi.A
$^{137}\text{Ba}(\gamma,n)^{136}\text{Ba}$	-6949	38	-6905.63	0.07	1.1	U			Phi		60Ge01
$^{136}\text{Ba}(d,p)^{137}\text{Ba}$	4680	15	4681.06	0.07	0.1	U			ANL		70Vo04
$^{136}\text{Ba}(n,\gamma)^{137}\text{Ba}$	ave.	6905.63	0.07	6905.63	0.07	0.0	1	100	99	$^{137}\text{Ba}$	average
$^{136}\text{Ce}(n,\gamma)^{137}\text{Ce}$	7481.3	0.4	7481.53	0.16	0.6	-					81Ko.A Z
	7481.58	0.17			-0.3	-			Bdn		06Fi.A
	ave.	7481.54	0.16			0.0	1	100	100	$^{137}\text{Ce}$	average
$^{137}\text{Te}(\beta^-)^{137}\text{I}$	7030	300	7053	9	0.1	U					85Sa15
	6925	130			1.0	U			Bwg		87Gr.A
$^{137}\text{I}(\beta^-)^{137}\text{Xe}$	5880	60	6027	8	2.5	U			Bwg		87Gr.A
$^{137}\text{Xe}(\beta^-)^{137}\text{Cs}$	4140	70	4162.2	0.4	0.3	U					64On03
	4150	100			0.1	U					68Ho22
$^{137}\text{Cs}(\beta^-)^{137}\text{Ba}$	1173.29	0.84	1175.63	0.17	2.8	U					68Wo02 *
	1175.55	0.26			0.3	2					78Ch22 *
	1175.69	0.23			-0.3	2					83Be18 *
$^{137}\text{Ce}(\beta^+)^{137}\text{La}$	1222.1	1.6				2					81Ar.A *
$^{137}\text{Pr}(\beta^+)^{137}\text{Ce}$	2702	10	2717	8	1.5	1	66	66	$^{137}\text{Pr}$		73Bu17
$^{137}\text{Nd}(\beta^+)^{137}\text{Pr}$	3497	40	3617	14	3.0	B					73Bu18 *
	3690	54			-1.3	U					85Af.A *
$^{137}\text{Pm}^m(\beta^+)^{137}\text{Nd}$	5690	130	5660	50	-0.3	-			IRS		83Al06 *
	5650	60			0.1	-			Dbn		95Ve08 *
	ave.	5660	50			0.0	1	71	70	$^{137}\text{Pm}^m$	average
$^{137}\text{Sm}(\beta^+)^{137}\text{Pm}^m$	5900	70	5900	50	0.0	1	53	30	$^{137}\text{Pm}^m$	Dbn	95Ve08
* $^{137}\text{Ce}-u$	$M - A = -85665(29)$ keV for mixture gs+m at 254.29 keV										Nub16b **
* $^{137}\text{Pm}-u$	$M - A = -74079(28)$ keV for mixture gs+m at 150(50) keV										Nub16b **
* $^{137}\text{Sm}-u$	$M - A = -67932(28)$ keV for mixture gs+m at 180#50 keV										Nub16b **
* $^{137}\text{Sm}-^{133}\text{Cs}_{1,030}$	Might be a mixture of ground state and isomer say authors										00Be42 **
*	$D_M = 24447(14)$ $\mu\text{u}$ for mixture gs+m at 180#50 keV; $M - A = -67941(13)$ keV										Nub16b **
* $^{136}\text{Xe}(d,p)^{137}\text{Xe}$	F : error severely underestimated and value low, see excitation energies										AHW **
* $^{137}\text{Cs}(\beta^-)^{137}\text{Ba}$	$E_{\beta^-} = 511.63(0.84)$ $513.89(0.26)$ $514.03(0.23)$ to $^{137}\text{Ba}^m$ at 661.659 keV										Nub16b **
* $^{137}\text{Ce}(\beta^+)^{137}\text{La}$	$E_{\beta^+} = 189.5(1.6)$ to $5/2^+$ level at 10.59 keV										Ens079 **
* $^{137}\text{Nd}(\beta^+)^{137}\text{Pr}$	$E_{\beta^+} = 2400(40)$ $E_{\beta^+} = 2592(54)$ respectively, to $3/2^+$ level at 75.5 keV										Ens079 **
* $^{137}\text{Pm}^m(\beta^+)^{137}\text{Nd}$	$E_{\beta^+} = 4132(+150-115)$ $4110(60)$ respectively, to $11/2^-$ $^{137}\text{Nd}^m$ at 519.43 keV										Nub16b **
$^{138}\text{Sb}-u$	-58208	457				2			GT3	2.5	16Kn03
$^{138}\text{Te}-u$	-70940	247	-70528	4	1.1	o			GT1	1.5	04Ma.A
	-70583	106			0.2	o			GT2	2.5	08Kn.A
	-70591	131			0.2	U			GT2	2.5	08Su19
$\text{C}_{10} \text{H}_{18}-^{138}\text{Ba}$	235609	20	235603.4	0.3	-0.1	U			M17	2.5	66Be10
$\text{C}_{11} \text{H}_7-^{138}\text{Ba} \text{H}$	141649	51	141703.0	0.3	0.7	U			R07	1.5	68De17
$\text{C}_{11} \text{H}_6-^{138}\text{Ba}$	141701	30			0.0	U			R07	1.5	68De17
$\text{C}_{12} \text{H}_{10}-^{138}\text{Ba} \text{O}$	178106	15	178088.5	0.3	-0.8	U			R07	1.5	68De17
	178105	49			-0.2	U			R07	1.5	68De17



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_{11} \text{ }^{13}\text{C H}_9 - ^{138}\text{Ba O}$	173612	37	173618.3	0.3	0.1	U			R07	1.5	68De17
$C_{10} \text{ H}_{18} - ^{138}\text{Ce}$	234799	60	234862	5	0.3	U			R05	4.0	65De13
$C_{12} \text{ H}_{10} - ^{138}\text{Ce O}$	177382	46	177347	5	-0.2	U			R05	4.0	65De13
$C_9 \text{ }^{13}\text{C H}_{17} - ^{138}\text{Ce}$	230358	60	230392	5	0.1	U			R05	4.0	65De13
$^{138}\text{Pr}^m - u$	-88896	30	-88870	18	0.9	1	35	35 $^{138}\text{Pr}^m$	GS2	1.0	05Li24
$^{138}\text{Nd} - u$	-88060	130	-88050	12	0.1	o			GS1	1.0	00Ra23
	-88060	30			0.3	R			GS2	1.0	05Li24
$^{138}\text{Pm} - u$	-80242	141	-80452	30	-1.5	o			GS1	1.0	00Ra23 *
	-80454	35			0.1	1	72	72 $^{138}\text{Pm}$	GS2	1.0	05Li24 *
$^{138}\text{Sm} - u$	-76766	30	-76756	13	0.3	R			GS2	1.0	05Li24
$^{138}\text{Eu} - u$	-66291	30				2			GS2	1.0	05Li24
$^{138}\text{Te} - ^{130}\text{Xe}_{1.062}$	31945.3	4.7	31946	4	0.0	1	75	75 $^{138}\text{Te}$	JY1	1.0	12Ha25
$^{138}\text{Te} - ^{133}\text{Cs}_{1.038}$	27614.0	8.1	27613	4	-0.1	1	25	25 $^{138}\text{Te}$	CP1	1.0	13Va12
$^{138}\text{Xe} - ^{133}\text{Cs}_{1.038}$	12284.1	3.5	12287	3	0.9	1	74	74 $^{138}\text{Xe}$	MA8	1.0	09Ne11
$^{138}\text{Cs} - ^{133}\text{Cs}_{1.038}$	9158	14	9158	10	0.0	1	49	49 $^{138}\text{Cs}$	MA1	1.0	99Am05
$^{138}\text{Ba} - ^{133}\text{Cs}_{1.038}$	3388	14	3388.1	0.3	0.0	U			MA1	1.0	99Am05
$^{138}\text{Nd} - ^{133}\text{Cs}_{1.038}$	10093	14	10091	12	-0.2	-			MA5	1.0	00Be42
ave.	10091	13			0.0	1	96	96 $^{138}\text{Nd}$			average
$^{138}\text{Pm}^m - ^{133}\text{Cs}_{1.038}$	17721	14				2			MA5	1.0	00Be42
$^{138}\text{Sm} - ^{133}\text{Cs}_{1.038}$	21387	14	21385	13	-0.2	2			MA5	1.0	00Be42
$^{138}\text{I} - ^{136}\text{Xe}_{1.015}$	16903.7	6.4				2			CP1	1.0	12Va02
$^{138}\text{Xe} - ^{136}\text{Xe}_{1.015}$	8332.2	5.9	8324	3	-1.5	1	26	26 $^{138}\text{Xe}$	CP1	1.0	12Va02
$^{138}\text{Ba} \text{ }^{35}\text{Cl} - ^{136}\text{Ba} \text{ }^{37}\text{Cl}$	3621.1	0.6	3621.38	0.11	0.2	U			H49	2.5	10Mc04
$^{138}\text{Ba} - ^{137}\text{Ba}$	-582	2	-580.15	0.04	0.4	U			M17	2.5	66Be10
	-480	27			-2.5	U			R07	1.5	68De17
	-553	40			-0.5	U			R07	1.5	68De17
$^{138}\text{Ba} - ^{136}\text{Ba}$	676	3	671.27	0.09	-0.6	U			M17	2.5	66Be10
	658	98			0.1	U			R07	1.5	68De17
	628	43			0.7	U			R07	1.5	68De17
$^{138}\text{Ce} - ^{136}\text{Ce}$	-1040	47	-1141	5	-0.5	U			R05	4.0	65De13
	-1158	20			0.3	U			M17	2.5	66Be10
$^{138}\text{Ba H} - ^{137}\text{Ba}$	7399	88	7244.89	0.04	-1.2	U			R07	1.5	68De17
	7280	43			-0.5	U			R07	1.5	68De17
$^{137}\text{Ba}(n,\gamma)^{138}\text{Ba}$	8611.3	0.8	8611.72	0.04	0.5	U					68Ma35
	8611.72	0.04			0.0	1	100	99 $^{138}\text{Ba}$	MMn		90Is07 Z
	8611.5	0.15			1.5	U			Ltn		95Bo05
	8611.63	0.18			0.5	U			Bdn		06Fi.A
$^{137}\text{Ba}(d,p)^{138}\text{Ba}$	6398	15	6387.15	0.04	-0.7	U			ANL		70Vo04
$^{138}\text{I}(\beta^-)^{138}\text{Xe}$	7820	70	7992	7	2.5	U			Bwg		87Gr.A
$^{138}\text{Xe}(\beta^-)^{138}\text{Cs}$	2700	50	2915	10	4.3	B					72Mo33 *
	2830	80			1.1	U			Trs		78Wo15
$^{138}\text{Cs}^x(\text{IT})^{138}\text{Cs}$	40	23				2					82Au01 *
$^{138}\text{Cs}(\beta^-)^{138}\text{Ba}$	5350	80	5375	9	0.3	U			Trs		78Wo15
	5388	25			-0.5	-			Gsn		81De25
	5370	15			0.3	-			McG		84He.A
ave.	5375	13			0.0	1	51	51 $^{138}\text{Cs}$			average
$^{138}\text{La}(\epsilon)^{138}\text{Ba}$	1620	15	1742	3	8.2	B					56Tu17 *
$^{138}\text{La}(\beta^-)^{138}\text{Ce}$	994	10	1052	4	5.8	B					57G120 *
	1159	40			-2.7	U					70El.A *
	1052.7	4.3			-0.2	1	88	82 $^{138}\text{Ce}$			16Qu01 *
$^{138}\text{Pr}(\beta^+)^{138}\text{Ce}$	4437	10				2					71Af05
$^{138}\text{Pr}^m(\beta^+)^{138}\text{Ce}$	4801	20	4789	16	-0.6	1	67	65 $^{138}\text{Pr}^m$			64Fu08 *
$^{138}\text{Nd}(\beta^+)^{138}\text{Pr}$	2020	100	1116	16	-9.0	C					61Bo.B
$^{138}\text{Pm}(\beta^+)^{138}\text{Nd}$	7090	100	7078	29	-0.1	-			IRS		83Al06
	7080	60			0.0	-			Dbn		95Ve08
ave.	7080	50			-0.1	1	31	28 $^{138}\text{Pm}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{138}\text{Pm}^m(\beta^+)^{138}\text{Nd}$	7000	250	7108	17	0.4	U					81De38 *
* $^{138}\text{Pm}-u$	$M-A=-74730(130)$ keV for mixture gs+m at 30(30) keV										Nub16b **
* $^{138}\text{Pm}-u$	$M-A=-74927(28)$ keV for mixture gs+m at 30(30) keV										Nub16b **
* $^{138}\text{Xe}(\beta^-)^{138}\text{Cs}$	$E_{\beta^-}=2460(50), 2270(50)$ to $(1^- 2^-)$ level at 258.400, $1^-$ at 412.260 keV										Ens035 **
* $^{138}\text{Cs}^x(\text{IT})^{138}\text{Cs}$	Based on $^{138}\text{Cs}^m(\text{IT})=79.9$ keV										Nub16b **
* $^{138}\text{La}(\epsilon)^{138}\text{Ba}$	$L/K=1.40(0.25)$ to $2^+$ level at 1435.816 keV										Ens035 **
* $^{138}\text{La}(\beta^-)^{138}\text{Ce}$	$E_{\beta^-}=205(10) 370(40) 264.0(4.3)$ respectively, to $2^+$ level at 788.744 keV										Ens035 **
* $^{138}\text{Pr}^m(\beta^+)^{138}\text{Ce}$	$E_{\beta^+}=1650(20)$ to $7^-$ level at 2129.17 keV										Ens035 **
* $^{138}\text{Pm}^m(\beta^+)^{138}\text{Nd}$	$E_{\beta^+}=3900(200)$ to $5^-$ level at 1990.4, $6^+$ at 2134.3 and $(5^-)$ at 2221.8keV										Ens035 **
$^{139}\text{Te}-u$	-64541	333	-64633	4	-0.2	U			GT1	1.5	04Ma.A
$^{139}\text{Te}-^{130}\text{Xe}_{1.069}$	38515.7	3.8				2			JY1	1.0	12Ha25
$^{139}\text{I}-u$	-73838	102	-73507	4	1.3	U			GT2	2.5	08Kn.A
	-73567	130			0.2	U			GT2	2.5	08Su19
$\text{C}_6 \text{ }^{13}\text{C} \text{ O}_3 \text{ H}_6-^{139}\text{La}$	128474	41	128690.1	2.2	1.3	U			R05	4.0	65De13
$\text{C}_7 \text{ O}_3 \text{ H}_7-^{139}\text{La}$	133063	32	133160.3	2.2	0.8	U			R05	4.0	65De13
$\text{C}_6 \text{ N} \text{ O}_3 \text{ H}_5-^{139}\text{La}$	120496	21	120584.2	2.2	1.1	U			R05	4.0	65De13
$\text{C}_{12} \text{ H}_{11}-^{139}\text{La} \text{ O}$	184568	66	184801.9	2.2	0.9	U			R05	4.0	65De13
$\text{C}_{11} \text{ }^{13}\text{C} \text{ H}_{10}-^{139}\text{La} \text{ O}$	180100	58	180331.7	2.2	1.0	U			R05	4.0	65De13
$^{139}\text{Nd}-u$	-87840	79	-88046	30	-2.6	U			GS2	1.0	05Li24 *
$^{139}\text{Sm}-u$	-77704	30	-77703	12	0.0	R			GS2	1.0	05Li24
	-77711	30			0.3	R			GS2	1.0	05Li24 *
$^{139}\text{Eu}-u$	-70215	30	-70208	14	0.2	R			GS2	1.0	05Li24
$^{139}\text{Te}-^{133}\text{Cs}_{1.045}$	34185	17	34170	4	-0.9	U			CP1	1.0	13Va12
$^{139}\text{I}-^{133}\text{Cs}_{1.045}$	25296.1	4.3				2			CP1	1.0	13Va12
$^{139}\text{Xe}-^{133}\text{Cs}_{1.045}$	17594.9	2.3				2			MA8	1.0	09Ne11 *
$^{139}\text{Cs}-^{133}\text{Cs}_{1.045}$	12163	14	12167	3	0.3	U			MA1	1.0	99Am05
$^{139}\text{Ba}-^{133}\text{Cs}_{1.045}$	7649	14	7644.0	0.3	-0.4	U			MA1	1.0	99Am05
$^{139}\text{Pm}-^{133}\text{Cs}_{1.045}$	15604	15	15603	15	-0.1	1	95	95 $^{139}\text{Pm}$	MA5	1.0	00Be42
$^{139}\text{Sm}-^{133}\text{Cs}_{1.045}$	21101	14	21099	12	-0.1	2			MA5	1.0	00Be42
$^{139}\text{Eu}-^{133}\text{Cs}_{1.045}$	28597	16	28595	14	-0.1	2			MA5	1.0	00Be42
$^{139}\text{I}-^{136}\text{Xe}_{1.022}$	21333	31	21320	4	-0.4	U			CP1	1.0	12Va02
$^{139}\text{Xe}-^{136}\text{Xe}_{1.022}$	13618	12	13619.0	2.3	0.1	U			CP1	1.0	12Va02
$^{139}\text{La}-^{138}\text{La}$	-622	132	-759.0	2.7	-0.3	U			R05	4.0	65De13
$^{139}\text{La}-^{138}\text{Ce}$	485	74	370	5	-0.4	U			R05	4.0	65De13
$^{133}\text{Cs}-^{139}\text{Cs}_{.239} \text{ }^{131}\text{Cs}_{.761}$	-1774	24	-1771	4	0.1	F			P33	2.5	86Au02 *
$^{138}\text{Cs}^x-^{139}\text{Cs}_{.496} \text{ }^{137}\text{Cs}_{.504}$	770	40	800	25	0.3	U			P23	2.5	82Au01
$^{138}\text{Ba}(n,\gamma)^{139}\text{Ba}$	4723.4	0.7	4723.43	0.04	0.0	U					69Mo13
	4723.4	0.3			0.1	U					80Ba.A
	4723.43	0.04			0.0	1	100	99 $^{139}\text{Ba}$	MMn		90Is07 Z
	4723.20	0.14			1.6	U			Bdn		06Fi.A
$^{138}\text{Ba}(d,p)^{139}\text{Ba}$	2495	10	2498.86	0.04	0.4	U			MIT		64Sp12
	2496	15			0.2	U			Hei		67Wi08
	2493	10			0.6	U			ANL		70Vo04
$^{139}\text{La}(\gamma,n)^{138}\text{La}$	-8775	25	-8778.3	2.5	-0.1	U			Phi		60Ge01
$^{138}\text{La}(d,p)^{139}\text{La}$	6553	3	6553.8	2.5	0.3	-			Tal		71Du02
$^{139}\text{La}(d,t)^{138}\text{La}$	-2522	5	-2521.1	2.5	0.2	-			Tal		72La20
$^{138}\text{La}(d,p)^{139}\text{La}$	ave.	6553.4	6553.8	2.5	0.1	1	96	94 $^{138}\text{La}$			average
$^{139}\text{I}(\beta^-)^{139}\text{Xe}$	6815	100	7174	5	3.6	C			Bwg		87Gr.A
	6806	23			16.0	B			Bwg		92Gr06
$^{139}\text{Xe}(\beta^-)^{139}\text{Cs}$	5020	60	5056	4	0.6	U			Trs		78Wo15
	5062	22			-0.3	U			Bwg		92Gr06
$^{139}\text{Cs}(\beta^-)^{139}\text{Ba}$	4290	70	4213	3	-1.1	U			Trs		78Wo15
	4190	25			0.9	o			Gsn		80Bl.A
	4213	5			0.0	o			Gsn		81De25
	4214	4			-0.3	2			McG		84He.A
	4211	5			0.4	2			Gsn		92Pr04
$^{139}\text{Ba}(\beta^-)^{139}\text{La}$	2307	5	2312.5	2.0	1.1	-					75Fi07 *
	2336	25			-0.9	U			Gsn		81De25
	2316	4			-0.9	-			McG		84He.A
ave.	2312	3			0.0	1	42	41 $^{139}\text{La}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{139}\text{Ce}(\epsilon)^{139}\text{La}$	278	7	278	7	0.0	1	99	$^{99}\text{Ce}$			Averag *
$^{139}\text{Pr}(\beta^+)^{139}\text{Ce}$	2129	3	2129.1	3.0	0.0	1	100	$^{98}\text{Pr}$			81Ar.A
$^{139}\text{Nd}(\beta^+)^{139}\text{Pr}$	2787	50	2805	28	0.4	1	31	$^{30}\text{Nd}$			75Vy02 *
$^{139}\text{Pm}(\beta^+)^{139}\text{Nd}$	4450	100	4513	26	0.6	-					77De06 *
	4540	40			-0.7	-			IRS		83Al06
	4470	50			0.9	-			Dbn		95Ve08
ave.	4507	30			0.2	1	76	$^{70}\text{Nd}$			average
$^{139}\text{Sm}(\beta^+)^{139}\text{Pm}$	5430	150	5120	17	-2.1	U					82De06 *
	5510	150			-2.6	U			IRS		83Al06 *
$^{139}\text{Eu}(\beta^+)^{139}\text{Sm}$	6080	50	6982	17	18.0	C			Dbn		95Ve08 *
* $^{139}\text{Nd}-u$	$M-A=-81707(30)$ keV for mixture gs+m at 231.15 keV										
* $^{139}\text{Sm}-u$	$M-A=-71930(28)$ keV for $^{139}\text{Sm}^m$ at 457.40 keV										
* $^{139}\text{Xe}-^{133}\text{Cs}_{1.045}$	Typo in original paper, ratio should read 1.045 245 4357(175)										
* $^{133}\text{Cs}-^{139}\text{Cs}_{239}$ $^{131}\text{Cs}$ .	F : rejection based on line-shape analysis										
* $^{139}\text{Ba}(\beta^+)^{139}\text{La}$	$E_{\beta^+}=2141(5)$ to $5/2^+$ level at 165.8576 keV										
* $^{139}\text{Ce}(\epsilon)^{139}\text{La}$	Average pK=0.73(0.01) to $5/2^+$ level at 165.8576 keV in 10 references:										
*	pK=0.76 (0.04)										
*	pK=0.73 (0.01)										
*	pK=0.68 (0.02)										
*	pK=0.75 (0.01)										
*	pK=0.69 (0.02)										
*	pK=0.716(0.02)										
*	pK=0.78 (0.02)										
*	pK=0.726(0.010)										
*	pK=0.801(0.034)										
*	pK=0.705(0.020)										
* $^{139}\text{Nd}(\beta^+)^{139}\text{Pr}$	$E_{\beta^+}=1770(50)$ ; and $1170(50)$ from $^{139}\text{Nd}^m$ at 231.15 to $1/2^-$ level at 821.98										
* $^{139}\text{Pm}(\beta^+)^{139}\text{Nd}$	$E_{\beta^+}=3020(120)$ , $2990(100)$ to $3/2^+$ levels at 463.10, 402.77 keV										
* $^{139}\text{Sm}(\beta^+)^{139}\text{Pm}$	$E_{\beta^+}=4100(150)$ to $(1/2,3/2)^+$ level at 306.69 keV										
* $^{139}\text{Sm}(\beta^+)^{139}\text{Pm}$	$E_{\beta^+}=4735(+180-130)$ from $^{139}\text{Sm}^m$ at 457.40 to $^{139}\text{Pm}^m$ at 188.7 keV										
* $^{139}\text{Eu}(\beta^+)^{139}\text{Sm}$	$E_{\beta^+}=4600(50)$ to $^{139}\text{Sm}^m$ at 457.40 keV										
$^{140}\text{Te}-u$	-60827	225	-60740	70	0.3	U			GT1	1.5	04Ma.A
$^{140}\text{Te}-^{130}\text{Xe}_{1.077}$	43419	30	43180	70	-7.9	B			JY1	1.0	12Ha25
$^{140}\text{I}-u$	-68181	193	-68284	13	-0.4	o			GT1	1.5	04Ma.A
	-68463	102			0.7	o			GT2	2.5	08Kn.A
	-68273	130			0.0	o			GT2	2.5	08Su19
	-68202	186			-0.2	U			GT2	2.5	16Kn03
$^{140}\text{Xe}-u$	-78449	103	-78354.2	2.5	0.4	o			GT2	2.5	08Kn.A
	-78229	130			-0.4	U			GT2	2.5	08Su19
$\text{C}_{11} \text{H}_8-^{140}\text{Ce}$	157116	29	157153.8	1.7	0.3	U			R05	4.0	65De13
$\text{C}_{10} \text{H}_7-^{140}\text{Ce}$	152553	17	152683.6	1.7	1.9	U			R05	4.0	65De13
$\text{C}_{10} \text{N H}_6-^{140}\text{Ce}$	144599	35	144577.8	1.7	-0.2	U			R05	4.0	65De13
$\text{C}_{10} \text{N}_2 \text{H}_8-^{140}\text{Ce O}$	168207	48	168387.2	1.7	0.9	U			R05	4.0	65De13
$^{140}\text{Nd}-u$	-90448	30	-90456	4	-0.3	U			GS2	1.0	05Li24
$^{140}\text{Pm}^m-u$	-83532	30	-83503	14	1.0	1	22	$^{22}\text{Pm}^m$	GS2	1.0	05Li24
$^{140}\text{Sm}-u$	-81018	30	-81005	13	0.4	R			GS2	1.0	05Li24
$^{140}\text{Gd}-u$	-66326	30				2			GS2	1.0	05Li24
$^{140}\text{Te}-^{133}\text{Cs}_{1.053}$	38822	67				2			CP1	1.0	13Va12
$^{140}\text{I}-^{133}\text{Cs}_{1.053}$	31275	13				2			CP1	1.0	13Va12
$^{140}\text{Xe}-^{133}\text{Cs}_{1.053}$	21204.9	2.5				2			MA8	1.0	09Ne11
$^{140}\text{Cs}-^{133}\text{Cs}_{1.053}$	16837	14	16842	9	0.4	-			MA1	1.0	99Am05
	16857	14			-1.0	-			MA4	1.0	99Am05
ave.	16847	10			-0.5	1	79	$^{79}\text{Cs}$			average
$^{140}\text{Ba}-^{133}\text{Cs}_{1.053}$	10150	14	10166	9	1.1	1	37	$^{37}\text{Ba}$	MA1	1.0	99Am05

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{140}\text{Ce O}-^{133}\text{Cs}_{1,173}$	11269.2	2.7	11265.9	1.7	-1.2	1	40	40 $^{140}\text{Ce}$	MA8 1.0 17Ma.A
$^{140}\text{Nd O}-^{133}\text{Cs}_{1,173}$	15363.8	3.7				2			MA8 1.0 17Ma.A
$^{140}\text{Pm}^m-^{133}\text{Cs}_{1,053}$	16064	16	16056	14	-0.5	1	78	78 $^{140}\text{Pm}^m$	MA5 1.0 00Be42
$^{140}\text{Sm}-^{133}\text{Cs}_{1,053}$	18557	15	18554	13	-0.2	2			MA5 1.0 00Be42
$^{140}\text{Xe}-^{136}\text{Xe}_{1,029}$	17134	11	17122.1	2.5	-1.1	U			CP1 1.0 12Va02
$\text{C}_{11} \text{H}_9-^{140}\text{Ce}$	164956	40	164978.9	1.7	0.2	U			M17 2.5 66Be10
$^{140}\text{Ce}-^{139}\text{La}$	-1029	80	-912.4	1.9	0.4	U			R05 4.0 65De13
$\text{C}_{11} \text{H}_{10}-^{140}\text{Ce}$	172765	40	172803.9	1.7	0.4	U			M17 2.5 66Be10
$^{140}\text{Ce}-^{138}\text{Ce}$	-497	83	-542	5	-0.1	U			R05 4.0 65De13
	-543	8			0.0	U			M17 2.5 66Be10
$^{139}\text{Cs}-^{140}\text{Cs}_{,883} \ ^{131}\text{Cs}_{,118}$	-2280	40	-2275	8	0.1	U			P23 2.5 82Au01
$^{139}\text{Cs}-^{140}\text{Cs}_{,869} \ ^{132}\text{Cs}_{,132}$	-2210	40	-2240	8	-0.3	U			P23 2.5 82Au01
$^{138}\text{Ce}(\text{t,p})^{140}\text{Ce}$	8184	15	8166	5	-1.2	-			LAI 72Mu09
$^{140}\text{Ce}(\text{p,t})^{138}\text{Ce}$	-8167	20	-8166	5	0.0	-			Brk 77Sh06
$^{138}\text{Ce}(\text{t,p})^{140}\text{Ce}$	ave. 8178	12	8166	5	-1.0	1	16	16 $^{138}\text{Ce}$	average
$^{139}\text{La}(\text{n},\gamma)^{140}\text{La}$	5161.1	1.0	5160.98	0.04	-0.1	U			70Ju04
	5160	1			1.0	U			72Fu10
	5160.97	0.05			0.1	-			MMn 90Is09
	5161.00	0.10			-0.2	-			Bdn 06Fi.A
$^{139}\text{La}(\text{d,p})^{140}\text{La}$	2938	3	2936.41	0.04	-0.5	U			Tal 67Ke02
$^{139}\text{La}(\text{n},\gamma)^{140}\text{La}$	ave. 5160.98	0.04	5160.98	0.04	0.0	1	100	57 $^{139}\text{La}$	average
$^{140}\text{Ho}(\text{p})^{139}\text{Dy}$	1093.9	10.				3			99Ry04
$^{140}\text{Xe}(\beta^-)^{140}\text{Cs}$	4060	60	4064	9	0.1	U			Trs 78Wo15
$^{140}\text{Cs}(\beta^-)^{140}\text{Ba}$	6100	100	6219	10	1.2	U			Trs 78Wo15
	6235	25			-0.6	o			Gsn 80Bl.A
	6220	15			-0.1	o			Gsn 81De25
	6212	20			0.4	-			Gsn 92Pr04
	6199	25			0.8	-			Ida 93Gr17
	ave. 6207	16			0.8	1	40	21 $^{140}\text{Cs}$	average
$^{140}\text{Ba}(\beta^-)^{140}\text{La}$	1060	20	1047	8	-0.7	-			49Be36 *
	1050	20			-0.2	-			59Bo61 *
	1055	30			-0.3	-			65Bu07 *
	ave. 1055	13			-0.7	1	39	38 $^{140}\text{Ba}$	average
$^{140}\text{La}(\beta^-)^{140}\text{Ce}$	3760.2	2.0	3760.2	1.7	0.0	1	75	56 $^{140}\text{La}$	72Na04 *
$^{140}\text{Pr}(\beta^+)^{140}\text{Ce}$	3388	6				2			68Ab17
$^{140}\text{Nd}(\epsilon)^{140}\text{Pr}$	160	60	429	7	4.5	B			72Ba91
$^{140}\text{Pm}(\beta^+)^{140}\text{Nd}$	6080	100	6045	24	-0.3	U			75Ke09
	6090	40			-1.1	3			IRS 83Al06
	6020	30			0.8	3			Dbn 95Ve08
$^{140}\text{Pm}^m(\beta^+)^{140}\text{Nd}$	6484	70	6476	14	-0.1	U			75Ke09 *
$^{140}\text{Sm}(\epsilon)^{140}\text{Pm}$	3400	300	2758	27	-2.1	U			87De04
$^{140}\text{Eu}(\beta^+)^{140}\text{Sm}$	8400	400	8470	50	0.2	U			LBL 91Fi03 *
	8470	50				3			Dbn 95Ve08
$^{140}\text{Gd}(\beta^+)^{140}\text{Eu}$	4800	400	5200	60	1.0	U			LBL 91Fi03
$^{140}\text{Tb}(\beta^+)^{140}\text{Gd}$	11300	800				3			LBL 91Fi03 *
* $^{140}\text{Ba}(\beta^-)^{140}\text{La}$	$E_{\beta^-}=1022(20), 480(20)$ to $2^-$ level at 29.9641, $0^-$ at 581.07 keV								
* $^{140}\text{Ba}(\beta^-)^{140}\text{La}$	$E_{\beta^-}=1020(20), 830(50), 590(50)$ to $2^-$ level at 29.9641, $2^-$ at 162.6591, and $1^-$ at 467.653 keV								
* $^{140}\text{Ba}(\beta^-)^{140}\text{La}$	$E_{\beta^-}=1030(30), 1020(30)$ to $2^-$ level at 29.9641, $1^-$ at 43.844 keV								
* $^{140}\text{La}(\beta^-)^{140}\text{Ce}$	$E_{\beta^-}=2164(2)$ to $2^+$ level at 1596.237 keV								
* $^{140}\text{Pm}^m(\beta^+)^{140}\text{Nd}$	$E_{\beta^+}=3240(70)$ to $7^-$ level at 2221.4 keV								
* $^{140}\text{Eu}(\beta^+)^{140}\text{Sm}$	From $p^+$ . May be lower limit								
* $^{140}\text{Tb}(\beta^+)^{140}\text{Gd}$	Lower limit								
$^{141}\text{I}-u$	-64316	419	-64334	17	0.0	o			GT1 1.5 04Ma.A
	-64549	120			0.7	o			GT2 2.5 08Kn.A
	-64736	137			1.2	o			GT2 2.5 08Su19
	-64445	186			0.2	U			GT3 2.5 16Kn03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{141}\text{Xe}-u$	-73092	126	-73213	3	-0.4	o			GT2	2.5	08Kn.A
	-73560	136			1.0	U			GT2	2.5	08Su19
$^{141}\text{Ba}-u$	-85603.5	7.5	-85596	6	0.9	1	58	58 $^{141}\text{Ba}$	CP1	1.0	06Sa56
$\text{C}_{11}\text{H}_9-^{141}\text{Pr}$	162852	41	162766.9	1.8	-0.5	U			R05	4.0	65De13
$\text{C}_{10}\text{N H}_7-^{141}\text{Pr}$	150229	37	150190.8	1.8	-0.3	U			R05	4.0	65De13
$\text{C}_9\text{ }^{13}\text{C N H}_6-^{141}\text{Pr}$	145722	65	145720.6	1.8	0.0	U			R05	4.0	65De13
$^{141}\text{Pr}-u$	-92374	30	-92341.6	1.8	1.1	U			GS2	1.0	05Li24
$^{141}\text{Nd}-u$	-90401	30	-90385	4	0.5	U			GS2	1.0	05Li24
	-90365	30			-0.7	U			GS2	1.0	05Li24 *
$^{141}\text{Sm}-u$	-81496	62	-81518	9	-0.4	U			GS2	1.0	05Li24 *
$^{141}\text{Eu}-u$	-75048	42	-75068	14	-0.5	U			GS2	1.0	05Li24 *
$^{141}\text{Gd}-u$	-67881	30	-67874	21	0.2	2			GS2	1.0	05Li24
	-67867	30			-0.2	2			GS2	1.0	05Li24 *
$^{141}\text{Tb}-u$	-58552	113				2			GS2	1.0	05Li24 *
$^{141}\text{I}-^{133}\text{Cs}_{1.060}$	35887	17				2			CP1	1.0	13Va12
$^{141}\text{Xe}-^{133}\text{Cs}_{1.060}$	27008.1	3.1				2			MA8	1.0	09Ne11
$^{141}\text{Cs}-^{133}\text{Cs}_{1.060}$	20269	16	20266	10	-0.2	1	38	38 $^{141}\text{Cs}$	MA4	1.0	99Am05
$^{141}\text{Ba}-^{133}\text{Cs}_{1.060}$	14625	15	14624	6	0.0	-			MA1	1.0	99Am05
	14631	16			-0.4	-			MA4	1.0	99Am05
ave.	14628	11			-0.3	1	27	27 $^{141}\text{Ba}$			average
$^{141}\text{Pm}-^{133}\text{Cs}_{1.060}$	13776	15				2			MA5	1.0	00Be42
$^{141}\text{Sm}-^{133}\text{Cs}_{1.060}$	18692	14	18703	9	0.8	1	43	43 $^{141}\text{Sm}$	MA5	1.0	00Be42 *
$^{141}\text{Eu}-^{133}\text{Cs}_{1.060}$	25164	15	25153	14	-0.8	1	82	82 $^{141}\text{Eu}$	MA5	1.0	00Be42 *
$^{141}\text{Xe}-^{136}\text{Xe}_{1.037}$	23003	10	23006	3	0.3	U			CP1	1.0	12Va02
$^{141}\text{Cs}-^{136}\text{Xe}_{1.037}$	16277	22	16264	10	-0.6	1	20	20 $^{141}\text{Cs}$	CP1	1.0	12Va02
$^{139}\text{Cs}-^{141}\text{Cs}_{.789}\text{ }^{131}\text{Cs}_{.212}$	-3190	40	-3270	8	-0.8	U			P23	2.5	82Au01
$^{140}\text{Cs}-^{141}\text{Cs}_{.894}\text{ }^{131}\text{Cs}_{.107}$	-970	40	-1045	12	-0.7	U			P23	2.5	82Au01
$^{139}\text{Cs}-^{141}\text{Cs}_{.767}\text{ }^{132}\text{Cs}_{.234}$	-3210	40	-3183	8	0.3	U			P23	2.5	82Au01
$^{141}\text{Cs}(\beta^-n)^{140}\text{Ba}$	735	30	721	12	-0.5	1	15	9 $^{141}\text{Cs}$			84Kr.B
$^{140}\text{Ce}(n,\gamma)^{141}\text{Ce}$	5428.6	0.6	5428.14	0.10	-0.8	U			BNn		70Ge03 Z
	5428.01	0.20			0.7	-			Ptn		80Ba.A Z
	5428.19	0.12			-0.4	-			Bdn		06Fi.A
$^{140}\text{Ce}(d,p)^{141}\text{Ce}$	3210	10	3203.58	0.10	-0.6	U			MIT		64Sp12
	3202	15			0.1	U			Hei		67Wi08
ave.	5428.14	0.10	5428.14	0.10	0.0	1	100	64 $^{141}\text{Ce}$			average
$^{140}\text{Ce}(n,\gamma)^{141}\text{Ce}$	-9361	23	-9399	6	-1.6	U			Phi		60Ge01
$^{141}\text{Pr}(\gamma,n)^{140}\text{Pr}$	1177.4	8.	1177	7	-0.1	3					98Da03
$^{141}\text{Ho}(p)^{140}\text{Dy}$	1172.9	20.			0.2	3					99Ry04 *
$^{141}\text{Xe}(\beta^-)^{141}\text{Cs}$	6150	90	6280	10	1.4	U			Trs		78Wo15
$^{141}\text{Cs}(\beta^-)^{141}\text{Ba}$	5200	80	5255	10	0.7	U			Trs		78Wo15
	5264	15			-0.6	o			Gsn		80Bl.A *
	5252	15			0.2	o			Gsn		81De25
	5242	15			0.9	1	41	33 $^{141}\text{Cs}$	Gsn		92Pr04
$^{141}\text{Ba}(\beta^-)^{141}\text{La}$	3010	60	3199	7	3.2	B			Trs		78Wo15
	3208	35			-0.3	U			Gsn		81De25
	3217	20			-0.9	1	11	7 $^{141}\text{Ba}$	McG		84He.A
$^{141}\text{La}(\beta^-)^{141}\text{Ce}$	2430	30	2501	4	2.4	U					51Du19
	2502	4			-0.2	1	96	96 $^{141}\text{La}$	McG		84He.A
$^{141}\text{Ce}(\beta^-)^{141}\text{Pr}$	584	3	582.7	1.2	-0.4	-					50Fr58 *
	585	4			-0.6	-					52Ko27 *
	576.4	2.0			3.2	B					55Jo02 *
	581.4	2.0			0.7	-					68Be06 *
	582.2	2.6			0.2	-					79Ha09 *
ave.	582.5	1.3			0.2	1	83	48 $^{141}\text{Pr}$			average
$^{141}\text{Nd}(\beta^+)^{141}\text{Pr}$	1816	8	1823.0	2.8	0.9	2					73Bu21
	1824	3			-0.3	2					76Ga.A *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{141}\text{Pm}(\beta^+)^{141}\text{Nd}$	3730	60	3670	14	-1.0	U					70Ch29 *
	3640	70			0.4	U					75Ke09
$^{141}\text{Sm}(\beta^+)^{141}\text{Pm}$	4580	50	4589	16	0.2	U					77Ke03 *
	4463	60			2.1	U		IRS			83A106 *
	4524	80			0.8	U		IRS			93A103 *
$^{141}\text{Eu}(\beta^+)^{141}\text{Sm}$	6030	100	6008	14	-0.2	U					77De25 *
	5950	40			1.5	-		IRS			83A106 *
	6035	60			-0.4	U					85Af.A
	5550	100			4.6	B		IRS			93A103
	5980	40			0.7	-		Dbn			95Ve08 *
	ave.	5965	28		1.5	1	26	18 $^{141}\text{Eu}$			average
* $^{141}\text{Nd}-u$	$M-A=-83418(28)$ keV for $^{141}\text{Nd}^m$ at 756.51 keV										Nub16b **
* $^{141}\text{Sm}-u$	$M-A=-75825(28)$ keV for mixture gs+m at 175.9 keV										Nub16b **
* $^{141}\text{Eu}-u$	$M-A=-69858(28)$ keV for mixture gs+m at 96.45 keV										Nub16b **
* $^{141}\text{Gd}-u$	$M-A=-62840(28)$ keV for $^{141}\text{Gd}^m$ at 377.76 keV										Nub16b **
* $^{141}\text{Tb}-u$	$M-A=-54541(34)$ keV for mixture gs+m at 0#200 keV										Nub16b **
* $^{141}\text{Sm}-^{133}\text{Cs}_{1.060}$	$D_M=18694(14)$ and $D_M=18878(14)$ from $^{141}\text{Sm}^m$ at 175.9 keV										Nub16b **
* $^{141}\text{Eu}-^{133}\text{Cs}_{1.060}$	Slight (< 10%) isomeric contamination cannot be excluded										00Be42 **
* $^{141}\text{Ho}(p)^{140}\text{Dy}$	$E_p=1230(20)$ from $^{141}\text{Ho}^m$ at 66(2) keV										Nub16b **
* $^{141}\text{Cs}(\beta^-)^{141}\text{Ba}$	$E_{\beta^-}=5215(15)$ to $(5/2)^-$ level at 48.53 keV										Ens141 **
* $^{141}\text{Ce}(\beta^-)^{141}\text{Pr}$	$E_{\beta^-}=442(3)$ 444(4) 432(2) 436(2) 436.7(2.6) respectively, to $7/2^+$ level at 145.4434										Ens141 **
* $^{141}\text{Nd}(\beta^+)^{141}\text{Pr}$	Was erroneously quoted 77Ga.A in the 1993 tables										GAu **
* $^{141}\text{Pm}(\beta^+)^{141}\text{Nd}$	Original error 40 increased due to lack of information on calibration										GAu **
* $^{141}\text{Sm}(\beta^+)^{141}\text{Pm}$	$E_{\beta^+}=3180(50)$ , 3100(50) to $3/2^+$ level at 403.82, $(1/2)^+$ at 438.69 keV										Ens141 **
*	and $E_{\beta^+}=1670(70)$ , 1600(70) from $^{141}\text{Sm}^m$										77Ke03 **
*	at 175.9 to $11/2^-$ at 2091.71, $(9/2, 11/2, 13/2)^-$ at 2119.05										Ens141 **
* $^{141}\text{Sm}(\beta^+)^{141}\text{Pm}$	$E_{\beta^+}=3020(60)$ 32% to $3/2^+$ level at 403.82, 31% to $(1/2)^+$ 438.29 keV										Ens141 **
* $^{141}\text{Sm}(\beta^+)^{141}\text{Pm}$	$Q_{\beta^+}=4700(80)$ from $^{141}\text{Sm}^m$ at 175.9 keV										Nub16b **
* $^{141}\text{Eu}(\beta^+)^{141}\text{Sm}$	$E_{\beta^+}=4620(110)$ to $(5/2)^+$ level at 395.56 keV, and other $E_{\beta^+}$ (not given)										Ens141 **
* $^{141}\text{Eu}(\beta^+)^{141}\text{Sm}$	$E_{\beta^+}=4925(40)$ to $3/2^+$ level at 1.58 keV										Ens141 **
* $^{141}\text{Eu}(\beta^+)^{141}\text{Sm}$	$E_{\beta^+}=4960(40)$ to $3/2^+$ level at 1.58 keV										Ens141 **
$^{142}\text{I}-u$	-58798	268					2		GT1	1.5	04Ma.A
$^{142}\text{Xe}-u$	-70247	111	-70026.9	2.9	0.8	U			GT2	2.5	08Kn.A
$^{142}\text{Xe}-^{133}\text{Cs}_{1.068}$	30950.4	2.9					2		MA8	1.0	09Ne11
$^{142}\text{Cs}-^{133}\text{Cs}_{1.068}$	25270	16	25277	8	0.4	-			MA4	1.0	99Am05
	25304	23			-1.2	-			CP1	1.0	13Va12
	ave.	25281	13		-0.3	1	33	33 $^{142}\text{Cs}$			average
$^{142}\text{Ba}-^{133}\text{Cs}_{1.068}$	17410	15	17410	6	0.0	-			MA1	1.0	99Am05
	17420	16			-0.6	-			MA4	1.0	99Am05
	ave.	17415	11		-0.4	1	34	34 $^{142}\text{Ba}$			average
$^{142}\text{Ba}-u$	-83576.8	9.1	-83567	6	1.1	1	49	49 $^{142}\text{Ba}$	CP1	1.0	06Sa56
$\text{C}_{11} \text{H}_{10}-^{142}\text{Ce}$	169111	15	169000.4	2.7	-1.8	U			R05	4.0	65De13
	168955	40			0.5	U			M17	2.5	66Be10
	168955	40			0.5	U			M17	2.5	66Be10
$\text{C}_{10} \text{C} \text{H}_9-^{142}\text{Ce}$	164528	82	164530.2	2.7	0.0	U			R05	4.0	65De13
$\text{C}_{10} \text{N} \text{H}_8-^{142}\text{Ce}$	156558	42	156424.4	2.7	-0.8	U			R05	4.0	65De13
$\text{C}_{11} \text{H}_{10}-^{142}\text{Nd}$	170509	36	170521.4	1.5	0.1	U			R05	4.0	65De13
$\text{C}_{10} \text{N} \text{H}_8-^{142}\text{Nd}$	157870	43	157945.4	1.5	0.4	U			R05	4.0	65De13
$\text{C}_{10} \text{O} \text{H}_6-^{142}\text{Nd}$	134076	36	134135.9	1.5	0.4	U			R05	4.0	65De13
$\text{C}_{10} \text{C} \text{H}_9-^{142}\text{Nd}$	166021	32	166051.2	1.5	0.2	U			R05	4.0	65De13 *
$^{142}\text{Pm}-u$	-87136	30	-87110	25	0.9	-			GS2	1.0	05Li24
	ave.	-87124	27		0.5	1	89	89 $^{142}\text{Pm}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{142}\text{Sm}-^{133}\text{Cs}_{1,068}$	16173	14	16182	3	0.6	U			MA5	1.0	00Be42
$^{142}\text{Eu}^m-^{133}\text{Cs}_{1,068}$	24909	15	24910	13	0.1	2			MA5	1.0	00Be42
$^{142}\text{Eu}^m-u$	-76063	30	-76067	13	-0.1	R			GS2	1.0	05Li24
$^{142}\text{Gd}-u$	-71884	30				2			GS2	1.0	05Li24
$^{142}\text{Cs}-^{136}\text{Xe}_{1,044}$	21171	11	21168	8	-0.3	1	48	48 $^{142}\text{Cs}$	CP1	1.0	12Va02
$^{142}\text{Ce}-\text{C}_{11}\text{H}_9$	-161176	40	-161175.4	2.7	0.0	U			M17	2.5	66Be10
$^{142}\text{Nd}-\text{C}_{11}\text{H}_9$	-162665	30	-162696.4	1.5	-0.4	U			M17	2.5	66Be10
$^{142}\text{Ce}-^{140}\text{Ce}$	3818	3	3803.5	2.7	-1.9	U			M17	2.5	66Be10
$^{142}\text{Ce}-^{138}\text{Ce}$	3644	35	3261	6	-2.7	B			R05	4.0	65De13
$^{139}\text{Cs}-^{142}\text{Cs}_{,685}$ $^{132}\text{Cs}_{,316}$	-4840	40	-4858	6	-0.2	U			P23	2.5	82Au01
$^{140}\text{Cs}-^{142}\text{Cs}_{,789}$ $^{132}\text{Cs}_{,212}$	-2950	40	-2938	10	0.1	U			P23	2.5	82Au01
$^{141}\text{Cs}-^{142}\text{Cs}_{,794}$ $^{137}\text{Cs}_{,206}$	-580	40	-661	11	-0.8	U			P23	2.5	82Au01
$^{138}\text{Cs}^x-^{142}\text{Cs}_{,194}$ $^{137}\text{Cs}_{,806}$	550	40	589	25	0.4	U			P23	2.5	82Au01
$^{140}\text{Cs}-^{142}\text{Cs}_{,329}$ $^{139}\text{Cs}_{,671}$	260	40	300	9	0.4	U			P23	2.5	82Au01
$^{141}\text{Cs}-^{142}\text{Cs}_{,662}$ $^{139}\text{Cs}_{,338}$	-410	40	-520	10	-1.1	U			P23	2.5	82Au01
$^{141}\text{Cs}-^{142}\text{Cs}_{,496}$ $^{140}\text{Cs}_{,504}$	-640	40	-669	11	-0.3	U			P23	2.5	82Au01
	-663	19			-0.1	U			P33	2.5	86Au02
$^{142}\text{Ce}(\alpha)^{138}\text{Ba}$	1545	200	1303.5	2.5	-1.2	U					57Ri43
$^{140}\text{Ce}(\text{t,p})^{142}\text{Ce}$	4112	5	4117.9	2.5	1.2	1	25	20 $^{142}\text{Ce}$	LAI		72Mu09
$^{142}\text{Ce}(\text{p,t})^{140}\text{Ce}$	-4170	20	-4117.9	2.5	2.6	U			Osa		70Ya05
$^{142}\text{Nd}(\text{p,t})^{140}\text{Nd}$	-9150	20	-9352	4	-10.1	B			Osa		71Ya10 *
$^{142}\text{Ce}(\gamma,\text{n})^{141}\text{Ce}$	-7240	70	-7171.6	2.5	1.0	U			Phi		60Ge01
$^{142}\text{Ce}(\text{d,t})^{141}\text{Ce}$	-909	15	-914.4	2.5	-0.4	U			Mtr		72Le17
$^{141}\text{Pr}(\text{n},\gamma)^{142}\text{Pr}$	5843.14	0.10	5843.15	0.08	0.1	-			MMn		81Ke11 Z
	5843.16	0.12			-0.1	-			Bdn		06Fi.A
$^{141}\text{Pr}(\text{d,p})^{142}\text{Pr}$	3626	10	3618.58	0.08	-0.7	U			MIT		64Sp12
$^{141}\text{Pr}(\text{n},\gamma)^{142}\text{Pr}$	ave.	5843.15	0.08	5843.15	0.08	0.0	1	100	52 $^{141}\text{Pr}$		average
$^{142}\text{Xe}(\beta^-)^{142}\text{Cs}$	5040	100	5285	8	2.4	U			Trs		78Wo15
$^{142}\text{Cs}(\beta^-)^{142}\text{Ba}$	7230	70	7328	8	1.4	U			Trs		78Wo15
	7329	20			-0.1	o			Gsn		81De25
	7280	40			1.2	U			Bwg		87Gr.A
	7315	15			0.8	1	31	19 $^{142}\text{Cs}$	Gsn		92Pr04
$^{142}\text{Ba}(\beta^-)^{142}\text{La}$	2200	25	2182	8	-0.7	1	11	6 $^{142}\text{La}$			83Ch39
	2216	5			-6.8	C			McG		84He.A
$^{142}\text{La}(\beta^-)^{142}\text{Ce}$	4517	25	4509	6	-0.3	U					65Pr03
	4510	6			-0.2	1	95	94 $^{142}\text{La}$	McG		84He.A
$^{142}\text{Pr}(\beta^-)^{142}\text{Nd}$	2164	2	2162.5	1.4	-0.7	-					66Be12
	2158	3			1.5	-					75Ra09
	ave.	2162.2	1.7		0.2	1	72	52 $^{142}\text{Pr}$			average
$^{142}\text{Pm}(\beta^+)^{142}\text{Nd}$	4800	80	4808	24	0.1	R					60Ma.A
	4880	80			-0.9	R			IRS		83Al06
	4880	160			-0.5	U			LBL		91Fi03
$^{142}\text{Sm}(\beta^+)^{142}\text{Pm}$	2050	70	2156	24	1.5	1	12	11 $^{142}\text{Pm}$			60Ma.A
	2100	400			0.1	U			LBL		91Fi03
$^{142}\text{Eu}(\beta^+)^{142}\text{Sm}$	8000	300	7670	30	-1.1	U					75Ke08
	7400	100			2.7	U					82Gr.A
	7000	300			2.2	U			LBL		91Fi03
	7673	30			2				Dbn		94Po26
$^{142}\text{Eu}^m(\beta^+)^{142}\text{Sm}$	8150	100	8130	13	-0.2	U					75Ke08 *
	8174	50			-0.9	U			IRS		83Al06 *
	7480	100			6.5	B			IRS		93Al03 *
	8150	60			-0.3	U			Dbn		94Po26 *
$^{142}\text{Gd}(\beta^+)^{142}\text{Eu}$	4200	300	4350	40	0.5	U			LBL		91Fi03
$^{142}\text{Tb}(\beta^+)^{142}\text{Gd}$	10400	700				3			LBL		91Fi03
$^{142}\text{Dy}(\beta^+)^{142}\text{Tb}$	7100	200	6440#	200#	-3.3	D			LBL		91Fi03 *
* $\text{C}_{10}^{13}\text{C H}_9-^{142}\text{Nd}$	Original 1002055(32) is certainly a typo; rebuilt from M=141.907760(36)u										
* $^{142}\text{Nd}(\text{p,t})^{140}\text{Nd}$	Disagrees strongly with $^{140}\text{Nd}-u$										
* $^{142}\text{Eu}^m(\beta^+)^{142}\text{Sm}$	$E_{\beta^+}=4760(100)$ $4782(50)$ respectively, to $7^-$ level at 2372.1 keV										
* $^{142}\text{Eu}^m(\beta^+)^{142}\text{Sm}$	Measured half-life 73.4(0.5) s corresponds to $^{142}\text{Eu}^m$										
* $^{142}\text{Eu}^m(\beta^+)^{142}\text{Sm}$	$E_{\beta^+}=4756(60)$ to $7^-$ level at 2372.1 keV										
* $^{142}\text{Dy}(\beta^+)^{142}\text{Tb}$	Trends from Mass Surface TMS suggest $^{142}\text{Dy}$ 660 more bound										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{143}\text{I}-u$	-53849	495	-54350#	220#	-0.4	D			GT3	2.5	16Kn03 *
$^{143}\text{Xe}-u$	-64649	290	-64630	5	0.0	o			GT1	1.5	04Ma.A
	-64858	108			0.8	o			GT2	2.5	08Kn.A
	-64684	133			0.2	U			GT2	2.5	08Su19
	37008.7	5.0				2			MA8	1.0	09Ne11
$^{143}\text{Xe}-^{133}\text{Cs}_{1.075}$								GT2	2.5	08Kn.A	
$^{143}\text{Cs}-u$	-72771	117	-72653	8	0.4	U			GT2	2.5	08Kn.A
$^{143}\text{Cs}-^{133}\text{Cs}_{1.075}$	28985.6	8.5	28986	8	0.1	1	91	91 $^{143}\text{Cs}$	CP1	1.0	13Va12
$^{143}\text{Ba}-^{133}\text{Cs}_{1.075}$	22268	16	22264	7	-0.2	1	21	21 $^{143}\text{Ba}$	MA1	1.0	99Am05
$^{143}\text{Ba}-u$	-79375.0	8.5	-79375	7	0.0	1	73	73 $^{143}\text{Ba}$	CP1	1.0	06Sa56
$^{143}\text{La}-u$	-83918.1	8.7	-83921	8	-0.3	1	82	82 $^{143}\text{La}$	CP1	1.0	06Sa56
$\text{C}_{10} \text{ N H}_9 - ^{143}\text{Nd}$	163719	31	163679.4	1.5	-0.3	U			R05	4.0	65De13
$\text{C}_{10} \text{ O H}_7 - ^{143}\text{Nd}$	139814	42	139870.0	1.5	0.3	U			R05	4.0	65De13
$^{143}\text{Pm}-^{133}\text{Cs}_{1.075}$	12567	15	12577	3	0.7	U			MA5	1.0	00Be42
$^{143}\text{Sm}-^{133}\text{Cs}_{1.075}$	16268	15	16274	3	0.4	U			MA5	1.0	00Be42
$^{143}\text{Sm}-u$	-85347	30	-85365	3	-0.6	U			GS2	1.0	05Li24 *
$^{143}\text{Eu}-^{133}\text{Cs}_{1.075}$	21947	14	21938	12	-0.7	2			MA5	1.0	00Be42
$^{143}\text{Eu}-u$	-79706	30	-79701	12	0.2	R			GS2	1.0	05Li24
$^{143}\text{Gd}-u$	-73012	56	-73250	220	-4.2	C			GS2	1.0	05Li24 *
$^{143}\text{Tb}-u$	-64879	64	-64860	60	0.3	U			GS2	1.0	05Li24 *
$^{143}\text{Tb}-^{85}\text{Rb}_{1.682}$	83507	55				2			SH1	1.0	07Ra37 *
$^{143}\text{Dy}-^{85}\text{Rb}_{1.682}$	92364	14				2			SH1	1.0	07Ra37 *
$^{143}\text{Nd}-^{35}\text{Cl}-^{141}\text{Pr}-^{37}\text{Cl}$	5116	4	5111.6	1.5	-0.4	U			H21	2.5	70Ma05
$^{143}\text{Nd}-\text{C}_{11} \text{ H}_{10}$	-168422	30	-168430.4	1.5	-0.1	U			M17	2.5	66Be10
$^{143}\text{Nd}-^{142}\text{Nd}$	2322	46	2090.99	0.07	-1.3	U			R05	4.0	65De13
	2084	2			1.4	U			M17	2.5	66Be10
	-160594	30	-160605.4	1.5	-0.2	U			M17	2.5	66Be10
$^{141}\text{Cs}-^{143}\text{Cs}_{.493} \ ^{139}\text{Cs}_{.507}$	-230	40	-198	10	0.3	U			P23	2.5	82Au01
	-115	22			-1.5	U			P33	2.5	86Au02
$^{142}\text{Cs}-^{143}\text{Cs}_{.497} \ ^{141}\text{Cs}_{.504}$	647	15	657	9	0.3	U			P33	2.5	86Au02
$^{143}\text{Nd}(n,\alpha)^{140}\text{Ce}$	9699	15	9720.3	1.6	1.4	U			ILL		75Em.A
$^{143}\text{Nd}(p,t)^{141}\text{Nd}$	-7450	20	-7470	3	-1.0	U			Osa		71Ya10
$^{142}\text{Ce}(n,\gamma)^{143}\text{Ce}$	5145.9	0.5	5144.80	0.09	-2.2	U					76Ge02
	5144.78	0.15			0.1	-			Ptn		80Ba.A Z
	5144.81	0.12			-0.1	-			Bdn		06Fi.A
$^{142}\text{Ce}(d,p)^{143}\text{Ce}$	2945	15	2920.23	0.09	-1.7	U			Mtr		72Le17
$^{142}\text{Ce}(n,\gamma)^{143}\text{Ce}$	ave.	5144.80	0.09	5144.80	0.09	0.0	1	100	79 $^{142}\text{Ce}$		average
	$^{142}\text{Nd}(n,\gamma)^{143}\text{Nd}$	6123.62	0.08	6123.57	0.07	-0.6	-		MMn		82Is05 Z
$^{142}\text{Nd}(d,p)^{143}\text{Nd}$	6123.41	0.14			1.1	-			Bdn		06Fi.A
	3916	15	3899.00	0.07	-1.1	U			Kop		67Ch16
	3902	15			-0.2	U			Tal		67Ne04
	3902	15			-0.2	U			Hei		67Wi08
	ave.	6123.57	0.07	6123.57	0.07	0.0	1	100	79 $^{142}\text{Nd}$		average
$^{142}\text{Nd}(^3\text{He},d)^{143}\text{Pm}$	-1099	25	-1193.8	2.7	-3.8	B			Oak		71Wi04
$^{143}\text{Cs}(\beta^-)^{143}\text{Ba}$	-1195	5			0.2	1	29	29 $^{143}\text{Pm}$	McM		80St10 *
	6250	90	6262	10	0.1	o			Gsn		81De25 *
	6240	70			0.3	U			Bwg		87Gr.A
$^{143}\text{Ba}(\beta^-)^{143}\text{La}$	6270	25			-0.3	1	15	9 $^{143}\text{Cs}$	Gsn		92Pr04
	4240	50	4234	10	-0.1	U					79Sc11
	4259	40			-0.6	U			Gsn		81De25
	4210	70			0.3	U			Bwg		87Gr.A
$^{143}\text{La}(\beta^-)^{143}\text{Ce}$	3425	17	3435	8	0.6	1	20	18 $^{143}\text{La}$			84Is09 *
$^{143}\text{Ce}(\beta^-)^{143}\text{Pr}$	1460.6	2.	1461.6	1.9	0.5	1	87	77 $^{143}\text{Ce}$			77Ra18 *
$^{143}\text{Pr}(\beta^-)^{143}\text{Nd}$	932	2	934.0	1.4	1.0	-					49Fe18
	935	2			-0.5	-					76Ra33
	ave.	933.5	1.4			0.3	1	94	90 $^{143}\text{Pr}$		average
$^{143}\text{Pm}(\beta^+)^{143}\text{Nd}$	1000	70	1041.6	2.7	0.6	U					67Va01 *



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{143}\text{Sm}(\beta^+)^{143}\text{Pm}$	3492	30	3443	4	-1.6	U					66Be21
	3437	30			0.2	U			IRS		83A106
	3500	60			-0.9	U			IRS		93A103
	3461	40			-0.4	U			Dbn		94Po26
$^{143}\text{Eu}(\beta^+)^{143}\text{Sm}$	5100	50	5276	11	3.5	B					74Ch21
	5160	60			1.9	o			IRS		83Ve.A
	5240	70			0.5	o			IRS		83A106
	5250	80			0.3	U			IRS		93A103
	5236	30			1.3	R			Dbn		94Po26
$^{143}\text{Gd}(\beta^+)^{143}\text{Eu}$	6010	200				3			IRS		93A103 *
* $^{143}\text{I}-u$	Trends from Mass Surface TMS suggest $^{143}\text{I}$ 750 less bound										GAu **
* $^{143}\text{Sm}-u$	$M-A=-78746(28)$ keV for $^{143}\text{Sm}^m$ at 753.99 keV										Nub16b **
* $^{143}\text{Gd}-u$	$M-A=-67934(28)$ keV for mixture gs+m at 152.6 keV										Nub16b **
* $^{143}\text{Tb}-u$	$M-A=-60434(32)$ keV for mixture gs+m at 0#100 keV										Nub16b **
*	outweighed by next item before correcting for isomeric mixture										GAu **
* $^{143}\text{Tb}-^{85}\text{Rb}_{1.682}$	$M-A=-60419.5(7.8)$ keV for mixture gs+m at 0#100 keV										Nub16b **
* $^{143}\text{Dy}-^{85}\text{Rb}_{1.682}$	$D_M=92354(17)$ and $D_M=92705(14)$ for $^{143}\text{Dy}^m$ at 310.7 keV										Nub16b **
* $^{142}\text{Nd}(^3\text{He},d)^{143}\text{Pm}$	Based on $^{146}\text{Nd}(^3\text{He},d)^{147}\text{Pm}$ $Q=-87.6(0.9)$ keV										AHW **
* $^{143}\text{Cs}(\beta^-)^{143}\text{Ba}$	$E_{\beta^-}=6070(50)$ and $5847(100)$ to $3/2^-$ level at 228.83 keV										Ens123 **
* $^{143}\text{La}(\beta^-)^{143}\text{Ce}$	$E_{\beta^-}=3419(17)$ 64% to $3/2^-$ ground state, 29% to $7/2^-$ level at 18.9 keV										Ens123 **
* $^{143}\text{Ce}(\beta^-)^{143}\text{Pr}$	$E_{\beta^-}=1110(2)$ to $3/2^+$ level at 350.622 keV										Ens123 **
* $^{143}\text{Pm}(\beta^+)^{143}\text{Nd}$	$pK=0.806(0.023)$ to $3/2^-$ level at 742.05 keV, and $p^+ < 1 \times 10^{-6}$										Ens123 **
* $^{143}\text{Gd}(\beta^+)^{143}\text{Eu}$	$Q_{\beta^+}=6160(200)$ from $^{143}\text{Gd}^m$ at 152.6 keV										Nub16b **
$^{144}\text{Xe}-^{133}\text{Cs}_{1.083}$	41340.6	5.7				2			MA8	1.0	09Ne11
$^{144}\text{Cs}-^{133}\text{Cs}_{1.083}$	34488	33	34471	22	-0.5	1	43	43 $^{144}\text{Cs}$	CP1	1.0	13Va12
$^{144}\text{Ba}-^{133}\text{Cs}_{1.083}$	25347	15	25350	8	0.2	1	26	26 $^{144}\text{Ba}$	MA1	1.0	99Am05
$^{144}\text{Ba}-u$	-77045.3	9.1	-77045	8	0.0	1	71	71 $^{144}\text{Ba}$	CP1	1.0	06Sa56
$^{144}\text{La}-u$	-80337.1	19.3	-80354	14	-0.9	2			CP1	1.0	06Sa56
	-80373	20			0.9	2			GS3	1.0	12Ch19
$\text{C}_{10} \text{O H}_8-^{144}\text{Nd}$	147408	28	147422.0	1.5	0.1	U			R05	4.0	65De13
	147384	29			0.3	U			R05	4.0	65De13
$\text{C}_9 \text{ }^{13}\text{C N H}_9-^{144}\text{Nd}$	166777	28	166761.3	1.5	-0.1	U			R05	4.0	65De13
$\text{C}_{10} \text{ H}_8 \text{ O}-^{144}\text{Sm}$	145450	50	145508.5	1.7	0.3	U			R04	4.0	64De15
$\text{C}_9 \text{ }^{13}\text{C H}_9 \text{ N}-^{144}\text{Sm}$	164955	46	164847.8	1.7	-0.6	U			R04	4.0	64De15
$^{144}\text{Eu}-^{133}\text{Cs}_{1.083}$	21223	17	21215	12	-0.5	1	46	46 $^{144}\text{Eu}$	MA5	1.0	00Be42
$^{144}\text{Eu}-u$	-81117	30	-81180	12	-2.1	1	15	15 $^{144}\text{Eu}$	GS2	1.0	05Li24
$^{144}\text{Gd}-u$	-77037	30				2			GS2	1.0	05Li24
$^{144}\text{Tb}-u$	-66955	30				2			GS2	1.0	05Li24 *
$^{144}\text{Dy}-u$	-60746	33	-60730	8	0.5	U			GS2	1.0	05Li24
$^{144}\text{Dy}-^{85}\text{Rb}_{1.694}$	88697.7	7.7				2			SH1	1.0	07Ra37
$^{144}\text{Ho}-^{85}\text{Rb}_{1.694}$	101537.9	9.1				2			SH1	1.0	07Ra37
$^{144}\text{Nd } ^{35}\text{Cl}-^{142}\text{Nd } ^{37}\text{Cl}$	5329	3	5314.08	0.12	-1.2	U			H12	4.0	64Ba15
	5308	3			0.8	U			H21	2.5	70Ma05
$^{144}\text{Sm}-^{144}\text{Nd}$	1951	3	1913.5	0.9	-3.1	B			H19	4.0	64Mc11
	1911.9	1.1			0.6	-			H25	2.5	72Ba08
	1913.68	0.94			-0.2	-			SH1	1.0	11Go23
	ave.	1913.5	0.9		0.0	1	91	85 $^{144}\text{Sm}$			average
$^{144}\text{Nd}-^{143}\text{Nd}$	269	25	272.98	0.06	0.0	U			R05	4.0	65De13
	273	3			0.0	U			M17	2.5	66Be10
$^{144}\text{Nd}-^{142}\text{Nd}$	2366	3	2363.97	0.09	-0.3	U			M17	2.5	66Be10
$^{142}\text{Cs}-^{144}\text{Cs}_{.592} \text{ } ^{139}\text{Cs}_{.409}$	-60	40	-51	14	0.1	U			P23	2.5	82Au01
$^{143}\text{Cs}-^{144}\text{Cs}_{.745} \text{ } ^{140}\text{Cs}_{.255}$	-920	50	-891	17	0.2	U			P23	2.5	82Au01
$^{142}\text{Cs}-^{144}\text{Cs}_{.329} \text{ } ^{141}\text{Cs}_{.671}$	290	40	276	11	-0.1	U			P23	2.5	82Au01
$^{143}\text{Cs}-^{144}\text{Cs}_{.662} \text{ } ^{141}\text{Cs}_{.338}$	-651	21	-616	16	0.7	U			P33	2.5	86Au02

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{143}\text{Cs}-^{144}\text{Cs}_{.497}$ $^{142}\text{Cs}_{.504}$	-790	50	-690	13	0.8	U			P23	2.5	82Au01
$^{144}\text{Nd}(\alpha)^{140}\text{Ce}$	1882.4	30.	1903.2	1.6	0.7	U					61Ma05
	1882.4	20.			1.0	U					65Is01
$^{144}\text{Sm}(\alpha)^{141}\text{Sm}$	-8693	12	-8693	9	0.0	1	51	50	$^{141}\text{Sm}$	MSU	78Pa11
$^{142}\text{Ce}(\text{t,p})^{144}\text{Ce}$	3582	15	3560	3	-1.5	U			LAI		72Mu09
$^{142}\text{Nd}(\text{t,p})^{144}\text{Nd}$	5450	30	5458.82	0.09	0.3	U			Ald		72Ch11
$^{144}\text{Nd}(\text{p,t})^{142}\text{Nd}$	-5470	20	-5458.82	0.09	0.6	U			Osa		71Ya10
$^{144}\text{Sm}(\text{p,t})^{142}\text{Sm}$	-10649	15	-10639.9	2.7	0.6	U			Ham		73Oe02
$^{143}\text{Nd}(\text{n},\gamma)^{144}\text{Nd}$	7817.11	0.07	7817.04	0.05	-1.0	-			MMn		82Is05 Z
	7816.93	0.08			1.4	-			ILn		91Ro.A Z
	7816.94	0.23			0.4	U			Bdn		06Fi.A
$^{144}\text{Nd}(\text{d,t})^{143}\text{Nd}$	-1555	15	-1559.81	0.05	-0.3	U			Ors		73Ga01
$^{143}\text{Nd}(\text{n},\gamma)^{144}\text{Nd}$	ave. 7817.03	0.05	7817.04	0.05	0.1	1	99	61	$^{144}\text{Nd}$		average
$^{143}\text{Nd}(\alpha)^{144}\text{Pm}$	-804	5	-790.6	2.6	2.7	B			McM		80St10 *
$^{143}\text{Nd}(\alpha)^{144}\text{Pm}-^{142}\text{Nd}(\alpha)^{143}\text{Pm}$	402.7	1.6	403.2	1.5	0.3	1	91	49	$^{143}\text{Pm}$		75Ma04
$^{144}\text{Sm}(\text{t},\alpha)^{143}\text{Pm}$	13542	25	13520.0	2.7	-0.9	U			Ald		68Ha13
$^{144}\text{Sm}(\text{d,t})^{143}\text{Sm}$	-4262	10	-4262.5	2.3	0.0	U					72Ja28
$^{144}\text{Sm}(\text{p,d})^{143}\text{Sm}-^{148}\text{Gd}(\alpha)^{147}\text{Gd}$	-1536	2	-1536.0	2.0	0.0	1	100	100	$^{143}\text{Sm}$		86Ru04
$^{144}\text{Tm}(\text{p})^{143}\text{Er}$	1712.0	16.				3			ORp		05Gr32
$^{144}\text{Cs}(\beta^-)^{144}\text{Ba}$	8451	30	8496	20	1.5	o			Gsn		81De25
	8560	80			-0.8	-			Bwg		87Gr.A
	8462	35			1.0	-			Gsn		92Pr04
	ave. 8480	30			0.6	1	41	38	$^{144}\text{Cs}$		average
$^{144}\text{Ba}(\beta^-)^{144}\text{La}$	3055	70	3083	15	0.4	U			Bwg		87Gr.A
$^{144}\text{La}(\beta^-)^{144}\text{Ce}$	4300	100	5582	13	12.8	B					79Ik07
	5435	90			1.6	U			Bwg		87Gr.A
	5540	100			0.4	o			Kur		02Sh.B
	5540	100			0.4	U			Kur		02Sh16
$^{144}\text{Ce}(\beta^-)^{144}\text{Pr}$	315.6	1.5	318.6	0.8	2.0	3					66Da04
	320	1			-1.4	3					76Ra33
$^{144}\text{Pr}(\beta^-)^{144}\text{Nd}$	2996	3	2997.4	2.4	0.5	2					59Po77
	3000	4			-0.6	2					66Da04
$^{144}\text{Eu}(\beta^+)^{144}\text{Sm}$	6330	30	6346	11	0.5	-			IRS		83Al06
	6400	80			-0.7	U			IRS		93Al03
	6287	30			2.0	-			Dbn		94Po26
$^{144}\text{Sm}(\text{p,n})^{144}\text{Eu}$	-7110.0	30.	-7129	11	-0.6	-					65Me12
$^{144}\text{Eu}(\beta^+)^{144}\text{Sm}$	ave. 6315	17	6346	11	1.8	1	39	39	$^{144}\text{Eu}$		average
$^{144}\text{Gd}(\beta^+)^{144}\text{Eu}$	4300	400	3860	30	-1.1	U					70Ar04
* $^{144}\text{Tb}-u$	$M-A=-61971(28)$ keV for $^{144}\text{Tb}^m$ at 396.9 keV										
* $^{143}\text{Nd}(\alpha)^{144}\text{Pm}$	Based on $^{146}\text{Nd}(\alpha)^{147}\text{Pm}$ $Q=-87.6(0.9)$ keV										
$^{145}\text{Xe}-^{133}\text{Cs}_{1.090}$	47777	12				2			MA8	1.0	09Ne11
$^{145}\text{Cs}-^{133}\text{Cs}_{1.090}$	38588	12	38586	10	-0.1	-			MA8	1.0	08We02
	38583	17			0.2	-			CP1	1.0	13Va12
	ave. 38586	10			0.0	1	99	99	$^{145}\text{Cs}$		average
$^{145}\text{Ba}-u$	-72481.6	9.1				2			CP1	1.0	06Sa56
$^{145}\text{La}-u$	-78188.8	13.3	-78192	13	-0.2	1	98	98	$^{145}\text{La}$	1.0	06Sa56
$^{145}\text{Ce}-u$	-82771.8	92.2	-82730	40	0.4	1	16	16	$^{145}\text{Ce}$	1.0	06Sa56
$\text{C}_{10} \text{O H}_9-^{145}\text{Nd}$	152641	55	152760.7	1.5	0.5	U			R05	4.0	65De13
	152653	30			0.9	U			R05	4.0	65De13
$\text{C}_9 \text{ }^{13}\text{C O H}_8-^{145}\text{Nd}$	148231	31	148290.5	1.5	0.5	U			R05	4.0	65De13
$^{145}\text{Pm}-u$	-87255	30	-87244	3	0.4	U			GS2	1.0	05Li24
$^{145}\text{Sm}-u$	-86535	30	-86582.8	1.7	-1.6	U			GS2	1.0	05Li24
$^{145}\text{Eu}-^{133}\text{Cs}_{1.090}$	19338	17	19330	3	-0.5	U			MA5	1.0	00Be42
$^{145}\text{Gd}-u$	-78287	30	-78290	21	-0.1	-			GS2	1.0	05Li24
	-78294	30			0.1	-			GS2	1.0	05Li24 *
	ave. -78291	21			0.0	1	99	99	$^{145}\text{Gd}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{145}\text{Tb}-\text{u}$	-71134	266	-71270	120	-0.5	1	19	$^{19}\text{ }^{145}\text{Tb}$	GS2	1.0	05Li24 *
$^{145}\text{Dy}-\text{u}$	-62575	49	-62526	7	1.0	U			GS2	1.0	05Li24 *
$^{145}\text{Dy}-^{85}\text{Rb}_{1.706}$	87960.7	7.0				2			SH1	1.0	07Ra37 *
$^{145}\text{Ho}-^{85}\text{Rb}_{1.706}$	97754.1	8.0				2			SH1	1.0	07Ra37
$^{145}\text{Nd } ^{35}\text{Cl}_2-^{141}\text{Pr } ^{37}\text{Cl}_2$	10828	7	10821.0	1.5	-0.4	U			H21	2.5	70Ma05
$^{145}\text{Nd } ^{35}\text{Cl}-^{143}\text{Nd } ^{37}\text{Cl}$	5744	5	5709.42	0.26	-1.7	U			H12	4.0	64Ba15
	5703	4			0.6	U			H21	2.5	70Ma05
$^{145}\text{Nd}-^{144}\text{Nd}$	2582	21	2486.33	0.25	-1.1	U			R05	4.0	65De13
	2480	2			1.3	U			M17	2.5	66Be10
$^{145}\text{Nd}-^{143}\text{Nd}$	2862	40	2759.31	0.25	-0.6	U			R05	4.0	65De13
	2751	3			1.1	U			M17	2.5	66Be10
$^{142}\text{Cs}-^{145}\text{Cs}_{.490} \text{ } ^{139}\text{Cs}_{.511}$	240	50	150	9	-0.7	U			P23	2.5	82Au01
$^{144}\text{Cs}-^{145}\text{Cs}_{.828} \text{ } ^{139}\text{Cs}_{.173}$	450	50	415	21	-0.3	U			P23	2.5	82Au01
$^{143}\text{Cs}-^{145}\text{Cs}_{.592} \text{ } ^{140}\text{Cs}_{.409}$	-700	80	-610	10	0.5	U			P23	2.5	82Au01
$^{143}\text{Cs}-^{145}\text{Cs}_{.493} \text{ } ^{141}\text{Cs}_{.507}$	-310	40	-309	10	0.0	U			P23	2.5	82Au01
$^{144}\text{Cs}-^{145}\text{Cs}_{.662} \text{ } ^{142}\text{Cs}_{.338}$	320	18	319	21	0.0	1	21	$^{20}\text{ }^{144}\text{Cs}$	P33	2.5	86Au02
$^{144}\text{Cs}-^{145}\text{Cs}_{.497} \text{ } ^{143}\text{Cs}_{.503}$	600	40	616	21	0.2	U			P23	2.5	82Au01
$^{145}\text{Pm}(\alpha)^{141}\text{Pr}$	2303.6	40.	2323.3	2.9	0.5	U					62Nu01
$^{145}\text{Nd}(n,\alpha)^{142}\text{Ce}$	8706	30	8747.6	2.2	1.4	U			ILL		75Em04
$^{145}\text{Nd}(p,t)^{143}\text{Nd}$	-5100	20	-5090.56	0.24	0.5	U			Osa		71Ya10
$^{144}\text{Nd}(n,\gamma)^{145}\text{Nd}$	5755.3	0.7	5755.31	0.23	0.0	-					75Na.A
	5756.9	2.0			-0.8	U					77Mc09
	5755.26	0.25			0.2	-			Bdn		06Fi.A
$^{144}\text{Nd}(d,p)^{145}\text{Nd}$	3521	15	3530.75	0.23	0.6	U			Hei		67Wi08
	3538	15			-0.5	U			Ors		73Ga01
$^{144}\text{Nd}(n,\gamma)^{145}\text{Nd}$	ave. 5755.26	0.24	5755.31	0.23	0.2	1	95	$^{89}\text{ }^{145}\text{Nd}$			average
$^{144}\text{Nd}(^3\text{He},d)^{145}\text{Pm}$	-680	5	-685.0	2.5	-1.0	1	26	$^{25}\text{ }^{145}\text{Pm}$	McM		80St10 *
$^{144}\text{Nd}(^3\text{He},d)^{145}\text{Pm}-^{143}\text{Nd}(^3\text{He},d)^{144}\text{Pm}$	105.2	1.6	105.7	1.5	0.3	1	91	$^{57}\text{ }^{144}\text{Pm}$			75Ma04
$^{144}\text{Sm}(n,\gamma)^{145}\text{Sm}$	6757.1	0.3	6757.10	0.30	0.0	1	99	$^{92}\text{ }^{145}\text{Sm}$			79Wa22
$^{144}\text{Sm}(d,p)^{145}\text{Sm}$	4533	12	4532.53	0.30	0.0	U			Tal		65Ke09
	4547	15			-1.0	U			Kop		67Ch16
$^{144}\text{Sm}(^3\text{He},d)^{145}\text{Eu}$	-2184	4	-2178.5	2.7	1.4	-			Mun		82Sc25
	-2174	4			-1.1	-					84Ru.A
	ave. -2179.0	2.8			0.2	1	92	$^{91}\text{ }^{145}\text{Eu}$			average
$^{145}\text{Dy}(\epsilon p)^{144}\text{Gd}$	6000	500	6228	29	0.5	U					83La.A *
$^{145}\text{Tm}(p)^{144}\text{Er}$	1740.1	10.	1736	7	-0.4	3			ORp		98Ba13
	1732.1	10.			0.4	3			Arp		07Se06
$^{145}\text{Cs}(\beta^-)^{145}\text{Ba}$	7358	70	7462	12	1.5	U			Gsn		81De25
	7930	75			-6.2	C			Bwg		87Gr.A
	7865	50			-8.1	B			Gsn		92Pr04
$^{145}\text{Ba}(\beta^-)^{145}\text{La}$	4925	80	5319	15	4.9	C			Bwg		87Gr.A
$^{145}\text{La}(\beta^-)^{145}\text{Ce}$	4110	80	4230	40	1.5	1	19	$^{18}\text{ }^{145}\text{Ce}$	Bwg		87Gr.A
$^{145}\text{Ce}(\beta^-)^{145}\text{Pr}$	2490	100	2560	30	0.7	-					67Ho19 *
	2600	100			-0.4	-					80Ya07 *
	2530	50			0.6	-			Bwg		87Gr.A
	ave. 2540	40			0.6	1	68	$^{67}\text{ }^{145}\text{Ce}$			average
$^{145}\text{Pr}(\beta^-)^{145}\text{Nd}$	1805	10	1806	7	0.1	1	50	$^{50}\text{ }^{145}\text{Pr}$			59Dr.A
$^{145}\text{Pm}(\epsilon)^{145}\text{Nd}$	143	15	164.5	2.5	1.4	U					59Br65 *
	150	5			2.9	B					74To04 *
$^{145}\text{Sm}(\epsilon)^{145}\text{Pm}$	607	6	616.2	2.5	1.5	-					71My01 *
	622	5			-1.2	-					83Vo10 *
	ave. 616	4			0.1	1	44	$^{41}\text{ }^{145}\text{Pm}$			average
$^{145}\text{Eu}(\beta^+)^{145}\text{Sm}$	2710	15	2659.8	2.7	-3.3	B					68Ad04 *
	2647	12			1.1	U					83Sc28 *
$^{145}\text{Gd}(\beta^+)^{145}\text{Eu}$	5070	60	5065	20	-0.1	U					79Fi07
	5090	90			-0.3	o			IRS		83Ve.A *
	5070	80			-0.1	U			IRS		85Al13

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{145}\text{Gd}(\epsilon)^{145}\text{Eu}$	5000	70	5065	20	0.9	U					77Ho18
$^{145}\text{Tb}(\beta^+)^{145}\text{Gd}$	6700	200	6540	110	-0.8	-					86Ve.A *
	6400	150			0.9	-			IRS		93Al03
	ave.	6510	120		0.2	1	81	81	$^{145}\text{Tb}$		average
$^{145}\text{Dy}(\beta^+)^{145}\text{Tb}^m$	7300	200				3			IRS		93Al03
* $^{145}\text{Gd}-u$	$M - A = -72181(28)$ keV for $^{145}\text{Gd}^m$ at 749.1 keV										Nub16b **
* $^{145}\text{Tb}-u$	$M - A = -65881(28)$ keV for mixture gs+m at 760(220) keV										Nub16b **
* $^{145}\text{Dy}-u$	$M - A = -58230(30)$ keV for mixture gs+m at 118.2 keV										Nub16b **
* $^{145}\text{Dy}-^{85}\text{Rb}_{1.706}$	$D_M = 88054.7(6.8)$ $\mu\text{u}$ for mixture gs+m at 118.2 keV with ratio $R = 0.741(13)$										Nub16b **
* $^{144}\text{Nd}(\beta^+\text{He,d})^{145}\text{Pm}$	Based on $^{146}\text{Nd}(\beta^+\text{He,d})^{147}\text{Pm}$ $Q = -87.6(0.9)$ keV										AHW **
* $^{145}\text{Dy}(\epsilon\text{p})^{144}\text{Gd}$	As read from graph										AHW **
* $^{145}\text{Ce}(\beta^-)^{145}\text{Pr}$	$E_{\beta^-} = 1700(100)$ 1810(100) respectively, to $(3/2)^-$ level at 786.91; and other $E_{\beta^-}$										Ens092 **
* $^{145}\text{Pm}(\epsilon)^{145}\text{Nd}$	LM/K=0.85(0.03) to $3/2^-$ level at 67.167 keV										Ens092 **
* $^{145}\text{Pm}(\epsilon)^{145}\text{Nd}$	pK=0.554(0.025) to $5/2^-$ level at 72.486 keV, and other pK										Ens092 **
* $^{145}\text{Sm}(\epsilon)^{145}\text{Pm}$	pK=0.27(0.03) 0.35(0.025) respectively, to $3/2^+$ level at 492.31 keV										Ens092 **
* $^{145}\text{Eu}(\beta^+)^{145}\text{Sm}$	$E_{\beta^+} = 794(15)$ to $3/2^+$ level at 893.788 keV										Ens092 **
* $^{145}\text{Eu}(\beta^+)^{145}\text{Sm}$	pK=0.72(0.02) to $(5/2^-, 7/2^-)$ level at 2508.31 and $9^-$ at 2513.37 levels										Ens092 **
* $^{145}\text{Gd}(\beta^+)^{145}\text{Eu}$	$E_{\beta^+} = 2310(90)$ to $3/2^+$ level at 1758.03 keV, and other $E_{\beta^+}$										Ens092 **
* $^{145}\text{Tb}(\beta^+)^{145}\text{Gd}$	$E_{\beta^+} = 3300(200)$ to $(9/2^-)$ level at 2382.3(0.2) keV										Ens092 **
$^{146}\text{Xe}-^{133}\text{Cs}_{1.098}$	52332	26				2			MA8	1.0	09Ne11
$^{146}\text{Cs}-^{133}\text{Cs}_{1.098}$	44437.4	3.3	44436	3	-0.5	2			MA8	1.0	15At.A
	44421.8	9.2			1.5	2			CP1	1.0	13Va12
$^{146}\text{Ba}-u$	-69618	112	-69724	22	-0.4	o			GT2	2.5	08Kn.A
	-69963	141			0.7	U			GT2	2.5	08Su19
	-69717.5	23.7			-0.3	1	89	89	$^{146}\text{Ba}$	1.0	06Sa56
$^{146}\text{La}-u$	-74252	86	-74130	40	1.4	1	18	18	$^{146}\text{La}$	1.0	06Sa56 *
$^{146}\text{Ce}-u$	-81191.8	20.8	-81198	18	-0.3	-			CP1	1.0	06Sa56
	-81171	40			-0.7	-			GS3	1.0	12Ch19
	ave.	-81187	18		-0.6	1	90	90	$^{146}\text{Ce}$		average
$\text{C}_{12}\text{H}_2-^{146}\text{Nd}$	102453	31	102527.6	1.5	0.6	U			R05	4.0	65De13
$\text{C}_{10}\text{O H}_{10}-^{146}\text{Nd}$	160017	27	160042.4	1.5	0.2	U			R05	4.0	65De13
	159971	50			0.4	U			R05	4.0	65De13
$\text{C}_9\text{ }^{13}\text{C O H}_9-^{146}\text{Nd}$	155525	35	155572.2	1.5	0.3	U			R05	4.0	65De13
$^{146}\text{Pm}-u$	-85289	30	-85298	5	-0.3	U			GS2	1.0	05Li24
$^{146}\text{Eu}-^{133}\text{Cs}_{1.098}$	21029	15	21025	6	-0.3	1	19	19	$^{146}\text{Eu}$	1.0	00Be42
$^{146}\text{Tb}-u$	-72464	77	-72750	50	-3.7	C			GS2	1.0	05Li24 *
$^{146}\text{Dy}-u$	-67150	30	-67155	7	-0.2	U			GS2	1.0	05Li24
$^{146}\text{Dy}-^{85}\text{Rb}_{1.718}$	84390.0	7.2	84390	7	0.0	1	100	100	$^{146}\text{Dy}$	1.0	07Ra37
$^{146}\text{Ho}-^{133}\text{Cs}_{1.098}$	48797	10	48807	7	1.0	1	50	50	$^{146}\text{Ho}$	1.0	07Ra37
$^{146}\text{Ho}-^{85}\text{Rb}_{1.718}$	96549	10	96539	7	-1.0	1	50	50	$^{146}\text{Ho}$	1.0	07Ra37
$^{146}\text{Er}-^{85}\text{Rb}_{1.718}$	103960.4	9.2	103964	7	0.3	1	61	61	$^{146}\text{Er}$	1.0	07Ra37
$^{146}\text{Nd } ^{35}\text{Cl}-^{144}\text{Nd } ^{37}\text{Cl}$	6003	3	5979.75	0.27	-1.9	U			H12	4.0	64Ba15
	5966	4			1.4	U			H21	2.5	70Ma05
	5982.8	1.1			-1.1	U			H25	2.5	72Ba08
$^{146}\text{Nd}-^{145}\text{Nd}$	526	33	543.30	0.09	0.1	U			R05	4.0	65De13
	536	2			1.5	U			M17	2.5	66Be10
$^{146}\text{Nd}-^{144}\text{Nd}$	3147	36	3029.64	0.26	-0.8	U			R05	4.0	65De13
	3026	3			0.5	U			M17	2.5	66Be10
$^{145}\text{Cs}-^{146}\text{Cs}_{.828}$ $^{140}\text{Cs}_{.173}$	-580	80	-928	9	-1.7	U			P23	2.5	82Au01
$^{144}\text{Cs}-^{146}\text{Cs}_{.329}$ $^{143}\text{Cs}_{.671}$	320	50	336	21	0.1	U			P23	2.5	82Au01
$^{145}\text{Cs}-^{146}\text{Cs}_{.662}$ $^{143}\text{Cs}_{.338}$	-440	30	-565	10	-1.7	U			P33	2.5	86Au02
$^{145}\text{Cs}-^{146}\text{Cs}_{.497}$ $^{144}\text{Cs}_{.503}$	-730	30	-740	13	-0.1	U			P33	2.5	86Au02
$^{146}\text{Sm}(\alpha)^{142}\text{Nd}$	2529.5	20.	2528.8	2.8	0.0	U					64Nu02
	2622.0	30.			-3.1	B					66Fr11
	2524.2	4.			1.1	1	47	46	$^{146}\text{Sm}$		87Me08 Z

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{144}\text{Nd}(t,p)^{146}\text{Nd}$	4834	30	4838.75	0.24	0.2	U			Ald		72Ch11
$^{144}\text{Sm}(t,p)^{146}\text{Sm}$	6681	25	6691.6	2.8	0.4	U			Ald		66Bj01
$^{144}\text{Sm}(^3\text{He},p)^{146}\text{Eu}$	2797	12	2794	6	-0.2	1	25	24 $^{146}\text{Eu}$			84Ru.A
$^{144}\text{Sm}(^3\text{He},n)^{146}\text{Gd}$	977	30	980	4	0.1	U			Bld		79A107
$^{144}\text{Sm}(^{12}\text{C},^{10}\text{Be})^{146}\text{Gd}$	-18476	25	-18487	4	-0.4	U			MSU		80Pa07
$^{146}\text{Nd}(d,^3\text{He})^{145}\text{Pr}$	-3095	10	-3095	7	0.0	1	50	49 $^{145}\text{Pr}$	KVI		79Sa.A
$^{145}\text{Nd}(n,\gamma)^{146}\text{Nd}$	7565.28	0.10	7565.23	0.09	-0.5	-			MMn		82Is05
	7565.05	0.18			1.0	-			Bdn		06Fi.A
	ave.	7565.23	0.09		0.1	1	99	88 $^{146}\text{Nd}$			average
$^{146}\text{Sm}(^3\text{He},\alpha)^{145}\text{Sm}$	12161	5	12161.3	2.9	0.1	1	33	30 $^{146}\text{Sm}$			86Ru04
$^{146}\text{Tm}(p)^{145}\text{Er}$	895.2	8.	896	6	0.1	o			ORp		03Gi10
	896.2	8.			-0.1	3			Arp		05Ro40
	895.2	8.			0.1	3			ORp		06Ta08
$^{146}\text{Tm}^m(p)^{145}\text{Er}$	1197.3	5.	1199.3	1.0	0.4	U			Dap		93Li18
	1198.3	10.			0.1	o			ORp		01Ry01
	1200.3	8.			-0.1	U			Arp		05Ro40
	1199.3	1.				3			ORp		06Ta08
$^{146}\text{Tm}^m(p)^{145}\text{Er}^m$	994.5	4.				4			ORp		06Ta08
$^{146}\text{Tm}^n(p)^{145}\text{Er}^m$	1126.8	5.	1127.8	1.0	0.2	U			Dap		93Li18
	1127.8	10.			0.0	o			ORp		01Ry01
	1129.8	8.			-0.3	U			Arp		05Ro40
	1127.8	1.				5			ORp		06Ta08
$^{146}\text{Cs}(\beta^-)^{146}\text{Ba}$	9300	900	9637	21	0.4	o			Gsn		81De25
	9310	60			5.4	B			Bwg		87Gr.A
	9375	50			5.2	B			Gsn		92Pr04
$^{146}\text{Ba}(\beta^-)^{146}\text{La}$	4280	100	4100	30	-1.8	-			Gsn		81De25
	4030	50			1.5	-			Bwg		87Gr.A
	ave.	4080	40		0.5	1	56	45 $^{146}\text{La}$			average
$^{146}\text{La}(\beta^-)^{146}\text{Ce}$	6175	100	6590	30	4.1	B			Gsn		81De25
	6380	30			6.8	B			Trs		82Br23
	6620	70			-0.5	-			Bwg		87Gr.A
	6580	80			0.1	-					01Ko07
	ave.	6600	50		-0.3	1	43	37 $^{146}\text{La}$			average
$^{146}\text{Ce}(\beta^-)^{146}\text{Pr}$	1100	80	1050	30	-0.7	-					54Be10
	1050	100			0.0	-					67Ho19
	951	50			1.9	-					80Ya07
	1065	100			-0.2	-					81Eb01
	ave.	1010	40		1.0	1	80	76 $^{146}\text{Pr}$			average
$^{146}\text{Pr}(\beta^-)^{146}\text{Nd}$	4150	200	4240	30	0.5	U					54Be10
	4250	200			0.0	U					65Ra02
	4080	100			1.6	-					68Da13
	4140	100			1.0	-					78Ik03
	ave.	4110	70		1.9	1	24	24 $^{146}\text{Pr}$			average
$^{146}\text{Pm}(\beta^-)^{146}\text{Sm}$	1542	3				2					74Sc06
$^{146}\text{Eu}(\beta^+)^{146}\text{Sm}$	3871	10	3879	6	0.8	-					62Fu16
	3871	20			0.4	-					64Ta11
	3896	20			-0.9	-			Got		88Sa06
	ave.	3875	8		0.4	1	52	46 $^{146}\text{Eu}$			average
$^{146}\text{Gd}(\beta^+)^{146}\text{Eu}$	1757	30	1032	7	-24.2	B					70Ag01
	1300	200			-1.3	U					81Ka07
$^{146}\text{Tb}(\beta^+)^{146}\text{Gd}$	8240	150	8320	40	0.5	o			IRS		83Al06
	7910	150			2.7	U			IRS		93Al03
	8310	50			0.2	1	80	80 $^{146}\text{Tb}$	Dbn		94Po26
$^{146}\text{Dy}(\beta^+)^{146}\text{Tb}$	5160	100	5210	50	0.5	1	20	20 $^{146}\text{Tb}$	IRS		93Al03
* $^{146}\text{La}-u$	$D_M = -74182.5(30.6) \mu\text{u}$ for mixture gs+m at 130(130) keV; $M - A = -69100.6(28.5) \text{keV}$										
* $^{146}\text{Tb}-u$	$M - A = -67424(28) \text{keV}$ for mixture gs+m at 150#100 keV										
* $^{146}\text{Sm}(^3\text{He},\alpha)^{145}\text{Sm}$	$Q - Q(^{148}\text{Gd}(^3\text{He},\alpha)) = -567(5) \text{keV}$										
* $^{146}\text{Ba}(\beta^-)^{146}\text{La}$	$E_{\beta^-} = 3910(100)$ to $1^+$ level at 372.4 keV, and other $E_{\beta^-}$										
* $^{146}\text{La}(\beta^-)^{146}\text{Ce}$	$E_{\beta^-} = 5919(100)$ 6120(30) respectively, to $2^+$ level at 258.46 keV, and other $E_{\beta^-}$										
* $^{146}\text{La}(\beta^-)^{146}\text{Ce}$	$E_{\beta^-} = 6580(100)$ and 6320(80) to ground state and $2^+$ level at 258.46 keV										
* $^{146}\text{Ce}(\beta^-)^{146}\text{Pr}$	$E_{\beta^-} = 750(80)$ 700(100) 600(50) 715(100) respectively, to $1^+$ at 351.78 keV										
* $^{146}\text{Pr}(\beta^-)^{146}\text{Nd}$	$E_{\beta^-} = 3700(200)$ 3800(200) respectively, to $2^+$ level at 453.77 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* $^{146}\text{Pr}(\beta^-)^{146}\text{Nd}$	$E_{\beta^-}=4100(200), 3600(100), 2100(100)$ to ground state, $2^+$ 453.77, $2^+$ 1978.45 levels										
* $^{146}\text{Pr}(\beta^-)^{146}\text{Nd}$	$E_{\beta^-}=4150(150), 3700(100), 2160(100)$ to ground state, $2^+$ 453.77, $2^+$ 1978.45 levels										
* $^{146}\text{Pm}(\beta^-)^{146}\text{Sm}$	$E_{\beta^-}=795(3)$ to $2^+$ level at 747.2 keV										
* $^{146}\text{Eu}(\beta^+)^{146}\text{Sm}$	$E_{\beta^+}=2107(11) 2100(20)$ respectively, to $2^+$ level at 747.2 keV, and other $E_{\beta^+}$										
* $^{146}\text{Eu}(\beta^+)^{146}\text{Sm}$	$e/\beta^+$ to 2045.8 level										
* $^{146}\text{Gd}(\beta^+)^{146}\text{Eu}$	$E_{\beta^+}=350(30)$ to $1^-$ level at 384.79 keV										
* $^{146}\text{Gd}(\beta^+)^{146}\text{Eu}$	pK to 690.7 level, $p^+ < 1 \times 10^{-4}$ to 384.8 level, see $^{150}\text{Dy}(\beta^+)$										
* $^{146}\text{Tb}(\beta^+)^{146}\text{Gd}$	Reported half-life 24.1(0.5)s corresponds to $^{146}\text{Tb}^m$										
*	$Q_{\beta^-}=8060(100)$ keV from $^{146}\text{Tb}^m$ at estimated 150#100 keV										
$^{147}\text{Cs}-^{133}\text{Cs}_{1,105}$	48640	64	48737	9	1.5	U			MA8	1.0	08We02
	48737.1	9.0				2			MA8	1.0	15At.A
$^{147}\text{Ba}-\text{u}$	-64696.1	21.2				2			CP1	1.0	06Sa56
$^{147}\text{La}-\text{u}$	-71582.2	11.5				2			CP1	1.0	06Sa56
$^{147}\text{Ce}-\text{u}$	-77309.2	9.6	-77310	9	-0.1	1	92	92 $^{147}\text{Ce}$	CP1	1.0	06Sa56
$\text{C}_8 \text{H}_5 \text{N O}_2-^{147}\text{Sm}$	117197	40	117124.3	1.5	-0.5	U			R04	4.0	64De15
$\text{C}_9 \text{H}_7 \text{O}_2-^{147}\text{Sm}$	129703	17	129700.4	1.5	0.0	U			R04	4.0	64De15
$^{147}\text{Eu}-^{133}\text{Cs}_{1,105}$	21215	16	21227.9	2.8	0.8	U			MA5	1.0	00Be42
$^{147}\text{Tb}-\text{u}$	-75934	34	-75945	9	-0.3	U			GS2	1.0	05Li24 *
$^{147}\text{Tb}-^{133}\text{Cs}_{1,105}$	28533	12	28530	9	-0.2	1	53	53 $^{147}\text{Tb}$	SH1	1.0	07Ra37 *
$^{147}\text{Dy}-\text{u}$	-68909	30	-68917	10	-0.3	U			GS2	1.0	05Li24 *
	-68908	30			-0.3	U			GS2	1.0	05Li24 *
$^{147}\text{Dy}-^{133}\text{Cs}_{1,105}$	35558.3	9.5				2			SH1	1.0	07Ra37 *
$^{147}\text{Ho}-\text{u}$	-59944	30	-59858	5	2.9	B			GS2	1.0	05Li24
$^{147}\text{Ho}-^{133}\text{Cs}_{1,105}$	44613.7	7.8	44618	5	0.5	1	47	47 $^{147}\text{Ho}$	SH1	1.0	07Ra37
$^{147}\text{Ho}-^{85}\text{Rb}_{1,729}$	92661.6	7.4	92658	5	-0.5	1	53	53 $^{147}\text{Ho}$	SH1	1.0	07Ra37
$^{147}\text{Er}-^{133}\text{Cs}_{1,105}$	54452	42	54440	40	-0.3	o			SH1	1.0	07Ra37 *
$^{147}\text{Er}-^{85}\text{Rb}_{1,729}$	102480	41				2			SH1	1.0	07Ra37 *
$^{147}\text{Tm}-^{85}\text{Rb}_{1,729}$	113900	11	113895	7	-0.4	1	45	45 $^{147}\text{Tm}$	SH1	1.0	07Ra37
$^{147}\text{Eu}-^{142}\text{Sm}_{1,035}$	4516	17	4516	4	0.0	U			MA7	1.0	01Bo59
$^{147}\text{Sm } ^{35}\text{Cl}-^{145}\text{Nd } ^{37}\text{Cl}$	5305	4	5275.0	0.4	-1.9	U			H12	4.0	64Ba15
	5264	4			1.1	U			H21	2.5	70Ma05
$^{145}\text{Cs}-^{147}\text{Cs}_{,705} \ ^{140}\text{Cs}_{,296}$	-170	170	-644	11	-1.1	U			P23	2.5	82Au01
$^{144}\text{Cs}-^{147}\text{Cs}_{,490} \ ^{141}\text{Cs}_{,511}$	80	250	227	21	0.2	U			P23	2.5	82Au01
$^{145}\text{Cs}-^{147}\text{Cs}_{,493} \ ^{143}\text{Cs}_{,507}$	-87	22	-146	11	-1.1	U			P33	2.5	86Au02
$^{147}\text{Sm}(\alpha)^{143}\text{Nd}$	2292.5	10.	2311.0	0.4	1.8	U					62Si14 Z
	2296.7	5.			2.9	U					66Ma05 Z
	2300.8	5.			2.0	U					70Gu14 Z
	2310.5	0.5			0.9	1	50	28 $^{147}\text{Sm}$			16Ca.1
$^{147}\text{Eu}(\alpha)^{143}\text{Pm}$	2990.6	10.	2991	3	0.0	U					62Si14 Z
	2981.5	20.			0.5	U					64To04 Z
	2987.2	5.			0.7	1	37	22 $^{143}\text{Pm}$	DbA		67Go32 Z
$^{147}\text{Sm}(n,\alpha)^{144}\text{Nd}$	10114	8	10128.0	0.4	1.8	U			ILL		74Em01
$^{144}\text{Sm}(^{12}\text{C},^9\text{Be})^{147}\text{Gd}$	-17832	30	-17957.0	1.2	-4.2	B			MSU		80Pa07
	-17921	25			-1.4	U			Ors		85Be24
$^{144}\text{Sm}(^{14}\text{N},^{11}\text{Be})^{147}\text{Tb}$	-28280	50	-28537	8	-5.1	B			Hei		85Gy01
$^{147}\text{Sm}(p,t)^{145}\text{Sm}$	-6287	8	-6275.9	0.9	1.4	U			Min		72De47
$^{146}\text{Nd}(n,\gamma)^{147}\text{Nd}$	5292.19	0.15	5292.20	0.09	0.1	-			ILn		75Ro16 Z
	5292.19	0.11			0.1	-			Bdn		06Fi.A
$^{146}\text{Nd}(d,p)^{147}\text{Nd}$	3070	15	3067.63	0.09	-0.2	U			Hei		67Wi08
$^{146}\text{Nd}(n,\gamma)^{147}\text{Nd}$	ave.	5292.19	5292.20	0.09	0.1	1	99	89 $^{147}\text{Nd}$			average
$^{147}\text{Sm}(d,t)^{146}\text{Sm}$	-98	10	-84.1	2.8	1.4	U			McM		75Si03
$^{147}\text{Tb}(p)^{146}\text{Gd}$	-1945	18	-1946	9	0.0	1	23	19 $^{147}\text{Tb}$			87Sc.A
$^{147}\text{Tm}(p)^{146}\text{Er}$	1062.2	6.	1059	3	-0.6	o					82K103
	1058.2	3.3			0.1	1	94	55 $^{147}\text{Tm}$	Dap		93Se04 *
	1067.3	15.			-0.6	U			ORp		03Gi10
$^{147}\text{Tm}^m(p)^{146}\text{Er}$	1124.7	6.	1120	3	-0.7	2					84Ho.A
	1118.5	3.9			0.5	2			Dap		93Se04

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{147}\text{Ba}(\beta^-)^{147}\text{La}$	5750	50	6414	22	13.3	C			Bwg		87Gr.A
$^{147}\text{La}(\beta^-)^{147}\text{Ce}$	4945	55	5336	14	7.1	C			Bwg		87Gr.A
	5150	40			4.6	B			Kur		95Ik03
	5370	100			-0.3	o			Kur		02Sh.B
	5366	40			-0.8	U			Kur		09Ha.B
$^{147}\text{Ce}(\beta^-)^{147}\text{Pr}$	3290	40	3430	16	3.5	C			Bwg		87Gr.A
	3426	20			0.2	1	60	52 $^{147}\text{Pr}$	Kur		95Ik03
	3380	100			0.5	U			Kur		02Sh.B
$^{147}\text{Pr}(\beta^-)^{147}\text{Nd}$	2700	200	2703	16	0.0	U					64Ho03
	2790	100			-0.9	U					81Ya06 *
	2711	28			-0.3	-			Kur		95Ik03
ave.	2697	23			0.2	1	48	48 $^{147}\text{Pr}$			average
$^{147}\text{Nd}(\beta^-)^{147}\text{Pm}$	894.6	1.0	895.5	0.5	0.9	1	23	13 $^{147}\text{Pm}$			67Ca18 *
$^{147}\text{Pm}(\beta^-)^{147}\text{Sm}$	223.2	0.5	224.09	0.29	1.8	-					50La04
	224.3	1.3			-0.2	-					58Ha32
	224.5	0.4			-1.0	-					66Hs01
ave.	224.0	0.3			0.3	1	93	87 $^{147}\text{Pm}$			average
$^{147}\text{Eu}(\beta^+)^{147}\text{Sm}$	1767	10	1721.6	2.3	-4.5	B					67Ad03
	1723	3			-0.5	1	58	57 $^{147}\text{Eu}$			80Bu04
	1702	13			1.5	U					84Sc18 *
	1692	18			1.6	U					84Sc18
$^{147}\text{Gd}(\beta^+)^{147}\text{Eu}$	2185	5	2187.8	2.5	0.6	1	25	19 $^{147}\text{Eu}$			80Vy01 *
	2199	17			-0.7	U					84Sc18 *
$^{147}\text{Tb}(\beta^+)^{147}\text{Gd}$	4700	90	4614	8	-1.0	U					83Ve06 *
	4490	60			2.1	U			Got		85Ti01
	4560	50			1.1	U					Averag *
	4609	15			0.4	1	29	28 $^{147}\text{Tb}$	GSI		91Ke11 *
	4509	60			1.8	U			IRS		93Al03 *
$^{147}\text{Dy}(\beta^+)^{147}\text{Tb}$	6334	60	6547	12	3.5	B			IRS		83Al06 *
	6480	100			0.7	U			IRS		83Al18 *
	6334	60			3.5	C					85Af.A *
	6480	100			0.7	U			IRS		85Al08 *
* $^{147}\text{Tb}-u$	$M - A = -70707(28)$ keV for mixture gs+m at 50.6 keV										
* $^{147}\text{Tb}-^{133}\text{Cs}_{1.105}$	$D_M = 28574(12)$ $\mu\text{u}$ for mixture gs+m at 50.6 keV with ratio $R = 0.741(13)$										
* $^{147}\text{Dy}-u$	$M - A = -63437(28)$ keV for $^{147}\text{Dy}^m$ at 750.5 keV										
* $^{147}\text{Dy}-^{133}\text{Cs}_{1.105}$	$D_M = 35567(14)$ and $D_M = 36358.4(9.5)$ for $^{147}\text{Dy}^m$ at 750.5 keV										
* $^{147}\text{Er}-^{133}\text{Cs}_{1.105}$	$D_M = 54531(11)$ $\mu\text{u}$ for mixture gs+m at 100#50 keV with ratio $R = 0.741(13)$										
*	error due to excitation energy, use only next item										
* $^{147}\text{Er}-^{85}\text{Rb}_{1.729}$	$D_M = 102559.5(8.3)$ $\mu\text{u}$ for mixture gs+m at 100#50 keV with $R = 0.741(13)$										
* $^{147}\text{Tm}(p)^{146}\text{Er}$	$Q_p$ from $E_p = 1051.0(3.3)$ , no screening correction should be applied										
* $^{147}\text{Pr}(\beta^-)^{147}\text{Nd}$	$E_{\beta^-} = 2760(100)$ to 49.93, 1450(100) to 1310.7 and 1350.5 keV										
* $^{147}\text{Nd}(\beta^-)^{147}\text{Pm}$	$E_{\beta^-} = 803.5(1.0)$ to $5/2^+$ level at 91.1047 keV										
* $^{147}\text{Eu}(\beta^+)^{147}\text{Sm}$	$p^+ = 2.9(0.3) \times 10^{-3}$ to $3/2^-$ level at 197.284 keV										
* $^{147}\text{Eu}(\beta^+)^{147}\text{Sm}$	$pK = 0.724(0.026)$ to $(3/2^+, 5/2^+)$ level at 1548.634 keV										
* $^{147}\text{Gd}(\beta^+)^{147}\text{Eu}$	$E_{\beta^+} = 933(5)$ to $7/2^+$ level at 229.323 keV										
* $^{147}\text{Gd}(\beta^+)^{147}\text{Eu}$	$pK = 0.694(0.016)$ to 2073 level, recalculated by AHW										
* $^{147}\text{Tb}(\beta^+)^{147}\text{Gd}$	$E_{\beta^+} = 2460(80)$ to $3/2^+$ level at 1152.56 and $1/2^+$ at 1292.3 keV, reinterpreted										
* $^{147}\text{Tb}(\beta^+)^{147}\text{Gd}$	Average $\text{KLM}/\beta^+ = 2.03(0.15) \rightarrow E_{\beta^+} = 2190(50)$ from $^{147}\text{Tb}^m$ at 50.6(0.9) to $9/2^-$ level at 1397.00 from 3 references (no side-feeding correction applied):										
*	$p^+ = 0.32(0.07)$ gives $\text{KLM}/\beta^+ = 2.2(0.8)$										
*	$\text{KLM}/\beta^+ = 2.17(0.30)$										
*	$\beta^+/K = 0.59(0.05)$ gives $\text{KLM}/\beta^+ = 1.99(0.17)$										
* $^{147}\text{Tb}(\beta^+)^{147}\text{Gd}$	$Q_{\beta^+} = 4660(15)$ 4560(60) respectively, from $^{147}\text{Tb}^m$ at 50.6(0.9) keV										
* $^{147}\text{Dy}(\beta^+)^{147}\text{Tb}$	$E_{\beta^+} = 6012(60)$ from $^{147}\text{Dy}^m$ at 750.5 to $^{147}\text{Tb}^m$ at 50.6(0.9) keV										
* $^{147}\text{Dy}(\beta^+)^{147}\text{Tb}$	$Q_{\beta^+} = 7180(100)$ from $^{147}\text{Dy}^m$ at 750.5 to $^{147}\text{Tb}^m$ at 50.6(0.9) keV										
* $^{147}\text{Dy}(\beta^+)^{147}\text{Tb}$	$E_{\beta^+} = 6012(60)$ from $^{147}\text{Dy}^m$ at 750.5 to $^{147}\text{Tb}^m$ at 50.6(0.9) keV										
* $^{147}\text{Dy}(\beta^+)^{147}\text{Tb}$	$Q_{\beta^+} = 7180(100)$ from $^{147}\text{Dy}^m$ at 750.5 to $^{147}\text{Tb}^m$ at 50.6(0.9) keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{148}\text{Cs}-^{133}\text{Cs}_{1.113}$	54871	14				2			MA8	1.0	15At.A
$^{148}\text{La}-u$	-67320.6	20.9				2			CP1	1.0	06Sa56
$^{148}\text{Ce}-u$	-75578.2	13.0	-75576	12	0.2	1	85	$^{85}\text{ }^{148}\text{Ce}$	CP1	1.0	06Sa56
$^{148}\text{Pr}-u$	-77781	41	-77870	16	-2.2	B			CP1	1.0	06Sa56 *
$\text{C}_{12}\text{H}_4-^{148}\text{Nd}$	114261	34	114401.0	2.3	1.0	U			R05	4.0	65De13
$\text{C}_9\text{N O H}_{10}-^{148}\text{Nd}$	159186	59	159339.9	2.3	0.7	U			R05	4.0	65De13
$\text{C}_9\text{H}_8\text{O}_2-^{148}\text{Sm}$	137540	26	137600.5	1.5	0.6	U			R04	4.0	64De15
$\text{C}_9\text{H}_{10}\text{N O}-^{148}\text{Sm}$	161275	31	161409.9	1.5	1.1	U			R04	4.0	64De15
$\text{C}_8\text{ }^{13}\text{C H}_7\text{O}_2-^{148}\text{Sm}$	133030	60	133130.3	1.5	0.4	U			R04	4.0	64De15
$^{148}\text{Eu}-^{133}\text{Cs}_{1.113}$	23315	15	23321	11	0.4	1	51	$^{51}\text{ }^{148}\text{Eu}$	MA5	1.0	00Be42
$^{148}\text{Tb}-u$	-75692	41	-75725	13	-0.8	U			GS2	1.0	05Li24 *
$^{148}\text{Dy}-^{133}\text{Cs}_{1.113}$	32394	16	32382	9	-0.8	-			MA5	1.0	00Be42
	32380	14			0.1	-			SH1	1.0	07Ra37
	ave.	32386			-0.4	1	79	$^{79}\text{ }^{148}\text{Dy}$			average
$^{148}\text{Ho}-u$	-62201	100	-62260	90	-0.6	U			GS2	1.0	05Li24 *
$^{148}\text{Ho}-^{85}\text{Rb}_{1.741}$	91318	90				2			SH1	1.0	07Ra37 *
$^{148}\text{Er}-^{133}\text{Cs}_{1.113}$	49967	11				2			SH1	1.0	07Ra37
$^{148}\text{Tm}-^{133}\text{Cs}_{1.113}$	63616	11				2			SH1	1.0	07Ra37
$^{148}\text{Eu}-^{142}\text{Sm}_{1.042}$	6451	17	6446	11	-0.3	1	41	$^{38}\text{ }^{148}\text{Eu}$	MA7	1.0	01Bo59
$^{148}\text{Nd }^{35}\text{Cl}_2-^{144}\text{Nd }^{37}\text{Cl}_2$	12690	9	12706.4	1.8	0.7	U			H21	2.5	70Ma05
	12703.6	2.1			0.5	1	12	$^{11}\text{ }^{148}\text{Nd}$	H25	2.5	72Ba08
$^{148}\text{Sm }^{35}\text{Cl}_2-^{144}\text{Sm }^{37}\text{Cl}_2$	8710	10	8722.9	0.9	0.3	U			H12	4.0	64Ba15
	8721.4	2.6			0.2	U			H25	2.5	72Ba08
$^{148}\text{Nd }^{35}\text{Cl}-^{146}\text{Nd }^{37}\text{Cl}$	6740	5	6726.7	1.8	-0.7	U			H12	4.0	64Ba15
	6721	4			0.6	U			H21	2.5	70Ma05
	6723.8	2.7			0.4	U			H25	2.5	72Ba08
	6725.7	0.9			0.4	1	62	$^{61}\text{ }^{148}\text{Nd}$	H26	2.5	73Me28
$^{148}\text{Sm }^{35}\text{Cl}-^{146}\text{Nd }^{37}\text{Cl}$	4656	3	4656.6	0.5	0.1	U			H12	4.0	64Ba15
$^{148}\text{Sm}-^{147}\text{Sm}$	110	44	-75.05	0.28	-1.1	U			R04	4.0	64De15
$^{148}\text{Nd}-^{146}\text{Nd}$	3866	50	3776.6	1.8	-0.4	U			R05	4.0	65De13
	3773	3			0.5	U			M17	2.5	66Be10
$^{145}\text{Cs}-^{148}\text{Cs}_{.392}\text{ }^{143}\text{Cs}_{.608}$	-370	90	-519	11	-0.7	U			P33	2.5	86Au02
$^{148}\text{Sm}(\alpha)^{144}\text{Nd}$	2014.6	20.	1986.8	0.4	-1.4	U					70Gu14
	1987.3	0.5			-1.0	1	52	$^{26}\text{ }^{144}\text{Nd}$			16Ca.1
$^{148}\text{Eu}(\alpha)^{144}\text{Pm}$	2703.2	30.	2692	10	-0.4	1	11	$^{10}\text{ }^{148}\text{Eu}$			64To04
$^{148}\text{Gd}(\alpha)^{144}\text{Sm}$	3271.29	0.03	3271.29	0.03	0.0	1	100	$^{96}\text{ }^{148}\text{Gd}$			73Go29
$^{146}\text{Nd}(t,p)^{148}\text{Nd}$	4139	30	4143.0	1.7	0.1	U			Ald		72Ch11
$^{148}\text{Sm}(p,t)^{146}\text{Sm}$	-6011	8	-6000.8	2.8	1.3	1	12	$^{12}\text{ }^{146}\text{Sm}$	Min		72De47
	-6018	15			1.1	U			Ham		74Oe03
$^{148}\text{Gd}(p,t)^{146}\text{Gd}$	-7844	14	-7844	4	0.0	U			LAL		83Fl05
$^{148}\text{Gd}(p,t)^{146}\text{Gd}-^{65}\text{Cu}(\alpha)^{63}\text{Cu}$	1500	4	1500	4	0.1	1	90	$^{89}\text{ }^{146}\text{Gd}$	Liv		86Ma40
$^{148}\text{Nd}(d,^3\text{He})^{147}\text{Pr}$	-3726	40	-3759	16	-0.8	R			KVI		79Sa.A
$^{148}\text{Nd}(d,t)^{147}\text{Nd}$	-1072	4	-1075.3	1.7	-0.8	1	17	$^{17}\text{ }^{148}\text{Nd}$	McM		77St22
$^{147}\text{Sm}(n,\gamma)^{148}\text{Sm}$	8139.8	1.2	8141.23	0.26	1.2	U					69Re04 Z
	8141.1	1.5			0.1	U					70Bu19 Z
	8141.8	0.8			-0.7	-					71Gr37 Z
	8141.3	0.3			-0.2	-			Bdn		06Fi.A
$^{147}\text{Sm}(d,p)^{148}\text{Sm}$	5920	10	5916.66	0.26	-0.3	U			Tal		64Ke03
$^{148}\text{Sm}(d,t)^{147}\text{Sm}$	-1890	15	-1884.00	0.26	0.4	U			Kop		67Ve04
$^{147}\text{Sm}(n,\gamma)^{148}\text{Sm}$	ave.	8141.36	8141.23	0.26	-0.5	1	84	$^{51}\text{ }^{147}\text{Sm}$			average
$^{148}\text{Gd}(p,d)^{147}\text{Gd}$	-6755	5	-6759.1	1.2	-0.8	U					86Ru04
$^{148}\text{Gd}(p,d)^{147}\text{Gd}-^{148}\text{Sm}(\alpha)^{147}\text{Sm}$	-842	2	-842.5	1.2	-0.2	-					86Ru04
$^{148}\text{Gd}(d,t)^{147}\text{Gd}-^{148}\text{Sm}(\alpha)^{147}\text{Sm}$	-843	2			0.3	-					86Ru04
$^{148}\text{Gd}(^3\text{He},\alpha)^{147}\text{Gd}-^{148}\text{Sm}(\alpha)^{147}\text{Sm}$	-842	3			-0.2	-					86Ru04
$^{148}\text{Gd}(p,d)^{147}\text{Gd}-^{148}\text{Sm}(\alpha)^{147}\text{S}$	ave.	-842.4			-0.1	1	89	$^{86}\text{ }^{147}\text{Gd}$			average
$^{148}\text{Ba}(\beta^-)^{148}\text{La}$	5115	60				3			Bwg		90Gr10



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{148}\text{La}(\beta^-)^{148}\text{Ce}$	7310 140	7690 22	2.7	U			Trs		82Br23 *
	7255 55		7.9	B			Bwg		90Gr10
	7650 100		0.4	U			Kur		02Sh.B
	7732 70		-0.6	U			Kur		09Ha.B
$^{148}\text{Ce}(\beta^-)^{148}\text{Pr}$	2060 75	2137 13	1.0	U			Bwg		87Gr.A
	2140 14		-0.2	1	81	66 $^{148}\text{Pr}$	Kur		95Ik03
$^{148}\text{Pr}(\beta^-)^{148}\text{Nd}$	4800 200	4873 15	0.4	U					79Ik06
	4965 100		-0.9	U			Bwg		87Gr.A
	4890 50		-0.3	-					88Ka14
	4880 30		-0.2	-			Kur		95Ik03
	4930 100		-0.6	U			Kur		02Sh.B
$^{148}\text{Pm}(\beta^-)^{148}\text{Sm}$	ave. 4883 26		-0.4	1	34	34 $^{148}\text{Pr}$			average
	2480 15	2471 6	-0.6	R					62Sc04 *
$^{148}\text{Eu}(\beta^+)^{148}\text{Sm}$	2475 30		-0.1	U					63Ba31 *
	3122 30	3037 10	-2.8	U					63Ba32 *
$^{148}\text{Tb}(\beta^+)^{148}\text{Gd}$	3150 30		-3.8	B					70Ag01 *
	5630 80	5732 13	1.3	F					76Cr.B *
$^{148}\text{Dy}(\beta^+)^{148}\text{Tb}$	5835 70		-1.5	U					83Ve06 *
	5710 100		0.2	U			Got		85Sc09 *
	5390 100		3.4	B			Got		85Ti01 *
	5760 80		-0.3	U			IRS		93Al03 *
	5752 40		-0.5	1	10	10 $^{148}\text{Tb}$	GSI		95Ke05 *
	2660 60	2678 10	0.3	U					81Sc21
	2805 60		-2.1	U					81Sp03 *
	2700 60		-0.4	U			IRS		82Al.A
	2700 60		-0.4	U					82Ve.A
	2722 60		-0.7	U					83Ve06 *
$^{148}\text{Ho}^m(\beta^+)^{148}\text{Dy}$	2835 95		-1.7	U					84Ha.B *
	2740 60		-1.0	U			Got		85Sc09 *
$^{148}\text{Ho}^m(\beta^+)^{148}\text{Dy}$	2682 10		-0.4	1	92	86 $^{148}\text{Tb}$	GSI		95Ke05 *
	9400 250	10120# 130#	2.9	B			IRS		93Al03
* $^{148}\text{Pr}-u$	$D_M = -77739.3(30.6) \mu\text{u}$ for mixture gs+m at 76.80; $M - A = -72413.7(28.5) \text{ keV}$								Nub16b **
* $^{148}\text{Tb}-u$	$M - A = -70462(28) \text{ keV}$ for mixture gs+m at 90.1 keV								Nub16b **
* $^{148}\text{Ho}-u$	$M - A = -57815(30) \text{ keV}$ for mixture gs+m at 250#100 keV								Nub16b **
*	outweighed by next item before correcting for isomeric mixture								GAu **
* $^{148}\text{Ho}-^{85}\text{Rb}_{1,741}$	$D_M = 91517.5(9.5) \mu\text{u}$ for mixture gs+m at 250#100 keV with $R = 0.74(15)$								Nub16b **
* $^{148}\text{La}(\beta^-)^{148}\text{Ce}$	$E_{\beta^-} = 5862(100)$ supposed to go to levels around $E = 1450(100) \text{ keV}$								90Gr10 **
* $^{148}\text{Pm}(\beta^-)^{148}\text{Sm}$	$E_{\beta^-} = 2460(20) 1020(15)$ to ground state, $1^-$ level at 1465.137 keV								Ens144 **
* $^{148}\text{Pm}(\beta^-)^{148}\text{Sm}$	$E_{\beta^-} = 2480(30) 1930(30) 1020(30)$ to ground state, $2^+$ at 550.255, $1^-$ at 1465.137 keV								Ens144 **
*	and $E_{\beta^-} = 400(30)$ from $^{148}\text{Pm}^m$ at 137.9 to $6^+$ level at 2194.061 keV								Nub16b **
* $^{148}\text{Eu}(\beta^+)^{148}\text{Sm}$	$E_{\beta^+} = 920(30)$ to 1180.261 keV $4^+$ level								Ens144 **
* $^{148}\text{Eu}(\beta^+)^{148}\text{Sm}$	$E_{\beta^+} = 540(30)$ to 1594.247 keV $5^-$ level, and other $E_{\beta^+}$								Ens144 **
* $^{148}\text{Tb}(\beta^+)^{148}\text{Gd}$	$E_{\beta^+} = 4610(80)$ assumed to ground state								76Cr.B **
*	$F$ : since $^{148}\text{Tb}$ ground state $2^-$ , transition to $^{148}\text{Gd}$ ground state weak								AHW **
* $^{148}\text{Tb}(\beta^+)^{148}\text{Gd}$	$E_{\beta^+} = 2210(70)$ from $^{148}\text{Tb}^m$ at 90.1 to $8^+$ level at 2693.35 keV and								Nub16b **
*	$E_{\beta^+} = 4560(80)$ mainly to $2^+$ at 784.433 keV. Conflicting, not used								Ens144 **
* $^{148}\text{Tb}(\beta^+)^{148}\text{Gd}$	$p^+ = 0.271(0.10) \rightarrow E_{\beta^+} = 1920(30)$ from $^{148}\text{Tb}^m$ at 90.1 to $8^+$ at 2693.35 keV								Ens144 **
*	but assuming 5(5)% side-feeding; see reference								90Sa32 **
* $^{148}\text{Tb}(\beta^+)^{148}\text{Gd}$	$KL/\beta^+ = 1.54(0.09)$ to 1863.42 level; yields $Q_{\beta^+} = 5295(45) \text{ keV}$								85Ti01 **
*	but assuming 7(7)% side-feeding; see 1990Sa32								AHW **
* $^{148}\text{Tb}(\beta^+)^{148}\text{Gd}$	$Q_{\beta^+} = 5700(80)$ ; and 5910(80) from $^{148}\text{Tb}^m$ at 90.1 keV								Nub16b **
* $^{148}\text{Tb}(\beta^+)^{148}\text{Gd}$	$Q_{\beta^+} = 5750(40)$ ; and 5846(50) from $^{148}\text{Tb}^m$ at 90.1 keV								Nub16b **
* $^{148}\text{Dy}(\beta^+)^{148}\text{Tb}$	$p^+ = 0.069(0.014)$ to $1^+$ level at 620.24 keV, recalculated $Q$								Ens144 **
* $^{148}\text{Dy}(\beta^+)^{148}\text{Tb}$	$E_{\beta^+} = 1040(60), 1120(60)$ to $1^+$ level at 620.24 keV								Ens144 **
* $^{148}\text{Dy}(\beta^+)^{148}\text{Tb}$	$p^+ = 0.055(0.015)$ to $1^+$ level at 620.24 keV								Ens144 **
* $^{148}\text{Dy}(\beta^+)^{148}\text{Tb}$	$\beta^+/K = 0.045(0.005)$ to $1^+$ level at 620.24 keV, gives $Q_{\beta^+} = 2680(30) \text{ keV}$								Ens144 **
* $^{148}\text{Dy}(\beta^+)^{148}\text{Tb}$	GSI average of $E_{\beta^+} = 1043(10)$ and 1036(10) of reference								91Ke11 **
*	to $1^+$ level at 620.24 keV								Ens144 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{149}\text{Ba}-u$	-57027	188				2			GT3	2.5	16Kn03
$^{149}\text{Ce}-u$	-71573.1	11.0				2			CP1	1.0	06Sa56
$^{149}\text{Pr}-u$	-76263.9	10.6				2			CP1	1.0	06Sa56
$\text{C}_8 \text{ }^{13}\text{C} \text{ H}_8 \text{ O}_2-^{149}\text{Sm}$	138597	29	138593.0	1.4	0.0	U			R04	4.0	64De15
$\text{C}_9 \text{ H}_{11} \text{ N O}-^{149}\text{Sm}$	166820	33	166872.6	1.4	0.4	U			R04	4.0	64De15
$\text{C}_8 \text{ }^{13}\text{C} \text{ H}_{10} \text{ N O}-^{149}\text{Sm}$	162408	46	162402.4	1.4	0.0	U			R04	4.0	64De15
$^{149}\text{Eu}-^{133}\text{Cs}_{1,120}$	23849	17	23831	4	-1.1	U			MA5	1.0	00Be42
$^{149}\text{Tb}-u$	-76730	32	-76746	4	-0.5	U			GS2	1.0	05Li24 *
$^{149}\text{Dy}-^{133}\text{Cs}_{1,120}$	33278	109	33219	10	-0.5	U			MA5	1.0	00Be42
$^{149}\text{Dy}-u$	-72698	30	-72675	10	0.8	U			GS2	1.0	05Li24 *
$^{149}\text{Ho}-u$	-66179	34	-66180	13	0.0	1	14	14 $^{149}\text{Ho}$	GS2	1.0	05Li24 *
$^{149}\text{Er}-u$	-57694	30				2			GS2	1.0	05Li24 *
$^{149}\text{Eu}-^{142}\text{Sm}_{1,049}$	6909	18	6888	5	-1.2	U			MA7	1.0	01Bo59
$^{149}\text{Dy}-^{142}\text{Sm}_{1,049}$	16249	16	16276	10	1.7	1	39	36 $^{149}\text{Dy}$	MA7	1.0	01Bo59
$^{149}\text{Sm} \text{ }^{35}\text{Cl}-^{147}\text{Sm} \text{ }^{37}\text{Cl}$	5257	4	5237.4	1.0	-1.2	U			H12	4.0	64Ba15
	5231	3			0.9	U			H21	2.5	70Ma05
	5239.8	0.8			-1.2	1	24	14 $^{147}\text{Sm}$	M21	2.5	75Ka25
$^{149}\text{Sm}-^{148}\text{Sm}$	2282	31	2362.4	1.0	0.6	U			R04	4.0	64De15
$^{149}\text{Sm}-^{147}\text{Sm}$	2320	60	2287.3	1.0	-0.1	U			R04	4.0	64De15
$^{149}\text{Gd}(\alpha)^{145}\text{Sm}$	3102.3	10.	3099	3	-0.3	-					65Ma51 Z
	3096.1	10.3			0.3	-			ORa		66Wi12 Z
	3099.1	5.			0.1	-			DbA		67Go32 Z
ave.	3099	4			0.1	1	56	53 $^{149}\text{Gd}$			average
$^{149}\text{Tb}(\alpha)^{145}\text{Eu}$	4074.4	3.	4077.9	2.2	1.2	-			DbA		67Go32 Z
	4073.8	7.			0.6	U			ORa		74To07 *
	4074.6	10.			0.3	U					81Ho.A Z
	4081.8	5.			-0.8	-			Bka		82Bo04 Z
	4082.8	4.			-1.2	-			Daa		96Pa01
ave.	4078.2	2.2			-0.1	1	95	86 $^{149}\text{Tb}$			average
$^{149}\text{Sm}(n,\alpha)^{146}\text{Nd}$	9429	4	9436.5	1.0	1.9	U			McM		67Oa01
	9421	15			1.0	U			ILL		75Em.A
$^{149}\text{Sm}(p,t)^{147}\text{Sm}$	-5532	8	-5530.2	0.9	0.2	U			Min		72De47
	-5532	7			0.3	U			McM		73Ga04
$^{148}\text{Nd}(n,\gamma)^{149}\text{Nd}$	5038.76	0.10	5038.79	0.07	0.3	2			ILn		76Pi04 Z
	5038.82	0.11			-0.3	2			Bdn		06Fi.A
$^{148}\text{Nd}(^3\text{He},d)^{149}\text{Pm}$	455	5	451.8	2.5	-0.6	1	24	13 $^{149}\text{Pm}$	McM		80St10 *
$^{149}\text{Sm}(d,^3\text{He})^{148}\text{Pm}$	-2064	6	-2066	6	-0.3	2					88No02
$^{148}\text{Sm}(n,\gamma)^{149}\text{Sm}$	5872.5	1.8	5870.8	0.9	-0.9	1	25	15 $^{148}\text{Sm}$			70Sm.A
	5850.8	0.6			33.3	C					82Ba15
$^{149}\text{Sm}(\gamma,n)^{148}\text{Sm}$	-5890	160	-5870.8	0.9	0.1	U			Phi		60Ge01
$^{148}\text{Sm}(d,p)^{149}\text{Sm}$	3656	15	3646.2	0.9	-0.7	U			Kop		67Ve04
$^{149}\text{Er}(\text{ep})^{148}\text{Dy}$	5758	900	6829	29	1.2	U					83La.A *
	7080	470			-0.5	U			LBL		89Fi01
$^{149}\text{La}(\beta^-)^{149}\text{Ce}$	6450	200				3			Kur		02Sh.B
$^{149}\text{Ce}(\beta^-)^{149}\text{Pr}$	4190	75	4369	14	2.4	U			Bwg		87Gr.A
	4380	60			-0.2	U			Kur		95Ik03
	4310	100			0.6	U			Kur		02Sh.B
$^{149}\text{Pr}(\beta^-)^{149}\text{Nd}$	3000	200	3336	10	1.7	U					67Va14
	3390	90			-0.6	U			Kur		95Ik03
$^{149}\text{Nd}(\beta^-)^{149}\text{Pm}$	1669	10	1688.8	2.5	2.0	U					64Go08 *
$^{149}\text{Pm}(\beta^-)^{149}\text{Sm}$	1072	2	1071.5	1.9	-0.3	1	88	87 $^{149}\text{Pm}$			60Ar05
	1062	2			4.7	B					78Re01
$^{149}\text{Eu}(\epsilon)^{149}\text{Sm}$	680	10	695	4	1.5	1	14	14 $^{149}\text{Eu}$			85Ad.A
$^{149}\text{Gd}(\epsilon)^{149}\text{Eu}$	1308	6	1314	4	1.0	1	48	30 $^{149}\text{Eu}$	Got		84Sc.B
$^{149}\text{Tb}(\beta^+)^{149}\text{Gd}$	3575	50	3638	4	1.3	U			Got		85Sc09 *
	3635	10			0.3	1	19	10 $^{149}\text{Tb}$	GSI		91Ke06 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{149}\text{Dy}(\beta^+)^{149}\text{Tb}$	3930	150	3793	9	-0.9	U					84A136 *	
	3925	65			-2.0	U			Got		90Sa32 *	
	3797	13			-0.3	1	50	46	$^{149}\text{Dy}$	GSI	91Ke11 *	
	3950	100			-1.6	U			IRS		93A103	
$^{149}\text{Ho}(\beta^+)^{149}\text{Dy}$	6043	50	6049	13	0.1	-			IRS		83A106 *	
	5330	100			7.2	C					84Ha.B	
	6000	90			0.5	U			IRS		93A103	
	6009	20			2.0	-			GSI		91Ke11 *	
	ave.	6014	19		1.9	1	48	32	$^{149}\text{Ho}$		average	
$^{149}\text{Er}(\epsilon)^{149}\text{Ho}$	8610	650	7900	30	-1.1	U			LBL		89Fi01 *	
* $^{149}\text{Tb}-u$	$M-A=-71456(28)$ keV for mixture gs+m at 35.78 keV										Nub16c **	
* $^{149}\text{Dy}-u$	$M-A=-65057(28)$ keV for $^{149}\text{Dy}^m$ at 2661.1 keV										Nub16b **	
* $^{149}\text{Ho}-u$	$M-A=-61621(28)$ keV for mixture gs+m at 48.80 keV										Nub16b **	
* $^{149}\text{Er}-u$	$M-A=-53000(28)$ keV for $^{149}\text{Er}^m$ at 741.8 keV										Nub16b **	
* $^{149}\text{Tb}(\alpha)^{145}\text{Eu}$	$E_\alpha=3999(7)$ from $^{149}\text{Tb}^m$ at 35.78 keV										Nub16c **	
* $^{148}\text{Nd}(^3\text{He},d)^{149}\text{Pm}$	Based on $^{146}\text{Nd}(^3\text{He},d)^{147}\text{Pm}$ $Q=-87.6(0.9)$ keV										AHW **	
* $^{149}\text{Er}(\epsilon p)^{148}\text{Dy}$	As read from graph; $Q=6500$ from $^{149}\text{Er}^m$ at 741.8 keV										Nub16b **	
* $^{149}\text{Nd}(\beta^-)^{149}\text{Pm}$	$E_{\beta^-}=1555(10)$ to $5/2^+$ level at 114.312 level										Ens046 **	
* $^{149}\text{Tb}(\beta^+)^{149}\text{Gd}$	$\beta^+/\text{K}=0.31(0.03)$ from $^{149}\text{Tb}^m$ at 35.78 to $9/2^-$ level at 795.82 keV										Ens046 **	
* $^{149}\text{Tb}(\beta^+)^{149}\text{Gd}$	$E_{\beta^+}=1853(10)$ from $^{149}\text{Tb}^m$ at 35.78 to $9/2^-$ level at 795.82 keV										Ens046 **	
* $^{149}\text{Dy}(\beta^+)^{149}\text{Tb}$	$E_{\beta^+}=1030(150)$ to 1728.31-1876.96 levels										Ens046 **	
* $^{149}\text{Dy}(\beta^+)^{149}\text{Tb}$	KL/ $\beta^+=29.4(10.6)$ , 14.5(3.7), 17.6(4.9) to 1883, 1735, 1728 levels										90Sa32 **	
* $^{149}\text{Dy}(\beta^+)^{149}\text{Tb}$	Original $Q=3812(10)$ from $E_{\beta^+}=1965(10)$ to 825.16 level corrected to $E_{\beta^+}=1950(13)$ for background subtraction										GAu **	
* $^{149}\text{Ho}(\beta^+)^{149}\text{Dy}$	$E_{\beta^+}=3930(50)$ to $9/2^-$ level at 1090.76 keV										Ens046 **	
* $^{149}\text{Ho}(\beta^+)^{149}\text{Dy}$	$E_{\beta^+}=3896(20)$ to $9/2^-$ level at 1090.76 keV										Ens046 **	
* $^{149}\text{Er}(\epsilon)^{149}\text{Ho}$	KLM/ $\beta^+=0.68(0.34)$ from $^{149}\text{Er}^m$ at 741.8 to 4699.7 level										Ens046 **	
$^{150}\text{Ba}-u$	-55309	371	-53570#	320#	1.9	D			GT3	2.5	16Kn03 *	
$^{150}\text{La}-u$	-60258	187				2			GT3	2.5	16Kn03	
$^{150}\text{Ce}-u$	-69618.6	13.1	-69616	13	0.2	1	92	92	$^{150}\text{Ce}$	1.0	06Sa56	
$^{150}\text{Pr}-u$	-73322.9	10.6	-73324	10	-0.1	1	83	83	$^{150}\text{Pr}$	1.0	06Sa56	
$\text{C}_{12}\text{H}_6-^{150}\text{Nd}$	126194	43	126048.7	1.4	-0.8	U			R05	4.0	65De13	
$\text{C}_8\text{ }^{13}\text{C N O H}_{11}-^{150}\text{Nd}$	166439	34	166517.3	1.4	0.6	U			R05	4.0	65De13	
$\text{C}_9\text{ N O H}_{12}-^{150}\text{Nd}$	170931	46	170987.5	1.4	0.3	U			R05	4.0	65De13	
$\text{C}_{12}\text{H}_6-^{150}\text{Sm}$	129810	140	129668.0	1.4	-0.3	U			R04	4.0	64De15	
$\text{C}_8\text{ }^{13}\text{C H}_{11}\text{ N O}-^{150}\text{Sm}$	170029	25	170136.6	1.4	1.1	U			R04	4.0	64De15	
$\text{C}_9\text{ H}_{12}\text{ N O}-^{150}\text{Sm}$	174612	47	174606.8	1.4	0.0	U			R04	4.0	64De15	
$^{150}\text{Tb}^m-u$	-75850	30	-75840	28	0.3	1	89	89	$^{150}\text{Tb}^m$	1.0	05Li24	
$^{150}\text{Ho}-^{133}\text{Cs}_{1,128}$	40150	29	40149	15	-0.1	-			MA5	1.0	00Be42	
ave.	40132	21			0.8	1	53	53	$^{150}\text{Ho}$		average	
$^{150}\text{Ho}-u$	-66504	40	-66502	15	0.1	U			GS2	1.0	05Li24 *	
$^{150}\text{Er}-u$	-62060	30	-62084	18	-0.8	1	38	38	$^{150}\text{Er}$	1.0	05Li24	
$^{150}\text{Nd } ^{35}\text{Cl}_2-^{146}\text{Nd } ^{37}\text{Cl}_2$	13654	9	13679.2	1.1	1.1	U			H21	2.5	70Ma05	
	13672.5	1.8			1.5	U			H25	2.5	72Ba08	
$^{150}\text{Nd } ^{35}\text{Cl}-^{148}\text{Nd } ^{37}\text{Cl}$	7006	4	6952.5	2.0	-3.3	B			H12	4.0	64Ba15	
	6939	4			1.4	U			H21	2.5	70Ma05	
$^{150}\text{Sm } ^{35}\text{Cl}-^{148}\text{Sm } ^{37}\text{Cl}$	5452	8	5403.3	1.0	-1.5	U			H12	4.0	64Ba15	
	5400	4			0.3	U			H21	2.5	70Ma05	
	5404.8	0.6			-1.0	1	41	26	$^{148}\text{Sm}$	M21	2.5	75Ka25
$^{150}\text{Nd}-^{150}\text{Sm}$	3633	4	3619.33	0.21	-0.9	U			H19	4.0	64Mc11	
	3617.0	1.2			0.8	U			H25	2.5	72Ba08	
	3619.33	0.21			0.0	1	100	100	$^{150}\text{Nd}$	JY1	1.0	10Ko28
$^{150}\text{Sm}-^{149}\text{Sm}$	149	30	90.8	0.4	-0.5	U			R04	4.0	64De15	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{150}\text{Nd}-^{148}\text{Nd}$	3860	46	4002.4	2.0	0.8	U			R05	4.0	65De13
	3988	3			1.9	U			M17	2.5	66Be10
$^{150}\text{Sm}-^{148}\text{Sm}$	2430	50	2453.2	1.0	0.1	U			R04	4.0	64De15
$^{150}\text{Nd}-^{146}\text{Nd}$	7719	67	7779.0	1.0	0.2	U			R05	4.0	65De13
$^{150}\text{Gd}(\alpha)^{146}\text{Sm}$	2804.9	10.	2807	6	0.2	-					62Si14
	2792.6	18.			0.8	-					65Og01
	ave.	2802	9		0.6	1	45	39 $^{150}\text{Gd}$			average
$^{150}\text{Tb}(\alpha)^{146}\text{Eu}$	3585.5	5.	3587	5	0.3	1	92	80 $^{150}\text{Tb}$	DbA		67Go32 Z
$^{150}\text{Dy}(\alpha)^{146}\text{Gd}$	4345.8	5.	4351.3	1.5	1.1	-			DbA		67Go32 Z
	4349.5	5.			0.3	-			GSa		79Ho10 Z
	4351.3	3.			0.0	-			Bka		82Bo04 *
	4352.5	2.1			-0.6	-			Ora		82De11 Z
	ave.	4351.3	1.5			0.0	1	99	92 $^{150}\text{Dy}$		average
$^{148}\text{Nd}(\text{t,p})^{150}\text{Nd}$	3935	30	3932.6	1.9	-0.1	U			Ald		72Ch11
$^{148}\text{Sm}(\text{t,p})^{150}\text{Sm}$	5372	25	5375.7	0.9	0.1	U			Ald		66Bj01
$^{150}\text{Sm}(\text{p,t})^{148}\text{Sm}$	-5379	8	-5375.7	0.9	0.4	U			Min		72De47
	-5378	15			0.2	U			Ham		74Oe03
$^{150}\text{Nd}(\text{d},^3\text{He})^{149}\text{Pr}$	-4501	10	-4436	10	6.5	C			KVI		79Sa.A
$^{150}\text{Nd}(\text{d,t})^{149}\text{Nd}$	-1122	10	-1118.4	1.9	0.4	U			McM		73Bu02
$^{149}\text{Sm}(\text{n},\gamma)^{150}\text{Sm}$	7984.9	0.6	7986.7	0.4	3.0	B					69Re04 Z
	7986.7	1.5			0.0	-					70Bu19 Z
	7986.7	0.4			0.0	-			Bdn		06Fi.A
$^{149}\text{Sm}(\text{d,p})^{150}\text{Sm}$	5764	4	5762.2	0.4	-0.5	U			Tal		64Ke03
$^{150}\text{Sm}(\text{d,t})^{149}\text{Sm}$	-1738	15	-1729.5	0.4	0.6	U			Kop		67Ve04
$^{149}\text{Sm}(\text{n},\gamma)^{150}\text{Sm}$	ave.	7986.7	0.4	7986.7	0.4	0.0	1	95	79 $^{149}\text{Sm}$		average
	$^{150}\text{Lu}(\text{p})^{149}\text{Yb}$	1269.6	4.	1269.6	2.3	0.0	3				84Ho.A
	1269.6	4.			0.0	3			Dap		93Se04
$^{150}\text{Lu}^m(\text{p})^{149}\text{Yb}$	1269.6	4.			0.0	3			ORp		03Gi10
	1303.8	15.	1291	5	-0.8	o			ORp		00Gi01
	1285.6	8.			0.7	3			ORp		03Gi10
	1294.7	6.			-0.5	3			Arp		03Ro21
	3010	90	3454	14	4.9	C			Bwg		87Gr.A
$^{150}\text{Ce}(\beta^-)^{150}\text{Pr}$	3480	40			-0.7	1	13	8 $^{150}\text{Ce}$	Kur		95Ik03
	5690	80	5379	9	-3.9	C			Bwg		87Gr.A
$^{150}\text{Pr}(\beta^-)^{150}\text{Nd}$	5386	26			-0.3	1	12	12 $^{150}\text{Pr}$	Kur		95Ik03
	5290	100			0.9	U			Kur		02Sh.B
	3454	20				2					77Ho09
$^{150}\text{Pm}(\beta^-)^{150}\text{Sm}$	2222	25	2259	6	1.5	U					65Gu03 *
$^{150}\text{Eu}(\beta^+)^{150}\text{Sm}$	978	10	972	4	-0.6	-					63Yo07 *
	968	4			0.9	-					65Gu03 *
	ave.	969	4			0.6	1	91	53 $^{150}\text{Eu}$		average
$^{150}\text{Tb}(\beta^+)^{150}\text{Gd}$	4720	40	4658	8	-1.5	U					68Wi21
	4670	15			-0.8	1	31	20 $^{150}\text{Tb}$			76Cr.B
	4760	50			-2.0	U					77Ha31 *
	4620	60			0.6	U					83Ve06
$^{150}\text{Tb}^m(\beta^+)^{150}\text{Gd}$	5040	100	5119	27	0.8	U			IRS		93Al03
$^{150}\text{Dy}(\beta^+)^{150}\text{Tb}$	1760	40	1796	8	0.9	U					81Ka07 *
$^{150}\text{Ho}(\beta^+)^{150}\text{Dy}$	6980	150	7364	14	2.6	U					84Al36 *
	6560	100			8.0	B			IRS		93Al03
$^{150}\text{Ho}(\epsilon)^{150}\text{Dy}$	7400	200			-0.2	U					98Ag.A
	7372	27			-0.3	1	29	27 $^{150}\text{Ho}$			00Ca.A
	7444	126			-0.6	U					01Ro35
$^{150}\text{Ho}^m(\beta^+)^{150}\text{Dy}$	7360	50				2			IRS		83Al06 *
	6575	75	7360	50	10.5	C					84Ha.B *
	6625	120			6.1	B			Got		85Sc09 *
	6900	130			3.5	B			Got		90Sa32 *
	7060	80			3.7	C			IRS		93Al03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{150}\text{Er}(\beta^+)^{150}\text{Ho}$	4010	80	4115	14	1.3	o					82No08 *
	4105	75			0.1	U					84Ha.B *
	4108	15			0.4	1	82	62 $^{150}\text{Er}$	GSI		91Ke11 *
* $^{150}\text{Ba}-u$	Trends from Mass Surface TMS suggest $^{150}\text{Ba}$ 1620 less bound										WgM168**
* $^{150}\text{Ho}-u$	$M-A=-61948(28)$ keV for mixture gs+m at $-0(50)$ keV										Nub16b **
* $^{150}\text{Dy}(\alpha)^{146}\text{Gd}$	Recalibrated as in reference										91Ry01 **
* $^{150}\text{Eu}(\beta^+)^{150}\text{Sm}$	$E_{\beta^+}=1242(25)$ from $^{150}\text{Eu}^m$ at 41.7 keV										Nub16b **
* $^{150}\text{Eu}(\beta^-)^{150}\text{Gd}$	$Q_{\beta^-}=1020(10)$ 1010(4) respectively, from $^{150}\text{Eu}^m$ at 41.7 keV										Nub16b **
* $^{150}\text{Tb}(\beta^+)^{150}\text{Gd}$	$E_{\beta^+}=1655(80)$ to $2^+$ at 2091.623, and other $E_{\beta^+}$ . Orig. error 35 increased										Ens136 **
* $^{150}\text{Dy}(\beta^+)^{150}\text{Tb}$	$p^+=2(10)\times 10^{-3}$ to $1^+$ 397.2 level combined with lower limit through $^{146}\text{Gd}(\epsilon)$										Ens136 **
* $^{150}\text{Ho}(\beta^+)^{150}\text{Dy}$	$E_{\beta^+}=4550(150)$ to 1395.0 and 1456.8 levels										82No08 **
* $^{150}\text{Ho}^m(\beta^+)^{150}\text{Dy}$	$E_{\beta^+}=3940(50)$ to $8^+$ level at 2401.6 keV										Ens136 **
* $^{150}\text{Ho}^m(\beta^+)^{150}\text{Dy}$	$p^+=0.56(0.02)$ 0.58(0.07) respectively to $8^+$ level at 2401.6 keV										Ens136 **
* $^{150}\text{Ho}^m(\beta^+)^{150}\text{Dy}$	$Q_{\beta^+}=6819(+117-100)$ from $p^+$ to 2401.6 level; could be raised 140 if 4% feeding of higher Dy levels										90Sa32 **
* $^{150}\text{Er}(\beta^+)^{150}\text{Ho}$	$p^+=0.36(0.04)$ 0.39(0.04) to $1^+$ level at 476.16 keV										90Sa32 **
* $^{150}\text{Er}(\beta^+)^{150}\text{Ho}$	$E_{\beta^+}=2610(15)$ to $1^+$ level at 476.16 keV										Ens136 **
$^{151}\text{La}-u$	-58734	397	-57230	470	2.5	C				1.5	04Ma.A
	-57231	187				2				2.5	16Kn03
$^{151}\text{Ce}-u$	-65727.8	19.0				2				1.0	06Sa56
$^{151}\text{Pr}-u$	-71697.5	14.3	-71691	13	0.5	1	77	77 $^{151}\text{Pr}$	CP1	1.0	06Sa56
$\text{C}_{12}\text{H}_7-^{151}\text{Eu}$	134920	37	134918.4	1.4	0.0	U			R04	4.0	64De15
$\text{C}_{10}\text{H}_{15}\text{O}-^{151}\text{Eu}$	192490	70	192433.2	1.4	-0.2	U			R04	4.0	64De15
$^{151}\text{Eu}-^{85}\text{Rb}_{1.776}$	76520	15	76518.3	1.4	-0.1	U			MA5	1.0	00Be42
$^{151}\text{Tb}-u$	-76866	43	-76891	4	-0.6	U			GS2	1.0	05Li24 *
$^{151}\text{Dy}-u$	-73809	30	-73809	4	0.0	U			GS2	1.0	05Li24 *
$^{151}\text{Ho}-u$	-68323	33	-68302	9	0.6	U			GS2	1.0	05Li24 *
$^{151}\text{Er}-u$	-62528	30	-62551	18	-0.8	2			GS2	1.0	05Li24 *
	-62540	30			-0.4	2			GS2	1.0	05Li24 *
$^{151}\text{Eu}\ ^{35}\text{Cl}-^{149}\text{Sm}\ ^{37}\text{Cl}$	5620.3	2.6	5615.6	0.7	-0.7	U			H25	2.5	72Ba08
$^{151}\text{Eu}-^{150}\text{Sm}$	2800	60	2574.7	0.6	-0.9	U			R04	4.0	64De15
$^{151}\text{Eu}(\alpha)^{147}\text{Pm}$	1960	30	1964.5	1.1	0.1	U					07Be48
	1948.9	8.6			1.8	U					14Ca13
$^{151}\text{Gd}(\alpha)^{147}\text{Sm}$	2670.8	30.	2652.7	2.9	-0.6	U					65Si06
$^{151}\text{Tb}(\alpha)^{147}\text{Eu}$	3499.6	5.	3496	4	-0.6	1	58	49 $^{151}\text{Tb}$	DbA		67Go32
$^{151}\text{Dy}(\alpha)^{147}\text{Gd}$	4175.5	5.	4179.6	2.6	0.8	2					67Go32 Z
	4181.1	3.			-0.5	2					82Bo04 Z
$^{151}\text{Ho}(\alpha)^{147}\text{Tb}$	4696.3	5.	4695.0	1.8	-0.3	2			GSa		79Ho10 *
	4695.8	3.			-0.3	2			Bka		82Bo04 *
	4693.8	3.			0.4	2			Ora		82De11 *
	4694.9	5.			0.0	2			Daa		96Pa01 *
$^{151}\text{Eu}(n,\alpha)^{148}\text{Pm}$	7870	20	7859	6	-0.5	U			ILL		74Em01
$^{151}\text{Sm}(p,t)^{149}\text{Sm}$	-5100	4	-5101.4	0.4	-0.3	U			McM		73Ga04
$^{151}\text{Eu}(p,t)^{149}\text{Eu}$	-5872	5	-5873	4	-0.1	1	57	56 $^{149}\text{Eu}$	Min		75Ta12
$^{150}\text{Nd}(n,\gamma)^{151}\text{Nd}$	5334.55	0.2	5334.55	0.10	0.0	-			ILn		76Pi13 Z
	5334.55	0.11			0.0	-			Bdn		06Fi.A
$^{150}\text{Nd}(d,p)^{151}\text{Nd}$	3084	15	3109.98	0.10	1.7	U			Tal		67Ne08
$^{150}\text{Nd}(n,\gamma)^{151}\text{Nd}$	ave.	5334.55	5334.55	0.10	0.0	1	100	100 $^{151}\text{Nd}$			average
$^{150}\text{Nd}(^3\text{He,d})^{151}\text{Pm}$	1503	5	1502	4	-0.2	1	80	80 $^{151}\text{Pm}$	McM		80St10 *
$^{150}\text{Sm}(n,\gamma)^{151}\text{Sm}$	5596.5	1.8	5596.46	0.11	0.0	U					70Sm.A *
	5596.	1.5			0.3	U					71Gr22
	5596.42	0.20			0.2	-			ILn		86Va08 Z
	5596.44	0.13			0.1	-			Bdn		06Fi.A
$^{150}\text{Sm}(d,p)^{151}\text{Sm}$	3369	16	3371.89	0.11	0.2	U			Tal		65Ke09
$^{150}\text{Sm}(n,\gamma)^{151}\text{Sm}$	ave.	5596.43	5596.46	0.11	0.2	1	100	62 $^{150}\text{Sm}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{151}\text{Eu}(\gamma,n)^{150}\text{Eu}$	-8040	110	-7932	6	1.0	U			Phi		60Ge01
$^{151}\text{Eu}(p,d)^{150}\text{Eu}$	-5721	9	-5707	6	1.5	1	47	47 $^{150}\text{Eu}$			82So.B
$^{151}\text{Yb}(\epsilon p)^{150}\text{Er}$	9000	300				2			ORa		86To12 *
$^{151}\text{Lu}(p)^{150}\text{Yb}$	1239.2	3.	1241.0	1.8	0.6	o			Dap		82Ho04
	1241.0	2.8			0.0	3			Dap		93Se04
	1241.3	3.0			-0.1	3			ORp		03Gi10
	1240.3	4.0			0.2	3			Man		15Ta12
$^{151}\text{Lu}^m(p)^{150}\text{Yb}$	1318.8	10.	1294	4	-2.5	B			Dap		98Ba.B
	1293.6	4.				3			Man		15Ta12
$^{151}\text{Ce}(\beta^-)^{151}\text{Pr}$	5270	100	5555	21	2.8	U			Kur		02Sh.B
$^{151}\text{Pr}(\beta^-)^{151}\text{Nd}$	4170	75	4163	12	-0.1	U			Bwg		90Gr10
	4136	40			0.7	-			Ida		93Gr17 *
	4210	30			-1.6	-			Kur		95Ik03
	ave.	4183	24			-0.8	1	24	23 $^{151}\text{Pr}$		average
$^{151}\text{Nd}(\beta^-)^{151}\text{Pm}$	2510	50	2443	4	-1.3	U					73Se12 *
	2480	50			-0.7	U			Kur		95Ik03
$^{151}\text{Pm}(\beta^-)^{151}\text{Sm}$	1195	10	1190	4	-0.5	1	20	20 $^{151}\text{Pm}$			64Be10 *
$^{151}\text{Sm}(\beta^-)^{151}\text{Eu}$	75.9	0.6	76.6	0.5	1.1	1	80	59 $^{151}\text{Eu}$			59Ac28
$^{151}\text{Gd}(\epsilon)^{151}\text{Eu}$	463	3	464.1	2.8	0.4	1	86	85 $^{151}\text{Gd}$			83Vo10 *
$^{151}\text{Tb}(\beta^+)^{151}\text{Gd}$	2562	5	2565	4	0.6	-					77Cr05 *
	2566	12			-0.1	-					84Sc18 *
	ave.	2563	5			0.6	1	66	51 $^{151}\text{Tb}$		average
	$^{151}\text{Ho}(\beta^+)^{151}\text{Dy}$	5080	50	5130	9	1.0	U			IRS	
	5100	80			0.4	U			IRS		93Al03
$^{151}\text{Er}(\beta^+)^{151}\text{Ho}$	5130	110	5356	18	2.1	U					98Fo06
$^{151}\text{Tm}(\beta^+)^{151}\text{Er}$	6025	145	7494	25	10.1	C					84Ha.B *
	7074	50			8.4	F			GSI		91Ke11 *
$^{151}\text{Tm}^m(\text{IT})^{151}\text{Tm}$	96.4	7.0	94	6	-0.4	o					97Da07 *
$^{151}\text{Lu}^m(\text{IT})^{151}\text{Lu}$	77	5	53	4	-4.9	B			Dap		99Bi14
* $^{151}\text{Tb}-u$	$M-A=-71551(28)$ keV for mixture gs+m at 99.53 keV										Nub16b **
* $^{151}\text{Ho}-u$	$M-A=-63622(28)$ keV for mixture gs+m at 41.0 keV										Nub16b **
* $^{151}\text{Er}-u$	$M-A=-55670(28)$ keV for $^{151}\text{Er}^m$ at 2586.0 keV										Nub16b **
* $^{151}\text{Ho}(\alpha)^{147}\text{Tb}$	$E=4523.8(5,Z)$ to $^{147}\text{Tb}^m$ at 50.6(0.9); 4610.8(5,Z) from $^{151}\text{Ho}^m$ 41.0(0.2)										Nub16b **
* $^{151}\text{Ho}(\alpha)^{147}\text{Tb}$	$E=4521.5(3,Z)$ to $^{147}\text{Tb}^m$ at 50.6(0.9); 4611.5(3,Z) from $^{151}\text{Ho}^m$ 41.0(0.2)										Nub16b **
* $^{151}\text{Ho}(\alpha)^{147}\text{Tb}$	$E=4521.2(3,Z)$ to $^{147}\text{Tb}^m$ at 50.6(0.9); 4607.2(4,Z) from $^{151}\text{Ho}^m$ 41.0(0.2)										Nub16b **
* $^{151}\text{Ho}(\alpha)^{147}\text{Tb}$	$E_\alpha=4521(5,Z)$ to $^{147}\text{Tb}^m$ at 50.6(0.9)										Nub16b **
* $^{150}\text{Nd}(\text{He,d})^{151}\text{Pm}$	Based on $^{146}\text{Nd}(\text{He,d})^{147}\text{Pm}$ $Q=-87.6(0.9)$ keV										AHW **
* $^{150}\text{Sm}(n,\gamma)^{151}\text{Sm}$	$E(\gamma)=5591.7(1.8)$ to $3/2^-$ level at 4.821 keV										Ens091 **
* $^{151}\text{Yb}(\epsilon p)^{150}\text{Er}$	$E_p$ estimated 7300(300) to levels around 1700 keV										GAu **
*	"Statistical p's originate from $11/2^-$ isomer."										86To12 **
* $^{151}\text{Pr}(\beta^-)^{151}\text{Nd}$	Two highest $Q_{\beta^-}=4135(50),4137(40)$ keV										AHW **
* $^{151}\text{Nd}(\beta^-)^{151}\text{Pm}$	$E_{\beta^-}=2260(90)$ 2100(90) 1210(50) to $3/2^+$ level at 255.692, $1/2^+$ at 426.451, and $5/2^+$ at 1297.682 keV										Ens091 **
*											Ens091 **
* $^{151}\text{Pm}(\beta^-)^{151}\text{Sm}$	$E_{\beta^-}=1190(10)$ to ground state and $3/2^-$ level at 4.821 keV, and other $E_{\beta^-}$										Ens091 **
* $^{151}\text{Gd}(\epsilon)^{151}\text{Eu}$	$pK=0.652(0.007)$ to $(5/2^-,7/2^-)$ level at 353.64 keV										Ens091 **
* $^{151}\text{Tb}(\beta^+)^{151}\text{Gd}$	$E_{\beta^+}=700(5)$ $p^+=104(5)\times 10^{-4}$ respectively, to $1/2^-$ level at 839.320 keV, and other $E_{\beta^+}$										Ens091 **
* $^{151}\text{Ho}(\beta^+)^{151}\text{Dy}$	$E_{\beta^+}=3530(50)$ to $9/2^-$ level at 527.40 keV										Ens091 **
* $^{151}\text{Tm}(\beta^+)^{151}\text{Er}$	$p^+=0.71(0.02)$ to $9/2^-$ level at 801.52 keV										Ens091 **
* $^{151}\text{Tm}(\beta^+)^{151}\text{Er}$	F : lower limit: positrons escape from detector; $E_{\beta^+}=5250(50)$ to 801.6 level										91Ke11 **
* $^{151}\text{Tm}^m(\text{IT})^{151}\text{Tm}$	Only $\alpha$ -decay energies are used										GAu **
$^{152}\text{Pr}-u$	-68447.1	19.9				2			CP1	1.0	06Sa56
$\text{C}_{12}\text{H}_8-^{152}\text{Sm}$	142764	32	142861.2	1.3	0.8	U			R04	4.0	64De15
	142867.0	5.0			-0.5	U			M22	2.5	75Ka25
$^{152}\text{Eu}-u$	-78347	50	-78248.8	1.4	2.0	U			GS2	1.0	05Li24 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_{12} H_8 -^{152}Gd$	142870	50	142801.4	1.3	-0.3	U			R04	4.0	64De15
$^{152}Gd O - C_{14}$	-85290.7	3.5	-85286.6	1.3	0.8	U			TG1	1.5	11Ke03
$^{152}Tb-u$	-76212	159	-75920	40	1.9	U			GS2	1.0	05Li24 *
$^{152}Dy-u$	-75278	30	-75275	5	0.1	U			GS2	1.0	05Li24
$^{152}Ho-u$	-68248	58	-68283	13	-0.6	U			GS2	1.0	05Li24 *
$^{152}Er-u$	-64962	30	-64950	9	0.4	U			GS2	1.0	05Li24
$^{152}Tm-u$	-55524	58	-55520	60	0.0	1	100	100 $^{152}Tm$	GS2	1.0	05Li24
$^{152}Sm \ ^{35}Cl_2 - ^{148}Sm \ ^{37}Cl_2$	10802	10	10810.2	1.1	0.3	U			H21	2.5	70Ma05
	10810.8	2.0			-0.1	U			H25	2.5	72Ba08
	10807.9	1.4			0.7	U			M21	2.5	75Ka25
$^{152}Sm \ ^{35}Cl - ^{150}Sm \ ^{37}Cl$	5429	4	5407.0	0.7	-1.4	U			H12	4.0	64Ba15
	5396	4			1.1	o			H21	2.5	70Ma05
	5402.7	0.8			2.1	1	11	8 $^{150}Sm$	M21	2.5	75Ka25
$^{152}Gd - ^{152}Sm$	59.80	0.19	59.78	0.19	-0.1	1	98	72 $^{152}Sm$	SH1	1.0	11El02
$^{152}Sm - ^{151}Eu$	95	42	-117.8	0.7	-1.3	U			R04	4.0	64De15
$^{152}Sm - ^{150}Sm$	2563	31	2456.8	0.7	-0.9	U			R04	4.0	64De15
$^{152}Gd(\alpha) ^{148}Sm$	2197.9	30.	2204.4	1.0	0.2	U					61Ma05
$^{152}Dy(\alpha) ^{148}Gd$	3728.0	8.	3727	4	-0.2	2					65Ma51 Z
	3726.0	5.			0.1	2			DbA		67Go32 Z
$^{152}Ho(\alpha) ^{148}Tb$	4506.9	3.	4507.4	1.3	0.2	-			Bka		82Bo04 *
	4508.0	2.			-0.3	-			Ora		82De11 Z
	4505.8	3.			0.5	-					82To14
	4507.9	3.			-0.2	-					87St.A Z
	ave.	4507.3	1.3		0.0	1	100	95 $^{152}Ho$			average
$^{152}Er(\alpha) ^{148}Dy$	4935.2	5.	4934.3	1.6	-0.2	-			GSa		79Ho10
	4934.6	3.			-0.1	-			Bka		82Bo04 Z
	4934.3	2.			0.0	-			Ora		82De11 Z
	ave.	4934.5	1.6		-0.1	1	100	85 $^{152}Er$			average
$^{150}Nd(t,p) ^{152}Nd$	4125	30	4130	24	0.2	1	66	66 $^{152}Nd$	Ald		72Ch11
$^{150}Sm(t,p) ^{152}Sm$	5376	25	5372.3	0.6	-0.1	U			Ald		66Bj01
$^{152}Sm(p,t) ^{150}Sm$	-5378	8	-5372.3	0.6	0.7	U			Min		72De47
	-5376	4			0.9	U			McM		73Ga04
	-5379	15			0.4	U			Ham		74Oe03
$^{151}Sm(n,\gamma) ^{152}Sm$	8257.6	0.8	8257.6	0.6	0.1	1	58	41 $^{151}Sm$			71Gr22 Z
$^{151}Sm(p,\gamma) ^{152}Eu$	5604	4	5600.9	0.5	-0.8	U					75Jo.A
$^{151}Eu(n,\gamma) ^{152}Eu$	6306.70	0.10	6306.72	0.10	0.2	1	99	60 $^{152}Eu$	ILn		85Vo15 Z
	6307.11	0.14			-2.8	C			Bdn		06Fi.A
$^{152}Gd(d,t) ^{151}Gd$	-2338	10	-2332.3	2.9	0.6	U			Kop		67Tj01
$^{152}Pr(\beta^-) ^{152}Nd$	6350	120	6390	30	0.3	U			Kur		95Ik03
$^{152}Nd(\beta^-) ^{152}Pm$	1088	27	1105	19	0.6	-					93Sh23
	1120	30			-0.5	-			Kur		95Ik03
	ave.	1102	20		0.1	1	85	51 $^{152}Pm$			average
$^{152}Pm(\beta^-) ^{152}Sm$	3600	200	3508	26	-0.5	U					71Da19
	3520	150			-0.1	U					72Wa04
	3400	200			0.5	U					75Wi08
	3500	100			0.1	-					77Ya07
	3500	40			0.2	-			Kur		95Ik03
	ave.	3500	40		0.2	1	49	49 $^{152}Pm$			average
$^{152}Pm^m(\beta^-) ^{152}Sm$	3603	100	3650	80	0.5	2					71Da19 *
	3753	150			-0.7	2					72Wa04 *
$^{152}Eu(\beta^+) ^{152}Sm$	1871	5	1874.3	0.7	0.7	U					58A199 *
	1866	5			1.7	U					62Lo10 *
	1870.8	2.			1.8	-					72Sv02 *
	1872.8	1.5			1.0	-					77Mi.A *
	ave.	1872.1	1.2		1.9	1	33	26 $^{152}Eu$			average
$^{152}Eu(\beta^-) ^{152}Gd$	1809	10	1818.7	0.7	1.0	U					58A199 *
	1827	7			-1.2	U					60La04 *
	1836	20			-0.9	U					60Sc14 *
	1806	4			3.2	B					69An18 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{152}\text{Tb}(\beta^+)^{152}\text{Gd}$	3990	40				2					76Cr.B *
$^{152}\text{Ho}(\beta^+)^{152}\text{Dy}$	6690	100	6513	13	-1.8	U			IRS		83A106 *
	6270	140			1.7	U					Averag *
	6225	90			3.2	B			IRS		93A103 *
$^{152}\text{Tm}(\beta^+)^{152}\text{Er}$	8820	240	8780	50	-0.2	U					16Na02
$^{152}\text{Tm}^m(\beta^+)^{152}\text{Er}$	6850	110	8680	240	16.6	C					84Ha.B *
	8680	240				2					16Na02
$^{152}\text{Yb}(\beta^+)^{152}\text{Tm}$	5465	195	5450	140	-0.1	-			Got		90Sa.A
	5434	200			0.1	-			GSI		04Na.A *
	ave.	5450	140		0.0	1	100	100	$^{152}\text{Yb}$		average
* $^{152}\text{Eu}-u$	$M-A=-72915(35)$ keV for mixture gs+m+r at 45.5998 and 147.86 keV										Nub16b **
* $^{152}\text{Tb}-u$	$M-A=-70740(29)$ keV for mixture gs+n at 501.74 keV										Nub16b **
* $^{152}\text{Ho}-u$	$M-A=-63492(28)$ keV for mixture gs+m at 160(1) keV										Nub16b **
* $^{152}\text{Ho}(\alpha)^{148}\text{Tb}$	$E_\alpha=4389.1(3,Z)$ ; and $4455.1(3,Z)$ from $^{152}\text{Ho}^m$ to $^{148}\text{Tb}^m$										82Bo04 **
*	combined with $^{152}\text{Ho}^m(\text{IT})$ - $^{148}\text{Tb}^m(\text{IT})=160(1)$ - $90.1(0.3)$ keV										87St.A **
* $^{152}\text{Pm}^m(\beta^-)^{152}\text{Sm}$	$E_{\beta^-}=1800(100)$ $1950(150)$ respectively, to $5^-$ level at 1803.94 keV										Ens13b **
* $^{152}\text{Eu}(\beta^+)^{152}\text{Sm}$	$E_{\beta^+}=895(5)$ $890(5)$ respectively, from $^{152}\text{Eu}^m$ at 45.5998 keV										Nub16b **
* $^{152}\text{Eu}(\beta^+)^{152}\text{Sm}$	$E_{\beta^+}=727(2)$ $729(1.5)$ respectively, to $2^+$ level at 121.7818 keV										Ens13b **
* $^{152}\text{Eu}(\beta^-)^{152}\text{Gd}$	$Q_{\beta^-}=1855(10)$ from $^{152}\text{Eu}^m$ at 45.5998 keV										Nub16b **
* $^{152}\text{Eu}(\beta^-)^{152}\text{Gd}$	$E_{\beta^-}=1483(7)$ to $2^+$ level at 344.2790 keV										Ens13b **
* $^{152}\text{Eu}(\beta^-)^{152}\text{Gd}$	$E_{\beta^-}=1840(30)$ $1490(20)$ $1072(20)$ to ground state, $2^+$ at 344.2790, $4^+$ at 755.3961 keV										Ens13b **
* $^{152}\text{Eu}(\beta^-)^{152}\text{Gd}$	$Q_{\beta^-}=1852(4)$ from $^{152}\text{Eu}^m$ at 45.5998 keV										Nub16b **
* $^{152}\text{Tb}(\beta^+)^{152}\text{Gd}$	$E_{\beta^+}=2830(15)$ $8(4)\%$ to ground state, $5.2(1)\%$ to $2^+$ level at 344.2790 keV										Ens13b **
* $^{152}\text{Ho}(\beta^+)^{152}\text{Dy}$	$E_{\beta^+}=3390(100)$ from $^{152}\text{Ho}^m$ at 160(1) to $8^+$ level at 2437.42 keV										Ens13b **
* $^{152}\text{Ho}(\beta^+)^{152}\text{Dy}$	From adopted $\text{KLM}/\beta^+=0.97(0.13)$										AHW **
*	from $^{152}\text{Ho}^m$ at 160(1) to $8^+$ level at 2437.42 keV										Ens13b **
*	after extra 3(2)% side-feeding correction; see reference										90Sa32 **
*	$p^+=0.52(0.04)/.967$ gives $\text{KLM}/\beta^+=0.86(0.14)$										85Sc09 **
*	$\text{KLM}/\beta^+=1.12(0.10)$ after 0.967(0.008) side-feeding correction										90Sa32 **
* $^{152}\text{Ho}(\beta^+)^{152}\text{Dy}$	$Q_{\beta^+}=6270(90)$ ; and $6330(100)$ from $^{152}\text{Ho}^m$ at 160(1) keV										Nub16b **
* $^{152}\text{Tm}^m(\beta^+)^{152}\text{Er}$	$p^+=0.64(0.02)$ to $8^+$ level at 2183.3 keV										Ens13b **
* $^{152}\text{Yb}(\beta^+)^{152}\text{Tm}$	As reported in reference										11Es03 **
$^{153}\text{Pr}-u$	-66110.5	15.3	-66096	13	0.9	-			CP1	1.0	06Sa56
	-66065	40			-0.8	-			CP1	1.0	12Va02 *
	ave.	-66105	14		0.6	1	80	80	$^{153}\text{Pr}$		average
$^{153}\text{Pr}-^{80}\text{Kr}_{1,913}$	93906	40	93872	13	-0.8	1	10	10	$^{153}\text{Pr}$	1.0	12Va02
$^{153}\text{Pr}-^{86}\text{Kr}_{1,779}$	92958	40	92927	13	-0.8	1	10	10	$^{153}\text{Pr}$	1.0	12Va02
$^{153}\text{Nd}-u$	-72283.3	5.2	-72282.1	2.9	0.2	1	32	32	$^{153}\text{Nd}$	1.0	12Va02 *
$^{153}\text{Nd}-^{80}\text{Kr}_{1,913}$	87687.9	4.7	87687	3	-0.2	1	42	36	$^{153}\text{Nd}$	1.0	12Va02
$^{153}\text{Nd}-^{86}\text{Kr}_{1,779}$	86740.7	5.3	86741.6	2.9	0.2	1	31	31	$^{153}\text{Nd}$	1.0	12Va02
$^{153}\text{Pm}-u$	-75833	23	-75844	10	-0.5	1	18	18	$^{153}\text{Pm}$	1.0	12Va02 *
$^{153}\text{Pm}-^{80}\text{Kr}_{1,913}$	84139	23	84125	10	-0.6	1	18	18	$^{153}\text{Pm}$	1.0	12Va02
$^{153}\text{Pm}-^{86}\text{Kr}_{1,779}$	83192	23	83180	10	-0.5	1	18	18	$^{153}\text{Pm}$	1.0	12Va02
$\text{C}_{12}\text{H}_9-^{153}\text{Eu}$	149103	18	149188.2	1.4	1.2	U			R04	4.0	64De15
$\text{C}_{11}\text{H}_8-^{153}\text{Eu}$	144606	30	144718.0	1.4	0.9	U			R04	4.0	64De15
$\text{C}_9\text{H}_{16}\text{O}-^{153}\text{Eu}$	201934	38	202232.9	1.4	2.0	U			R04	4.0	64De15
$^{153}\text{Eu}-^{85}\text{Rb}_{1,800}$	80021	16	80015.5	1.4	-0.3	U			MA5	1.0	00Be42
$^{153}\text{Ho}-u$	-69814	37	-69793	5	0.6	U			GS2	1.0	05Li24 *
$^{153}\text{Er}-u$	-64942	30	-64916	10	0.9	U			GS2	1.0	05Li24
$^{153}\text{Eu}-^{35}\text{Cl}-^{151}\text{Eu}-^{37}\text{Cl}$	4334	4	4330.29	0.18	-0.4	U			H21	2.5	70Ma05 *
$^{138}\text{La}\text{O}-^{153}\text{Eu}$	-19266	123	-19205	4	0.1	U			R05	4.0	65De13
$^{153}\text{Eu}\text{O}-\text{C}_{14}$	-83849.6	5.8	-83848.3	1.4	0.1	U			TG1	1.5	11Ke03
$^{153}\text{Eu}-^{152}\text{Sm}$	1544	42	1498.0	0.7	-0.3	U			R04	4.0	64De15
$^{153}\text{Eu}-^{151}\text{Eu}$	1567	33	1380.18	0.17	-1.4	U			R04	4.0	64De15
$^{153}\text{Dy}(\alpha)^{149}\text{Gd}$	3560.0	8.	3559	4	-0.1	-					65Ma51 Z
	3554.9	5.			0.8	-			DbA		67Go32 Z
	ave.	3556	4		0.7	1	69	48	$^{153}\text{Dy}$		average
$^{153}\text{Ho}(\alpha)^{149}\text{Tb}$	4052.3	5.	4052	4	-0.1	2					68Go.C *
	4051.0	5.			0.1	2			ORa		71To01 *



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{153}\text{Er}(\alpha)^{149}\text{Dy}$	4799.8	10.	4802.4	1.4	0.3	U			81Ho.A	
	4804.5	3.			-0.7	-	Bka		82Bo04 Z	
	4802.0	2.			0.2	-	Ora		82De11 Z	
	4802.8	3.			-0.1	-			87Sc.A Z	
	4799.7	4.			0.6	-	Daa		96Pa01	
ave.	4802.4	1.4			0.0	1	100	97 $^{153}\text{Er}$	average	
$^{153}\text{Tm}(\alpha)^{149}\text{Ho}$	5252.3	5.	5248.3	1.5	-0.8	U	GSa		79Ho10 *	
	5246.1	3.			0.7	-	Bka		82Bo04 *	
	5249.2	2.			-0.4	-	Ora		82De11 *	
	5247.7	3.			0.2	o			87Sc.A *	
	5247.7	3.			0.2	-			88Sc.A	
ave.	5249.5	5.			-0.2	U	Daa		96Pa01	
	5248.1	1.5			0.1	1	100	53 $^{149}\text{Ho}$	average	
$^{153}\text{Gd}(\text{n},\alpha)^{150}\text{Sm}$	9790	30	9815.0	0.6	0.8	U	ILL		81Wa31	
$^{153}\text{Eu}(\text{p},\text{t})^{151}\text{Eu}$	-6374	5	-6375.21	0.16	-0.2	U	Min		75Ta12	
$^{152}\text{Sm}(\text{n},\gamma)^{153}\text{Sm}$	5867.1	0.4	5868.40	0.13	3.3	B			69Re04 Z	
	5868.4	0.3			0.0	2			71Be41 Z	
	5868.4	0.7			0.0	U			82Ba15 Z	
	5868.40	0.15			0.0	2	Bdn		06Fi.A	
$^{152}\text{Sm}(\text{d},\text{p})^{153}\text{Sm}$	3645	12	3643.83	0.13	-0.1	U	Tal		65Ke09	
$^{152}\text{Eu}(\text{n},\gamma)^{153}\text{Eu}$	8550.28	0.12	8550.28	0.12	0.0	1	100	87 $^{153}\text{Eu}$	ILn	85Vo15 Z
$^{153}\text{Eu}(\gamma,\text{n})^{152}\text{Eu}$	-8650	130	-8550.28	0.12	0.8	U	Phi		60Ge01	
$^{152}\text{Gd}(\text{n},\gamma)^{153}\text{Gd}$	6247.27	0.35	6246.95	0.13	-0.9	-	ILn		85Vo15 Z	
	6246.89	0.14			0.4	-	ILn		93Sp.A	
	6247.48	0.21			-2.5	B	Bdn		06Fi.A	
$^{152}\text{Gd}(\text{d},\text{p})^{153}\text{Gd}$	4015	10	4022.39	0.13	0.7	U	Kop		67Tj01	
ave.	6246.94	0.13	6246.95	0.13	0.1	1	99	74 $^{152}\text{Gd}$	average	
$^{152}\text{Gd}(\beta^-\text{He},\text{d})^{153}\text{Tb}$	-1634	30	-1598	4	1.2	U	McM		76St10	
$^{153}\text{Pr}(\beta^-)^{153}\text{Nd}$	5720	100	5762	12	0.4	U	Kur		02Sh.B	
$^{153}\text{Nd}(\beta^-)^{153}\text{Pm}$	3336	25	3318	9	-0.7	1	14	13 $^{153}\text{Pm}$	Ida	93Gr17
	3260	100			0.6	U			Kur	02Sh.B
$^{153}\text{Pm}(\beta^-)^{153}\text{Sm}$	1777	50	1912	9	2.7	U			62Ko10 *	
	1863	15			3.3	B			Ida	93Gr17
$^{153}\text{Sm}(\beta^-)^{153}\text{Eu}$	810	10	807.5	0.7	-0.2	U			54Gr19	
	795	10			1.3	U			54Le08	
	820	10			-1.2	U			55Ma62	
	825	10			-1.7	U			56Du31	
	792	10			1.6	U			57Jo24	
$^{153}\text{Tb}(\beta^+)^{153}\text{Gd}$	1573	5	1569	4	-0.8	1	59	59 $^{153}\text{Tb}$	78Cr02 *	
$^{153}\text{Dy}(\beta^+)^{153}\text{Tb}$	2171	2	2170.4	1.9	-0.3	1	93	52 $^{153}\text{Dy}$	78Gr13 *	
$^{153}\text{Ho}(\beta^+)^{153}\text{Dy}$	4153	50	4131	6	-0.4	o		IRS	83Al06 *	
	4160	60			-0.5	U		IRS	93Al03	
$^{153}\text{Lu}^m(\text{IT})^{153}\text{Lu}$	80	5				4			97Ir01	
* $^{153}\text{Pr}-\text{u}$	Represents frequency ratio $^{153}\text{Pr}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.99430250(26)$								WgM124**	
* $^{153}\text{Nd}-\text{u}$	Represents frequency ratio $^{153}\text{Pr}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.994342.931(34)$								WgM124**	
* $^{153}\text{Pm}-\text{u}$	Represents frequency ratio $^{153}\text{Pm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.99436601(15)$								WgM124**	
* $^{153}\text{Ho}-\text{u}$	$M - A = -64997(28)$ keV for mixture gs+m at 68.7 keV								Nub16b **	
* $^{153}\text{Eu} \text{ } ^{35}\text{Cl} - ^{151}\text{Eu} \text{ } ^{37}\text{Cl}$	Increased by 5 for systematic difference of H21 with later data								AHW **	
* $^{153}\text{Ho}(\alpha)^{149}\text{Tb}$	$E_\alpha = 4013.1(5, Z)$ from $^{153}\text{Ho}^m$ at 68.7 keV								Nub16b **	
* $^{153}\text{Ho}(\alpha)^{149}\text{Tb}$	$E_\alpha = 3910(5)$ to $^{149}\text{Tb}^m$ at 35.78 keV								Nub16b **	
* $^{153}\text{Tm}(\alpha)^{149}\text{Ho}$	$E_\alpha = 5114.2(5, Z)$ $5108.2(3, Z)$ $5111.2(2, Z)$ respectively, contain a 8% $^{153}\text{Tm}(\alpha)^{149}\text{Ho} - ^{153}\text{Tm}^m(\alpha)^{149}\text{Ho} = 48.80(0.20) - 43.2(0.2) = 5.6(0.3)$ keV lower $^{153}\text{Tm}^m(\alpha)^{149}\text{Ho}^m$ branch, corrected thus +0.5keV								Nub16b **	
*									GAu **	
* $^{153}\text{Tm}(\alpha)^{149}\text{Ho}$	$E_\alpha = 5110.6(3, Z)$ ; and $5103.6(4, Z)$ for lower $^{153}\text{Tm}^m(\alpha)$ branch								87Sc.A **	
* $^{153}\text{Pm}(\beta^-)^{153}\text{Sm}$	$E_{\beta^-} = 1650(50)$ to $3/2^-$ level at 127.298 keV, and other $E_{\beta^-}$								Ens062 **	
* $^{153}\text{Tb}(\beta^+)^{153}\text{Gd}$	$E_{\beta^+} = 339(5)$ to $3/2^+$ level at 212.0078 keV								Ens062 **	
* $^{153}\text{Dy}(\beta^+)^{153}\text{Tb}$	$E_{\beta^+} = 886(2)$ to $9/2^-$ level at 262.831 keV								Ens062 **	
* $^{153}\text{Ho}(\beta^+)^{153}\text{Dy}$	$E_{\beta^+} = 2835(50)$ to $9/2^-$ level at 295.84 keV								Ens062 **	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{154}\text{Ce}-u$	-56404	619	-56060#	220#	0.2	D			GT3	2.5	16Kn03 *
$\text{C}_{12} \text{H}_{10}-^{154}\text{Sm}$	155830	29	156034.2	1.6	1.8	U			R04	4.0	64De15
	156035.7	4.0			-0.2	U			M22	2.5	75Ka25
$\text{C}_{12} \text{H}_{10}-^{154}\text{Gd}$	157149	40	157376.9	1.3	1.4	U			R04	4.0	64De15
$\text{C}_{11} \text{}^{13}\text{C} \text{H}_9-^{154}\text{Gd}$	152550	110	152906.7	1.3	0.8	U			R04	4.0	64De15
$\text{C}_{10} \text{}^{13}\text{C}_2 \text{H}_8-^{154}\text{Gd}$	148030	90	148436.5	1.3	1.1	U			R04	4.0	64De15
$\text{C}_{10} \text{H}_6 \text{N}_2-^{154}\text{Gd}$	131980	240	132224.8	1.3	0.3	U			R04	4.0	64De15
$^{154}\text{Gd}-^{138}\text{La} \text{O}$	19005	80	18841	4	-0.5	U			R05	4.0	65De13
$^{154}\text{Tb}-u$	-75376	115	-75320	50	0.5	R			GS2	1.0	05Li24 *
$^{154}\text{Dy}-^{133}\text{Cs}_{1.158}$	33903	19	33916	8	0.7	1	18	18 $^{154}\text{Dy}$	MA5	1.0	00Be42 *
$^{154}\text{Ho}-u$	-69350	83	-69393	9	-0.5	U			GS2	1.0	05Li24 *
$^{154}\text{Tm}-u$	-58479	49	-58430	15	1.0	U			GS2	1.0	05Li24 *
$^{154}\text{Sm} \text{}^{35}\text{Cl}_2-^{150}\text{Sm} \text{}^{37}\text{Cl}_2$	10832.9	5.2	10834.2	1.1	0.1	U			M21	2.5	75Ka25
$^{154}\text{Sm} \text{}^{35}\text{Cl}-^{152}\text{Sm} \text{}^{37}\text{Cl}$	5480	4	5427.2	0.9	-3.3	B			H12	4.0	64Ba15
	5417	4			1.0	U			H21	2.5	70Ma05
	5427.2	0.4			0.0	1	80	78 $^{154}\text{Sm}$	M21	2.5	75Ka25
$^{154}\text{Gd} \text{}^{35}\text{Cl}-^{152}\text{Gd} \text{}^{37}\text{Cl}$	4019.5	2.	4024.69	0.23	1.0	U			H25	2.5	72Ba08
	4016	30			0.1	U			H12	4.0	64Ba15
$^{154}\text{Sm}-^{154}\text{Gd}$	1338.2	3.8	1342.8	0.9	0.5	U			H25	2.5	72Ba08
	1342.8	0.8			0.0	1	21	21 $^{154}\text{Sm}$	M21	2.5	75Ka25
$^{154}\text{Sm}-\text{C}_{12} \text{H}_9$	-148211.0	8.0	-148209.1	1.6	0.1	U			M21	2.5	75Ka25
$^{139}\text{La} \text{O}-^{154}\text{Gd}$	-19616	55	-19600.0	2.3	0.1	U			R05	4.0	65De13
$^{154}\text{Sm}-^{153}\text{Eu}$	1082	42	979.1	1.2	-0.6	U			R04	4.0	64De15
$^{154}\text{Sm}-^{152}\text{Sm}$	2664	43	2477.1	0.9	-1.1	U			R04	4.0	64De15
$^{154}\text{Gd}-^{152}\text{Gd}$	1400	50	1074.58	0.22	-1.6	U			R04	4.0	64De15
$^{154}\text{Gd} \text{O}-\text{C}_{15}$	-84207.4	5.9	-84212.0	1.3	-0.5	U			TG1	1.5	09Ke.A
	-84206.6	4.3			-0.8	U			TG1	1.5	11Ke03
$^{154}\text{Dy}(\alpha)^{150}\text{Gd}$	2946.4	5.	2945	5	-0.3	1	93	82 $^{154}\text{Dy}$	DbA		67Go32 Z
$^{154}\text{Ho}(\alpha)^{150}\text{Tb}$	4041.3	5.	4041	4	0.0	2					68Go.C Z
	4041.7	5.			0.0	2			ORa		74Sc19 Z
$^{154}\text{Ho}^m(\alpha)^{150}\text{Tb}^m$	3819.2	10.	3823	5	0.4	-			ORa		71To01 Z
	3824.0	5.			-0.1	-			ORa		74Sc19 Z
	ave.	3823			0.1	1	100	89 $^{154}\text{Ho}^m$			average
$^{154}\text{Er}(\alpha)^{150}\text{Dy}$	4280.5	5.	4279.7	2.6	-0.2	-					68Go.C Z
	4279.5	3.			0.1	-			Bka		82Bo04 Z
	ave.	4279.8			0.0	1	98	92 $^{154}\text{Er}$			average
$^{154}\text{Tm}(\alpha)^{150}\text{Ho}$	5096.9	5.1	5093.8	2.6	-0.6	2			GSa		79Ho10 Z
	5092.7	3.			0.4	2			Bka		82Bo04 Z
$^{154}\text{Tm}^m(\alpha)^{150}\text{Ho}^m$	5174.8	5.	5171.8	1.6	-0.6	3			GSa		79Ho10 Z
	5170.8	3.			0.3	3			Bka		82Bo04 Z
	5171.8	2.1			0.0	3			Ora		82De11 Z
$^{154}\text{Yb}(\alpha)^{150}\text{Er}$	5473.4	5.	5474.3	1.7	0.2	-			GSa		79Ho10 Z
	5474.7	2.			-0.2	-			Ora		82De11 Z
	5473.4	4.			0.2	-			Daa		96Pa01
	ave.	5474.3			0.0	1	100	100 $^{154}\text{Yb}$			average
$^{152}\text{Sm}(\text{t,p})^{154}\text{Sm}$	5361	25	5353.4	0.8	-0.3	U			Ald		66Bj01
$^{154}\text{Sm}(\text{p,t})^{152}\text{Sm}$	-5357	8	-5353.4	0.8	0.4	U			Min		72De47
	-5353	15			0.0	U			Ham		74Oe03
$^{154}\text{Gd}(\text{p,t})^{152}\text{Gd}$	-6660	5	-6659.88	0.21	0.0	U			Min		73Oo01
$^{154}\text{Sm}(\text{d},^3\text{He})^{153}\text{Pm}$	-3623	25	-3603	9	0.8	-					76Su.B
$^{154}\text{Sm}(\text{t},\alpha)^{153}\text{Pm}$	10748	20	10718	9	-1.5	-			LAl		78Bu18
$^{154}\text{Sm}(\text{d},^3\text{He})^{153}\text{Pm}$	ave.	-3592	-3603	9	-0.7	1	34	33 $^{153}\text{Pm}$			average
$^{153}\text{Eu}(\text{n},\gamma)^{154}\text{Eu}$	6442.2	0.3	6442.22	0.24	0.1	-			ILn		87Ba52 Z
	6442.2	0.4			0.0	-			Bdn		06Fi.A
	ave.	6442.20			0.1	1	99	85 $^{154}\text{Eu}$			average
$^{153}\text{Gd}(\text{n},\gamma)^{154}\text{Gd}$	8895.25	0.30	8894.72	0.17	-1.8	-			ILn		85Vo15 Z
	8894.47	0.20			1.3	-			ILn		93Sp.A Z

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{154}\text{Gd}(d,t)^{153}\text{Gd}$		-2642	10	-2637.49	0.17	0.5	U		Kop		67Tj01
$^{153}\text{Gd}(n,\gamma)^{154}\text{Gd}$	ave.	8894.71	0.17	8894.72	0.17	0.1	1	98	74 $^{153}\text{Gd}$		average
$^{154}\text{Pr}(\beta^-)^{154}\text{Nd}$		7720	100				5		Kur		02Sh.B *
$^{154}\text{Nd}(\beta^-)^{154}\text{Pm}$		2687	25				4		Ida		93Gr17
$^{154}\text{Pm}^m(\text{IT})^{154}\text{Pm}$		-210	70	20	12	3.3	B				72Ta13 *
		20	12				3				90So08 *
$^{154}\text{Pm}(\beta^-)^{154}\text{Sm}$		3900	200	3940	50	0.2	U				71Da28
		4396	180			-2.5	U				72Ta13
		3880	200			0.3	U				74Ya07 *
$^{154}\text{Pm}^m(\beta^-)^{154}\text{Sm}$		3900	200	3960	40	0.3	U				71Da28 *
		4190	170			-1.3	U				72Ta13 *
		3940	50			0.5	2				73Pr05 *
		3940	200			0.1	U				74Ya07 *
		4056	100			-0.9	2		Ida		93Gr17
$^{154}\text{Eu}(\beta^-)^{154}\text{Gd}$		1978	5	1967.8	0.8	-2.0	U				60La04 *
		1967	2			0.4	1	14	12 $^{154}\text{Eu}$		77Ra08 *
		1975	3			-2.4	U				81Bu.A *
$^{154}\text{Tb}(\beta^+)^{154}\text{Gd}$		3562	50	3550	50	-0.2	2				70Ag03 *
$^{154}\text{Ho}(\beta^+)^{154}\text{Dy}$		5700	80	5755	10	0.7	U		IRS		83Al06 *
		5750	80			0.1	U		IRS		93Al03
$^{154}\text{Ho}^m(\beta^+)^{154}\text{Dy}$		5994	100	5997	28	0.0	o		IRS		83Al.A *
		6070	80			-0.9	1	12	11 $^{154}\text{Ho}^m$		93Al03
$^{154}\text{Tm}^m(\beta^+)^{154}\text{Er}$		8234	150	8250	50	0.1	U		Dbn		94Po26 *
$^{154}\text{Lu}(\beta^+)^{154}\text{Yb}$		7556	450	10220#	200#	5.9	C				84Ha.B *
$^{154}\text{Lu}^m(\text{IT})^{154}\text{Lu}$		58.7	9.3	60	12	0.1	o		Ara		97Da07 *
* $^{154}\text{Ce}-u$	Trends from Mass Surface TMS suggest $^{154}\text{Ce}$ 320 less bound										
* $^{154}\text{Tb}-u$	$M - A = -70142(43)$ keV for mixture gs+m+n at 12(7) and 200#150 keV										
* $^{154}\text{Dy}-^{133}\text{Cs}_{1,158}$	No contamination observed, but contamination by $^{154}\text{Tb}$ cannot be excluded										
* $^{154}\text{Ho}-u$	$M - A = -64478(28)$ keV for mixture gs+m at 243(28) keV										
* $^{154}\text{Tm}-u$	$M - A = -54438(32)$ keV for mixture gs+m at 70(50) keV										
* $^{154}\text{Pr}(\beta^-)^{154}\text{Nd}$	$E_{\beta^-} = 7490(100)$ to $4^+$ level at 233.2 keV										
* $^{154}\text{Pm}^m(\text{IT})^{154}\text{Pm}$	Only use the two $Q^-$ 's to $^{154}\text{Sm}$ , see below										
* $^{154}\text{Pm}^m(\text{IT})^{154}\text{Pm}$	Supported by reference 30(10) keV										
* $^{154}\text{Pm}(\beta^-)^{154}\text{Sm}$	$E_{\beta^-} = 2410(180)$ to $3^-$ level at 1986.59 keV										
* $^{154}\text{Pm}(\beta^-)^{154}\text{Sm}$	$E_{\beta^-} = 2400(200), 1850(200)$ to $2^+$ levels at 1440.04, 2069.07 keV										
* $^{154}\text{Pm}^m(\beta^-)^{154}\text{Sm}$	$E_{\beta^-} = 3270, 3090, 2810$ (all 170) to $921.345 1^-, 1099.26 0^+, 1475.81 1^-$										
* $^{154}\text{Pm}^m(\beta^-)^{154}\text{Sm}$	$E_{\beta^-} = 2810(170)$ to $1^-$ level at 1475.81 keV, and other $E_{\beta^-}$										
* $^{154}\text{Pm}^m(\beta^-)^{154}\text{Sm}$	$E_{\beta^-} = 3950(50) 3010(80)$ to ground state, $1^-$ level at 921.345 keV										
* $^{154}\text{Pm}^m(\beta^-)^{154}\text{Sm}$	$E_{\beta^-} = 3000(200), 1900(200), 1800(200)$ to $1^-$ level at 921.345, $2^+$ at 2069.07, and $(1,2^+)$ at 2139.82 keV										
* $^{154}\text{Eu}(\beta^-)^{154}\text{Gd}$	$E_{\beta^-} = 1855(5) 1844(2)$ respectively, to $2^+$ level at 123.0709 keV										
* $^{154}\text{Eu}(\beta^-)^{154}\text{Gd}$	$E_{\beta^-} = 257(3)$ to $2^-$ level at 1719.5593 keV, and other $E_{\beta^-}$										
* $^{154}\text{Tb}(\beta^+)^{154}\text{Gd}$	$E_{\beta^+} = 2540(50) 1860(50)$ to ground state and $0^+$ level at 680.6673 keV										
* $^{154}\text{Ho}(\beta^+)^{154}\text{Dy}$	$E_{\beta^+} = 4340(80)$ to $2^+$ level at 334.34 keV										
* $^{154}\text{Ho}^m(\beta^+)^{154}\text{Dy}$	$E_{\beta^+} = 2500(100)$ to $7^+$ level at 2472.40 keV										
* $^{154}\text{Tm}^m(\beta^+)^{154}\text{Er}$	$E_{\beta^+} = 4882(150)$ to $8^+$ level 2329.5 keV										
* $^{154}\text{Lu}(\beta^+)^{154}\text{Yb}$	$p^+ = 0.75(0.05) Q = 5710(450)$ from $^{154}\text{Lu}^m$ at 200#150 to $8^+$ level at 2046.2 keV										
* $^{154}\text{Lu}^m(\text{IT})^{154}\text{Lu}$	Use only their $Q_{\alpha}$ 's										
$^{155}\text{Pr}-u$	-59492	31	-59491	18	0.0	1	35	35 $^{155}\text{Pr}$	CP1	1.0	12Va02 *
$^{155}\text{Pr}-^{80}\text{Kr}_{1,938}$	102571	33	102569	18	-0.1	1	31	31 $^{155}\text{Pr}$	CP1	1.0	12Va02
$^{155}\text{Pr}-^{86}\text{Kr}_{1,802}$	101588	32	101589	18	0.0	1	33	33 $^{155}\text{Pr}$	CP1	1.0	12Va02
$^{155}\text{Nd}-u$	-66866	17	-66864	10	0.1	1	33	33 $^{155}\text{Nd}$	CP1	1.0	12Va02 *
$^{155}\text{Nd}-^{80}\text{Kr}_{1,938}$	95197	17	95195	10	-0.1	1	34	33 $^{155}\text{Nd}$	CP1	1.0	12Va02

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{155}\text{Nd}-^{86}\text{Kr}_{1.802}$	94215	17	94215	10	0.0	1	33	33 $^{155}\text{Nd}$	CP1	1.0	12Va02
$^{155}\text{Pm}-\text{u}$	-71863.8	8.8	-71863	5	0.1	1	33	33 $^{155}\text{Pm}$	CP1	1.0	12Va02 *
$^{155}\text{Pm}-^{80}\text{Kr}_{1.938}$	90197.8	8.6	90196	5	-0.2	1	36	34 $^{155}\text{Pm}$	CP1	1.0	12Va02
$^{155}\text{Pm}-^{86}\text{Kr}_{1.802}$	89216.0	8.8	89217	5	0.1	1	33	33 $^{155}\text{Pm}$	CP1	1.0	12Va02
$^{155}\text{Sm}-\text{u}$	-75357	24	-75352.9	1.6	0.2	U			CP1	1.0	12Va02 *
$^{155}\text{Sm}-^{80}\text{Kr}_{1.938}$	86704	24	86706.4	2.1	0.1	U			CP1	1.0	12Va02
$^{155}\text{Sm}-^{86}\text{Kr}_{1.802}$	85722	24	85726.7	1.6	0.2	U			CP1	1.0	12Va02
$\text{C}_{12} \text{H}_{11}-^{155}\text{Gd}$	163530	70	163445.6	1.3	-0.3	U			R04	4.0	64De15
$\text{C}_{11} \text{}^{13}\text{C} \text{H}_{10}-^{155}\text{Gd}$	158921	42	158975.4	1.3	0.3	U			R04	4.0	64De15
$\text{C}_{10} \text{}^{13}\text{C}_2 \text{H}_9-^{155}\text{Gd}$	154450	140	154505.2	1.3	0.1	U			R04	4.0	64De15
$\text{C}_{10} \text{H}_7 \text{N}_2-^{155}\text{Gd}$	138213	38	138293.4	1.3	0.5	U			R04	4.0	64De15
$^{155}\text{Gd}-^{139}\text{La} \text{O}$	21252	32	21356.4	2.3	0.8	U			R05	4.0	65De13
$^{155}\text{Tb}-\text{u}$	-76431	30	-76490	11	-2.0	U			GS2	1.0	05Li24
$^{155}\text{Dy}-\text{u}$	-74227	30	-74242	10	-0.5	U			GS2	1.0	05Li24
$^{155}\text{Ho}-\text{u}$	-70867	30	-70896	19	-1.0	1	39	39 $^{155}\text{Ho}$	GS2	1.0	05Li24
$^{155}\text{Er}-\text{u}$	-66785	30	-66784	7	0.0	U			GS2	1.0	05Li24
$^{155}\text{Tm}-\text{u}$	-60814	33	-60790	11	0.7	U			GS2	1.0	05Li24 *
$^{155}\text{Gd} \text{}^{35}\text{Cl}_3-^{149}\text{Sm} \text{}^{37}\text{Cl}_3$	14282.4	6.3	14288.8	0.9	0.4	U			M21	2.5	75Ka25
$^{155}\text{Gd} \text{}^{35}\text{Cl}-^{153}\text{Eu} \text{}^{37}\text{Cl}$	4345.4	2.4	4342.9	0.8	-0.4	U			H25	2.5	72Ba08
$^{155}\text{Gd}-^{138}\text{La} \text{O}$	20558	49	20597	4	0.2	U			R05	4.0	65De13
$^{155}\text{Gd}-^{154}\text{Gd}$	1480	60	1756.40	0.20	1.2	U			R04	4.0	64De15
$^{155}\text{Gd} \text{O}-\text{C}_{15}$	-82452.8	5.0	-82455.6	1.3	-0.4	o			TG1	1.5	09Ke.A
	-82452.2	2.6			-0.9	1	11	11 $^{155}\text{Gd}$	TG1	1.5	11Ke03
$^{155}\text{Er}(\alpha)^{151}\text{Dy}$	4118.3	5.				3			ORa		74To07 Z
$^{155}\text{Tm}(\alpha)^{151}\text{Ho}$	4578.3	10.3	4572	5	-0.6	3			ORa		71To01 *
	4568.1	10.			0.4	3			ORa		71To01 *
	4570.1	8.			0.2	3					92Ha10 *
$^{155}\text{Yb}(\alpha)^{151}\text{Er}$	5344.1	5.	5338.8	2.1	-1.1	3			GSa		79Ho10
	5336.6	5.			0.4	3			Bka		82Bo04 Z
	5344.2	5.			-1.1	3					87Ka.A
	5331.8	4.			1.7	3			ORa		91To08
	5340.1	4.			-0.3	3			Daa		96Pa01
$^{155}\text{Lu}(\alpha)^{151}\text{Tm}$	5796.9	5.	5802.8	2.6	1.2	5					89Ho12 *
	5797.9	5.			1.0	5			ORa		91To08
	5805.1	5.			-0.5	5			Daa		96Pa01
	5811.2	5.			-1.7	5			Ara		97Da07
$^{155}\text{Lu}^m(\alpha)^{151}\text{Tm}^m$	5723.0	10.	5730.6	2.8	0.7	6					89Ho12
	5727.1	5.			0.7	6			ORa		91To08
	5732.2	5.			-0.3	6			Daa		96Pa01
	5734.2	5.			-0.7	6			Ara		97Da07
$^{155}\text{Lu}^n(\alpha)^{151}\text{Tm}$	7574.9	15.	7584	3	0.2	U					89Ho12 *
	7586.2	5.			-0.5	o			Daa		96Pa01 *
$^{155}\text{Gd}(n,\alpha)^{152}\text{Sm}$	8331	6	8339.1	0.3	1.4	U			McM		69Be17
$^{155}\text{Gd}(p,t)^{153}\text{Gd}$	-6850	7	-6848.17	0.25	0.3	U			McM		73Lo08
	-6853	5			1.0	U			Min		73Oo01
$^{154}\text{Sm}(n,\gamma)^{155}\text{Sm}$	5806.8	0.6	5806.96	0.27	0.3	2					82Ba15 Z
	5807.0	0.3			-0.1	2			ILn		82Sc03 Z
$^{154}\text{Sm}(d,p)^{155}\text{Sm}$	3584	12	3582.39	0.27	-0.1	U			Tal		65Ke09
$^{154}\text{Eu}(n,\gamma)^{155}\text{Eu}$	8151.3	0.4	8151.3	0.4	0.0	1	100	98 $^{155}\text{Eu}$	ILn		86Pr03
$^{154}\text{Gd}(n,\gamma)^{155}\text{Gd}$	6435.11	0.30	6435.24	0.18	0.4	-			ILn		86Sc25 Z
	6435.29	0.23			-0.2	-			Bdn		06Fi.A
$^{154}\text{Gd}(d,p)^{155}\text{Gd}$	4217	10	4210.68	0.18	-0.6	U			Kop		67Tj01
$^{155}\text{Gd}(d,t)^{154}\text{Gd}$	-190	10	-178.01	0.18	1.2	U			Kop		67Tj01
$^{154}\text{Gd}(n,\gamma)^{155}\text{Gd}$	ave. 6435.22	0.18	6435.24	0.18	0.1	1	99	73 $^{154}\text{Gd}$			average
$^{155}\text{Ta}(p)^{154}\text{Hf}$	1776	10	1453	15	-32.3	B			Arp		99Uu01 *
	1453	15				3			Jya		07Pa27

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{155}\text{Nd}(\beta^-)^{155}\text{Pm}$	4222	150	4656	10	2.9	U			Ida		93Gr17
$^{155}\text{Pm}(\beta^-)^{155}\text{Sm}$	3224	30	3251	5	0.9	U			Ida		93Gr17
$^{155}\text{Sm}(\beta^-)^{155}\text{Eu}$	1634	15	1627.3	1.2	-0.4	U					63Kr04 *
	1624	15			0.2	U					65Fu13 *
	1607	25			0.8	U			Ida		93Gr17
$^{155}\text{Eu}(\beta^-)^{155}\text{Gd}$	252	5	251.8	0.9	0.0	U					54Le08
	245	5			1.4	U					58G156
	245	5			1.4	U					59Am16
$^{155}\text{Dy}(\beta^+)^{155}\text{Tb}$	2099	6	2094.5	1.9	-0.8	2					63Pe13 *
	2094	2			0.2	2					80Bu04 *
$^{155}\text{Ho}(\beta^+)^{155}\text{Dy}$	3102	20	3116	17	0.7	1	69	61 $^{155}\text{Ho}$			72To07 *
$^{155}\text{Lu}^m(\text{IT})^{155}\text{Lu}$	23.0	6.2	21	4	-0.3	5					96Pa01
	19.9	6.2			0.2	5					97Da07
$^{155}\text{Lu}^n(\text{IT})^{155}\text{Lu}$	1781	2				5					96Pa01
* $^{155}\text{Pr}-u$	Represents frequency ratio $^{155}\text{Pr}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.98142559(20)$										WgM124**
* $^{155}\text{Nd}-u$	Represents frequency ratio $^{155}\text{Nd}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.98147230(11)$										WgM124**
* $^{155}\text{Pm}-u$	Represents frequency ratio $^{155}\text{Pm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.981503965(56)$										WgM124**
* $^{155}\text{Sm}-u$	Represents frequency ratio $^{155}\text{Sm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.98152610(15)$										WgM124**
* $^{155}\text{Tm}-u$	$M - A = -56627(28)$ keV for mixture gs+m at 41(6) keV										Nub16b **
* $^{155}\text{Tm}(\alpha)^{151}\text{Ho}$	First assigned to $^{156}\text{Tm}^m$ but belonging to $^{155}\text{Tm}$ ground state										94To10 **
* $^{155}\text{Tm}(\alpha)^{151}\text{Ho}$	Doublet from ground state and isomer, less than 5 keV apart										90Po13 **
* $^{155}\text{Lu}(\alpha)^{151}\text{Tm}$	Original value $E=5656(6)$ ( $Q=5806.1$ ) recalibrated										79Ho10 **
* $^{155}\text{Lu}^n(\alpha)^{151}\text{Tm}$	Original value $E=7408(10)$ recalibrated										81Ho.A **
* $^{155}\text{Lu}^n(\alpha)^{151}\text{Tm}$	Replaced by authors' value for $^{155}\text{Lu}^n(\text{IT})$										AHW **
* $^{155}\text{Ta}(p)^{154}\text{Hf}$	$E_p=1765(10)$ for ( $11/2^-$ ) state; ground state may be $1/2^+$ , slightly lower										99Uu01 **
*	1776 keV proton not observed in coincidence with feeding $\alpha$										07Pa27 **
* $^{155}\text{Sm}(\beta^-)^{155}\text{Eu}$	$E_{\beta^-}=1530(15)$ $E_{\beta^-}=1520(15)$ respectively, to $5/2^-$ level at 104.334 keV										Ens051 **
* $^{155}\text{Dy}(\beta^+)^{155}\text{Tb}$	$E_{\beta^+}=850(6)$ $845(2)$ respectively, to $5/2^-$ level at 226.918 keV, and other $E_{\beta^+}$										Ens051 **
* $^{155}\text{Ho}(\beta^+)^{155}\text{Dy}$	$E_{\beta^+}=1840(20)$ to $3/2^+$ level at 240.196 keV										Ens051 **
$^{156}\text{Pm}-u$	-68883.4	6.9	-68883	4	0.1	1	32	32 $^{156}\text{Pm}$	CP1	1.0	12Va02 *
$^{156}\text{Pm}-^{80}\text{Kr}_{1.950}$	94181.7	6.4	94180	4	-0.2	1	38	35 $^{156}\text{Pm}$	CP1	1.0	12Va02
$^{156}\text{Pm}-^{86}\text{Kr}_{1.814}$	93269.1	6.8	93270	4	0.1	1	33	33 $^{156}\text{Pm}$	CP1	1.0	12Va02
$\text{C}_{12}\text{H}_{12}-^{156}\text{Gd}$	171923	44	171769.8	1.3	-0.9	U			R04	4.0	64De15
$\text{C}_{11}\text{}^{13}\text{C}\text{H}_{11}-^{156}\text{Gd}$	167384	43	167299.6	1.3	-0.5	U			R04	4.0	64De15
$\text{C}_{10}\text{}^{13}\text{C}_2\text{H}_{10}-^{156}\text{Gd}$	162810	60	162829.4	1.3	0.1	U			R04	4.0	64De15
$\text{C}_{10}\text{H}_8\text{N}_2-^{156}\text{Gd}$	146661	38	146617.7	1.3	-0.3	U			R04	4.0	64De15
$^{156}\text{Tb}-u$	-75165	40	-75246	4	-2.0	U			GS2	1.0	05Li24 *
$\text{C}_{10}\text{H}_8\text{N}_2-^{156}\text{Dy}$	145130	100	144464.2	1.3	-1.7	U			R04	4.0	64De15
$^{156}\text{Dy}-^{133}\text{Cs}_{1.173}$	35195.1	4.8	35188.9	1.3	-1.3	1	7	7 $^{156}\text{Dy}$	MA8	1.0	17Ma.A
$^{156}\text{Ho}-u$	-70107	122	-70290	60	-1.5	o			GS1	1.0	00Ra23 *
$^{156}\text{Ho}^n-u$	-70107	30				2			GS2	1.0	05Li24 *
$^{156}\text{Er}-u$	-68907	30	-68934	26	-0.9	1	78	78 $^{156}\text{Er}$	GS2	1.0	05Li24
$^{156}\text{Tm}-u$	-61044	30	-61014	15	1.0	U			GS2	1.0	05Li24
$^{156}\text{Yb}-u$	-57202	30	-57183	10	0.6	U			GS2	1.0	05Li24
$^{156}\text{Gd}\text{}^{35}\text{Cl}-^{154}\text{Gd}\text{}^{37}\text{Cl}$	4199	5	4207.27	0.22	0.4	U			H12	4.0	64Ba15
	4206	10			0.1	U			H21	2.5	70Ma05
	4204.8	1.4			0.7	U			H25	2.5	72Ba08
	4203.0	1.0			1.7	U			M21	2.5	75Ka25
$^{156}\text{Dy}-^{156}\text{Gd}$	2153.47	0.11	2153.48	0.11	0.0	1	100	92 $^{156}\text{Dy}$	SH1	1.0	11El05
$^{156}\text{Gd}-^{139}\text{La}\text{O}$	20618	71	20857.1	2.3	0.8	U			R05	4.0	65De13
$^{156}\text{Gd}-^{155}\text{Gd}$	-584	33	-499.23	0.07	0.6	U			R04	4.0	64De15
$^{156}\text{Gd}\text{O}-\text{C}_{15}$	-82946.5	5.8	-82954.8	1.3	-1.0	o			TG1	1.5	09Ke.A
	-82945.6	3.6			-1.7	U			TG1	1.5	11Ke03
$^{156}\text{Er}(\alpha)^{152}\text{Dy}$	3109.9	70.	3481	25	5.3	C					95Ka.A
$^{156}\text{Tm}(\alpha)^{152}\text{Ho}$	4341.6	10.	4345	7	0.4	-			ORa		71To10
	4345.6	10.			0.0	-					81Ga36
ave.	4344	7			0.2	1	98	94 $^{156}\text{Tm}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{156}\text{Tm}^m(\alpha)^{152}\text{Ho}$	4737.5	10.	*			F			ORa		71To01 *
$^{156}\text{Yb}(\alpha)^{152}\text{Er}$	4813.6	10.	4810	4	-0.4	-					77Ha48
	4809.6	10.			0.0	-			GSa		79Ho10
	4810.6	4.			-0.2	-			Daa		96Pa01
ave.	4811	4			-0.3	1	98	83	$^{156}\text{Yb}$		average
$^{156}\text{Lu}(\alpha)^{152}\text{Tm}$	5593.7	10.	5596	3	0.2	U			GSa		79Ho10
	5592.7	5.			0.6	2			DbA		92Po14
	5597.9	4.			-0.5	2			Daa		96Pa01
$^{156}\text{Lu}^m(\alpha)^{152}\text{Tm}^m$	5713.7	5.	5711.5	2.6	-0.4	3			GSa		79Ho10 Z
	5709.7	5.			0.4	3			DbA		92Po14
	5709.7	8.			0.2	3					92Ha10
	5711.7	4.			-0.1	3			Daa		96Pa01
$^{156}\text{Hf}(\alpha)^{152}\text{Yb}$	6033.0	10.	6029	4	-0.4	-			GSa		79Ho10
	6027.9	4.			0.2	-			Daa		96Pa01
ave.	6029	4			0.0	1	100	100	$^{156}\text{Hf}$		average
$^{156}\text{Hf}^m(\alpha)^{152}\text{Yb}$	8009.8	15.	7988	4	-1.5	U			GSa		81Ho.A
	7987.2	4.			0.1	o			Daa		96Pa01 *
$^{154}\text{Sm}(t,p)^{156}\text{Sm}$	4556	25	4566	9	0.4	1	12	11	$^{156}\text{Sm}$		66Bj01
$^{154}\text{Eu}(t,p)^{156}\text{Eu}$	6003	4	6005	3	0.6	1	71	70	$^{156}\text{Eu}$		84La06 *
$^{154}\text{Gd}(t,p)^{156}\text{Gd}$	6495.1	3.6	6489.80	0.19	-1.5	U			McM		89Lo07
$^{156}\text{Gd}(p,t)^{154}\text{Gd}$	-6490	7	-6489.80	0.19	0.0	U			McM		73Lo08
	-6490	5			0.0	U			Min		73Oo01
$^{155}\text{Gd}(n,\gamma)^{156}\text{Gd}$	8536.8	0.5	8536.35	0.07	-0.9	U			ILn		82Ba28
	8536.39	0.07			-0.6	-			MMn		82Is05 Z
	8536.04	0.19			1.6	-			Bdn		06Fi.A
$^{155}\text{Gd}(d,p)^{156}\text{Gd}$	6319	10	6311.78	0.07	-0.7	U			Kop		67Tj01
$^{156}\text{Gd}(d,t)^{155}\text{Gd}$	-2287	10	-2279.12	0.07	0.8	U			Kop		67Tj01
$^{155}\text{Gd}(n,\gamma)^{156}\text{Gd}$	ave.	8536.35	0.07	8536.35	0.07	0.0	1	100	59	$^{155}\text{Gd}$	average
$^{155}\text{Gd}(\alpha,t)^{156}\text{Tb}-^{158}\text{Gd}()^{159}\text{Tb}$	-821.9	3.6	-822	4	0.0	1	100	100	$^{156}\text{Tb}$	McM	75Bu02
$^{156}\text{Dy}(d,t)^{155}\text{Dy}$	-3184	10	-3188	10	-0.4	1	92	92	$^{155}\text{Dy}$	Kop	70Gr46
$^{156}\text{Ta}(p)^{155}\text{Hf}$	1028.6	13.	1020	4	-0.7	o			Dap		92Pa05
	1013.6	5.			1.2	o			Dap		96Pa01
	1017.9	5.			0.4	3			Dap		11Da12
$^{156}\text{Ta}^m(p)^{155}\text{Hf}$	1110.2	12.	1114	7	0.3	3			Dap		93Li34
	1115.2	8.			-0.2	3			Dap		96Pa01
$^{156}\text{Nd}(\beta^-)^{156}\text{Pm}$	3690	200				2			Kur		02Sh.B *
$^{156}\text{Pm}(\beta^-)^{156}\text{Sm}$	5155	35	5197	9	1.2	U			Stu		90He11
	5110	100			0.9	U			Kur		02Sh.B
$^{156}\text{Sm}(\beta^-)^{156}\text{Eu}$	721	10	722	8	0.1	-					63Gu04 *
	721	15			0.1	-					65Wi08 *
ave.	721	8			0.1	1	90	89	$^{156}\text{Sm}$		average
$^{156}\text{Eu}(\beta^-)^{156}\text{Gd}$	2430	10	2452	3	2.2	-					62Ew01
	2460	10			-0.8	-					63Th02
	2450	15			0.2	-					64Pe17
	2478	20			-1.3	U					67Va23
ave.	2446	6			1.0	1	28	28	$^{156}\text{Eu}$		average
$^{156}\text{Tb}(\beta^+)^{156}\text{Gd}$	3570	50	2444	4	-22.5	B					70Ag02 *
$^{156}\text{Ho}(\beta^+)^{156}\text{Dy}$	4400	400	5050	60	1.6	U					76Gr20 *
	5050	90			0.0	o					02Iz01
	5050	60				2					04Iz02 *
$^{156}\text{Er}(\beta^+)^{156}\text{Ho}$	1670	70	1270	60	-5.8	B					82Vy06 *
$^{156}\text{Tm}(\beta^+)^{156}\text{Er}$	7458	50	7377	27	-1.6	1	29	22	$^{156}\text{Er}$	Dbn	94Po26 *
	7390	100			-0.1	U					95Ga.A
$^{156}\text{Hf}^m(\text{IT})^{156}\text{Hf}$	1959	1				2					96Pa01
* $^{156}\text{Pm}-u$	Represents frequency ratio $^{156}\text{Pm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.975190689(43)$										
* $^{156}\text{Tb}-u$	$M-A = -69968(32)$ keV for mixture gs+m+n at 54(3) and 88.4 keV										
* $^{156}\text{Ho}-u$	$M-A = -65230(100)$ keV for mixture gs+m+n at 52.37 and 170(70) keV										
* $^{156}\text{Ho}^n-u$	Assuming high spin isomer is favored										
* $^{156}\text{Tm}^m(\alpha)^{152}\text{Ho}$	F : originally $E_\alpha = 4460(10)$ to $^{152}\text{Ho}^m$ at 160(1), reassigned to $^{155}\text{Tm}$										
* $^{156}\text{Hf}^m(\alpha)^{152}\text{Yb}$	Replaced by authors' value for $^{156}\text{Hf}^m(\text{IT})$										
* $^{154}\text{Eu}(t,p)^{156}\text{Eu}$	$Q = 5569(4)$ to $3^-$ level at 434.23 keV										
* $^{156}\text{Nd}(\beta^-)^{156}\text{Pm}$	Trends from Mass Surface TMS suggest $^{156}\text{Nd}$ 70 less bound										
											GAu **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>156</sup> Sm( $\beta^-$ ) <sup>156</sup> Eu	$E_{\beta^-}$ =430(10) 430(15) respectively, to 1 <sup>+</sup> level at 291.3037 keV										Ens12a **
* <sup>156</sup> Tb( $\beta^+$ ) <sup>156</sup> Gd	$E_{\beta^+}$ =2640(50) from <sup>156</sup> Tb <sup>II</sup> at 88.4 to ground state										Nub16b **
* <sup>156</sup> Ho( $\beta^+$ ) <sup>156</sup> Dy	$E_{\beta^+}$ =1800(50) to levels around 1600										Ens12a **
* <sup>156</sup> Ho( $\beta^+$ ) <sup>156</sup> Dy	Original error 20 is for statistics only, increased by evaluator										GAu **
* <sup>156</sup> Er( $\beta^+$ ) <sup>156</sup> Ho	$p^+$ =0.0036(0.0017) to 2 <sup>-</sup> level at 82.23 keV, reanalyzed										Ens12a **
* <sup>156</sup> Tm( $\beta^+$ ) <sup>156</sup> Er	$E_{\beta^+}$ =6091(50) to 2 <sup>+</sup> level at 344.53 keV										Ens12a **
<sup>157</sup> Nd-u	-60614	47	-60614	27	0.0	1	32	32 <sup>157</sup> Nd	CP1	1.0	12Va02 *
<sup>157</sup> Nd- <sup>80</sup> Kr <sub>1.963</sub>	103537	46	103536	27	0.0	1	34	34 <sup>157</sup> Nd	CP1	1.0	12Va02
<sup>157</sup> Nd- <sup>86</sup> Kr <sub>1.826</sub>	102610	46	102611	27	0.0	1	34	34 <sup>157</sup> Nd	CP1	1.0	12Va02
<sup>157</sup> Pm-u	-66880	13	-66879	8	0.1	1	33	33 <sup>157</sup> Pm	CP1	1.0	12Va02 *
<sup>157</sup> Pm- <sup>80</sup> Kr <sub>1.963</sub>	97273	13	97271	8	-0.1	1	34	33 <sup>157</sup> Pm	CP1	1.0	12Va02
<sup>157</sup> Pm- <sup>86</sup> Kr <sub>1.826</sub>	96346	13	96346	8	0.0	1	33	33 <sup>157</sup> Pm	CP1	1.0	12Va02
<sup>157</sup> Sm-u	-71582.2	8.3	-71581	5	0.1	1	33	33 <sup>157</sup> Sm	CP1	1.0	12Va02 *
<sup>157</sup> Sm- <sup>80</sup> Kr <sub>1.963</sub>	92570.0	8.0	92569	5	-0.2	1	36	34 <sup>157</sup> Sm	CP1	1.0	12Va02
<sup>157</sup> Sm- <sup>86</sup> Kr <sub>1.826</sub>	91643.0	8.3	91644	5	0.1	1	33	33 <sup>157</sup> Sm	CP1	1.0	12Va02
C <sub>10</sub> H <sub>9</sub> N <sub>2</sub> - <sup>157</sup> Gd	152720	60	152605.4	1.3	-0.5	U			R04	4.0	64De15
C <sub>9</sub> <sup>13</sup> C H <sub>8</sub> N <sub>2</sub> - <sup>157</sup> Gd	148170	70	148135.2	1.3	-0.1	U			R04	4.0	64De15
C <sub>10</sub> H <sub>5</sub> O <sub>2</sub> - <sup>157</sup> Gd	105080	60	104986.5	1.3	-0.4	U			R04	4.0	64De15
<sup>157</sup> Ho-u	-71724	30	-71748	25	-0.8	1	71	71 <sup>157</sup> Ho	GS2	1.0	05Li24
<sup>157</sup> Er-u	-68084	30	-68077	28	0.2	1	90	90 <sup>157</sup> Er	GS2	1.0	05Li24
<sup>157</sup> Tm-u	-63027	30				2			GS2	1.0	05Li24
<sup>157</sup> Yb-u	-57389	30	-57351	12	1.3	U			GS2	1.0	05Li24
<sup>157</sup> Lu-u	-49842	31	-49856	13	-0.5	1	17	17 <sup>157</sup> Lu	GS2	1.0	05Li24 *
<sup>157</sup> Gd <sup>35</sup> Cl- <sup>155</sup> Gd <sup>37</sup> Cl	4318	4	4288.18	0.19	-1.9	U			H12	4.0	64Ba15
	4287	3			0.2	U			H21	2.5	70Ma05
	4289.0	0.7			-0.5	U			M21	2.5	75Ka25
	4288.83	0.66			-0.4	U			H41	2.5	85Dy04
<sup>157</sup> Gd- <sup>156</sup> Gd	1860	60	1837.31	0.16	-0.1	U			R04	4.0	64De15
<sup>157</sup> Gd O-C <sub>15</sub>	-81114.2	5.4	-81117.5	1.3	-0.4	o			TG1	1.5	09Ke.A
	-81113.6	3.3			-0.8	U			TG1	1.5	11Ke03
<sup>157</sup> Yb( $\alpha$ ) <sup>153</sup> Er	4622.0	7.	4622	6	0.0	-					77Ha48
	4623.0	10.			-0.1	-			GSa		79Ho10
	ave.	4622	6		-0.1	1	99	96 <sup>157</sup> Yb			average
<sup>157</sup> Lu( $\alpha$ ) <sup>153</sup> Tm	5097.2	5.	5107.9	2.9	2.1	o			DbA		91Le15 *
	5096.2	20.			0.6	U			Bka		91To09 *
	5111.5	5.			-0.7	o			DbA		92Po14 *
<sup>157</sup> Lu <sup>m</sup> ( $\alpha$ ) <sup>153</sup> Tm	5128.9	10.	5128.8	2.0	0.0	U			IRa		79Al16 Z
	5131.8	5.			-0.6	-			GSa		79Ho10 Z
	5133.7	5.			-0.9	-			ORa		83To01 Z
	5128.9	5.			0.0	o			DbA		91Le15
	5118.7	5.			2.0	-			Bka		91To09
	5125.8	6.			0.5	-					92Ha10
	5132.0	5.			-0.6	-			DbA		92Po14
	5127.9	4.			0.2	-			Daa		96Pa01
	ave.	5128.4	2.1		0.2	1	99	54 <sup>153</sup> Tm			average
<sup>157</sup> Hf( $\alpha$ ) <sup>153</sup> Yb	5869.4	10.	5880	3	1.0	3					73Ea01 Z
	5884.1	5.			-0.8	3			GSa		79Ho10 Z
	5879.1	4.			0.2	3			Daa		96Pa01
<sup>157</sup> Ta( $\alpha$ ) <sup>153</sup> Lu <sup>m</sup>	6277.2	4.	6275	8	-0.6	o			Ara		97Ir01 *
<sup>157</sup> Ta <sup>m</sup> ( $\alpha$ ) <sup>153</sup> Lu	6381.9	10.	6377	4	-0.5	3			GSa		79Ho10
	6375.8	4.			0.2	3			Daa		96Pa01 *
<sup>157</sup> Ta <sup>n</sup> ( $\alpha$ ) <sup>153</sup> Lu	7946.9	8.	7948	8	0.0	o			Daa		96Pa01 *
<sup>155</sup> Gd(t,p) <sup>157</sup> Gd	6417.8	2.9	6414.43	0.16	-1.2	U			McM		89Lo07
<sup>157</sup> Gd(p,t) <sup>155</sup> Gd	-6414	7	-6414.43	0.16	-0.1	U			McM		73Lo08
	-6417	5			0.5	U			Min		73Oo01

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{156}\text{Gd}(n,\gamma)^{157}\text{Gd}$	6359.6	0.8	6359.88	0.15	0.3	U					70Bo29
	6360	1			-0.1	U					71Gr42
	6359.80	0.15			0.5	o					87Sp.A Z
	6359.86	0.15			0.1	1	99	57 $^{156}\text{Gd}$	ILn		03Bo25
$^{157}\text{Gd}(\gamma,n)^{156}\text{Gd}$	-6350	80	-6359.88	0.15	-0.1	U			Phi		60Ge01
$^{156}\text{Gd}(d,p)^{157}\text{Gd}$	4136	10	4135.31	0.15	-0.1	U			Kop		67Tj01
$^{157}\text{Gd}(d,t)^{156}\text{Gd}$	-112	10	-102.65	0.15	0.9	U			Kop		67Tj01
$^{156}\text{Gd}(\alpha,t)^{157}\text{Tb}-^{158}\text{Gd}(\alpha)^{159}\text{Tb}$	-616.2	2.0	-614.3	0.8	0.9	1	16	9 $^{159}\text{Tb}$	McM		75Bu02
$^{156}\text{Dy}(d,p)^{157}\text{Dy}$	4748	10	4742	5	-0.6	-			Tal		68Be.A
	4753	10			-1.1	-			Kop		70Gr46
	ave. 4750	7			-1.2	1	53	52 $^{157}\text{Dy}$			average
$^{157}\text{Ta}(p)^{156}\text{Hf}$	925.0	17.	935	10	0.6	o			Dap		96Pa01
	933.0	7.			0.2	o			Ara		97Ir01 *
$^{157}\text{Pm}(\beta^-)^{157}\text{Sm}$	4360	100	4381	8	0.2	U			Kur		02Sh.B
$^{157}\text{Sm}(\beta^-)^{157}\text{Eu}$	2700	200	2781	6	0.4	U					73Ka23 *
	2734	50			0.9	U			Ida		93Gr17
$^{157}\text{Eu}(\beta^-)^{157}\text{Gd}$	1350	20	1365	4	0.7	U					64Sh21 *
	1370	20			-0.3	U					66Fu05 *
$^{157}\text{Tb}(\epsilon)^{157}\text{Gd}$	62.4	0.6	60.04	0.30	-3.9	B					67Na08 *
	62.2	0.6			-3.6	B					83Be42 *
	60.0	0.3			0.1	1	98	93 $^{157}\text{Tb}$			92Ra18 *
$^{157}\text{Ho}(\beta^+)^{157}\text{Dy}$	2540	50	2592	24	1.0	1	23	22 $^{157}\text{Ho}$			72To05 *
$^{157}\text{Er}(\beta^+)^{157}\text{Ho}$	3470	80	3420	30	-0.6	1	18	10 $^{157}\text{Er}$			75Al.A
	3805	100			-3.9	B			Dbn		94Po26 *
$^{157}\text{Tm}(\beta^+)^{157}\text{Er}$	4480	100	4700	40	2.2	B			IRS		93Al03
	4482	100			2.2	B			Dbn		94Po26
$^{157}\text{Yb}(\beta^+)^{157}\text{Tm}$	5074	100	5290	30	2.1	U			Dbn		94Po26
$^{157}\text{Lu}^m(\text{IT})^{157}\text{Lu}$	32	2	20.9	2.0	-5.5	B			Dba		91Le15
	21	2			0.0	1	100	83 $^{157}\text{Lu}$	Dba		92Po14 *
$^{157}\text{Ta}^m(\text{IT})^{157}\text{Ta}$	22	5				3					97Ir01
$^{157}\text{Ta}^n(\text{IT})^{157}\text{Ta}^m$	1571	7				3			Daa		96Pa01
* $^{157}\text{Nd}-u$	Represents frequency ratio $^{157}\text{Nd}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.96892546(29)$										WgM124**
* $^{157}\text{Pm}-u$	Represents frequency ratio $^{157}\text{Pm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.968964141(81)$										WgM124**
* $^{157}\text{Sm}-u$	Represents frequency ratio $^{157}\text{Sm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.968993178(51)$										WgM124**
* $^{157}\text{Lu}-u$	$M-A = -46417(28)$ keV for mixture gs+m at 20.9(2.0) keV										Nub16b **
* $^{157}\text{Lu}(\alpha)^{153}\text{Tm}$	$E_\alpha = 4925(5)$ to $^{153}\text{Tm}^m$ at 43.2(0.2) keV										Nub16b **
* $^{157}\text{Lu}(\alpha)^{153}\text{Tm}$	$E_\alpha = 4924(20)$ to $^{153}\text{Tm}^m$ at 43.2(0.2) keV										Nub16b **
* $^{157}\text{Lu}(\alpha)^{153}\text{Tm}$	$E_\alpha = 4939(5)$ to $^{153}\text{Tm}^m$ at 43.2(0.2) keV; replaced by $^{157}\text{Lu}^m(\text{IT})$										Nub16b **
* $^{157}\text{Ta}(\alpha)^{153}\text{Lu}^m$	Replaced by $^{153}\text{Lu}^m(\text{IT})$										AHW **
* $^{157}\text{Ta}^m(\alpha)^{153}\text{Lu}$	Reassigned										97Ir01 **
* $^{157}\text{Ta}^n(\alpha)^{153}\text{Lu}$	Replaced by authors' value for $^{157}\text{Ta}^n(\text{IT})$										AHW **
* $^{157}\text{Ta}(p)^{156}\text{Hf}$	Use instead $^{157}\text{Ta}^m(\text{IT})$										AHW **
* $^{157}\text{Sm}(\beta^-)^{157}\text{Eu}$	$E_{\beta^-} = 2400(200)$ to $5/2^-$ level at 197.863 and $3/2^+$ at 394.334 keV										Ens162 **
* $^{157}\text{Eu}(\beta^-)^{157}\text{Gd}$	$E_{\beta^-} = 870(30)$ 910(20) respectively, to $3/2^+$ level at 474.630 keV, and other $E_{\beta^-}$										Ens162 **
* $^{157}\text{Tb}(\epsilon)^{157}\text{Gd}$	LK=2.65(0.20); original value 66(6) recalculated										92Ha03 **
* $^{157}\text{Tb}(\epsilon)^{157}\text{Gd}$	LK=2.69(0.20); original value 62.9(0.7) recalculated										85Vo09 **
* $^{157}\text{Ho}(\beta^+)^{157}\text{Dy}$	$E_{\beta^+} = 1180(50)$ to $5/2^-$ level at 341.118 keV										Ens162 **
* $^{157}\text{Er}(\beta^+)^{157}\text{Ho}$	$E_{\beta^+} = 2525(100)$ to ground state yielding 3547(100), rather 24% to $(3/2^+)$										94Po26 **
*	level at 174.55 keV, 15% to $5/2^-$ at 391.32 keV $\rightarrow$ +258 keV										Ens162 **
* $^{157}\text{Lu}^m(\text{IT})^{157}\text{Lu}$	Derived from $^{157}\text{Lu}^m(\alpha)-^{157}\text{Lu}(\alpha)$ difference										92Po14 **
$^{158}\text{Pm}-u$	-63436	25	-63435	14	0.0	1	33	33 $^{158}\text{Pm}$	CP1	1.0	12Va02 *
$^{158}\text{Pm}-^{80}\text{Kr}_{1.975}$	101720	25	101718	14	-0.1	1	33	33 $^{158}\text{Pm}$	CP1	1.0	12Va02
$^{158}\text{Pm}-^{86}\text{Kr}_{1.837}$	100773	25	100773	14	0.0	1	33	33 $^{158}\text{Pm}$	CP1	1.0	12Va02
$^{158}\text{Sm}-u$	-70049.2	9.5	-70049	5	0.0	1	31	31 $^{158}\text{Sm}$	CP1	1.0	12Va02 *



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference		
$^{158}\text{Sm}-^{80}\text{Kr}_{1,975}$	95106.5	9.1	95104	5	-0.2	1	34	32	$^{158}\text{Sm}$	CP1	1.0	12Va02	
$^{158}\text{Sm}-^{86}\text{Kr}_{1,837}$	94159.3	9.4	94159	5	0.0	1	31	31	$^{158}\text{Sm}$	CP1	1.0	12Va02	
$^{158}\text{Eu}-u$	-72208	25	-72201	11	0.3	1	19	19	$^{158}\text{Eu}$	CP1	1.0	12Va02	*
$^{158}\text{Eu}-^{80}\text{Kr}_{1,975}$	92949	25	92952	11	0.1	1	20	19	$^{158}\text{Eu}$	CP1	1.0	12Va02	
$^{158}\text{Eu}-^{86}\text{Kr}_{1,837}$	92001	25	92007	11	0.2	1	19	19	$^{158}\text{Eu}$	CP1	1.0	12Va02	
$\text{C}_{10}\text{H}_6\text{O}_2-^{158}\text{Gd}$	112444	33	112667.8	1.3	1.7	U				R04	4.0	64De15	
$\text{C}_{10}\text{H}_6\text{O}_2-^{158}\text{Dy}$	112870	100	112364.8	2.5	-1.3	U				R04	4.0	64De15	
$^{158}\text{Ho}-u$	-71101	67	-71055	29	0.7	R				GS2	1.0	05Li24	*
$^{158}\text{Er}-u$	-70220	110	-70107	27	1.0	U				GS1	1.0	00Ra23	
	-70107	30			0.0	1	81	81	$^{158}\text{Er}$	GS2	1.0	05Li24	
$^{158}\text{Tm}-u$	-63080	110	-63020	27	0.5	U				GS1	1.0	00Ra23	
	-63020	30			0.0	1	81	81	$^{158}\text{Tm}$	GS2	1.0	05Li24	
$^{158}\text{Yb}-^{142}\text{Sm}_{1,113}$	34252	22	34248	9	-0.2	1	16	14	$^{158}\text{Yb}$	MA7	1.0	01Bo59	
$^{158}\text{Lu}-u$	-50720	30	-50684	16	1.2	R				GS2	1.0	05Li24	
$^{158}\text{Gd}-^{35}\text{Cl}-^{156}\text{Gd}-^{37}\text{Cl}$	4956	4	4931.19	0.19	-1.6	U				H12	4.0	64Ba15	
	4929	3			0.3	U				H21	2.5	70Ma05	
	4926.2	1.4			1.4	U				H25	2.5	72Ba08	
	4930.8	0.7			0.2	U				M21	2.5	75Ka25	
	4930.13	1.36			0.3	U				H41	2.5	85Dy04	
$^{158}\text{Dy}-^{35}\text{Cl}-^{156}\text{Dy}-^{37}\text{Cl}$	3081.4	3.3	3080.7	2.6	-0.1	U				H25	2.5	72Ba08	
$^{158}\text{Gd}-^{157}\text{Gd}$	392	48	143.78	0.07	-1.3	U				R04	4.0	64De15	
$^{158}\text{Gd}\text{O}-\text{C}_{15}$	-80968.3	5.4	-80973.7	1.3	-0.7	o				TG1	1.5	09Ke.A	
	-80967.8	3.2			-1.2	U				TG1	1.5	11Ke03	
$^{158}\text{Gd}\text{O}-\text{C}_{14}$	-80964.7	8.2			-0.7	U				TG1	1.5	11Ke03	
$^{158}\text{Yb}(\alpha)^{154}\text{Er}$	4174.9	10.	4170	7	-0.5	-						77Ha48	
	4164.6	12.			0.4	-						92Ha10	
	ave.	4171	8		-0.1	1	80	71	$^{158}\text{Yb}$			average	
$^{158}\text{Lu}(\alpha)^{154}\text{Tm}$	4792.2	10.	4790	5	-0.2	3				IRa		79Al16	Z
	4789.5	5.			0.1	3				ORa		83To01	Z
$^{158}\text{Hf}(\alpha)^{154}\text{Yb}$	5406.0	5.	5404.8	2.7	-0.2	-				GSa		79Ho10	Z
	5401.4	5.			0.7	-				ORa		83To01	Z
	5406.1	4.			-0.3	-				Daa		96Pa01	
	ave.	5404.8	2.7		0.0	1	100	100	$^{158}\text{Hf}$			average	
$^{158}\text{Ta}(\alpha)^{154}\text{Lu}$	6124.4	8.	6124	4	-0.1	9				Daa		96Pa01	
	6123.3	5.			0.1	9				Ara		97Da07	
$^{158}\text{Ta}^m(\alpha)^{154}\text{Lu}^m$	6208.5	6.	6205.1	2.8	-0.6	10				GSa		79Ho10	
	6203.4	4.			0.4	10				Daa		96Pa01	
	6205.4	5.			-0.1	10				Ara		97Da07	
$^{158}\text{Ta}^n(\alpha)^{154}\text{Lu}^n$	8869.0	11.3				11				Jya		14Ca03	
$^{158}\text{W}(\alpha)^{154}\text{Hf}$	6600.4	30.	6613	3	0.4	U				GSa		81Ho10	*
	6609.7	30.			0.1	U				Daa		96Pa01	
	6612.7	3.				3				Ara		00Ma95	
$^{158}\text{W}^m(\alpha)^{154}\text{Hf}$	8495.5	30.	8502	7	0.2	U				GSa		89Ho12	
	8506.8	24.			-0.2	U				Daa		96Pa01	
	8501.6	7.				3				Ara		00Ma95	
$^{158}\text{Gd}(\text{p,t})^{156}\text{Gd}$	-5818	5	-5815.47	0.16	0.5	U				Min		73Oo01	
$^{158}\text{Dy}(\text{p,t})^{156}\text{Dy}$	-7535	15	-7539.2	2.4	-0.3	U				Pri		77Ko04	
$^{158}\text{Gd}(\text{t},\alpha)^{157}\text{Eu}-^{156}\text{Gd}()^{155}\text{Eu}$	-512	5	-514	4	-0.4	1	69	67	$^{157}\text{Eu}$	LAL		79Bu05	
$^{157}\text{Gd}(\text{n},\gamma)^{158}\text{Gd}$	7937.39	0.07	7937.39	0.06	0.0	-				MMn		82Is05	Z
	7937.39	0.17			0.0	-				Bdn		06Fi.A	
$^{157}\text{Gd}(\text{d,p})^{158}\text{Gd}$	5724	10	5712.82	0.06	-1.1	U				Kop		67Tj01	
	5706	5			1.4	U				Tal		71Sh04	
$^{158}\text{Gd}(\text{d,t})^{157}\text{Gd}$	-1688	10	-1680.16	0.06	0.8	U				Kop		67Tj01	
$^{157}\text{Gd}(\text{n},\gamma)^{158}\text{Gd}$	ave.	7937.39	0.06	7937.39	0.06	0.0	1	100	58	$^{158}\text{Gd}$		average	
$^{158}\text{Gd}(\text{d,t})^{157}\text{Gd}-^{159}\text{Tb}()^{158}\text{Tb}$	195.0	1.5	195.6	0.6	0.4	1	18	18	$^{158}\text{Tb}$	McM		84Bu14	
$^{157}\text{Gd}(\alpha,\text{t})^{158}\text{Tb}-^{158}\text{Gd}()^{159}\text{Tb}$	-198.3	1.0	-195.6	0.6	2.7	o				McM		75Bu02	
	-196.6	1.0			1.0	1	41	39	$^{158}\text{Tb}$	McM		84Bu14	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{158}\text{Tb}(p,d)^{157}\text{Tb}$	-4560.3	4.2	-4554.0	1.0	1.5	U			Pri		85Al02 *
$^{158}\text{Dy}(d,t)^{157}\text{Dy}$	-2804	10	-2797	5	0.7	-			Tal		68Be.A
	-2804	10			0.7	-			Kop		70Gr46
	ave.	-2804	7		1.0	1	53	48 $^{157}\text{Dy}$			average
$^{158}\text{Pm}(\beta^-)^{158}\text{Sm}$	6120	100	6161	14	0.4	o			Kur		02Sh.A
	6085	80			1.0	o			Kur		07Ha57
	6080	80			1.0	U			Kur		10Ha.A
$^{158}\text{Sm}(\beta^-)^{158}\text{Eu}$	1999	15	2005	10	0.4	1	48	42 $^{158}\text{Eu}$	Ida		93Gr17
$^{158}\text{Eu}(\beta^-)^{158}\text{Gd}$	3550	120	3434	10	-1.0	U					65Sc19 *
	3440	100			-0.1	U					66Da06 *
$^{158}\text{Tb}(\epsilon)^{158}\text{Gd}$	1237.542	0.018	1218.9	1.0	*****	F					83Ra25 *
	1220	13			-0.1	U					87Br33
	1222.1	3.			-1.1	U					85Vo13 *
$^{158}\text{Tb}(\beta^-)^{158}\text{Dy}$	952	10	936.7	2.5	-1.5	U					68Sc04 *
	933	6			0.6	1	17	14 $^{158}\text{Dy}$			85Vo03 *
$^{158}\text{Ho}(\beta^+)^{158}\text{Dy}$	4350	100	4220	27	-1.3	U					61Bo24 *
	4230	30			-0.3	2					68Ab14 *
$^{158}\text{Er}(\beta^+)^{158}\text{Ho}$	1710	40	880	40	-20.7	F					82Vy06 *
$^{158}\text{Tm}(\beta^+)^{158}\text{Er}$	6530	100	6600	30	0.7	-			IRS		93A103
	6624	60			-0.4	-			Dbn		94Po26 *
	ave.	6600	50		0.0	1	37	19 $^{158}\text{Er}$			average
$^{158}\text{Lu}(\epsilon)^{158}\text{Yb}$	8960	200	8798	17	-0.8	U					95Ga.A
* $^{158}\text{Pm}-u$	Represents frequency ratio $^{158}\text{Pm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.96280782(15)$										WgM124**
* $^{158}\text{Sm}-u$	Represents frequency ratio $^{158}\text{Sm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.962848141(58)$										WgM124**
* $^{158}\text{Eu}-u$	Represents frequency ratio $^{158}\text{Eu}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.96286130(15)$										WgM124**
* $^{158}\text{Ho}-u$	$M-A=66148(29)$ keV for mixture gs+m+n at 67.199 and 180#70 keV										Nub16b **
* $^{158}\text{W}(\alpha)^{154}\text{Hf}$	Original value $E=6450(30)$ ( $Q=6617.8$ ) recalibrated to $E=6433(30)$ keV										89Ho12 **
* $^{158}\text{Tb}(p,d)^{157}\text{Tb}$	$Q-Q(^{158}\text{Gd}(p,d))=1152.5(4.2)$ keV										AHW **
* $^{158}\text{Eu}(\beta^-)^{158}\text{Gd}$	$E_{\beta^-}=2520(120)$ 2430(100) respectively, to $2^-$ level at 1023.6974 keV										Ens043 **
*	and $3^-$ level at 1041.6376 keV, and other $E_{\beta^-}$										Ens043 **
* $^{158}\text{Tb}(\epsilon)^{158}\text{Gd}$	$pK=0.00009(2)$ to $2^+$ level at 1187.143, recalculated $Q$										Ens043 **
*	$F : pK < 0.00002$										87Br33 **
* $^{158}\text{Tb}(\epsilon)^{158}\text{Gd}$	$pL=0.689(0.01)$ to $2^+$ level at 1187.143 keV, recalculated $Q$										Ens043 **
* $^{158}\text{Tb}(\beta^-)^{158}\text{Dy}$	$E_{\beta^-}=853(10)$ 834(6) respectively, to $2^+$ level at 98.9180 keV										Ens043 **
* $^{158}\text{Ho}(\beta^+)^{158}\text{Dy}$	$E_{\beta^+}=780(80)$ to 2436-2605 levels; originally assigned to $^{158}\text{Er}(\beta^+)$ ;										Ens043 **
*	reinterpreted by evaluator										AHW **
* $^{158}\text{Ho}(\beta^+)^{158}\text{Dy}$	$E_{\beta^+}=2890(20)$ , 700(60) to 317.139-637.712 and 2436.52-2605.96 levels, and										Ens043 **
*	$E_{\beta^+}=1300(30)$ , 1850(25) keV from $^{158}\text{Ho}^m$ at 67.199 to 1920.43-1940.75										Nub16b **
*	and 1441.75 levels; $E_{\beta^+}=700(60)$ was originally assigned to $^{158}\text{Er}(\beta^+)$ ;										68Ab14 **
*	reinterpreted by evaluator										AHW **
* $^{158}\text{Er}(\beta^+)^{158}\text{Ho}$	$p^+ = 0.3(0.1)$ from annih. $\gamma$ coinc. to 146.90 level										96Go06 **
*	$F : Q_i(1550)$ from upper limit on $p^+$										75Bu.A **
* $^{158}\text{Tm}(\beta^+)^{158}\text{Er}$	$E_{\beta^+}=5410(60)$ to $2^+$ level at 192.15 keV										Ens07a **
$^{159}\text{Pm}-u$	-60715	18	-60714	11	0.1	1	36	36 $^{159}\text{Pm}$	CP1	1.0	12Va02 *
$^{159}\text{Pm}-^{80}\text{Kr}_{1.988}$	105529	19	105527	11	-0.1	1	32	32 $^{159}\text{Pm}$	CP1	1.0	12Va02
$^{159}\text{Pm}-^{86}\text{Kr}_{1.849}$	104567	19	104567	11	0.0	1	32	32 $^{159}\text{Pm}$	CP1	1.0	12Va02
$^{159}\text{Sm}-u$	-66784	11	-66783	6	0.1	1	34	34 $^{159}\text{Sm}$	CP1	1.0	12Va02 *
$^{159}\text{Sm}-^{80}\text{Kr}_{1.988}$	99459	11	99458	6	-0.1	1	34	33 $^{159}\text{Sm}$	CP1	1.0	12Va02
$^{159}\text{Sm}-^{86}\text{Kr}_{1.849}$	98498	11	98498	6	0.0	1	34	34 $^{159}\text{Sm}$	CP1	1.0	12Va02
$^{159}\text{Eu}-u$	-70899	10	-70900	5	-0.1	1	22	22 $^{159}\text{Eu}$	CP1	1.0	12Va02 *
$^{159}\text{Eu}-^{80}\text{Kr}_{1.988}$	95344	10	95340	5	-0.4	1	23	21 $^{159}\text{Eu}$	CP1	1.0	12Va02
$^{159}\text{Eu}-^{86}\text{Kr}_{1.849}$	94382	10	94381	5	-0.1	1	22	22 $^{159}\text{Eu}$	CP1	1.0	12Va02
$\text{C}_9 \text{ }^{13}\text{C} \text{ H}_6 \text{ O}_2 - ^{159}\text{Tb}$	114840	50	114780.3	1.3	-0.3	U			R04	4.0	64De15
$\text{C}_{10} \text{ H}_7 \text{ O}_2 - ^{159}\text{Tb}$	119238	25	119250.5	1.3	0.1	U			R04	4.0	64De15

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{159}\text{Dy}-u$	-74285	30	-74254.0	1.6	1.0	U			GS2	1.0	05Li24
$^{159}\text{Ho}-u$	-72365	71	-72281	3	1.2	U			GS2	1.0	05Li24 *
$^{159}\text{Er}-u$	-69290	30	-69309	4	-0.6	U			GS2	1.0	05Li24
$^{159}\text{Tm}-u$	-65025	30				2			GS2	1.0	05Li24
$^{159}\text{Yb}-^{142}\text{Sm}_{1.120}$	35035	24	35026	19	-0.4	2			MA7	1.0	01Bo59
$^{159}\text{Yb}-u$	-59960	30	-59945	19	0.5	R			GS2	1.0	05Li24
$^{159}\text{Lu}-u$	-53420	61	-53360	40	0.9	2			GS2	1.0	05Li24 *
$^{159}\text{Hf}-u$	-46044	32	-46004	18	1.2	R			GS2	1.0	05Li24
$^{159}\text{Tb } ^{35}\text{Cl}_2-^{155}\text{Gd } ^{37}\text{Cl}_2$	8625.64	1.03	8624.4	0.8	-0.5	1	11	7 $^{159}\text{Tb}$	H41	2.5	85Dy04
$^{159}\text{Tb } ^{35}\text{Cl}-^{157}\text{Gd } ^{37}\text{Cl}$	4333.3	1.2	4336.2	0.8	1.0	U			H25	2.5	72Ba08
	4337.01	0.61			-0.5	1	28	18 $^{159}\text{Tb}$	H41	2.5	85Dy04
$^{159}\text{Lu}(\alpha)^{155}\text{Tm}$	4534.3	10.	4490	40	-0.8	R			IRa		80Al14
	4531.3	10.			-0.8	R					92Ha10
$^{159}\text{Hf}(\alpha)^{155}\text{Yb}$	5221.2	10.	5225.1	2.7	0.4	U					73Ea01 Z
	5226.2	5.			-0.2	4			GSa		79Ho10 Z
	5223.0	5.			0.4	4			ORa		83To01 Z
	5219.6	6.			0.9	4					92Ha10
	5229.8	5.			-0.9	4			Daa		96Pa01
$^{159}\text{Ta}(\alpha)^{155}\text{Lu}^m$	5658.6	5.	5660	7	0.2	o			Daa		96Pa01 *
	5661.7	5.			-0.4	o			Ara		97Da07 *
$^{159}\text{Ta}^m(\alpha)^{155}\text{Lu}$	5745.8	6.	5745	3	-0.2	4			GSa		79Ho10
	5743.8	5.			0.2	4			Daa		96Pa01
	5744.8	5.			0.0	4			Ara		97Da07
$^{159}\text{W}(\alpha)^{155}\text{Hf}$	6444.5	6.	6450	4	1.0	3			GSa		81Ho10 *
	6440.3	5.1			2.0	o			Daa		92Pa05
	6454.7	5.			-0.8	3			Daa		96Pa01
$^{159}\text{Re}^m(\alpha)^{155}\text{Ta}$	6951.2	26.7	6969	23	0.7	R			Daa		07Pa27
$^{157}\text{Gd}(t,p)^{159}\text{Gd}$	5398.9	2.3	5398.80	0.11	0.0	U			McM		89Lo07
$^{158}\text{Gd}(n,\gamma)^{159}\text{Gd}$	5942	1	5943.21	0.08	1.2	U					71Gr42
	5943.07	0.15			0.9	-			ILn		87Sp.A Z
	5943.1	0.2			0.5	-			Dbn		03Gr13
	5943.32	0.12			-0.9	-			BNn		03Gr27
$^{158}\text{Gd}(d,p)^{159}\text{Gd}$	3717	10	3718.64	0.08	0.2	U			Kop		67Tj01
$^{158}\text{Gd}(n,\gamma)^{159}\text{Gd}$	ave. 5943.20	0.08	5943.21	0.08	0.1	1	100	91 $^{159}\text{Gd}$			average
$^{158}\text{Gd}(\alpha,t)^{159}\text{Tb}$	-13686.6	10.	-13682.1	0.8	0.5	U			McM		75Bu02
$^{158}\text{Gd}(\alpha,t)^{159}\text{Tb}-^{164}\text{Dy}()^{165}\text{Ho}$	-85.7	2.2	-88.2	1.1	-1.1	1	23	10 $^{165}\text{Ho}$	McM		84Bu14
$^{159}\text{Tb}(\gamma,n)^{158}\text{Tb}$	-8141	39	-8133.0	0.6	0.2	U			Phi		60Ge01
$^{159}\text{Tb}(d,t)^{158}\text{Tb}$	-1870	15	-1875.8	0.6	-0.4	U			Tal		70Jo22
$^{159}\text{Tb}(d,t)^{158}\text{Tb}-^{164}\text{Dy}()^{163}\text{Dy}$	-474.3	1.0	-474.9	0.6	-0.6	1	41	39 $^{158}\text{Tb}$	McM		84Bu14
$^{158}\text{Dy}(d,p)^{159}\text{Dy}$	4608	10	4606.5	2.6	-0.1	U			Tal		68Be.A
	4600	10			0.7	U			Kop		70Gr46
$^{159}\text{Re}^m(p)^{158}\text{W}$	1816.4	20.	1809	17	-0.4	4			Dap		06Jo10
$^{159}\text{Pm}(\beta^-)^{159}\text{Sm}$	5460	140	5653	12	1.4	o			Kur		07Ha57
	5430	140			1.6	U			Kur		10Ha.A
$^{159}\text{Sm}(\beta^-)^{159}\text{Eu}$	3840	100	3836	7	0.0	o			Kur		02Sh.A
	3805	65			0.5	o			Kur		07Ha57
	3800	65			0.5	U			Kur		10Ha.A
$^{159}\text{Eu}(\beta^-)^{159}\text{Gd}$	2600	50	2518	4	-1.6	U					65Iw01 *
$^{159}\text{Gd}(\beta^-)^{159}\text{Tb}$	969.0	1.5	970.9	0.8	1.3	1	26	16 $^{159}\text{Tb}$			77Bo.A
$^{159}\text{Dy}(\epsilon)^{159}\text{Tb}$	365.9	1.3	365.2	1.2	-0.5	1	80	62 $^{159}\text{Dy}$			68My.A *
$^{159}\text{Ho}(\beta^+)^{159}\text{Dy}$	1837.6	6.	1837.6	2.7	0.0	2					79Ad08 *
	1837.6	3.			0.0	2					82Vy02 *
$^{159}\text{Er}(\beta^+)^{159}\text{Ho}$	2768.5	2.0				3					84Ka.A *
	2810	100	2768.5	2.0	-0.4	U			IRS		93Al03
$^{159}\text{Tm}(\beta^+)^{159}\text{Er}$	3400	300	3991	28	2.0	U					75St07
	3850	100			1.4	U			IRS		93Al03
	3670	100			3.2	B			Dbn		94Po26

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{159}\text{Yb}(\beta^+)^{159}\text{Tm}$	5050	200	4730	30	-1.6	U			IRS		93Al03
	4554	150			1.2	U			Dbn		94Po26 *
$^{159}\text{Lu}(\beta^+)^{159}\text{Yb}$	5850	150	6130	40	1.9	U			IRS		93Al03
	5803	150			2.2	U			Dbn		94Po26
$^{159}\text{Ta}^m(\text{IT})^{159}\text{Ta}$	63.7	5.2				4			Ara		97Da07
* $^{159}\text{Pm}-u$	Represents frequency ratio $^{159}\text{Pm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.95673359(11)$										WgM124**
* $^{159}\text{Sm}-u$	Represents frequency ratio $^{159}\text{Sm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.956770122(65)$										WgM124**
* $^{159}\text{Eu}-u$	Represents frequency ratio $^{159}\text{Eu}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.956794898(63)$										WgM124**
* $^{159}\text{Ho}-u$	$M - A = -67304(28)$ keV for mixture gs+m at 205.91 keV										Nub16b **
* $^{159}\text{Lu}-u$	$M - A = -49710(28)$ keV for mixture gs+m at 100#80 keV										Nub16b **
* $^{159}\text{Ta}(\alpha)^{155}\text{Lu}^m$	Replaced by $^{155}\text{Lu}^m(\text{IT})$										AHW **
* $^{159}\text{W}(\alpha)^{155}\text{Hf}$	Original value $E_\alpha = 6299(6)$ recalibrated to $E_\alpha = 6282(6)$ keV										89Ho12 **
* $^{159}\text{Eu}(\beta^-)^{159}\text{Gd}$	$E_{\beta^-} = 2350(50)$ to $7/2^-$ level at 227.412 level, and other $E_{\beta^-}$										Ens121 **
* $^{159}\text{Dy}(\epsilon)^{159}\text{Tb}$	From intensity of feeding $5/2^-$ level at 363.5449 keV										Ens121 **
* $^{159}\text{Ho}(\beta^+)^{159}\text{Dy}$	$E_{\beta^+} = 506(6)$ $506(3)$ respectively, to $5/2^-$ level at 309.593 keV										Ens121 **
* $^{159}\text{Er}(\beta^+)^{159}\text{Ho}$	$E_{\beta^+} = 1122(3)$ to $13/2^+$ level at 624.5 keV, and other $E_{\beta^+}$										Ens121 **
* $^{159}\text{Yb}(\beta^+)^{159}\text{Tm}$	$E_{\beta^+} = 3366(150)$ to $7/2^-$ level at 166.17 keV										Ens121 **
$^{160}\text{Sm}-u$	-64666	11	-64665	6	0.1	1	34	34 $^{160}\text{Sm}$	CP1	1.0	12Va02 *
$^{160}\text{Sm}-^{80}\text{Kr}_{2,000}$	102581	11	102579	6	-0.2	1	34	33 $^{160}\text{Sm}$	CP1	1.0	12Va02
$^{160}\text{Sm}-^{86}\text{Kr}_{1,860}$	101599	11	101600	6	0.0	1	34	34 $^{160}\text{Sm}$	CP1	1.0	12Va02
$^{160}\text{Eu}-u$	-68150	17	-68149	10	0.1	1	36	36 $^{160}\text{Eu}$	CP1	1.0	12Va02 *
$^{160}\text{Eu}-^{80}\text{Kr}_{2,000}$	99096	18	99095	10	-0.1	1	32	32 $^{160}\text{Eu}$	CP1	1.0	12Va02
$^{160}\text{Eu}-^{86}\text{Kr}_{1,860}$	98115	18	98115	10	0.0	1	32	32 $^{160}\text{Eu}$	CP1	1.0	12Va02
$\text{C}_{12}\text{H}_{16}-^{160}\text{Gd}$	198150	50	198139.0	1.4	-0.1	U			R04	4.0	64De15
$\text{C}_{12}\text{H}_{16}-^{160}\text{Dy}$	200050	70	199997.3	0.8	-0.2	U			R04	4.0	64De15
$^{160}\text{Er}-u$	-70916	30	-70923	26	-0.2	-			GS2	1.0	05Li24
	ave.	-70914	27		-0.3	1	95	95 $^{160}\text{Er}$			average
$^{160}\text{Tm}-u$	-64773	127	-64740	40	0.3	U			GS1	1.0	00Ra23 *
	-64755	39			0.5	1	89	89 $^{160}\text{Tm}$	GS2	1.0	05Li24 *
$^{160}\text{Yb}-^{142}\text{Sm}_{1,127}$	33120	20	33124	8	0.2	1	18	15 $^{160}\text{Yb}$	MA7	1.0	01Bo59
$^{160}\text{Yb}-u$	-62440	120	-62440	8	0.0	U			GS1	1.0	00Ra23
	-62438	30			-0.1	U			GS2	1.0	05Li24
$^{160}\text{Yb}-^{133}\text{Cs}_{1,203}$	51301.8	8.4	51301	8	-0.1	1	85	85 $^{160}\text{Yb}$	MA8	1.0	17Ma.A
$^{160}\text{Lu}-u$	-53967	61				2			GS2	1.0	05Li24 *
$^{160}\text{Hf}-u$	-49334	30	-49317	10	0.6	U			GS2	1.0	05Li24
$^{160}\text{Gd}\ ^{35}\text{Cl}_2-^{156}\text{Gd}\ ^{37}\text{Cl}_2$	10831.70	1.27	10831.2	0.8	-0.2	U			H41	2.5	85Dy04
$^{160}\text{Gd}\ ^{35}\text{Cl}-^{158}\text{Gd}\ ^{37}\text{Cl}$	5890	5	5900.0	0.8	0.5	U			H12	4.0	64Ba15
	5899	3			0.1	U			H21	2.5	70Ma05
	5900.0	0.5			0.0	-			M21	2.5	75Ka25
	5899.88	0.96			0.1	-			H41	2.5	85Dy04
	ave.	5900.0	1.1		0.0	1	51	35 $^{160}\text{Gd}$			average
$^{160}\text{Dy}\ ^{35}\text{Cl}-^{158}\text{Dy}\ ^{37}\text{Cl}$	3731.8	2.3	3738.8	2.4	1.2	1	18	18 $^{158}\text{Dy}$	H25	2.5	72Ba08
$^{160}\text{Gd}-^{160}\text{Dy}$	1854.5	0.8	1858.3	1.3	1.9	1	41	35 $^{160}\text{Gd}$	H25	2.5	72Ba08
$^{160}\text{Gd}\ \text{O}-\text{C}_{15}$	-78020.1	5.8	-78023.8	1.4	-0.4	o			TG1	1.5	09Ke.A
	-78019.9	3.6			-0.7	U			TG1	1.5	11Ke03
$^{160}\text{Hf}(\alpha)^{156}\text{Yb}$	4892.2	10.	4901.9	2.6	0.9	-					73Ea01 Z
	4905.0	5.			-0.6	-			GSa		79Ho10 Z
	4904.0	5.			-0.4	-			ORa		83To01 Z
	4901.8	6.			0.0	-					92Ha10
	4902.8	10.			-0.1	-					95Hi12
	4900.8	6.			0.2	-			Daa		96Pa01
	ave.	4902.4	2.6		-0.2	1	99	82 $^{160}\text{Hf}$			average
$^{160}\text{Ta}(\alpha)^{156}\text{Lu}$	5449.5	5.	5451	5	0.3	3			Daa		96Pa01
	5456.6	10.			-0.6	3			Jya		09Ha42
$^{160}\text{Ta}^m(\alpha)^{156}\text{Lu}^m$	5550.9	5.	5548.5	3.0	-0.5	4			GSa		79Ho10 Z
	5538.7	6.			1.6	4					92Ha10
	5552.1	5.			-0.7	4			Daa		96Pa01
	5551.0	10.			-0.3	4			Jya		09Ha42
$^{160}\text{W}(\alpha)^{156}\text{Hf}$	6072.1	10.	6066	5	-0.6	-			GSa		79Ho10

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{160}\text{W}(\alpha)^{156}\text{Hf}$		6063.9	5.	6066	5	0.3	–		Daa		96Pa01	
	ave.	6066	5			0.0	1	100	$^{160}\text{W}$		average	
$^{160}\text{Re}(\alpha)^{156}\text{Ta}$		6704.9	16.	6698	4	–0.4	o		Daa		92Pa05	
		6711.1	16.			–0.8	o		Daa		96Pa01	
		6697.7	4.				4		Daa		11Da12	
$^{158}\text{Gd}(\text{t,p})^{160}\text{Gd}$		4912.0	2.2	4913.0	0.7	0.5	U		McM		89Lo07	
$^{160}\text{Gd}(\text{p,t})^{158}\text{Gd}$		–4919	5	–4913.0	0.7	1.2	U		Min		73Oo01	
$^{160}\text{Dy}(\text{p,t})^{158}\text{Dy}$		–6924	5	–6926.2	2.3	–0.4	–		Min		73Oo01	
		–6925.1	3.4			–0.3	–		McM		88Bu08 *	
	ave.	–6924.8	2.8			–0.5	1	64	$^{158}\text{Dy}$		average	
$^{160}\text{Gd}(\text{t},\alpha)^{159}\text{Eu}-^{158}\text{Gd}()^{157}\text{Eu}$		–666	5	–668	4	–0.4	1	69	$^{159}\text{Eu}$	LAI	79Bu05	
$^{160}\text{Gd}(\text{d,t})^{159}\text{Gd}$		–1200	10	–1194.4	0.7	0.6	U		Kop		67Tj01	
$^{159}\text{Tb}(\text{n},\gamma)^{160}\text{Tb}$		6375.45	0.3	6375.21	0.13	–0.8	–				74Ke01 Z	
		6375.13	0.15			0.5	–		Bdn		06Fi.A	
$^{159}\text{Tb}(\text{d,p})^{160}\text{Tb}$		4165	20	4150.65	0.13	–0.7	U		MIT		64Sp12	
		4153	5			–0.5	U		Tal		67St14	
	ave.	6375.19	0.13	6375.21	0.13	0.1	1	99	$^{160}\text{Tb}$		average	
$^{159}\text{Tb}(\text{n},\gamma)^{160}\text{Tb}$		–2339	10	–2319.7	1.4	1.9	U		Tal		68Be.A	
$^{160}\text{Dy}(\text{d,t})^{159}\text{Dy}$		–2323	10			0.3	U		Kop		70Gr46	
$^{160}\text{Re}(\text{p})^{159}\text{W}$		1269.1	6.	1267	7	–0.3	o		Dap		92Pa05	
		1271	9			–0.4	o		Dap		96Pa01 *	
		1272.2	6.			–0.9	R		Dap		11Da12	
$^{160}\text{Eu}(\beta^-)^{160}\text{Gd}$		3900	300	4461	10	1.9	U				73Da05	
		4200	200			1.3	U				73Mo18	
		4705	60			–4.1	B		Kur		07Ha57	
		4695	60			–3.9	C		Kur		10Ha.A	
		4480	35			–0.5	U		Kur		14Ha38	
$^{160}\text{Tb}(\beta^-)^{160}\text{Dy}$		1838	10	1836.5	1.2	–0.2	U				57Na03 *	
		1827	10			0.9	U				59Gr93 *	
		1825	10			1.1	U				63Wu01 *	
$^{160}\text{Ho}(\beta^+)^{160}\text{Dy}$		3290	15				2				66Av03 *	
$^{160}\text{Er}(\epsilon)^{160}\text{Ho}$		420	150	319	29	–0.7	U				82Vy06 *	
$^{160}\text{Tm}(\beta^+)^{160}\text{Er}$		5600	300	5760	40	0.5	U				75St12 *	
		5890	100			–1.3	1	16	$^{160}\text{Tm}$	IRS	93A103	
$^{160}\text{Lu}(\beta^+)^{160}\text{Yb}$		7210	240	7890	60	2.8	U				83Ge08	
		7340	100			5.5	C		IRS		83Vi.A	
		7300	100			5.9	B		IRS		93A103	
* $^{160}\text{Sm}-\text{u}$	Represents frequency ratio $^{160}\text{Sm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.950775181(67)$										WgM124**	
* $^{160}\text{Eu}-\text{u}$	Represents frequency ratio $^{160}\text{Eu}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.95079589(10)$										WgM124**	
* $^{160}\text{Tm}-\text{u}$	$M - A = -60300(110)$ keV for mixture gs+m at 70(20) keV										Nub16b **	
* $^{160}\text{Tm}-\text{u}$	$M - A = -60283(28)$ keV for mixture gs+m at 70(20) keV										Nub16b **	
* $^{160}\text{Lu}-\text{u}$	$M - A = -50270(28)$ keV for mixture gs+m at 0#100 keV										Nub16b **	
* $^{160}\text{Dy}(\text{p,t})^{158}\text{Dy}$	$Q - Q(^{164}\text{Dy}(\text{p,t})) = -1477.9(3.4)$ , see $^{164}\text{Dy}(\text{p,t})$										AHW **	
* $^{160}\text{Re}(\text{p})^{159}\text{W}$	Corrected : Ame2003 assumed $E_p = 1271(9)$ thus $Q_p = 1279.1(9.)$ keV										WgM105**	
* $^{160}\text{Tb}(\beta^-)^{160}\text{Dy}$	$E_{\beta^-} = 870(10)$ 858(10) 868(10) respectively, to $8^+$ level at 966.85 keV, and other $E_{\beta^-}$										Ens059 **	
* $^{160}\text{Ho}(\beta^+)^{160}\text{Dy}$	$E_{\beta^+} = 570(15)$ to $4^+$ level at 1694.37 keV; and 1045(15) from										Ens059 **	
*	$^{160}\text{Ho}^m$ at 59.98 to $1^-$ level at 1285.602 and $3^-$ at 1286.711 keV										Nub16b **	
* $^{160}\text{Er}(\epsilon)^{160}\text{Ho}$	pK=0.795(0.2) to $1^+$ level at 67.11 keV										Ens059 **	
* $^{160}\text{Tm}(\beta^+)^{160}\text{Er}$	$E_{\beta^+} = 3700(300)$ to 854.4–1007.95 levels, reassigned by evaluator										Ens059 **	
$^{161}\text{Sm}-\text{u}$	–60841	13	–60840	7	0.1	1	32	32	$^{161}\text{Sm}$	CP1	1.0	12Va02 *
$^{161}\text{Sm}-^{80}\text{Kr}_{2.013}$	107493	12	107491	7	–0.2	1	38	37	$^{161}\text{Sm}$	CP1	1.0	12Va02
$^{161}\text{Sm}-^{86}\text{Kr}_{1.872}$	106496	13	106497	7	0.1	1	32	32	$^{161}\text{Sm}$	CP1	1.0	12Va02
$^{161}\text{Eu}-\text{u}$	–66336	19	–66336	11	0.0	1	35	35	$^{161}\text{Eu}$	CP1	1.0	12Va02 *
$^{161}\text{Eu}-^{80}\text{Kr}_{2.013}$	101996	19	101995	11	0.0	1	35	34	$^{161}\text{Eu}$	CP1	1.0	12Va02

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference		
$^{161}\text{Eu}-^{86}\text{Kr}_{1.872}$	101000	20	101001	11	0.0	1	31	31	$^{161}\text{Eu}$	CP1	1.0	12Va02	
$\text{C}_{13}\text{H}_5-^{161}\text{Dy}$	112246	25	112186.1	0.8	-0.6	U			R04	4.0	64De15		
$^{161}\text{Tm}-u$	-66451	30				2			GS2	1.0	05Li24	*	
$^{161}\text{Yb}-^{142}\text{Sm}_{1.134}$	34071	19	34065	16	-0.3	2			MA7	1.0	01Bo59		
$^{161}\text{Yb}-u$	-62120	110	-62093	16	0.2	U			GS1	1.0	00Ra23		
	-62107	30			0.5	R			GS2	1.0	05Li24		
$^{161}\text{Lu}-u$	-56428	30				2			GS2	1.0	05Li24		
$^{161}\text{Hf}-u$	-49733	30	-49721	24	0.4	1	65	65	$^{161}\text{Hf}$	GS2	1.0	05Li24	
$^{161}\text{Dy}\ ^{35}\text{Cl}-^{159}\text{Tb}\ ^{37}\text{Cl}$	4535.0	1.0	4535.3	1.3	0.1	1	25	22	$^{159}\text{Tb}$	H25	2.5	72Ba08	
$^{161}\text{Hf}(\alpha)^{157}\text{Yb}$	4717.0	10.	4682	24	-0.7	-						73Ea01	Z
	4725.2	10.			-0.9	-						82Sc15	Z
	4724.2	5.			-0.8	-			ORa			83To01	Z
	4716.4	7.			-0.7	-						92Ha10	
	4721.5	10.			-0.8	-						95Hi12	
	ave.	4721	3		-0.8	1	23	19	$^{161}\text{Hf}$			average	
$^{161}\text{Ta}^m(\alpha)^{157}\text{Lu}^m$	5278.9	5.	5277	6	-0.5	F			GSa			79Ho10	*
	5280.4	5.			-0.8	F						92Ha10	
	5271.2	7.			0.8	F			Daa			96Pa01	*
	5282.5	7.			-0.9	F			Jya			05Sc22	*
	5273.2	6.			0.5	1	94	56	$^{161}\text{Ta}^m$	Jya		12Th13	
$^{161}\text{W}(\alpha)^{157}\text{Hf}$	5923.4	5.	5923	4	-0.1	4			GSa			79Ho10	Z
	5922.4	5.			0.1	4			Daa			96Pa01	
$^{161}\text{Re}^m(\alpha)^{157}\text{Ta}^m$	6439.3	10.	6430	4	-0.9	2			GSa			79Ho10	
	6425.0	6.			0.8	2			Daa			96Pa01	
	6432.1	7.			-0.3	2			Ara			97Ir01	
$^{161}\text{Os}(\alpha)^{157}\text{W}$	7065.9	12.				3						10Bi03	
$^{161}\text{Os}(\alpha)^{157}\text{W}^p$	6748.0	30.				4						10Bi03	
$^{161}\text{Dy}(p,t)^{159}\text{Dy}$	-6546	5	-6549.5	1.4	-0.7	-			Min			73Oo01	
	-6547.9	2.5			-0.6	-			McM			88Bu08	*
	ave.	-6547.5	2.2		-0.9	1	40	38	$^{159}\text{Dy}$			average	
$^{160}\text{Gd}(n,\gamma)^{161}\text{Gd}$	5635.4	1.0				2						71Gr42	
$^{160}\text{Gd}(d,p)^{161}\text{Gd}$	3411	10	3410.8	1.0	0.0	U			Kop			67Tj01	
$^{160}\text{Gd}(\alpha,t)^{161}\text{Tb}-^{158}\text{Gd}()^{159}\text{Tb}$	678.0	1.0	677.0	0.7	-1.0	1	56	27	$^{160}\text{Gd}$	McM		75Bu02	
$^{160}\text{Tb}(n,\gamma)^{161}\text{Tb}$	7696.3	0.6	7696.6	0.6	0.6	1	84	74	$^{161}\text{Tb}$			75He.C	
$^{160}\text{Dy}(n,\gamma)^{161}\text{Dy}$	6451.5	2.	6454.39	0.08	1.4	U						77Be03	
	6454.40	0.09			-0.1	-			ILn			86Sc16	Z
	6454.34	0.14			0.3	-			Bdn			06Fi.A	
$^{160}\text{Dy}(d,p)^{161}\text{Dy}$	4231	10	4229.82	0.08	-0.1	U			Tal			68Be.A	
	4237	10			-0.7	U			Kop			70Gr46	
$^{161}\text{Dy}(d,t)^{160}\text{Dy}$	-205	10	-197.16	0.08	0.8	U			Kop			70Gr46	
$^{160}\text{Dy}(n,\gamma)^{161}\text{Dy}$	ave.	6454.38	0.08	6454.39	0.08	0.1	1	100	94	$^{160}\text{Dy}$		average	
$^{160}\text{Dy}(^3\text{He,d})^{161}\text{Ho}-^{164}\text{Dy}()^{165}\text{Ho}$	-1406.5	2.0	-1406.5	2.0	0.0	1	100	100	$^{161}\text{Ho}$	McM		75Bu02	
$^{161}\text{Re}(p)^{160}\text{W}$	1199.5	6.	1197	5	-0.4	1	79	79	$^{161}\text{Re}$	Ara		97Ir01	
$^{161}\text{Re}^m(p)^{160}\text{W}$	1323.3	7.	1321	5	-0.3	o			Ara			97Ir01	*
$^{161}\text{Sm}(\beta^-)^{161}\text{Eu}$	5065	130	5120	12	0.4	o			Kur			07Ha57	
	5050	130			0.5	U			Kur			10Ha.A	
$^{161}\text{Eu}(\beta^-)^{161}\text{Gd}$	3705	60	3714	11	0.2	o			Kur			07Ha57	
	3705	60			0.2	U			Kur			10Ha.A	
	3722	35			-0.2	U			Kur			14Ha38	
$^{161}\text{Gd}(\beta^-)^{161}\text{Tb}$	1977	30	1955.8	1.4	-0.7	U						66Zy02	*
$^{161}\text{Tb}(\beta^-)^{161}\text{Dy}$	584	6	594.2	1.3	1.7	U						63Ko08	
	590	10			0.4	U						64Fu11	
$^{161}\text{Er}(\beta^+)^{161}\text{Ho}$	2050	40	1996	9	-1.4	U						65Gr35	*
	1980	18			0.9	R						84Ka.A	*
$^{161}\text{Tm}(\beta^+)^{161}\text{Er}$	3100	200	3303	29	1.0	U						75Ad08	*
	3180	100			1.2	U			IRS			93Al03	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{161}\text{Yb}(\beta^+)^{161}\text{Tm}$	3850	250	4060	30	0.8	U					81Ad02	
	3585	200			2.4	U			Dbn		94Po26	
$^{161}\text{Lu}(\beta^+)^{161}\text{Yb}$	5300	100	5280	30	-0.2	o			IRS		83Vi.A	
	5300	100			-0.2	U			IRS		93AI03	
	5255	150			0.1	U			Dbn		94Po26 *	
$^{161}\text{Re}^m(\text{IT})^{161}\text{Re}$	123.8	1.3	123.7	1.3	-0.1	1	99	78	$^{161}\text{Re}^m$		97Ir01	
* $^{161}\text{Sm}-\text{u}$	Represents frequency ratio $^{161}\text{Sm}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.944844874(74)$										WgM124**	
* $^{161}\text{Eu}-\text{u}$	Represents frequency ratio $^{161}\text{Eu}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.94487714(11)$										WgM124**	
* $^{161}\text{Tm}-\text{u}$	$M - A = -61895(28)$ keV for mixture gs+m at 7.51 keV										Nub16b **	
* $^{161}\text{Ta}^m(\alpha)^{157}\text{Lu}^m$	F : above 4 items are not unambiguously assigned										WgM151**	
* $^{161}\text{Dy}(\text{p,t})^{159}\text{Dy}$	$Q - Q(^{164}\text{Dy}(\text{p,t})) = -1100.7(2.5)$ keV										AHW **	
* $^{161}\text{Re}^m(\text{p})^{160}\text{W}$	Replaced by author's result for $^{161}\text{Re}^m(\text{IT})^{161}\text{Re}$										AHW **	
* $^{161}\text{Gd}(\beta^-)^{161}\text{Tb}$	$E_{\beta^-} = 1560(30)$ mainly to $7/2^-$ level at 417.228 keV										Ens11b **	
* $^{161}\text{Er}(\beta^+)^{161}\text{Ho}$	$E_{\beta^+} = 820(40)$ 748(18) respectively, to $1/2^+$ level at 211.15 keV										Ens11b **	
* $^{161}\text{Tm}(\beta^+)^{161}\text{Er}$	$E_{\beta^+} = 1800(100)$ to several levels around $7/2^-$ one at 266.44 keV										Ens11b **	
* $^{161}\text{Lu}(\beta^+)^{161}\text{Yb}$	$E_{\beta^+} = 3866(150)$ to 367.28 level										Ens11b **	
$\text{C}_{13}\text{H}_6-^{162}\text{Dy}$	120115	19	120146.0	0.8	0.4	U			R04	4.0	64De15	
$\text{C}_{12}\text{H}_4\text{N}-^{162}\text{Er}$	105590	70	105587.2	0.9	0.0	U			R04	4.0	64De15	
$\text{C}_{13}\text{H}_6-^{162}\text{Er}$	118430	170	118163.2	0.9	-0.4	U			R04	4.0	64De15	
$^{162}\text{Tm}-\text{u}$	-65942	55	-65999	28	-1.0	R			GS2	1.0	05Li24 *	
$^{162}\text{Yb}-^{142}\text{Sm}_{1.141}$	32524	19	32525	16	0.1	2			MA7	1.0	01Bo59	
$^{162}\text{Yb}-\text{u}$	-64210	110	-64226	16	-0.1	U			GS1	1.0	00Ra23	
	-64223	30			-0.1	R			GS2	1.0	05Li24	
$^{162}\text{Lu}-\text{u}$	-56758	234	-56720	80	0.2	o			GS1	1.0	00Ra23 *	
	-56781	190			0.3	2			GS2	1.0	05Li24 *	
$^{162}\text{Hf}-\text{u}$	-52756	30	-52785	10	-1.0	U			GS2	1.0	05Li24	
$^{162}\text{Er}^{35}\text{Cl}_2-^{158}\text{Gd}^{37}\text{Cl}_2$	10577.5	2.7	10575.5	1.2	-0.3	U			H25	2.5	72Ba08	
$^{162}\text{Dy}^{35}\text{Cl}-^{160}\text{Dy}^{37}\text{Cl}$	4555	6	4551.03	0.12	-0.2	U			H12	4.0	64Ba15	
	4552.1	1.1			-0.4	U			H25	2.5	72Ba08	
$^{162}\text{Er}^{35}\text{Cl}-^{160}\text{Gd}^{37}\text{Cl}$	4674.6	1.9	4675.5	1.3	0.2	U			H25	2.5	72Ba08	
$^{162}\text{Er}-^{162}\text{Dy}$	1982.79	0.32	1982.8	0.3	0.0	1	100	100	$^{162}\text{Er}$	SH1	1.0	11El04
$^{161}\text{Dy}^{37}\text{Cl}-^{162}\text{Dy}^{35}\text{Cl}$	-3080	70	-2815.19	0.09	0.9	U			R04	4.0	64De15	
$^{162}\text{Dy}-^{161}\text{Dy}$	150	70	-134.92	0.06	-1.0	U			R04	4.0	64De15	
	78	23			-2.3	U			R04	4.0	64De15	
	22	40			-1.0	U			R04	4.0	64De15	
$^{162}\text{Hf}(\alpha)^{158}\text{Yb}$	4417.2	10.	4416	5	-0.1	-					82Sc15	
	4420.4	10.3			-0.4	-			ORa		83To01	
	4414.2	9.			0.2	-					92Ha10	
	4416.3	10.3			0.0	-					95Hi12	
ave.	4417	5			-0.1	1	95	81	$^{162}\text{Hf}$		average	
$^{162}\text{Ta}(\alpha)^{158}\text{Lu}$	5003.8	10.	5010	50	0.1	4					86Ru05	
	5007.9	5.			0.0	4					92Ha10	
$^{162}\text{W}(\alpha)^{158}\text{Hf}$	5669.9	10.	5678.3	2.4	0.8	U					73Ea01 Z	
	5668.0	10.			1.0	U			ORa		75To05 Z	
	5677.5	5.			0.2	-			GSa		81Ho10 Z	
	5674.5	4.			0.9	-			Ora		82De11 Z	
	5681.6	5.			-0.6	-			Daa		96Pa01	
	5681.5	5.1			-0.6	-			Jya		15Li24	
ave.	5678.3	2.4			0.0	1	100	100	$^{162}\text{W}$		average	
$^{162}\text{Re}(\alpha)^{158}\text{Ta}$	6240.3	5.				8			Ara		97Da07	
$^{162}\text{Re}^m(\alpha)^{158}\text{Ta}^m$	6274.2	6.	6274	3	0.0	9			GSa		79Ho10	
	6278.3	6.			-0.7	9			Daa		96Pa01	
	6271.1	5.			0.6	9			Ara		97Da07	
	6256	16			1.1	U			Jya		16Ca15 *	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{162}\text{Os}(\alpha)^{158}\text{W}$	6778.8	30.	6767	3	-0.4	U			GSa		89Ho12 *	
	6785.8	10.			-1.8	U			ORa		96Bi07	
	6767.4	3.				4			Ara		00Ma95	
	6781.7	13.			-1.1	U			Jya		04Jo12	
$^{160}\text{Gd}(\text{t,p})^{162}\text{Gd}$	3999.5	3.8				2			McM		89Lo07	
$^{160}\text{Dy}(\text{t,p})^{162}\text{Dy}$	6169.5	1.9	6169.59	0.10	0.0	U			McM		88Bu08 *	
$^{162}\text{Dy}(\text{p,t})^{160}\text{Dy}$	-6168	5	-6169.59	0.10	-0.3	U			Min		73Oo01	
	-6169.7	2.1			0.1	U			McM		88Bu08 *	
$^{162}\text{Er}(\text{p,t})^{160}\text{Er}$	-7944	55	-7931	24	0.2	R			Win		74De31 *	
$^{161}\text{Dy}(\text{n},\gamma)^{162}\text{Dy}$	8196.99	0.06	8196.99	0.06	0.1	1	100	88	$^{161}\text{Dy}$	MMn	82Is05 Z	
	8193	3			1.3	U			Bdn		06Fi.A	
$^{161}\text{Dy}(\text{d,p})^{162}\text{Dy}$	5969	10	5972.43	0.06	0.3	U			Tal		67Ba34	
	5981	10			-0.9	U			Kop		70Gr46	
$^{162}\text{Dy}(\text{d,t})^{161}\text{Dy}$	-1944	10	-1939.77	0.06	0.4	U			Kop		70Gr46	
	-1943	10			0.3	U			Tal		77Be03	
$^{161}\text{Dy}(\text{}^3\text{He,d})^{162}\text{Ho}-^{164}\text{Dy}()^{165}\text{Ho}$	-945.3	3.0	-945.3	3.0	0.0	1	100	100	$^{162}\text{Ho}$	McM	75Bu02	
$^{162}\text{Er}(\text{d,t})^{161}\text{Er}$	-2952	10	-2947	9	0.5	2			Kop		69Tj01	
$^{162}\text{Eu}(\beta^-)^{162}\text{Gd}$	5575	60	5580	40	0.0	o			Kur		07Ha57	
	5585	60			-0.1	o			Kur		10Ha.A	
	5577	35				3			Kur		14Ha38	
$^{162}\text{Gd}(\beta^-)^{162}\text{Tb}$	1442	100	1400	40	-0.5	R					70Ch02 *	
$^{162}\text{Tb}(\beta^-)^{162}\text{Dy}$	2448	100	2510	40	0.6	2					66Fu08 *	
	2523	50			-0.3	2					66Sc24 *	
	2528	80			-0.3	2					77Ka08 *	
	2220	50	2140	3	-1.6	U					69Ak01	
$^{162}\text{Ho}(\beta^+)^{162}\text{Dy}$	4840	50	4857	26	0.3	2					63Ab02	
$^{162}\text{Tm}(\beta^+)^{162}\text{Er}$	4705	70			2.2	2					74De47 *	
	4900	100			-0.4	2			IRS		93Al03	
	4892	50			-0.7	2			Dbn		94Po26 *	
	6740	270	6990	80	0.9	U					83Ge08	
	6990	120			0.0	o			IRS		83Vi.A	
$^{162}\text{Lu}(\beta^+)^{162}\text{Yb}$	6960	100			0.3	R			IRS		93Al03	
	7111	150			-0.8	R			Dbn		94Po26 *	
	$^{162}\text{Tm}-\text{u}$ $M-A=-61359(28)$ keV for mixture gs+m at 130(40) keV											
	$^{162}\text{Lu}-\text{u}$ $M-A=-52730(130)$ keV for mixture gs+m+n at 120#200 and 300#200 keV											
$^{162}\text{Lu}-\text{u}$ $M-A=-52751(28)$ keV for mixture gs+m+n at 120#200 and 300#200 keV												
$^{162}\text{Re}^m(\alpha)^{158}\text{Ta}^m$ $E_{\alpha}=6037(16)$ keV to $10^+$ level 66 keV above the $9^+ ^{158}\text{Ta}^m$												
$^{162}\text{Os}(\alpha)^{158}\text{W}$ Original value $E=6640(20)$ ( $Q=6808.4$ ) recalibrated												
$^{160}\text{Dy}(\text{t,p})^{162}\text{Dy}$ $Q-Q(^{162}\text{Dy}(\text{t,p}))=722.3(1.9)$ keV												
$^{162}\text{Dy}(\text{p,t})^{160}\text{Dy}$ $Q-Q(^{164}\text{Dy}(\text{p,t}))=-722.5(2.1)$ keV												
$^{162}\text{Er}(\text{p,t})^{160}\text{Er}$ Not resolved peak. Original uncertainty 28 increased to 51 keV and added systematic error 21 keV												
$^{162}\text{Gd}(\beta^-)^{162}\text{Tb}$ $E_{\beta^-}=1000(100)$ to $1^+$ level at 442.11 keV												
$^{162}\text{Tb}(\beta^-)^{162}\text{Dy}$ $E_{\beta^-}=1300(100)$ 1375(50) 1380(80) respectively, to $2^-$ level at 1148.232 keV												
$^{162}\text{Tm}(\beta^+)^{162}\text{Er}$ $E_{\beta^+}=2110(70)$ to $2^-$ level at 1572.84 keV												
$^{162}\text{Tm}(\beta^+)^{162}\text{Er}$ $E_{\beta^+}=3768(50)$ to $2^+$ level at 102.04 keV												
$^{162}\text{Lu}(\beta^+)^{162}\text{Yb}$ $E_{\beta^+}=6006(150)$ to ground state and $2^+$ level at 166.8, unknown intensity ratio												
$^{163}\text{Gd}-\text{u}$	-65824	16	-65823	9	0.1	1	32	32	$^{163}\text{Gd}$	CP1	1.0	12Va02 *
$^{163}\text{Gd}-^{80}\text{Kr}_{2,038}$	104600	16	104598	9	-0.1	1	32	32	$^{163}\text{Gd}$	CP1	1.0	12Va02
$^{163}\text{Gd}-^{86}\text{Kr}_{1,895}$	103569	15	103570	9	0.0	1	36	36	$^{163}\text{Gd}$	CP1	1.0	12Va02
$\text{C}_{13}\text{H}_7-^{163}\text{Dy}$	125906	36	126038.3	0.8	0.9	U			R04	4.0	64De15	
$^{163}\text{Tm}-\text{u}$	-67327	30	-67342	6	-0.5	U			GS2	1.0	05Li24	
$^{163}\text{Yb}-^{142}\text{Sm}_{1,148}$	33686	19	33685	16	-0.1	2			MA7	1.0	01Bo59	
$^{163}\text{Yb}-\text{u}$	-63663	30	-63660	16	0.1	R			GS2	1.0	05Li24	



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{163}\text{Lu}-u$	-58730	110	-58820	30	-0.8	U			GS1	1.0	00Ra23	
	-58821	30				2			GS2	1.0	05Li24	
$^{163}\text{Hf}-u$	-52911	30	-52887	27	0.8	1	79	79	$^{163}\text{Hf}$	GS2	1.0	05Li24
$^{163}\text{Ta}-u$	-45855	54	-45660	40	3.6	C			GS2	1.0	05Li24	
$^{163}\text{Dy } ^{35}\text{Cl}-^{161}\text{Dy } ^{37}\text{Cl}$	5200	60	4747.90	0.11	-1.9	U			R04	4.0	64De15	
	4746	3			0.3	U			H23	2.5	70Wh01	
	4744.7	1.2			1.1	U			H25	2.5	72Ba08	
$^{163}\text{Ho O}-^{163}\text{Dy O}$	2.72	0.77	3.043	0.020	0.3	U			TG1	1.5	15Sc13	
$^{163}\text{Ho}-^{163}\text{Dy}$	3.042	0.037	3.043	0.020	0.0	1	31	17	$^{163}\text{Ho}$	SH2	1.0	15E103
$^{162}\text{Dy } ^{37}\text{Cl}-^{163}\text{Dy } ^{35}\text{Cl}$	-5069	42	-4882.82	0.08	1.1	U			R04	4.0	64De15	
$^{163}\text{Dy}-^{162}\text{Dy}$	2164	35	1932.71	0.05	-1.7	U			R04	4.0	64De15	
	1985	38			-0.3	U			R04	4.0	64De15	
	2174	40			-1.5	U			R04	4.0	64De15	
$^{163}\text{Dy O}-\text{C}_{15}$	-76349.06	0.86	-76348.5	0.8	0.4	1	41	41	$^{163}\text{Dy}$	TG1	1.5	15Sc13
$^{163}\text{Ho O}-\text{C}_{15}$	-76346.61	0.97	-76345.5	0.8	0.8	1	32	32	$^{163}\text{Ho}$	TG1	1.5	15Sc13
$^{163}\text{Ta}(\alpha)^{159}\text{Lu}$	4741.5	15.	4749	5	0.5	3					83Sc18	*
	4746.7	10.			0.2	3					86Ru05	
	4751.8	7.			-0.4	3					92Ha10	
	5520.3	5.	5520	50	0.0	5					73Ea01	Z
$^{163}\text{W}(\alpha)^{159}\text{Hf}$	5518.1	5.			0.0	5			GSa		79Ho10	Z
	5519.9	3.			0.0	5			Ora		82De11	Z
	5525.9	10.3			-0.1	U					84Sc06	*
	5518.7	6.			0.0	5			Daa		96Pa01	
	6017.9	5.	6012	8	-1.2	o			Ara		97Da07	*
$^{163}\text{Re}(\alpha)^{159}\text{Ta}$	6067.2	6.	6068	3	0.2	3			GSa		79Ho10	
	6067.2	7.			0.1	3			Daa		96Pa01	
	6069.2	5.			-0.2	3			Ara		97Da07	
$^{163}\text{Os}(\alpha)^{159}\text{W}$	6674.1	30.	6677	8	0.1	4			GSa		81Ho10	
	6678.2	10.			-0.1	4			ORa		96Bi07	
	6676.2	19.			0.1	4			Daa		96Pa01	
	6674.1	30.			0.1	4			Jya		13Dr06	*
	5986.3	1.5	5986.21	0.08	-0.1	U			McM		88Bu08	*
$^{161}\text{Dy}(\text{t,p})^{163}\text{Dy}$	-5985	5	-5986.21	0.08	-0.2	U			Min		73Oo01	
	-5987.1	2.2			0.4	U			McM		88Bu08	*
$^{162}\text{Dy}(\text{n},\gamma)^{163}\text{Dy}$	6270.98	0.06	6271.01	0.05	0.5	-			MMn		82Is05	Z
	6271.00	0.09			0.1	-			ILn		89Sc31	Z
	6271.14	0.13			-1.0	-			Bdn		06Fi.A	
$^{163}\text{Dy}(\gamma,\text{n})^{162}\text{Dy}$	-6320	110	-6271.01	0.05	0.4	U			Phi		60Ge01	
$^{162}\text{Dy}(\text{d,p})^{163}\text{Dy}$	4049	5	4046.44	0.05	-0.5	U			Tal		67Sc05	
	4045	10			0.1	U			Kop		70Gr46	
	-14	5	-13.78	0.05	0.0	U					67Ba34	
$^{163}\text{Dy}(\text{d,t})^{162}\text{Dy}$	-27	10			1.3	U			Kop		70Gr46	
	ave.	6271.01	0.05	6271.01	0.05	0.1	1	100	100	$^{162}\text{Dy}$		average
$^{162}\text{Dy}(\text{d},\text{p})^{163}\text{Dy}$	-734.3	1.0	-734.2	0.8	0.1	1	68	56	$^{165}\text{Ho}$	McM	75Bu02	
$^{162}\text{Dy}(\text{d},\text{p})^{163}\text{Dy}$	4682	10	4680	5	-0.2	1	21	21	$^{163}\text{Er}$	Kop	69Tj01	
$^{163}\text{Eu}(\beta^-)^{163}\text{Gd}$	4828	70	4830	70	0.0	o			Kur		07Ha57	*
	4813	70			0.2	o			Kur		10Ha.A	*
	4829	65				2			Kur		14Ha38	
$^{163}\text{Gd}(\beta^-)^{163}\text{Tb}$	3170	70	3282	9	1.6	o			Kur		07Ha57	
	3120	70			2.3	o			Kur		10Ha.A	
	3187	40			2.4	U			Kur		14Ha38	
$^{163}\text{Tb}(\beta^-)^{163}\text{Dy}$	1684	50	1785	4	2.0	U					66Fu08	*
	1721	100			0.6	U					71Ka22	*
$^{163}\text{Ho}(\epsilon)^{163}\text{Dy}$	2.83	0.05	2.834	0.019	0.1	-					82An19	*
	2.65	0.20			0.9	U					83Ba32	
	2.84	0.10			-0.1	U					84La.A	*
	2.56	0.05			5.5	B					85Ha12	*
	2.60	0.03			7.8	B					86Ya17	
	2.561	0.020			13.7	B					92Ha15	
	2.54	0.03			9.8	C					93Bo.A	*
	2.71	0.10			1.2	U					94Ya07	
	2.800	0.050			0.7	-					97Ga12	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{163}\text{Ho}(\epsilon)^{163}\text{Dy}$	2.849	0.030	2.834	0.019	-0.5	-					14Ra.1 *	
	ave.	2.835	0.023		0.0	1	69	39	$^{163}\text{Ho}$		average	
$^{163}\text{Er}(\beta^+)^{163}\text{Ho}$	1210	6	1211	5	0.1	1	58	58	$^{163}\text{Er}$		63Pe16	
$^{163}\text{Tm}(\beta^+)^{163}\text{Er}$	2439	3				2					82Vy07 *	
	2360	100	2439	3	0.8	U			IRS		93Al03	
$^{163}\text{Yb}(\beta^+)^{163}\text{Tm}$	3370	100	3430	16	0.6	U					75Ad09 *	
$^{163}\text{Lu}(\beta^+)^{163}\text{Yb}$	4860	170	4510	30	-2.1	U					83Ge08	
	4600	200			-0.5	o			IRS		83Vi.A	
	4600	200			-0.5	U			IRS		93Al03	
$^{163}\text{Re}^m(\text{IT})^{163}\text{Re}$	115.1	4.0	120	5	1.2	o			Ara		97Da07 *	
* $^{163}\text{Gd}-u$	Represents frequency ratio $^{163}\text{Gd}^{++}/(\text{C}_{12}\text{H}_4)^+ = 0.933275822(91)$											
* $^{163}\text{Ta}-u$	$M - A = -42644(28)$ keV for mixture gs+m at 140#(18#) keV											
* $^{163}\text{Ta}(\alpha)^{159}\text{Lu}$	Original assignment to 13 s $^{164}\text{Ta}$ changed to $^{163}\text{Ta}$											
* $^{163}\text{W}(\alpha)^{159}\text{Hf}$	Originally assigned to $^{166}\text{Re}$ , re-assigned in reference											
*	original $E_\alpha = 5372$ recalibrated using their $^{168}\text{Os}-^{170}\text{Os}$ results											
* $^{163}\text{Re}(\alpha)^{159}\text{Ta}$	Replaced by author's value for $^{159}\text{Ta}^m(\text{IT})$											
* $^{163}\text{Os}(\alpha)^{159}\text{W}$	Error not given, estimated by evaluator											
* $^{161}\text{Dy}(\text{t,p})^{163}\text{Dy}$	$Q - Q(^{162}\text{Dy}(\text{t,p})) = -539.1(1.5)$ keV											
* $^{163}\text{Dy}(\text{p,t})^{161}\text{Dy}$	$Q - Q(^{164}\text{Dy}(\text{p,t})) = -539.9(2.2)$ keV											
* $^{163}\text{Eu}(\beta^-)^{163}\text{Gd}$	$E_{\beta^-} = 4690(70)$ to $^{163}\text{Gd}^m$ at 137.8 keV											
* $^{163}\text{Eu}(\beta^-)^{163}\text{Gd}$	$E_{\beta^-} = 4675(70)$ to $^{163}\text{Gd}^m$ at 137.8 keV											
* $^{163}\text{Tb}(\beta^-)^{163}\text{Dy}$	$E_{\beta^-} = 800(50)$ to $1/2^+$ level at 884.2943 keV											
* $^{163}\text{Tb}(\beta^-)^{163}\text{Dy}$	$E_{\beta^-} = 1300(100)$ to $3/2^-$ level at 421.8439 keV											
* $^{163}\text{Ho}(\epsilon)^{163}\text{Dy}$	Original 2.58(0.10) from partial T=40(12) $\times 10^{+3}$ y, re-evaluated by authors											
* $^{163}\text{Ho}(\epsilon)^{163}\text{Dy}$	Original value 2.82(+0.11-0.08)											
* $^{163}\text{Ho}(\epsilon)^{163}\text{Dy}$	Original value 2.60(0.03) corrected to 2.561(0.020) for dynamic effects											
*	error 0.020 is statistical only											
* $^{163}\text{Ho}(\epsilon)^{163}\text{Dy}$	Original $2616 < Q_i(2694 \text{ eV } 68\% \text{ CL for charge } 66^+ Q_{\beta^+}$ ,											
*	corrected to $2511 < Q_{\beta^+} < 2572 \text{ eV } 68\% \text{ CL}$											
* $^{163}\text{Ho}(\epsilon)^{163}\text{Dy}$	"Preliminary $Q_\epsilon = 2.849(0.005)$ "; syst error estimated 0.030 by evaluator											
* $^{163}\text{Tm}(\beta^+)^{163}\text{Er}$	$E_{\beta^+} = 884(3)$ to $1/2^+$ level at 540.56 keV, and other $E_{\beta^+}$											
* $^{163}\text{Yb}(\beta^+)^{163}\text{Tm}$	$E_{\beta^+} = 1400(100)$ to $5/2^-$ level at 947.29 keV											
* $^{163}\text{Re}^m(\text{IT})^{163}\text{Re}$	Redundant with $^{167}\text{Ir}(\alpha)^{163}\text{Re}$ in same paper											
$\text{C}_{13} \text{H}_8 - ^{164}\text{Dy}$	133320	38	133419.8	0.8	0.7	U			R04	4.0	64De15	
$\text{C}_{12} ^{13}\text{C} \text{H}_7 - ^{164}\text{Dy}$	128920	34	128949.6	0.8	0.2	U			R04	4.0	64De15	
$\text{C}_{12} \text{H}_6 \text{N} - ^{164}\text{Er}$	120876	39	120816.8	0.8	-0.4	U			R04	4.0	64De15	
$^{164}\text{Tm}-u$	-66440	30	-66457	26	-0.6	1	76	76	$^{164}\text{Tm}$	GS2	1.0	05Li24 *
$^{164}\text{Yb}-^{142}\text{Sm}_{1.155}$	32429	19	32434	16	0.3	2			MA7	1.0	01Bo59	
$^{164}\text{Yb}-u$	-65690	104	-65505	16	1.8	U			GS1	1.0	00Ra23	
	-65493	30			-0.4	R			GS2	1.0	05Li24	
$^{164}\text{Lu}-u$	-58750	110	-58660	30	0.8	U			GS1	1.0	00Ra23	
	-58661	30				2			GS2	1.0	05Li24	
$^{164}\text{Hf}-u$	-55620	110	-55629	17	-0.1	U			GS1	1.0	00Ra23	
	-55596	30			-1.1	1	32	32	$^{164}\text{Hf}$	GS2	1.0	05Li24
$^{164}\text{Ta}-u$	-46466	30				2			GS2	1.0	05Li24	
$^{164}\text{Dy} ^{35}\text{Cl} - ^{162}\text{Dy} ^{37}\text{Cl}$	5347	5	5326.41	0.11	-1.0	U			H12	4.0	64Ba15	
	5589	19			-3.5	B			R04	4.0	64De15	
	5321	3			0.7	U			H23	2.5	70Wh01	
	5326.5	0.9			0.0	U			H25	2.5	72Ba08	
$^{164}\text{Er} ^{35}\text{Cl} - ^{162}\text{Er} ^{37}\text{Cl}$	3373.3	1.3	3370.5	0.4	-0.8	U			H25	2.5	72Ba08	
$^{164}\text{Er} - ^{164}\text{Dy}$	26.92	0.12	26.92	0.12	0.0	1	100	100	$^{164}\text{Er}$	SH1	1.0	11El08
$^{164}\text{Dy} ^{35}\text{Cl} - ^{161}\text{Dy} ^{37}\text{Cl}$	5610	48	5191.49	0.13	-2.2	U			R04	4.0	64De15	
$^{163}\text{Dy} ^{37}\text{Cl} - ^{164}\text{Dy} ^{35}\text{Cl}$	-3360	50	-3393.70	0.10	-0.2	U			R04	4.0	64De15	
$^{164}\text{Dy} - ^{163}\text{Dy}$	392	48	443.59	0.07	0.3	U			R04	4.0	64De15	
	540	25			-1.0	U			R04	4.0	64De15	
	446	28			0.0	U			R04	4.0	64De15	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{164}\text{Er}-^{162}\text{Er}$	556	48	420.4	0.4	-0.7	U			R04	4.0	64De15	
$^{164}\text{W}(\alpha)^{160}\text{Hf}$	5281.7	5.	5278.3	2.0	-0.7	-					73Ea01 Z	
	5274.7	5.			0.7	-			ORa		75To05 Z	
	5268.7	10.			1.0	U					78Sc26 *	
	5279.0	5.			-0.1	-			GSa		79Ho10	
	5279.2	3.			-0.3	-			Ora		82De11 Z	
	5283.0	8.			-0.6	U					84Sc06 *	
	5277.0	6.			0.2	-			Daa		96Pa01	
	ave.	5278.6	2.0			-0.2	1	99	81 $^{164}\text{W}$			average
	$^{164}\text{Re}(\alpha)^{160}\text{Ta}$	5922.7	10.	5926	5	0.4	4			GSa		79Ho10
		5928.9	7.			-0.4	4			Daa		96Pa01
5924.7		10.			0.2	4			Jya		09Ha42	
$^{164}\text{Re}^m(\alpha)^{160}\text{Ta}^m$	5763.8	10.			5	5		Jya		09Ha42		
$^{164}\text{Os}(\alpha)^{160}\text{W}$	6478.3	20.	6479	5	0.1	U			GSa		81Ho10	
	6473.2	10.			0.6	-			ORa		96Bi07	
	6479.4	7.			0.0	-			Daa		96Pa01	
	ave.	6477	6			0.4	1	80	80 $^{164}\text{Os}$		average	
$^{164}\text{Ir}^m(\alpha)^{160}\text{Re}^m$	7052.3	10.2				6			Jya		14Dr02	
$^{162}\text{Dy}(\text{l,p})^{164}\text{Dy}$	5447.3	1.9	5447.33	0.08	0.0	U			McM		88Bu08 *	
$^{164}\text{Dy}(\text{p,t})^{162}\text{Dy}$	-5450	5	-5447.33	0.08	0.5	U			Min		73Oo01	
$^{164}\text{Er}(\text{p,t})^{162}\text{Er}$	-7262	10	-7269.2	0.3	-0.7	U			Min		73Oo01	
$^{164}\text{Dy}(\text{t},\alpha)^{163}\text{Tb}$	11153	4				2			McM		92Ga15 *	
$^{163}\text{Dy}(\text{n},\gamma)^{164}\text{Dy}$	7658.11	0.07	7658.11	0.07	0.0	1	100	84 $^{164}\text{Dy}$	MMn		82Is05 Z	
	7658.90	0.06			-13.1	C					99Fo.A	
	7655.0	0.9			3.5	C			Bdn		06Fi.A	
$^{163}\text{Dy}(\text{d,p})^{164}\text{Dy}$	5434	5	5433.55	0.07	-0.1	U			Tal		64Sh06	
	5441	10			-0.7	U			Kop		70Gr46	
$^{164}\text{Dy}(\text{d,t})^{163}\text{Dy}$	-1407	10	-1400.88	0.07	0.6	U			Kop		70Gr46	
	-1407	10			0.6	U			Kop		70Gr46	
$^{163}\text{Dy}(\text{}^3\text{He,d})^{164}\text{Ho}-^{164}\text{Dy}(\text{}^1\text{H})^{165}\text{Ho}$	-331.6	1.4	-330.7	1.1	0.6	1	67	67 $^{164}\text{Ho}$	McM		75Bu02 *	
$^{164}\text{Er}(\text{d,t})^{163}\text{Er}$	-2593	10	-2589	5	0.4	1	21	21 $^{163}\text{Er}$	Kop		69Tj01	
$^{164}\text{Ir}^m(\text{p})^{163}\text{Os}$	1828	8	1824	6	-0.5	o			Jyp		01Ke05	
	1818	14			0.4	5			Arp		02Ma61	
	1825	6			-0.2	5			Jyp		14Dr02	
	6430	70	6390	50	-0.5	o			Kur		07Ha57	
$^{164}\text{Eu}(\beta^-)^{164}\text{Gd}$	6440	70			-0.7	o			Kur		10Ha.A	
	6393	50				3			Kur		14Ha38	
	3890	100				2					71Gu18 *	
$^{164}\text{Tb}(\beta^-)^{164}\text{Dy}$	990	30	961.4	1.4	-1.0	U					54Br96	
$^{164}\text{Ho}(\beta^-)^{164}\text{Er}$	965	20			-0.2	U					66Se07	
	3985	20	4039	24	2.7	B					67Vr04 *	
$^{164}\text{Tm}(\beta^+)^{164}\text{Er}$	3989	50			1.0	1	24	24 $^{164}\text{Tm}$	IRS		94Po26 *	
	6390	140	6380	30	-0.1	U					83Ge08	
$^{164}\text{Lu}(\beta^+)^{164}\text{Yb}$	6250	90			1.4	o			IRS		83Vi.A	
	6290	90			0.9	U			IRS		93Al03 *	
	6255	120			1.0	U			Dbn		94Po26 *	
* $^{164}\text{Tm}-\text{u}$	$M-A=-61884(28)$ keV for mixture gs+m at 10(6) keV											
* $^{164}\text{W}(\alpha)^{160}\text{Hf}$	Originally assigned to $^{168}\text{Re}$											
* $^{164}\text{W}(\alpha)^{160}\text{Hf}$	Originally assigned to $^{167}\text{Re}$ , re-assigned in reference											
*	original $E_\alpha=5136$ recalibrated using their $^{168}\text{Os}-^{170}\text{Os}$ results											
* $^{162}\text{Dy}(\text{t,p})^{164}\text{Dy}$	$Q-Q(^{160}\text{Dy}(\text{t,p}))=-722.3(1.9)$ keV, see $^{162}\text{Dy}(\text{p,t})$											
* $^{164}\text{Dy}(\text{t},\alpha)^{163}\text{Tb}$	$Q-Q(^{162}\text{Dy}(\text{t},\alpha))=-123(4)+54-584=-653(4)$ keV											
* $^{163}\text{Dy}(\text{}^3\text{He,d})^{164}\text{Ho}-^{164}\text{Dy}(\text{}^1\text{H})^{165}\text{H}$	See erratum											
* $^{164}\text{Tb}(\beta^-)^{164}\text{Dy}$	$E_{\beta^-}=1700(100)$ to $4^+$ level at 2194.44 and $4^+$ at 2205.63 keV, and other $E_{\beta^-}$											
* $^{164}\text{Tm}(\beta^+)^{164}\text{Er}$	$E_{\beta^+}=2940(20)$ 29 to ground state 10 to $2^+$ level at 91.38 keV											
* $^{164}\text{Tm}(\beta^+)^{164}\text{Er}$	$E_{\beta^+}=2944(50)$ 29 to ground state 10 to $2^+$ level at 91.38 keV											
* $^{164}\text{Lu}(\beta^+)^{164}\text{Yb}$	$Q_{\beta^+}=6250(90)$ partly to $2^+$ level at 123.31 keV											
* $^{164}\text{Lu}(\beta^+)^{164}\text{Yb}$	$E_{\beta^+}=5191(120)$ partly to $2^+$ level at 123.31 keV											

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_{13} H_9 - {}^{165}Ho$	140043	29	140097.2	1.1	0.5	U	R04	4.0	64De15
$C_{12} H_7 N - {}^{165}Ho$	127537	28	127521.2	1.1	-0.1	U	R04	4.0	64De15
$C_{11} {}^{13}C H_6 N - {}^{165}Ho$	122970	50	123051.0	1.1	0.4	U	R04	4.0	64De15
${}^{165}Tm - {}^{142}Sm_{1.162}$	30970	20	30975	4	0.2	U	MA7	1.0	01Bo59
${}^{165}Yb - u$	-64721	30	-64730	28	-0.3	1	90	90 ${}^{165}Yb$	GS2 1.0 05Li24
${}^{165}Lu - u$	-60602	30	-60593	28	0.3	1	90	90 ${}^{165}Lu$	GS2 1.0 05Li24
${}^{165}Hf - u$	-55360	140	-55430	30	-0.5	U			GS1 1.0 00Ra23
	-55433	30				2			GS2 1.0 05Li24
${}^{165}Ta - u$	-49191	30	-49220	15	-1.0	1	25	25 ${}^{165}Ta$	GS2 1.0 05Li24
${}^{165}W - u$	-41720	30	-41719	27	0.0	1	80	80 ${}^{165}W$	GS2 1.0 05Li24
${}^{165}Ho {}^{35}Cl - {}^{163}Dy {}^{37}Cl$	4539	4	4541.3	0.9	0.2	U			H23 2.5 70Wh01
${}^{165}W(\alpha) {}^{161}Hf$	5031.0	5.	5029	30	0.0	-			ORa 75To05 Z
	5034.2	10.			-0.1	-			84Sc06 *
	ave. 5032	4			-0.1	1	36	20 ${}^{165}W$	average
${}^{165}Re(\alpha) {}^{161}Ta$	5629.6	6.	5694	6	10.8	B			Jya 05Sc22
	5694.3	6.1				6			Jya 12Th13
${}^{165}Re^m(\alpha) {}^{161}Ta^m$	5631.7	10.	5660.9	2.8	2.9	U			78Sc26 *
	5643.0	10.			1.8	U			GSa 81Ho10
	5664.5	4.			-0.9	-			Ora 82De11 *
	5655.4	5.			1.1	-			Daa 96Pa01 *
	5657.4	5.			0.7	o			Jya 05Sc22
	5657.4	6.			0.6	-			Jya 12Th13
	ave. 5660.2	2.8			0.2	1	99	55 ${}^{165}Re^m$	average
${}^{165}Os(\alpha) {}^{161}W$	6354.3	20.	6335	6	-0.9	5			Ora 78Ca11
	6317.4	10.			1.8	5			GSa 81Ho10
	6342.1	7.			-0.9	5			Daa 96Pa01
	6342.1	30.			-0.2	U			Jya 13Dr06 *
${}^{165}Ir^m(\alpha) {}^{161}Re^m$	6882.1	7.	6879	6	-0.4	1	70	48 ${}^{165}Ir^m$	Ara 97Da07
${}^{163}Dy(t,p) {}^{165}Dy$	4890.6	2.9	4892.28	0.09	0.6	U			McM 88Bu08 *
${}^{164}Dy(n,\gamma) {}^{165}Dy$	5716.36	0.20	5715.96	0.05	-2.0	U			ILn 79Br25 Z
	5715.96	0.06			0.0	2			MMn 82Is05 Z
	5715.70	0.30			0.9	U			ILn 90Ka21 Z
	5715.95	0.12			0.1	2			Bdn 06Fi.A
${}^{164}Dy(d,p) {}^{165}Dy$	3488	5	3491.39	0.05	0.7	U			Tal 64Sh13
	3496	10			-0.5	U			Kop 70Gr46
${}^{164}Dy({}^3He,d) {}^{165}Ho$	717.3	10.	726.5	0.8	0.9	U			McM 75Bu02
${}^{165}Ho(\gamma,n) {}^{164}Ho$	-8160	80	-7988.8	1.1	2.1	U			Phi 60Ge01
	-7987	2			-0.9	1	33	33 ${}^{164}Ho$	MMn 85Ts01
${}^{165}Ho(d,t) {}^{164}Ho$	-1730	15	-1731.6	1.1	-0.1	U			Tal 70Jo11
${}^{164}Er(n,\gamma) {}^{165}Er$	6650.1	0.6	6650.0	0.6	-0.1	1	96	94 ${}^{165}Er$	70Bo29 Z
${}^{164}Er(d,p) {}^{165}Er$	4431	10	4425.5	0.6	-0.6	U			Kop 69Tj01
${}^{164}Er(\alpha,t) {}^{165}Tm - {}^{168}Er() {}^{169}Tm$	-1298.0	2.0	-1297.3	1.5	0.3	1	59	47 ${}^{165}Tm$	McM 75Bu02
${}^{165}Ir^m(p) {}^{164}Os$	1717.5	7.	1721	6	0.4	1	72	52 ${}^{165}Ir^m$	Ara 97Da07
${}^{165}Eu(\beta^-) {}^{165}Gd$	5800	120	5730	70	-0.6	o			Kur 07Ha57
	5800	120			-0.6	o			Kur 10Ha.A
	5729	65				4			Kur 14Ha38
${}^{165}Gd(\beta^-) {}^{165}Tb$	4113	65				3			Kur 14Ha38
${}^{165}Dy(\beta^-) {}^{165}Ho$	1305	20	1286.4	0.8	-0.9	U			59Bo52
	1285	10			0.1	U			63Pe11
${}^{165}Er(\epsilon) {}^{165}Ho$	370	10	377.4	1.0	0.7	U			63Ry01
	371	6			1.1	U			63Zy01
${}^{165}Tm(\beta^+) {}^{165}Er$	1591.3	2.0	1592.0	1.5	0.3	1	59	53 ${}^{165}Tm$	82Vy03 *
${}^{165}Yb(\beta^+) {}^{165}Tm$	2762	20	2634	27	-6.4	B			67Pa04 *
${}^{165}Lu(\beta^+) {}^{165}Yb$	4250	140	3850	40	-2.8	B			83Ge08
	3920	80			-0.8	o			83Vi.A
	3920	80			-0.8	1	20	10 ${}^{165}Yb$	IRS 93Al03
* ${}^{165}W(\alpha) {}^{161}Hf$	Originally assigned to ${}^{168}Re$ , re-assigned in reference								
*	original $E_\alpha=4894$ recalibrated using their ${}^{168}Os-{}^{170}Os$ results								
* ${}^{165}Re^m(\alpha) {}^{161}Ta^m$	Originally assigned to ${}^{166}Re$								
* ${}^{165}Re^m(\alpha) {}^{161}Ta^m$	Originally assigned to ${}^{166}Re$								
* ${}^{165}Re^m(\alpha) {}^{161}Ta^m$	Due to a high spin isomer								
* ${}^{165}Os(\alpha) {}^{161}W$	Error not given, estimated by evaluator								

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>163</sup> Dy(t,p) <sup>165</sup> Dy	$Q - Q(^{162}\text{Dy}(t,p)) = -556.6(2.9)$ keV										AHW **
* <sup>165</sup> Tm( $\beta^+$ ) <sup>165</sup> Er	$E_{\beta^+} = 272(2)$ to $1/2^-$ level at 297.371 keV										Ens066 **
* <sup>165</sup> Yb( $\beta^+$ ) <sup>165</sup> Tm	$E_{\beta^+} = 1580(20)$ to $7/2^-$ level at 160.47 keV										Ens066 **
C <sub>12</sub> H <sub>8</sub> N- <sup>166</sup> Er	135376	29	135375.2	1.3	0.0	U			R04	4.0	64De15
	135420	60			-0.2	U			R04	4.0	64De15
C <sub>13</sub> H <sub>10</sub> - <sup>166</sup> Er	147740	60	147951.3	1.3	0.9	U			R04	4.0	64De15
<sup>166</sup> Lu-u	-60157	108	-60140	30	0.1	U			GS1	1.0	00Ra23 *
	-60141	32				2			GS2	1.0	05Li24 *
<sup>166</sup> Hf-u	-57860	110	-57820	30	0.4	U			GS1	1.0	00Ra23
	-57820	30				2			GS2	1.0	05Li24
<sup>166</sup> Ta-u	-49488	30				2			GS2	1.0	05Li24
<sup>166</sup> W-u	-44957	30	-44969	10	-0.4	1	12	12 <sup>166</sup> W	GS2	1.0	05Li24
<sup>166</sup> Er <sup>35</sup> Cl- <sup>164</sup> Er <sup>37</sup> Cl	4040.9	1.4	4041.7	1.2	0.2	U			H25	2.5	72Ba08
<sup>166</sup> Er- <sup>164</sup> Er	1214	46	1091.6	1.2	-0.7	U			R04	4.0	64De15
	1110	80			-0.1	U			R04	4.0	64De15
<sup>166</sup> W( $\alpha$ ) <sup>162</sup> Hf	4856.0	5.	4856	4	0.0	-			ORa		75To05
	4855.0	10.			0.1	-			GSa		79Ho10 Z
	4858.3	8.2			-0.3	-					89Hi04
ave.	4856	4			-0.1	1	97	78 <sup>166</sup> W			average
<sup>166</sup> Re( $\alpha$ ) <sup>162</sup> Ta	5461.8	10.				5					78Sc26 *
	5574.5	3.	5460	50	-2.3	U			Ora		82De11 *
	5637.0	13.			-3.5	B			Bea		92Me10 *
	5669.9	10.			-4.2	B			Daa		96Pa01 *
<sup>166</sup> Os( $\alpha$ ) <sup>162</sup> W	6148.5	20.	6143	3	-0.3	U					77Ca23
	6129.0	6.			2.2	-			GSa		81Ho10
	6148.5	6.			-0.9	-			Daa		96Pa01
	6148.4	5.1			-1.1	-			Jya		15Li24
ave.	6143	3			0.0	1	100	100 <sup>166</sup> Os			average
<sup>166</sup> Ir( $\alpha$ ) <sup>162</sup> Re	6702.8	20.	6722	6	1.0	U			GSa		81Ho10
	6724.3	6.			-0.3	7			Ara		97Da07
	6713.1	13.			0.7	7			Jya		04Ke06 *
<sup>166</sup> Ir <sup>m</sup> ( $\alpha$ ) <sup>162</sup> Re <sup>m</sup>	6718.2	11.	6719	4	0.0	8			Daa		96Pa01 *
	6723.3	5.			-0.9	8			Ara		97Da07
	6706.9	8.2			1.4	8			Jya		04Ke06
<sup>166</sup> Pt( $\alpha$ ) <sup>162</sup> Os	7285.9	15.				5			ORa		96Bi07
<sup>164</sup> Dy(t,p) <sup>166</sup> Dy	4276.4	4.4	4277.7	0.4	0.3	U			McM		88Bu08 *
<sup>166</sup> Er(p,t) <sup>164</sup> Er	-6641	5	-6644.0	1.1	-0.6	U			Min		73Oo01
<sup>165</sup> Dy(n, $\gamma$ ) <sup>166</sup> Dy	7043.5	0.4				3					83Ke.A
<sup>165</sup> Ho(n, $\gamma$ ) <sup>166</sup> Ho	6243.69	0.06	6243.640	0.020	-0.8	U			MMn		82Is05 Z
	6243.64	0.02			0.0	1	100	77 <sup>166</sup> Ho	MMn		84Ke15 Z
	6243.68	0.13			-0.3	U			Bdn		06Fi.A
<sup>165</sup> Ho(d,p) <sup>166</sup> Ho	4025	7	4019.074	0.020	-0.8	U			Tal		65St06
<sup>166</sup> Er(d,t) <sup>165</sup> Er	-2218	10	-2218.5	1.3	-0.1	U			Kop		69Tj01
<sup>166</sup> Ir(p) <sup>165</sup> Os	1152.0	8.0				6			Ara		97Da07
<sup>166</sup> Ir <sup>m</sup> (p) <sup>165</sup> Os	1324.1	8.	1323	10	-0.1	o			Ara		97Da07 *
<sup>166</sup> Eu( $\beta^-$ ) <sup>166</sup> Gd	7322	300				3			Kur		14Ha38
<sup>166</sup> Tb( $\beta^-$ ) <sup>166</sup> Dy	4830	100	4700	70	-1.3	o			Kur		02Sh.A
	4695	70			0.1	o			Kur		07Ha57
	4700	70				4			Kur		10Ha.A
<sup>166</sup> Dy( $\beta^-$ ) <sup>166</sup> Ho	483	5	486.5	0.9	0.7	U					60He09 *
<sup>166</sup> Ho( $\beta^-$ ) <sup>166</sup> Er	1859	3	1854.7	0.9	-1.4	-					63Fu17
	1857	3			-0.8	-					66Da04
	1854.7	1.5			0.0	-					74Gr41
	1851.6	2.0			1.6	-					83Ra.A
ave.	1854.7	1.0			0.1	1	78	54 <sup>166</sup> Er			average
<sup>166</sup> Tm( $\beta^+$ ) <sup>166</sup> Er	3043	20	3038	12	-0.3	2					61Gr33 *
	3031	20			0.3	2					61Zy02 *
	3039	20			-0.1	2					63Pr13 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{166}\text{Yb}(\epsilon)^{166}\text{Tm}$	280	40	293	14	0.3	U					Averag *
$^{166}\text{Lu}(\beta^+)^{166}\text{Yb}$	5480	160	5570	30	0.6	U					74De09 *
$^{166}\text{Ir}^m(\text{IT})^{166}\text{Ir}$	171.5	6.1				7			Ara		97Da07
* $^{166}\text{Lu}-u$	$M - A = -56010(100)$ keV for mixture gs+m+n at 34.37 and 43.0 keV										Nub16b **
* $^{166}\text{Lu}-u$	$M - A = -55995(28)$ keV for mixture gs+m+n at 34.37 and 43.0 keV										Nub16b **
* $^{166}\text{Re}(\alpha)^{162}\text{Ta}$	Originally assigned to $^{167}\text{Re}$										AHW **
* $^{166}\text{Re}(\alpha)^{162}\text{Ta}$	Assignment uncertain, no other obvious attribution										AHW **
* $^{166}\text{Re}(\alpha)^{162}\text{Ta}$	Originally assigned to $^{167}\text{Re}$										AHW **
* $^{166}\text{Re}(\alpha)^{162}\text{Ta}$	Assignment tentative, may be $^{165}\text{Re}$										92Me10 **
* $^{166}\text{Re}(\alpha)^{162}\text{Ta}$	Correlated to a $^{170}\text{Ir}$ 6003 line; assignment uncertain										GAU **
* $^{166}\text{Ir}(\alpha)^{162}\text{Re}$	All $Q_\alpha$ of reference increased by 7 keV for calibration error										04Ke06 **
* $^{166}\text{Ir}^m(\alpha)^{162}\text{Re}^m$	Correlated with $E_\alpha=6123$ of $^{162}\text{Re}^m$										96Pa01 **
* $^{164}\text{Dy}(t,p)^{166}\text{Dy}$	$Q - Q(^{162}\text{Dy}(t,p)) = -1170.8(4.4)$ keV										AHW **
* $^{166}\text{Ir}^m(p)^{165}\text{Os}$	Replaced by author's value for $^{166}\text{Ir}^m(\text{IT})^{166}\text{Ir}$										97Da07 **
* $^{166}\text{Dy}(\beta^-)^{166}\text{Ho}$	$E_{\beta^-} = 402(5)$ to $1^-$ level at 82.47 keV, and other $E_{\beta^-}$										Ens084 **
* $^{166}\text{Tm}(\beta^+)^{166}\text{Er}$	$E_{\beta^+} = 1940(20)$ 1928(20) 1936(20) respectively, to $2^+$ level at 80.5776 keV										Ens084 **
* $^{166}\text{Yb}(\epsilon)^{166}\text{Tm}$	Average $pK=0.712(0.038)$ to $1^+$ level at 82.298 keV from 2 references:										Ens084 **
*	$pK=0.74(0.05)$ to 82.298 level										63Ja06 **
*	$pK=0.675(0.059)$ to 82.298 level										73De22 **
* $^{166}\text{Lu}(\beta^+)^{166}\text{Yb}$	$E_{\beta^+} = 2225(160)$ to $(6^-, 7^-)$ level at 2233.36 keV										Ens084 **
$C_{13} \text{H}_{11} - ^{167}\text{Er}$	153840	130	154021.2	1.3	0.3	U			R04	4.0	64De15
	154040.4	6.2			-0.8	U			M23	4.0	79Ha32
$C_{12} \text{H}_9 \text{N} - ^{167}\text{Er}$	141480	27	141445.2	1.3	-0.3	U			R04	4.0	64De15
	141520	50			-0.4	U			R04	4.0	64De15
$^{167}\text{Lu}-u$	-61730	34				2			GS2	1.0	05Li24 *
$^{167}\text{Hf}-u$	-57490	110	-57400	30	0.8	U			GS1	1.0	00Ra23
	-57400	30				2			GS2	1.0	05Li24
$^{167}\text{Ta}-u$	-51870	120	-51910	30	-0.3	U			GS1	1.0	00Ra23
	-51907	30				2			GS2	1.0	05Li24
$^{167}\text{W}-u$	-45175	30	-45194	20	-0.6	R			GS2	1.0	05Li24
$^{167}\text{Er} \text{ } ^{35}\text{Cl} - ^{165}\text{Ho} \text{ } ^{37}\text{Cl}$	4666	3	4676.2	1.0	1.4	U			H23	2.5	70Wh01
	4679.5	1.2			-1.1	U			H25	2.5	72Ba08
$^{167}\text{Er} - ^{166}\text{Er}$	1722	31	1755.10	0.19	0.3	U			R04	4.0	64De15
$^{167}\text{W}(\alpha)^{163}\text{Hf}$	4661.9	20.	4741	28	1.6	-					89Me02
	4671.1	13.			1.4	-					91Me05
	ave.	4668	11		1.4	1	32	21 $^{163}\text{Hf}$			average
$^{167}\text{Re}(\alpha)^{163}\text{Ta}^m$	5138.3	12.				12			Bea		92Me10
$^{167}\text{Re}^m(\alpha)^{163}\text{Ta}$	5408.8	3.	5407.0	2.9	-0.6	4			Ora		82De11 *
	5397.5	10.			0.9	4			ChR		84Sc06 *
	5392.4	12.			1.2	4			Bea		92Me10
$^{167}\text{Os}(\alpha)^{163}\text{W}$	5983.6	5.	5980	50	0.0	6			GSa		81Ho10 Z
	5978.7	2.			0.1	6			Ora		82De11 Z
	5996.9	5.			-0.3	6			Daa		96Pa01
	5979.5	5.			0.0	6			Bka		02Ro17
$^{167}\text{Ir}(\alpha)^{163}\text{Re}$	6507.1	5.	6504.9	2.6	-0.4	2			Ara		97Da07
	6504.0	3.			0.3	2			Jya		05Sc22
$^{167}\text{Ir}^m(\alpha)^{163}\text{Re}^m$	6543.0	10.	6561	3	1.7	2			GSa		81Ho10
	6567.6	11.			-0.6	2			Daa		96Pa01
	6567.6	5.			-1.4	2			Ara		97Da07
	6551.2	7.2			1.3	2			Jya		04Ke06
	6561.5	6.			-0.1	2			Jya		05Sc22
$^{167}\text{Pt}(\alpha)^{163}\text{Os}$	7159.8	10.	7160	50	-0.1	5			ORa		96Bi07
	7150.6	10.			0.1	5			Jya		04Ke06
$^{167}\text{Er}(p,t)^{165}\text{Er}$	-6427	6	-6430.4	1.3	-0.6	U			Min		73Oo01
	-6430	5			-0.1	U					75St08
$^{166}\text{Er}(n,\gamma)^{167}\text{Er}$	6436.35	0.50	6436.46	0.18	0.2	-					70Bo29 Z
	6436.51	0.40			-0.1	-					70Mi01 Z
	6436.46	0.22			0.0	-			Bdn		06Fi.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{167}\text{Er}(\gamma,n)^{166}\text{Er}$	-6560	80	-6436.46	0.18	1.5	U			Phi		60Ge01
$^{166}\text{Er}(d,p)^{167}\text{Er}$	4209	10	4211.89	0.18	0.3	U			Tal		68Ha10
	4214	10			-0.2	U			Kop		69Tj01
$^{167}\text{Er}(d,t)^{166}\text{Er}$	-189	12	-179.23	0.18	0.8	U			Kop		69Bu01
$^{166}\text{Er}(n,\gamma)^{167}\text{Er}$	ave. 6436.46	0.18	6436.46	0.18	0.0	1	99	53 $^{167}\text{Er}$			average
$^{166}\text{Er}(\alpha,t)^{167}\text{Tm}-^{168}\text{Er}()^{169}\text{Tm}$	-666.5	1.0	-666.4	1.0	0.1	1	99	99 $^{167}\text{Tm}$	McM		75Bu02
$^{167}\text{Ir}(p)^{166}\text{Os}$	1070.5	6.	1070	4	-0.1	-					97Da07
	1068.5	6.			0.3	-			Jyp		05Sc22 *
	ave. 1069	4			0.1	1	77	77 $^{167}\text{Ir}$			average
$^{167}\text{Ir}^m(p)^{166}\text{Os}$	1245.5	7.	1246	4	0.0	o					97Da07 *
$^{167}\text{Dy}(\beta^-)^{167}\text{Ho}$	2350	60				3					77Tu01 *
$^{167}\text{Ho}(\beta^-)^{167}\text{Er}$	970	20	1011	5	2.0	U					68Fu07
$^{167}\text{Yb}(\beta^+)^{167}\text{Tm}$	1954	4	1953	4	-0.2	1	90	89 $^{167}\text{Yb}$			77Kr.A *
$^{167}\text{Lu}(\beta^+)^{167}\text{Yb}$	3130	100	3090	30	-0.4	U					64Ag.A *
$^{167}\text{W}(\beta^+)^{167}\text{Ta}$	5620	270	6250	30	2.3	U			Got		89Me02
$^{167}\text{Ir}^m(\text{IT})^{167}\text{Ir}$	175.3	2.2	175.5	2.1	0.1	1	94	70 $^{167}\text{Ir}^m$	Ara		97Da07
* $^{167}\text{Lu}-u$	$M-A=-57501(28)$ keV for mixture gs+m at 0#30 keV										Nub16b **
* $^{167}\text{Re}^m(\alpha)^{163}\text{Ta}$	Original assignment to $^{168}\text{Re}$ changed in reference										92Me10 **
* $^{167}\text{Re}^m(\alpha)^{163}\text{Ta}$	Original assignment to $^{168}\text{Re}^m$ changed in reference										92Me10 **
*	original $E_\alpha=5250$ recalibrated using their $^{168}\text{Os}-^{170}\text{Os}$ results										GAu **
* $^{167}\text{Ir}(p)^{166}\text{Os}$	$E_p=1062(6)$ ; also $E_p=1248(7)$ from $^{167}\text{Ir}^m$										05Sc22 **
* $^{167}\text{Ir}^m(p)^{166}\text{Os}$	Replaced by author's value for $^{167}\text{Ir}^m(\text{IT})^{167}\text{Ir}$										97Da07 **
* $^{167}\text{Dy}(\beta^-)^{167}\text{Ho}$	$E_{\beta^-}=1780(60)$ to $3/2^-$ level at 569.69 keV										Ens008 **
* $^{167}\text{Yb}(\beta^+)^{167}\text{Tm}$	$E_{\beta^+}=639(4)$ to $7/2^-$ level at 292.820 keV										Ens008 **
* $^{167}\text{Lu}(\beta^+)^{167}\text{Yb}$	$E_{\beta^+}=2060(100)$ to $5/2^+$ level at 29.658, $7/2^-$ at 78.671 keV										Ens008 **
$\text{C}_{13} \text{H}_{12}-^{168}\text{Er}$	161543.3	5.1	161524.2	1.3	-0.9	U			M23	4.0	79Ha32
$\text{C}_{12} \text{H}_{10} \text{N}-^{168}\text{Er}$	148884	44	148948.1	1.3	0.4	U			R04	4.0	64De15
$\text{C}_{11} \text{}^{13}\text{C} \text{H}_9 \text{N}-^{168}\text{Er}$	144524	29	144477.9	1.3	-0.4	U			R04	4.0	64De15
$\text{C}_{12} \text{H}_{10} \text{N}-^{168}\text{Yb}$	147010	100	147435.2	1.3	1.1	U			R04	4.0	64De15
$^{168}\text{Lu}-u$	-61217	70	-61260	40	-0.7	R			GS2	1.0	05Li24 *
$^{168}\text{Hf}-u$	-59560	104	-59430	30	1.2	U			GS1	1.0	00Ra23
	-59432	30				2			GS2	1.0	05Li24
$^{168}\text{Ta}-u$	-52020	110	-51950	30	0.6	U			GS1	1.0	00Ra23
	-51953	30				2			GS2	1.0	05Li24
$^{168}\text{W}-u$	-48181	30	-48195	14	-0.5	1	23	23 $^{168}\text{W}$	GS2	1.0	05Li24
$^{168}\text{Yb} \text{}^{35}\text{Cl}_2-^{164}\text{Dy} \text{}^{37}\text{Cl}_2$	10612.8	8.7	10608.9	1.2	-0.2	U			H27	2.5	74Ba90
$^{168}\text{Er} \text{}^{35}\text{Cl}-^{166}\text{Er} \text{}^{37}\text{Cl}$	5037	50	5027.28	0.24	-0.1	U			R08	1.5	69De19
	5026	3			0.2	U			H23	2.5	70Wh01
	5028.9	1.5			-0.4	U			H25	2.5	72Ba08
$^{168}\text{Yb}-^{168}\text{Er}$	1512.91	0.27	1512.91	0.27	0.0	1	100	99 $^{168}\text{Yb}$	SH1	1.0	11El04
$^{168}\text{Er}-^{167}\text{Er}$	284	31	322.07	0.13	0.3	U			R04	4.0	64De15
	320.9	4.3			0.1	U			M24	2.5	79Ha32
$^{168}\text{W}(\alpha)^{164}\text{Hf}$	4506.5	12.	4500	11	-0.5	1	87	68 $^{164}\text{Hf}$			91Me05
$^{168}\text{Re}(\alpha)^{164}\text{Ta}$	5063	13				3			Bea		92Me10 *
$^{168}\text{Os}(\alpha)^{164}\text{W}$	5819.0	3.	5815.6	2.7	-1.1	-			Ora		82De11 Z
	5800.4	8.			1.9	-					84Sc06
	5812.7	8.			0.4	-					95Hi02
	ave. 5816.3	2.7			-0.2	1	99	80 $^{168}\text{Os}$			average
$^{168}\text{Ir}(\alpha)^{164}\text{Re}$	6410.9	5.	6381	9	-5.9	B			Ora		82De11
	6379.2	15.			0.1	5			Daa		96Pa01
	6382.2	10.			-0.1	5			Jya		09Ha42
$^{168}\text{Ir}^m(\alpha)^{164}\text{Re}^m$	6477.5	8.	6476	6	-0.1	6			Daa		96Pa01
	6474.4	10.			0.2	6			Jya		09Ha42 *
$^{168}\text{Pt}(\alpha)^{164}\text{Os}$	6990.8	20.	6990	3	-0.1	U			GSa		81Ho10
	6998.9	10.			-0.9	U			ORa		96Bi07
	6986.7	8.			0.4	o			Jya		04Ke06
	6989.7	3.1				2			Jya		09Go16

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{168}\text{Er}(p,t)^{166}\text{Er}$	-5723	6	-5725.97	0.21	-0.5	U			Min		73Oo01
$^{168}\text{Yb}(p,t)^{166}\text{Yb}$	-7647	7				2			Min		73Oo01
$^{167}\text{Er}(n,\gamma)^{168}\text{Er}$	7771.43	0.40	7771.31	0.12	-0.3	-					70Mi01 Z
	7771.05	0.20			1.3	-			ILn		79Br25 Z
	7771.6	1.0			-0.3	U			ORn		84Ka22
	7771.0	0.5			0.6	U					85Va.A
	7771.45	0.16			-0.9	-			Bdn		06Fi.A
$^{167}\text{Er}(d,p)^{168}\text{Er}$	5541	6	5546.74	0.12	1.0	U			Tal		67Ha25
$^{168}\text{Er}(d,t)^{167}\text{Er}$	-1523	10	-1514.08	0.12	0.9	U			Kop		69Tj01
$^{167}\text{Er}(n,\gamma)^{168}\text{Er}$	ave. 7771.31	0.12	7771.31	0.12	0.0	1	100	67 $^{168}\text{Er}$			average
$^{167}\text{Er}(\alpha,t)^{168}\text{Tm}-^{168}\text{Er}()^{169}\text{Tm}$	-262.3	1.5	-262.3	1.5	0.0	1	100	100 $^{168}\text{Tm}$	McM		75Bu02
$^{168}\text{Yb}(d,t)^{167}\text{Yb}$	-2797	12	-2805	4	-0.7	1	11	11 $^{167}\text{Yb}$	Kop		66Bu16
$^{168}\text{Ho}(\beta^-)^{168}\text{Er}$	2740	100	2930	30	1.9	U					73Ka07 *
	2930	30				2					90Ch37
$^{168}\text{Lu}(\beta^+)^{168}\text{Yb}$	4475	80	4510	40	0.5	2					70Ch28 *
	4493	100			0.2	2					72Ch44 *
	4500	80			0.2	2			IRS		83Vi.A
* $^{168}\text{Lu}-u$	$M-A=-56922(28)$ keV for mixture gs+m at 202.81 keV										Nub16b **
* $^{168}\text{Re}(\alpha)^{164}\text{Ta}$	$E_\alpha=4833(13)$ to level at 111.5 keV										Ens089 **
* $^{168}\text{Ir}^m(\alpha)^{164}\text{Re}^m$	$E_\alpha=6320(10), 6260(10)$ to ground state and level at 69 keV										09Ha42 **
* $^{168}\text{Ho}(\beta^-)^{168}\text{Er}$	$E_{\beta^-}=1900(100)$ to $2^+$ level at 821.17 and $3^+$ at 895.79 keV										Ens108 **
* $^{168}\text{Lu}(\beta^+)^{168}\text{Yb}$	$E_{\beta^+}=1230(80)$ to 2222.37 level										Ens108 **
* $^{168}\text{Lu}(\beta^+)^{168}\text{Yb}$	$E_{\beta^+}=1470(100)$ from $^{168}\text{Lu}^m$ at 202.81 to $4^+$ level at 2203.84 keV										Ens108 **
$\text{C}_{12}\text{H}_{11}\text{N}-^{169}\text{Tm}$	154920	60	154931.0	0.9	0.0	U			R04	4.0	64De15
$^{169}\text{Lu}-u$	-62362	31	-62356	3	0.2	U			GS2	1.0	05Li24 *
$^{169}\text{Hf}-u$	-58741	30				2			GS2	1.0	05Li24
$^{169}\text{Ta}-u$	-53960	110	-53990	30	-0.3	U			GS1	1.0	00Ra23
	-53989	30				2			GS2	1.0	05Li24
$^{169}\text{W}-u$	-48195	30	-48221	17	-0.9	1	31	31 $^{169}\text{W}$	GS2	1.0	05Li24
$^{169}\text{Re}-u$	-41203	63	-41234	12	-0.5	U			GS2	1.0	05Li24 *
$^{169}\text{Tm}^{35}\text{Cl}_2-^{165}\text{Ho}^{37}\text{Cl}_2$	9793.0	1.1	9790.5	1.1	-0.9	1	17	11 $^{165}\text{Ho}$	H25	2.5	72Ba08
$^{169}\text{Tm}^{35}\text{Cl}-^{167}\text{Er}^{37}\text{Cl}$	5107	3	5114.3	1.2	1.0	U			H23	2.5	70Wh01
	5113.2	1.1			0.4	1	19	15 $^{167}\text{Er}$	H25	2.5	72Ba08
$^{169}\text{Re}(\alpha)^{165}\text{Ta}^m$	4989.3	12.				5			Bea		92Me10 *
$^{169}\text{Re}^m(\alpha)^{165}\text{Ta}$	5189.1	3.	5189	3	-0.1	1	99	75 $^{165}\text{Ta}$	Ora		82De11
	5191.1	10.			-0.2	U			ChR		84Sc06 *
	5184.0	10.			0.5	U			Bea		92Me10
$^{169}\text{Os}(\alpha)^{165}\text{W}$	5717.6	4.	5713	3	-1.0	2			Ora		82De11
	5699.2	8.			1.7	2					84Sc06 *
	5713	8			0.1	2					95Hi02 *
	5711.5	8.			0.2	2			Daa		96Pa01
$^{169}\text{Ir}(\alpha)^{165}\text{Re}$	6150.8	8.	6141	4	-1.2	5			Ara		99Po09
	6138.5	4.			0.6	5			Jya		05Sc22
	6165	14			-1.7	U			Jya		12Th13
$^{169}\text{Ir}^m(\alpha)^{165}\text{Re}^m$	6276.0	3.	6266.5	2.9	-3.2	B			Ora		82De11 Z
	6258.4	10.			0.8	U			GSa		84Sc.A
	6267.6	9.			-0.1	-			Daa		96Pa01
	6254.3	5.			2.4	B			Ara		99Po09
	6265.6	3.			0.3	-			Jya		05Sc22
	6268	14			-0.1	U			Jya		12Th13
	ave. 6265.8	2.9			0.3	1	99	54 $^{169}\text{Ir}^m$			average
$^{169}\text{Pt}(\alpha)^{165}\text{Os}$	6840.2	15.	6858	5	1.2	U			GSa		81Ho10
	6860.7	23.			-0.1	U			Daa		96Pa01
	6853.5	8.2			0.5	o			Jya		04Ke06
	6857.6	5.1				6			Jya		09Go16



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{168}\text{Er}(n,\gamma)^{169}\text{Er}$	6002.5	0.7	6003.25	0.15	1.1	U					70Bo29 Z
	6003.5	0.3			-0.8	2					70Mu15 Z
	6003.16	0.18			0.5	2			Bdn		06Fi.A
$^{168}\text{Er}(d,p)^{169}\text{Er}$	3773	12	3778.68	0.15	0.5	U			Tal		68Ha10
	3781	10			-0.2	U			Kop		69Tj01
$^{168}\text{Er}(\alpha,t)^{169}\text{Tm}$	-14244.8	10.	-14240.9	1.1	0.4	U			McM		75Bu02
$^{169}\text{Tm}(\gamma,n)^{168}\text{Tm}$	-8110	50	-8033.6	1.5	1.5	U			Phi		60Ge01
$^{169}\text{Tm}(d,t)^{168}\text{Tm}$	-1775	6	-1776.4	1.5	-0.2	U			Pit		73Ko06
$^{168}\text{Yb}(n,\gamma)^{169}\text{Yb}$	6866.8	0.4	6866.98	0.15	0.4	2					68Mi08 Z
	6867.2	0.4			-0.6	2					68Sh12 Z
	6866.97	0.18			0.0	2			Bdn		06Fi.A
$^{168}\text{Yb}(d,p)^{169}\text{Yb}$	4636	12	4642.41	0.15	0.5	U			Kop		66Bu16
$^{169}\text{Dy}(\beta^-)^{169}\text{Ho}$	3200	300				3			LBL		90Ch34
$^{169}\text{Ho}(\beta^-)^{169}\text{Er}$	2070	100	2126	20	0.6	U					63Mi17 *
$^{169}\text{Er}(\beta^-)^{169}\text{Tm}$	343.8	3.	352.1	1.1	2.8	B					56Bi30 *
	347.8	5.			0.9	U					65Du02 *
$^{169}\text{Yb}(\epsilon)^{169}\text{Tm}$	913	12	897.6	1.1	-1.3	U					86Ad07 *
	900	100			0.0	U					87Sa53 *
$^{169}\text{Lu}(\beta^+)^{169}\text{Yb}$	2293	3				3					77Bo31
$^{169}\text{Hf}(\beta^+)^{169}\text{Lu}$	3365	200	3368	28	0.0	U					69Ar23 *
	3250	90			1.3	U					73Me09 *
* $^{169}\text{Lu}-u$	$M-A=-58075(28)$ keV for mixture gs+m at 29.0 keV										Nub16b **
* $^{169}\text{Re}-u$	$M-A=-38293(29)$ keV for mixture gs+m at 175(13) keV										Nub16b **
* $^{169}\text{Re}(\alpha)^{165}\text{Ta}^m$	$E_\alpha=4871(12)$ , and a stronger $E_\alpha=4700(12)$										92Me10 **
* $^{169}\text{Re}(\alpha)^{165}\text{Ta}$	Original $E_\alpha=5050$ recalibrated using their $^{168}\text{Os}-^{170}\text{Os}$ results										GAu **
* $^{169}\text{Os}(\alpha)^{165}\text{W}$	Used to recalibrate other results in same reference										GAu **
* $^{169}\text{Os}(\alpha)^{165}\text{W}$	$E_\alpha=5578(8)$ , $5536(10)$ to ground state, $(3/2^-)$ level at 43 keV										Ens066 **
* $^{169}\text{Ho}(\beta^-)^{169}\text{Er}$	$E_{\beta^-}=1200(100)$ to $5/2^-$ level at 853.0 and $7/2^-$ at 941.04 keV										Ens089 **
* $^{169}\text{Er}(\beta^-)^{169}\text{Tm}$	$E_{\beta^-}=340(2)$ $344(4)$ respectively, 55% to ground state, 45% to $3/2^+$ level at 8.41 keV										Ens089 **
* $^{169}\text{Yb}(\epsilon)^{169}\text{Tm}$	From decay rates to $(5/2)^+$ level at 781.796, $(7/2^+)$ 878.35 of same band										Ens089 **
* $^{169}\text{Yb}(\epsilon)^{169}\text{Tm}$	$pK=0.812(0.029)$ to $9/2^-$ level at 472.88 keV										Ens089 **
* $^{169}\text{Hf}(\beta^+)^{169}\text{Lu}$	$E_{\beta^+}=1850(200)$ to $7/2^-$ level at 492.88 keV										Ens089 **
* $^{169}\text{Hf}(\beta^+)^{169}\text{Lu}$	$K/\beta^+=5.2(1.0)$ to $7/2^-$ level at 492.88 keV										Ens089 **
$\text{C}_{12} \text{H}_{12} \text{N}-^{170}\text{Er}$	161210	70	161503.7	1.7	1.0	U			R04	4.0	64De15
$\text{C}_{12} \text{H}_{12} \text{N}-^{170}\text{Yb}$	161831	43	162207.146	0.011	2.2	U			R04	4.0	64De15
$\text{C}_{11} \text{H}_8 \text{O N}-^{170}\text{Yb}$	125370	150	125821.636	0.011	0.8	U			R04	4.0	64De15
$\text{C}_{11} ^{13}\text{C} \text{H}_{11} \text{N}-^{170}\text{Yb}$	157320	210	157736.949	0.011	0.5	U			R04	4.0	64De15
$^{170}\text{Yb}-^{129}\text{Xe}_{1,318}$	60266.078	0.012	60266.070	0.009	-0.7	1	58	53 $^{170}\text{Yb}$	FS1	1.0	12Ra34
$^{170}\text{Yb}-^{132}\text{Xe}_{1,288}$	58215.483	0.013	58215.493	0.009	0.7	1	50	47 $^{170}\text{Yb}$	FS1	1.0	12Ra34
$^{170}\text{Lu}-u$	-61529	42	-61521	18	0.2	R			GS2	1.0	05Li24 *
$^{170}\text{Hf}-u$	-60400	104	-60390	30	0.1	U			GS1	1.0	00Ra23
	-60391	30				2			GS2	1.0	05Li24
$^{170}\text{Ta}-u$	-53810	104	-53830	30	-0.1	U			GS1	1.0	00Ra23
	-53825	30				2			GS2	1.0	05Li24
$^{170}\text{W}-u$	-50710	110	-50769	14	-0.5	U			GS1	1.0	00Ra23
	-50755	30			-0.5	1	22	22 $^{170}\text{W}$	GS2	1.0	05Li24
$^{170}\text{Re}-u$	-41782	30	-41775	25	0.2	-			GS2	1.0	05Li24
	ave. -41780	28			0.2	1	80	80 $^{170}\text{Re}$			average
$^{170}\text{Os}-u$	-36454	31	-36421	10	1.1	1	11	11 $^{170}\text{Os}$	GS2	1.0	05Li24
$^{170}\text{Er} ^{35}\text{Cl}-^{168}\text{Er} ^{37}\text{Cl}$	6073	31	6044.6	1.6	-0.6	U			R08	1.5	69De19
	6040	3			0.6	U			H23	2.5	70Wh01
	6046.9	1.8			-0.5	1	12	9 $^{170}\text{Er}$	H25	2.5	72Ba08
$^{170}\text{Yb} ^{35}\text{Cl}-^{168}\text{Yb} ^{37}\text{Cl}$	3806.0	7.6	3828.2	1.3	1.2	U			H27	2.5	74Ba90

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{170}\text{Er}-^{168}\text{Er}$	3450	70	3094.5	1.6	-1.3	U			R04	4.0	64De15
$^{170}\text{Yb}-^{168}\text{Yb}$	910	200	878.1	1.3	0.0	U			R04	4.0	64De15
$^{170}\text{Os}(\alpha)^{166}\text{W}$	5533.5	10.	5536.9	2.7	0.3	-			ORa		72To06 Z
	5541.8	4.1			-1.2	-			Ora		82De11 Z
	5523.2	8.			1.7	-					84Sc06 *
	5533.4	8.			0.4	-					95Hi02
	5537.5	5.			-0.1	-			Bka		02Ro17
	ave.	5537.1	2.7		-0.1	1	99	89 $^{170}\text{Os}$			average
$^{170}\text{Ir}(\alpha)^{166}\text{Re}^p$	5955.4	10.				7			Bka		02Ro17
$^{170}\text{Ir}^m(\alpha)^{166}\text{Re}$	6175.4	10.	6272	10	9.7	B					78Sc26 *
	6172.7	5.			19.9	B			Ora		82De11 *
	6147.9	10.			12.4	B			Daa		96Pa01 *
	6229.9	11.			3.9	B			Daa		96Pa01 *
	6272.4	10.				6			Jya		07Ha45 *
$^{170}\text{Pt}(\alpha)^{166}\text{Os}$	6703.0	8.	6707	3	0.5	-			GSa		81Ho10
	6705.0	10.			0.2	-					82En03
	6708.1	6.			-0.1	-			ORa		96Bi07
	6711.2	11.			-0.3	-			Jya		97Uu01
	6723.5	14.			-1.1	-			Bka		01Ro.B
	6707.1	7.			0.0	-			Jya		04Ke06
	ave.	6708	3		-0.1	1	84	84 $^{170}\text{Pt}$			average
$^{170}\text{Au}(\alpha)^{166}\text{Ir}$	7174.1	11.	7177	15	0.3	o			Jya		02Ke.C
	7170.0	12.			0.6	U			Jya		04Ke06
$^{170}\text{Au}^m(\alpha)^{166}\text{Ir}^m$	7277.5	6.	7285	12	0.2	o			Jya		02Ke.C
	7226.3	15.			1.2	U			Ara		02Ma61
	7278.5	9.			0.1	U			Jya		04Ke06
$^{170}\text{Er}(p,\alpha)^{167}\text{Ho}$	7036	5				2			NDm		83Ta.A
$^{170}\text{Er}(^{18}\text{O},^{20}\text{Ne})^{168}\text{Dy}$	4710	140				2					98Lu08
$^{170}\text{Er}(p,t)^{168}\text{Er}$	-4785	5	-4778.3	1.5	1.3	U			Min		73Oo01
$^{170}\text{Yb}(p,t)^{168}\text{Yb}$	-6861	6	-6842.9	1.2	3.0	B			Min		73Oo01
$^{170}\text{Er}(d,^3\text{He})^{169}\text{Ho}$	-3107	20				2					76Su.A
$^{170}\text{Er}(d,t)^{169}\text{Er}$	-1010	10	-999.7	1.5	1.0	U			Kop		69Tj01
$^{169}\text{Tm}(n,\gamma)^{170}\text{Tm}$	6595.	2.5	6591.96	0.17	-1.2	U					66Sh03
	6592.1	1.5			-0.1	U					70Or.A
	6591.7	0.9			0.3	U			BNn		96Ho12 Z
	6591.95	0.17			0.0	1	99	79 $^{169}\text{Tm}$	Bdn		06Fi.A
$^{169}\text{Tm}(d,p)^{170}\text{Tm}$	4420	20	4367.39	0.17	-2.6	U			CIT		66Ry01
	4369	15			-0.1	U			Tal		66Sh03
$^{170}\text{Yb}(d,t)^{169}\text{Yb}$	-2211	12	-2200.4	1.2	0.9	U			Kop		66Bu16
$^{170}\text{Au}(p)^{169}\text{Pt}$	1473.8	15.	1472	12	-0.1	o			Jyp		02Ke.C
	1471.7	12.				7			Jyp		04Ke06
$^{170}\text{Au}^m(p)^{169}\text{Pt}$	1749.5	8.	1751	5	0.2	o			Jyp		02Ke.C
	1745.4	10.			0.6	7			Arp		02Ma61
	1753.5	6.			-0.4	7			Jyp		04Ke06
$^{170}\text{Ho}(\beta^-)^{170}\text{Er}$	3870	50				2					78Tu04
$^{170}\text{Ho}^m(\beta^-)^{170}\text{Er}$	3970	60				2					78Tu04
$^{170}\text{Tm}(\beta^-)^{170}\text{Yb}$	970	2	968.1	0.8	-1.0	-					54Po26
	967.3	1.			0.8	-					69Va17 *
	ave.	967.8	0.9		0.3	1	80	80 $^{170}\text{Tm}$			average
$^{170}\text{Lu}(\beta^+)^{170}\text{Yb}$	3467	20	3458	17	-0.5	2					60Dz02
	3410	50			1.0	2					65Ha30
* $^{170}\text{Lu}-u$	$M-A=-57267(29)$ keV for mixture gs+m at 92.91 keV										Nub16b **
* $^{170}\text{Os}(\alpha)^{166}\text{W}$	Used to recalibrate other results in same reference										GAu **
* $^{170}\text{Ir}^m(\alpha)^{166}\text{Re}$	$E_\alpha=6029.8(10,Z)$ $6027.2(5,Z)$ $6003(10)$ most probably to low levels in $^{166}\text{Re}$										GAu **
* $^{170}\text{Ir}^m(\alpha)^{166}\text{Re}$	Correlated with $^{166}\text{Re}$ $E_\alpha=5533$ keV										96Pa01 **
* $^{170}\text{Ir}^m(\alpha)^{166}\text{Re}$	$E_\alpha=5951(10)$ to level at 175, $6007(10)$ to 122, $6053(10)$ to 75 keV										07Ha45 **
* $^{170}\text{Tm}(\beta^-)^{170}\text{Yb}$	$E_{\beta^-}=883(1)$ to $2^+$ level at 84.25468 keV										Ens02b **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$C_{11}^{13C H_{12} N-^{171}Yb}$	164140	80	163997.709	0.014	-0.4	U			R04	4.0	64De15
$C_{10}^{H_7 O N_2-^{171}Yb}$	119640	270	119506.337	0.014	-0.1	U			R04	4.0	64De15
$^{171}Yb-^{129}Xe_{1,326}$	62592.096	0.012	62592.096	0.012	0.0	1	100	100	$^{171}Yb$	1.0	12Ra34
$^{171}Lu-u$	-62132	41	-62081.3	2.0	1.2	U			GS2	1.0	05Li24 *
$^{171}Hf-u$	-59570	104	-59510	30	0.6	U			GS1	1.0	00Ra23 *
	-59508	31				2			GS2	1.0	05Li24 *
$^{171}Ta-u$	-55550	104	-55520	30	0.3	U			GS1	1.0	00Ra23
	-55524	30				2			GS2	1.0	05Li24
$^{171}W-u$	-50650	110	-50550	30	0.9	U			GS1	1.0	00Ra23
	-50549	30				2			GS2	1.0	05Li24
$^{171}Re-u$	-44284	30				2			GS2	1.0	05Li24
$^{171}Os-u$	-36796	30	-36825	19	-1.0	-			GS2	1.0	05Li24
	ave.	-36801	21		-1.1	1	81	81	$^{171}Os$		average
$^{171}Yb^{35}Cl_2-^{167}Er^{37}Cl_2$	10178.0	1.7	10177.6	1.3	-0.1	U			H27	2.5	74Ba90
$^{171}Yb^{35}Cl-^{169}Tm^{37}Cl$	5055	3	5063.3	0.9	1.1	U			H23	2.5	70Wh01
	5061.9	1.7			0.3	U			H27	2.5	74Ba90
$^{171}Yb-^{170}Yb$	1220	60	1564.271	0.015	1.4	U			R04	4.0	64De15
$^{171}Os(\alpha)^{167}W$	5365.8	10.	5371	4	0.5	-			ORa		72To06
	5365.8	10.			0.5	-					78Sc26
	5393.4	15.			-1.4	-					79Ha10
	5367.9	8.			0.4	-					95Hi02 *
	5374.0	9.			-0.3	-			Daa		96Pa01
	ave.	5371	4		0.1	1	99	90	$^{167}W$		average
$^{171}Ir(\alpha)^{167}Re^m$	5854.2	10.	5866	5	1.1	5			Bka		02Ro17 *
	5865.4	8.			0.0	5			Ara		11Ko.B *
	5871.6	7.2			-0.8	5			Anv		13An10
$^{171}Ir^m(\alpha)^{167}Re$	6159.2	3.	6161.1	2.3	0.6	11			Ora		82De11 *
	6159	5			0.4	11					92Sc16 *
	6180	11			-1.7	11			Daa		96Pa01 *
	6159.2	8.			0.2	11			Anv		10An01 *
	6172.4	8.			-1.4	11			Ara		11Ko.B *
$^{171}Pt(\alpha)^{167}Os$	6608.1	4.	6607	3	-0.2	7			Ora		81De22 Z
	6606.8	5.			0.1	7			GSa		81Ho10 Z
	6604.8	11.			0.2	7			Jya		97Uu01
	6600.6	15.			0.5	U			Anv		10An01
$^{171}Au^m(\alpha)^{167}Ir^m$	7163.9	6.	7164	4	0.1	-			Ara		97Da07
	7162.9	8.			0.2	-			Jya		04Ke06
	ave.	7164	5		0.2	1	69	39	$^{171}Au^m$		average
$^{171}Hg(\alpha)^{167}Pt$	7667.7	15.				6			Jya		04Ke06
$^{171}Yb(p,t)^{169}Yb$	-6599	5	-6590.1	1.2	1.8	U			Min		73Oo01
$^{170}Er(n,\gamma)^{171}Er$	5681.5	0.5	5681.6	0.4	0.2	-					71Al01
	5681.6	0.5			0.0	-			Bdn		06Fi.A
$^{170}Er(d,p)^{171}Er$	3450	10	3457.0	0.4	0.7	U			Tal		68Ha10
	3458	10			-0.1	U			Kop		69Tj01
$^{170}Er(n,\gamma)^{171}Er$	ave.	5681.6	0.4	5681.6	0.4	0.1	1	98	62	$^{171}Er$	average
$^{170}Er(\alpha,t)^{171}Tm-^{168}Er(^{169}Tm)$	817.9	1.0	817.6	0.9	-0.3	1	82	53	$^{170}Er$	McM	75Bu02
$^{170}Yb(n,\gamma)^{171}Yb$	6614.3	0.6	6614.208	0.014	-0.2	U					72Wa10 Z
	6616.6	0.4			-6.0	C			Bdn		06Fi.A
$^{170}Yb(d,p)^{171}Yb$	4390	12	4389.642	0.014	0.0	U			Kop		66Bu16
$^{171}Yb(d,t)^{170}Yb$	-359	12	-356.979	0.014	0.2	U			Kop		66Bu16
$^{170}Yb(\alpha,t)^{171}Lu-^{174}Yb(^{175}Lu)$	-1156.2	2.0	-1156.6	1.7	-0.2	1	73	61	$^{171}Lu$	McM	75Bu02
$^{171}Au(p)^{170}Pt$	1452.6	17.	1448	10	-0.3	2			Arp		99Po09
	1445.6	12.			0.2	2			Jyp		04Ke06
$^{171}Au^m(p)^{170}Pt$	1702.1	6.	1702	4	0.1	-					97Da07
	1704.1	6.			-0.3	-			Jyp		04Ke06
	ave.	1703	4		-0.1	1	77	61	$^{171}Au^m$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{171}\text{Ho}(\beta^-)^{171}\text{Er}$	3200	600				2			LBL		90Ch34
$^{171}\text{Er}(\beta^-)^{171}\text{Tm}$	1490	2	1491.3	1.3	0.7	1	39	38	$^{171}\text{Er}$		61Ar15 *
$^{171}\text{Tm}(\beta^-)^{171}\text{Yb}$	96.5	1.0	96.5	1.0	0.0	1	94	94	$^{171}\text{Tm}$		57Sm73
$^{171}\text{Lu}(\beta^+)^{171}\text{Yb}$	1479.3	3.	1478.4	1.9	-0.3	1	39	39	$^{171}\text{Lu}$		77Bo32 *
$^{171}\text{Re}(\beta^+)^{171}\text{W}$	5670	200	5840	40	0.8	U			Got		87Ru05
$^{171}\text{Au}^m(\text{IT})^{171}\text{Au}$	250	16	255	10	0.3	o					99Po09 *
* $^{171}\text{Lu}-u$	$M-A=-57840(33)$ keV for mixture gs+m at 71.13 keV										Nub16b **
* $^{171}\text{Hf}-u$	$M-A=-55480(100)$ keV for mixture gs+m at 21.93 keV										Nub16b **
* $^{171}\text{Hf}-u$	$M-A=-55420(28)$ keV for mixture gs+m at 21.93 keV										Nub16b **
* $^{171}\text{Os}(\alpha)^{167}\text{W}$	$E_\alpha=5241(8), 5166(8)$ to ground state and level at 79 keV										95Hi02 **
* $^{171}\text{Ir}(\alpha)^{167}\text{Re}^m$	Correlated with $E_\alpha=6412$ of $^{175}\text{Au}$										02Ro17 **
* $^{171}\text{Ir}(\alpha)^{167}\text{Re}^m$	Correlated with $E_\alpha=6430(8)$ of $^{175}\text{Au}$ and $6556(8)$ of $^{179}\text{Tl}$										11Ko.B **
* $^{171}\text{Ir}^m(\alpha)^{167}\text{Re}$	$E_\alpha=5925.2(3,Z) 5925(5) 5945(11) 5925(8)$ respectively, to $(11/2^-)$ level at 92 keV										92Sc16 **
*	$E_\alpha=5920$ correlated with $^{175}\text{Au}$ $E_\alpha=6438$ keV										02Ro17 **
* $^{171}\text{Ir}^m(\alpha)^{167}\text{Re}$	$E_\alpha=5938(8)$ to 92 level; correlated with $E_\alpha=6431(8)$ of $^{175}\text{Au}^m$										11Ko.B **
*	and $7194(8)$ of $^{179}\text{Tl}^m$										11Ko.B **
* $^{171}\text{Er}(\beta^-)^{171}\text{Tm}$	$E_{\beta^-}=1065(2)$ to $7/2^-$ level at 424.95 keV										Ens029 **
* $^{171}\text{Lu}(\beta^+)^{171}\text{Yb}$	$E_{\beta^+}=362(3)$ to $7/2^+$ level at 95.28 keV										Ens029 **
* $^{171}\text{Au}^m(\text{IT})^{171}\text{Au}$	Redundant; use only their $Q_p$										GAu **
$\text{C}_{10} \text{H}_6 \text{O}_2 \text{N}-^{172}\text{Yb}$	103560	60	103466.778	0.015	-0.4	U			R04	4.0	64De15
$^{172}\text{Yb}-^{132}\text{Xe}_{1.303}$	61272.578	0.013	61272.578	0.013	0.0	1	100	100	$^{172}\text{Yb}$	1.0	12Ra34
$^{172}\text{Hf}-u$	-60555	30	-60550	26	0.2	2			GS2	1.0	05Li24
$^{172}\text{Ta}-u$	-55105	30				2			GS2	1.0	05Li24
$^{172}\text{W}-u$	-52770	110	-52710	30	0.6	U			GS1	1.0	00Ra23
	-52708	30				2			GS2	1.0	05Li24
$^{172}\text{Re}-u$	-44702	221	-44590	40	0.5	U			GS1	1.0	00Ra23 *
	-44587	62			-0.1	1	46	46	$^{172}\text{Re}$	1.0	05Li24 *
$^{172}\text{Yb } ^{35}\text{Cl}_2-^{168}\text{Er } ^{37}\text{Cl}_2$	9906.7	1.7	9910.7	1.3	0.9	U			H27	2.5	74Ba90
$^{172}\text{Yb } ^{35}\text{Cl}-^{170}\text{Yb } ^{37}\text{Cl}$	4568.5	2.0	4569.52	0.07	0.2	U			H27	2.5	74Ba90
$^{172}\text{Yb}-^{171}\text{Yb}$	-50	230	55.142	0.018	0.1	U			R04	4.0	64De15
$^{172}\text{Os}(\alpha)^{168}\text{W}$	5226.8	10.	5224	7	-0.2	-					71Bo06
	5227.8	10.			-0.3	-			Daa		96Pa01
ave.	5227	7			-0.4	1	93	59	$^{168}\text{W}$		average
$^{172}\text{Ir}(\alpha)^{168}\text{Re}$	5990.6	10.				4					92Sc16 *
$^{172}\text{Ir}^m(\alpha)^{168}\text{Re}$	6129.3	3.	6129.2	2.6	0.0	4			Ora		82De11 *
	6161	20			-1.6	F			GSa		84Sc.A *
	6129.1	5.			0.0	4					92Sc16 *
	6123.0	12.			0.5	U			Daa		96Pa01 *
$^{172}\text{Pt}(\alpha)^{168}\text{Os}$	6464.8	4.	6463	4	-0.4	1	97	77	$^{172}\text{Pt}$		81De22 Z
	6474.8	15.			-0.8	U			Anv		09An20
$^{172}\text{Au}(\alpha)^{168}\text{Ir}$	6923.3	10.2				6			Jya		09Ha42
$^{172}\text{Au}^m(\alpha)^{168}\text{Ir}^m$	7023.6	10.	7034	6	1.0	7					93Se09
	7042.1	9.			-0.9	7			Daa		96Pa01
	7033.8	10.			0.0	7			Jya		09Ha42 *
$^{172}\text{Hg}(\alpha)^{168}\text{Pt}$	7525.3	12.	7524	6	-0.1	3					99Se14
	7536.5	16.			-0.8	o			Jya		04Ke06
	7523.3	7.2			0.1	3			Jya		09Sa27
$^{170}\text{Er}(\text{t,p})^{172}\text{Er}$	4034	4	4036	4	0.4	1	89	87	$^{172}\text{Er}$		80Sh14
$^{172}\text{Yb}(\text{p,t})^{170}\text{Yb}$	-6161	5	-6152.366	0.015	1.7	U			Min		73Oo01
$^{171}\text{Yb}(\text{n},\gamma)^{172}\text{Yb}$	8020.3	0.7	8019.953	0.017	-0.5	U					71Al14 Z
	8020.1	0.5			-0.3	U					75Gr32
	8019.67	0.35			0.8	U			ILn		85Ge02 Z
	8019.27	0.17			4.0	C			Bdn		06Fi.A
$^{171}\text{Yb}(\text{d,p})^{172}\text{Yb}$	5797	12	5795.387	0.017	-0.1	U			Kop		66Bu16
	5789	5			1.3	U			Tal		66Sh14

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{172}\text{Yb}(d,t)^{171}\text{Yb}$	-1772	12	-1762.724	0.017	0.8	U	Kop		66Bu16
$^{171}\text{Yb}(^3\text{He},d)^{172}\text{Lu}$	-792	34	-775.3	2.3	0.5	U	Roc		76El11
$^{171}\text{Yb}(\alpha,t)^{172}\text{Lu}-^{174}\text{Yb}(\alpha)^{175}\text{Lu}$	-791.9	2.0	-791.9	2.0	0.0	1	100	100	$^{172}\text{Lu}$ McM
$^{172}\text{Er}(\beta^-)^{172}\text{Tm}$	888	5	891	5	0.6	1	83	70	$^{172}\text{Tm}$
$^{172}\text{Tm}(\beta^-)^{172}\text{Yb}$	1870	10	1881	6	1.1	1	30	30	$^{172}\text{Tm}$
$^{172}\text{Hf}(\epsilon)^{172}\text{Lu}$	350	50	334	25	-0.3	R			79To18
$^{172}\text{Ta}(\beta+)^{172}\text{Hf}$	4920	180	5070	40	0.8	U			73Ca10
$^{172}\text{W}(\beta+)^{172}\text{Ta}$	3210	100	2230	40	-9.8	C			74Ca.A
* $^{172}\text{Re}-u$	$M-A=-41640(200)$ keV for mixture gs+m at 0#100 keV								
* $^{172}\text{Re}-u$	$M-A=-41533(28)$ keV for mixture gs+m at 0#100 keV								
* $^{172}\text{Ir}(\alpha)^{168}\text{Re}$	$E_\alpha=5510(10)$ to 89.7+123.2+136.3 level								
* $^{172}\text{Ir}^m(\alpha)^{168}\text{Re}$	$E_\alpha=5736$ followed by XK(Re), 128 M2 and 161 M3 $\gamma$ 's								
* $^{172}\text{Ir}^m(\alpha)^{168}\text{Re}$	F : first assigned to $^{173}\text{Ir}(\alpha)$ ; seen in neither nuclide in reference								
* $^{172}\text{Ir}^m(\alpha)^{168}\text{Re}$	$E_\alpha=5828.2(3,Z)$ 5828(5) 5822(12) respectively, to ( $8^+$ ) level at 162.1 keV								
* $^{172}\text{Au}^m(\alpha)^{168}\text{Ir}^m$	$E_\alpha=6870(10)$ 6800(10) to ground state and 70 keV level								
* $^{172}\text{Er}(\beta^-)^{172}\text{Tm}$	$E_{\beta^-}=278(5)$ to $1^+$ level at 610.062 keV								
* $^{172}\text{Ta}(\beta+)^{172}\text{Hf}$	$E_{\beta^+}=2480(180)$ to $4^-$ level at 1418.55 keV								
* $^{172}\text{W}(\beta+)^{172}\text{Ta}$	$E_{\beta^+}=1600(100)$ in coinc. with 459.2 keV $\gamma$ from 586.3 level								
$\text{C}_{14}\text{H}_5-^{173}\text{Yb}$	101030	70	100908.946	0.012	-0.4	U			R04 4.0 64De15
$\text{C}_{10}\text{H}_7\text{O}_2\text{N}-^{173}\text{Yb}$	109810	60	109462.254	0.012	-1.4	U			R04 4.0 64De15
$^{173}\text{Yb}-^{129}\text{Xe}_{1.341}$	65905.075	0.013	65905.080	0.010	0.3	1	60	56	$^{173}\text{Yb}$ FS1 1.0 12Ra34
$^{173}\text{Yb}-^{132}\text{Xe}_{1.311}$	63868.901	0.015	63868.895	0.010	-0.4	1	47	44	$^{173}\text{Yb}$ FS1 1.0 12Ra34
$^{173}\text{Hf}-u$	-59487	30				2			GS2 1.0 05Li24
$^{173}\text{Ta}-u$	-56270	104	-56250	30	0.2	U			GS1 1.0 00Ra23
	-56250	30				2			GS2 1.0 05Li24
$^{173}\text{W}-u$	-52340	104	-52310	30	0.3	U			GS1 1.0 00Ra23
	-52311	30				2			GS2 1.0 05Li24
$^{173}\text{Re}-u$	-46910	110	-46760	30	1.4	U			GS1 1.0 00Ra23
	-46757	30				2			GS2 1.0 05Li24
$^{173}\text{Os}-u$	-40169	30	-40192	16	-0.8	1	29	29	$^{173}\text{Os}$ GS2 1.0 05Li24
$^{173}\text{Ir}-u$	-32449	100	-32495	12	-0.5	U			GS2 1.0 05Li24
$^{173}\text{Yb } ^{35}\text{Cl}_2-^{169}\text{Tm } ^{37}\text{Cl}_2$	9898.3	1.2	9898.1	0.9	-0.1	U			H27 2.5 74Ba90
$^{173}\text{Yb } ^{35}\text{Cl}-^{171}\text{Yb } ^{37}\text{Cl}$	4827	4	4834.81	0.07	0.8	U			H23 2.5 70Wh01
	4835.3	1.6			-0.1	U			H27 2.5 74Ba90
$^{173}\text{Yb}-^{172}\text{Yb}$	1970	120	1829.556	0.017	-0.3	U			R04 4.0 64De15
$^{173}\text{Os}(\alpha)^{169}\text{W}$	5057.2	10.	5055	6	-0.2	-			71Bo06
	5055.2	7.			-0.1	-			GSa 84Sc.A
	ave.	6			-0.2	1	97	69	$^{169}\text{W}$ average
$^{173}\text{Ir}(\alpha)^{169}\text{Re}^m$	5544.4	10.	5541	10	-0.4	1	90	76	$^{169}\text{Re}^m$ 92Sc16
$^{173}\text{Ir}^m(\alpha)^{169}\text{Re}$	5930.4	5.	5941.8	2.5	2.3	4			67Si02
	5947.1	4.			-1.3	4			Ora 82De11
	5937	10			0.5	4			GSa 84Sc.A
	5944.8	5.			-0.6	4			92Sc16
	5951.9	13.			-0.8	4			Daa 96Pa01
	5927.3	20.			0.7	U			Ara 01Ko.B
$^{173}\text{Pt}(\alpha)^{169}\text{Os}$	6359.3	8.2	6350	50	-0.1	3			79Ha10
	6352.3	3.			0.1	3			Ora 81De22
	6382.9	10.			-0.6	o			GSa 84Sc.A
	6372.6	9.			-0.4	3			Daa 96Pa01
	6387.9	15.			-0.7	U			Anv 09An20
$^{173}\text{Au}(\alpha)^{169}\text{Ir}$	6830.2	6.	6836	5	1.0	4			Ara 99Po09
	6847.6	8.			-1.4	4			Ara 01Ko44
	6846.6	14.			-0.7	U			Ara 12Th13
$^{173}\text{Au}^m(\alpha)^{169}\text{Ir}^m$	6896.8	10.	6897	3	0.0	-			GSa 84Sc.A
	6909.1	9.			-1.3	-			Daa 96Pa01
	6891.6	4.			1.3	-			Ara 99Po09
	6900.8	6.			-0.6	-			Ara 01Ko44
	6899	15			-0.1	U			Jya 12Th13
	ave.	3			0.3	1	98	52	$^{173}\text{Au}^m$ average
$^{173}\text{Hg}(\alpha)^{169}\text{Pt}$	7382.0	11.3	7378	4	-0.4	7			99Se14

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{173}\text{Hg}(\alpha)^{169}\text{Pt}$	7362.5	15.4	7378	4	1.0	7			Jya		04Ke06
	7378.9	5.			-0.2	7					12Od01
$^{173}\text{Yb}(\text{p,t})^{171}\text{Yb}$	-5913	5	-5905.255	0.015	1.5	U			Min		73Oo01
$^{172}\text{Yb}(\text{n},\gamma)^{173}\text{Yb}$	6367.3	0.4	6367.097	0.015	-0.5	U					71Al01 Z
	6367.2	0.6			-0.2	U			Bdn		06Fi.A
$^{173}\text{Yb}(\gamma,\text{n})^{172}\text{Yb}$	-6500	80	-6367.097	0.015	1.7	U			Phi		60Ge01
$^{172}\text{Yb}(\text{d,p})^{173}\text{Yb}$	4145	12	4142.531	0.015	-0.2	U			Kop		66Bu16
$^{173}\text{Yb}(\text{d,t})^{172}\text{Yb}$	-114	12	-109.868	0.015	0.3	U			Kop		66Bu16
$^{172}\text{Yb}(\alpha,\text{t})^{173}\text{Lu}-^{174}\text{Yb}(\text{)}^{175}\text{Lu}$	-595.6	1.0	-595.6	1.0	0.0	1	100	100 $^{173}\text{Lu}$	McM		75Bu02
$^{173}\text{Tm}(\beta^-)^{173}\text{Yb}$	1260	50	1295	4	0.7	U					63Ku22
	1320	40			-0.6	U					63Or01
$^{173}\text{Lu}(\epsilon)^{173}\text{Yb}$	675	20	670.3	1.6	-0.2	U					73Ko13
$^{173}\text{Ta}(\beta^+)^{173}\text{Hf}$	3670	200	3020	40	-3.3	B					73Re03
$^{173}\text{W}(\beta^+)^{173}\text{Ta}$	4000	300	3670	40	-1.1	U					80Vi.A
$^{173}\text{Ir}-\text{u}$	$M - A = -30113(70)$ keV for mixture gs+m at 226(9) keV										Nub16b **
$^{173}\text{Ir}^m(\alpha)^{169}\text{Re}$	$E_\alpha = 5660.0(5,Z) 5676.2(4,Z) 5666(10) 5674(5) 5681(13)$ respectively,										AHW **
*	to $(11/2^-)$ level at 136.40 keV										Ens155 **
$\text{C}_{14} \text{H}_6 - ^{174}\text{Yb}$	108308	38	108082.645	0.012	-1.5	U			R04	4.0	64De15
$^{174}\text{Yb} - ^{129}\text{Xe}_{1,349}$	67318.179	0.011	67318.168	0.009	-1.0	1	72	68 $^{174}\text{Yb}$	FS1	1.0	12Ra34
$^{174}\text{Yb} - ^{132}\text{Xe}_{1,318}$	65191.113	0.017	65191.140	0.010	1.6	1	34	32 $^{174}\text{Yb}$	FS1	1.0	12Ra34
$^{174}\text{Ta}-\text{u}$	-55546	30				2			GS2	1.0	05Li24
$^{174}\text{W}-\text{u}$	-53940	104	-53920	30	0.2	U			GS1	1.0	00Ra23
	-53921	30				2			GS2	1.0	05Li24
$^{174}\text{Re}-\text{u}$	-46930	104	-46890	30	0.4	U			GS1	1.0	00Ra23
	-46885	30				2			GS2	1.0	05Li24
$^{174}\text{Os}-\text{u}$	-42880	110	-42937	11	-0.5	U			GS1	1.0	00Ra23
	-42919	30			-0.6	1	13	13 $^{174}\text{Os}$	GS2	1.0	05Li24
$^{174}\text{Ir}-\text{u}$	-33127	72	-33133	26	-0.1	R			GS2	1.0	05Li24 *
$^{174}\text{Yb} \text{ } ^{35}\text{Cl} - ^{172}\text{Yb} \text{ } ^{37}\text{Cl}$	5420	4	5431.00	0.07	1.1	U			H23	2.5	70Wh01
	5430.3	1.1			0.3	U			H27	2.5	74Ba90
$^{174}\text{Yb} - ^{173}\text{Yb}$	700	50	651.333	0.014	-0.2	U			R04	4.0	64De15
$^{174}\text{Hf}(\alpha)^{170}\text{Yb}$	2558.9	30.	2494.5	2.3	-2.1	U					61Ma05
$^{174}\text{Os}(\alpha)^{170}\text{W}$	4872.2	10.	4871	10	-0.2	1	90	78 $^{170}\text{W}$			71Bo06
$^{174}\text{Ir}(\alpha)^{170}\text{Re}$	5624.1	10.	5625	10	0.1	1	97	77 $^{174}\text{Ir}$			92Sc16 *
$^{174}\text{Ir}^m(\alpha)^{170}\text{Re}$	5817.6	6.	5817	4	-0.1	2					67Si02 *
	5816.4	5.			0.1	2					92Sc16 *
$^{174}\text{Pt}(\alpha)^{170}\text{Os}$	6176.3	10.	6183	3	0.7	2					79Ha10 Z
	6185.7	5.			-0.5	2			Ora		81De22 Z
	6182.4	5.1			0.2	2			Ara		04Go38
$^{174}\text{Au}(\alpha)^{170}\text{Ir}$	6700.3	10.	6699	7	-0.1	7			GSa		84Sc.A
	6698.3	10.			0.1	7			Daa		96Pa01 *
$^{174}\text{Au}^m(\alpha)^{170}\text{Ir}^m$	6683.9	20.	6784	8	5.0	B			GSa		83Sc24 *
	6778	10			0.6	7			GSa		84Sc.A *
	6793.5	13.			-0.7	7			Daa		96Pa01
$^{174}\text{Hg}(\alpha)^{170}\text{Pt}$	7235.6	11.	7233	6	-0.2	2			Jya		97Uu01
	7232.5	8.2			0.1	2					99Se14
	7231.5	14.3			0.1	2			Bka		01Ro.B
$^{174}\text{Yb}(\text{p,t})^{172}\text{Yb}$	-5359	5	-5349.906	0.015	1.8	U			Min		73Oo01
$^{173}\text{Yb}(\text{n},\gamma)^{174}\text{Yb}$	7464.63	0.06	7464.604	0.013	-0.4	U			MMn		82Is05 Z
	7464.58	0.35			0.1	U			ILn		87Ge01 Z
	7465.5	0.4			-2.2	U			Bdn		06Fi.A
$^{173}\text{Yb}(\text{d,p})^{174}\text{Yb}$	5239	12	5240.038	0.013	0.1	U			Kop		66Bu16
	5229	5			2.2	U			Tal		66Sh14
$^{174}\text{Yb}(\text{d,t})^{173}\text{Yb}$	-1218	12	-1207.375	0.013	0.9	U			Kop		66Bu16

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{173}\text{Yb}(\alpha, t)^{174}\text{Lu} - ^{174}\text{Yb}(\alpha)^{175}\text{Lu}$	-202.1	1.0	-202.1	1.0	0.0	1	100	100	$^{174}\text{Lu}$	McM	75Bu02	
$^{174}\text{Tm}(\beta^-)^{174}\text{Yb}$	3080	100	3080	40	0.0	2					64Ka16 *	
	3080	50			0.0	2					67Gu12 *	
$^{174}\text{Lu}(\beta^+)^{174}\text{Yb}$	1402	5	1374.3	1.6	-5.5	B					68K108	
$^{174}\text{Lu}(\epsilon)^{174}\text{Yb}$	1370	7			0.6	U					68Li01 *	
$^{174}\text{Ta}(\beta^+)^{174}\text{Hf}$	3845	80	4104	28	3.2	B					71Ch26 *	
* $^{174}\text{Ir} - \text{u}$	$M - A = -30761(36)$ keV for mixture gs+m at 192(11) keV										Nub16b **	
* $^{174}\text{Ir}(\alpha)^{170}\text{Re}$	$E_\alpha = 5275(10)$ to $(3^+)$ level at 224.7 keV										Ens02b **	
* $^{174}\text{Ir}^m(\alpha)^{170}\text{Re}$	$E_\alpha = 5478(6)$ to $(7^+)$ level at 210.32 keV										Ens02b **	
* $^{174}\text{Ir}^m(\alpha)^{170}\text{Re}$	$E_\alpha = 5478(5)$ 5316(10) to $(7^+)$ at 210.32 and 370.1 level										Ens02b **	
* $^{174}\text{Au}(\alpha)^{170}\text{Ir}$	$E_\alpha = 6538$ correlated with $^{170}\text{Ir}$ $E_\alpha = 5817$ keV										02Ro17 **	
*	and only this with $^{178}\text{Tl}$ $\alpha$ 's										02Ro17 **	
* $^{174}\text{Au}^m(\alpha)^{170}\text{Ir}^m$	$E_\alpha = 6530(20)$ to level above 76 keV										84Sc.A **	
* $^{174}\text{Au}^m(\alpha)^{170}\text{Ir}^m$	$E_\alpha = 6626, 6470, 6435$ to $^{170}\text{Ir}^m$ and levels above $^{170}\text{Ir}^m$ $(9^+)$ at 152.5,										Ens082 **	
*	$(7^-, 8^-, 9^-)$ at 190.56; last two $E_\alpha$ originally assigned to $^{175}\text{Au}$										01Ko.B **	
* $^{174}\text{Tm}(\beta^-)^{174}\text{Yb}$	$E_{\beta^-} = 1200(100)$ 1200(50) respectively, to $5^-$ level at 1884.674 keV, and other $E_{\beta^-}$										Ens998 **	
* $^{174}\text{Lu}(\epsilon)^{174}\text{Yb}$	No K capture to $2^-$ level at 1318.361 keV $\rightarrow Q_i(1380)$ ; and L capture										Ens998 **	
*	of $^{174}\text{Lu}^m$ at 170.83 to $^{174}\text{Yb}^m$ at 1518.148 keV $\rightarrow Q_{gs} > 1357$ keV										Nub16b **	
* $^{174}\text{Ta}(\beta^+)^{174}\text{Hf}$	$E_{\beta^+} = 2525(80)$ to $4^+$ level at 297.38 keV										Ens04a **	
$^{175}\text{Lu} \text{ } ^{37}\text{Cl} - ^{142}\text{Nd} \text{ } ^{35}\text{Cl}_2$	61249.5	2.5	61245.6	1.8	-0.6	U			H31	2.5	77So02	
$\text{C}_{14} \text{H}_7 - ^{175}\text{Lu}$	114121	37	113997.9	1.3	-0.8	U			R04	4.0	64De15	
$\text{C}_{13} \text{ } ^{13}\text{C} \text{H}_6 - ^{175}\text{Lu}$	109763	36	109527.7	1.3	-1.6	U			R04	4.0	64De15	
$^{175}\text{Ta} - \text{u}$	-56350	120	-56260	30	0.7	U			GS1	1.0	00Ra23	
	-56263	30				2			GS2	1.0	05Li24	
$^{175}\text{W} - \text{u}$	-53290	104	-53280	30	0.1	U			GS1	1.0	00Ra23	
	-53283	30				2			GS2	1.0	05Li24	
$^{175}\text{Re} - \text{u}$	-48630	104	-48620	30	0.1	U			GS1	1.0	00Ra23	
	-48619	30				2			GS2	1.0	05Li24	
$^{175}\text{Os} - \text{u}$	-43120	110	-43055	13	0.6	U			GS1	1.0	00Ra23	
	-43024	30			-1.0	1	18	18	$^{175}\text{Os}$	GS2	1.0	05Li24
$^{175}\text{Ir} - \text{u}$	-34353	1288	-35850	13	-0.5	U				2.5	91Br17	
	-35828	30			-0.7	1	20	20	$^{175}\text{Ir}$	GS2	1.0	05Li24
$^{175}\text{Lu} \text{ } ^{35}\text{Cl} - ^{173}\text{Yb} \text{ } ^{37}\text{Cl}$	5503	4	5511.2	1.3	0.8	U			H23	2.5	70Wh01	
	5507.3	1.4			1.1	1	14	14	$^{175}\text{Lu}$	H27	2.5	74Ba90
$^{175}\text{Lu} \text{ O} - \text{C}_{16}$	-64316.3	4.5	-64308.1	1.3	1.2	U			TG1	1.5	11Ke03	
$^{175}\text{Ir}(\alpha)^{171}\text{Re}$	5709.0	5.	5430	30	-55.6	B					67Si02 *	
	5709.2	5.			-55.7	B					92Sc16 *	
$^{175}\text{Pt}(\alpha)^{171}\text{Os}$	6179	5	6164	4	-3.1	C					79Ha10 *	
	6178.1	3.			-4.8	B					82De11 *	
	6164	4			-0.1	1	100	91	$^{175}\text{Pt}$	Ora	14Pe02 *	
$^{175}\text{Au}(\alpha)^{171}\text{Ir}$	6562.3	15.	6583	4	1.4	U			Bka		02Ro17 *	
	6580.7	8.2			0.3	6			Ara		11Ko.B *	
	6583.7	4.			-0.2	6			Anv		13An10	
$^{175}\text{Au}^m(\alpha)^{171}\text{Ir}^m$	6590.9	10.	6583	3	-0.7	10			Ora		75Ca06	
	6775.8	10.			-19.3	F			GSa		84Sc.A *	
	6588.8	9.			-0.6	10			Daa		96Pa01	
	6579.6	6.			0.6	10			Ara		01Ko44	
	6582.7	5.			0.1	10			Anv		10An01	
	6581.7	8.2			0.2	10			Ara		11Ko.B *	
$^{175}\text{Hg}(\alpha)^{171}\text{Pt}$	7020.7	20.	7072	5	2.5	o			GSa		83Sc24	
	7039.2	20.			1.7	U			GSa		84Sc.A	
	7071.0	24.			0.1	o			Daa		96Pa01	
	7058.7	11.			1.2	8			Jya		97Uu01	
	7075	5			-0.5	8			Daa		09Od01	
	7082.1	20.			-0.5	U			Anv		10An01	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{174}\text{Yb}(n,\gamma)^{175}\text{Yb}$	5822.35	0.07	5822.35	0.07	0.1	1	100	100 $^{175}\text{Yb}$	MMn		82Is05 Z
	5822.5	0.4			-0.4	U			Bdn		06Fi.A
$^{174}\text{Yb}(d,p)^{175}\text{Yb}$	3595	12	3597.79	0.07	0.2	U			Kop		66Bu16
$^{174}\text{Yb}(\alpha,t)^{175}\text{Lu}$	-14303	10	-14303.8	1.2	-0.1	U			McM		75Bu02
$^{175}\text{Lu}(\gamma,n)^{174}\text{Lu}$	-7880	80	-7666.7	1.0	2.7	U			Phi		60Ge01
$^{175}\text{Lu}(d,t)^{174}\text{Lu}$	-1400	10	-1409.5	1.0	-0.9	U			Tal		70Jo08
$^{174}\text{Hf}(n,\gamma)^{175}\text{Hf}$	6708.4	0.5	6708.5	0.4	0.2	-					71Al01 Z
	6708.8	0.6			-0.5	-			Bdn		06Fi.A
ave.	6708.6	0.4			-0.2	1	100	86 $^{175}\text{Hf}$			average
$^{175}\text{Tm}(\beta^-)^{175}\text{Yb}$	2385	50				2					66Wi04 *
$^{175}\text{Yb}(\beta^-)^{175}\text{Lu}$	466	3	470.0	1.2	1.3	-					55De18
	468	5			0.4	-					55Mi90
	471	3			-0.3	-					56Co13
	467	3			1.0	-					62Ba32
ave.	468.0	1.6			1.2	1	54	54 $^{175}\text{Lu}$			average
$^{175}\text{Hf}(\epsilon)^{175}\text{Lu}$	628	9	683.9	2.0	6.2	B					68Ja11 *
	650	20			1.7	U					69Jo16 *
	630	3			18.0	B					88Si22 *
* $^{175}\text{Ir}(\alpha)^{171}\text{Re}$	$E_\alpha=5392.8(5,Z)$ to 189.8 level										
* $^{175}\text{Ir}(\alpha)^{171}\text{Re}$	$E_\alpha=5393(5)$ to 189.8 level										
* $^{175}\text{Pt}(\alpha)^{171}\text{Os}$	$E_\alpha=6037(10), 5963.0(5,Z)$ to ground state, 76.4(0.5) level										
* $^{175}\text{Pt}(\alpha)^{171}\text{Os}$	$E_\alpha=5959.2(3,Z)$ to 76.4(0.5) level										
* $^{175}\text{Pt}(\alpha)^{171}\text{Os}$	$E_\alpha=6021(4), 5948(4)$ to ground state, 76.7(0.3) level										
* $^{175}\text{Au}(\alpha)^{171}\text{Ir}$	Analysis by AHW of data in Fig. 3 of reference										
* $^{175}\text{Au}(\alpha)^{171}\text{Ir}$	Correlated with $E_\alpha=6556(8)$ of $^{179}\text{Tl}$ and $5728(8)$ of $^{171}\text{Ir}$										
* $^{175}\text{Au}^m(\alpha)^{171}\text{Ir}^m$	$E_\alpha=6435(10)$ and $6470(20)$ to 190.0 and 152.7 levels										
* $^{175}\text{Au}^m(\alpha)^{171}\text{Ir}^m$	F : reassigned to $^{174}\text{Au}$ !										
* $^{175}\text{Au}^m(\alpha)^{171}\text{Ir}^m$	Correlated with $E_\alpha=7194(8)$ of $^{179}\text{Tm}$ and $5958(8)$ of $^{171}\text{Ir}^m$										
*	different method and different detectors as compared to 01Ko44										
* $^{175}\text{Tm}(\beta^-)^{175}\text{Yb}$	$E_{\beta^-}=1870(50)$ to $1/2^-$ level at 514.866 keV										
* $^{175}\text{Hf}(\epsilon)^{175}\text{Lu}$	pK=0.712(0.008) 0.740(0.015) 0.714(0.002) respectively,										
*	to $7/2^+$ level at 432.74 keV, and other capture ratios, recalculated										
	Ens04a **										
	Ens04a **										
$\text{C}_{14}\text{H}_8-^{176}\text{Yb}$	119980	46	120025.549	0.016	0.2	U			R04	4.0	64De15
$\text{C}_{13}\text{H}_6\text{N}-^{176}\text{Yb}$	107190	110	107449.489	0.016	0.6	U			R04	4.0	64De15
$^{176}\text{Yb}-^{129}\text{Xe}_{1,364}$	72453.619	0.016	72453.614	0.014	-0.3	1	75	73 $^{176}\text{Yb}$	FS1	1.0	12Ra34
$^{176}\text{Yb}-^{132}\text{Xe}_{1,333}$	70335.958	0.027	70335.973	0.014	0.5	1	28	27 $^{176}\text{Yb}$	FS1	1.0	12Ra34
$^{176}\text{Lu}\text{ }^{37}\text{Cl}-^{143}\text{Nd}\text{ }^{35}\text{Cl}_2$	61067.2	1.4	61069.1	1.8	0.5	1	27	15 $^{143}\text{Nd}$	H31	2.5	77So02
$\text{C}_{14}\text{H}_8-^{176}\text{Lu}$	119962	49	119908.4	1.3	-0.3	U			R04	4.0	64De15
$^{176}\text{Lu}\text{O}-\text{C}_{16}$	-62394.1	7.6	-62393.6	1.3	0.0	U			TG1	1.5	11Ke03
$^{176}\text{Hf}\text{O}-\text{C}_{16}$	-63668.5	9.8	-63675.5	1.6	-0.5	U			TG1	1.5	11Ke03
$^{176}\text{Ta}-\text{u}$	-55143	33				2			GS2	1.0	05Li24
$^{176}\text{W}-\text{u}$	-54420	104	-54370	30	0.5	U			GS1	1.0	00Ra23
	-54366	30				2			GS2	1.0	05Li24
$^{176}\text{Re}-\text{u}$	-48380	110	-48380	30	0.0	U			GS1	1.0	00Ra23
	-48377	30				2			GS2	1.0	05Li24
$^{176}\text{Os}-\text{u}$	-45150	110	-45190	30	-0.4	U			GS1	1.0	00Ra23
	-45194	30				2			GS2	1.0	05Li24
$^{176}\text{Ir}-\text{u}$	-36328	30	-36370	18	-1.4	1	36	36 $^{176}\text{Ir}$	GS2	1.0	05Li24
$^{176}\text{Yb}\text{ }^{35}\text{Cl}_2-^{172}\text{Yb}\text{ }^{37}\text{Cl}_2$	12088.9	2.4	12088.27	0.14	-0.1	U			H27	2.5	74Ba90
$^{176}\text{Yb}\text{ }^{35}\text{Cl}-^{174}\text{Yb}\text{ }^{37}\text{Cl}$	6652	3	6657.27	0.07	0.7	U			H23	2.5	70Wh01
	6656.3	1.4			0.3	U			H27	2.5	74Ba90
$^{176}\text{Hf}\text{ }^{35}\text{Cl}-^{174}\text{Hf}\text{ }^{37}\text{Cl}$	4106	16	4311.5	1.9	5.1	B			H24	2.5	73Ba40
	4314.21	0.86			-1.2	1	76	74 $^{174}\text{Hf}$	H37	2.5	77Sh12
$^{176}\text{Lu}-^{175}\text{Lu}$	1980	60	1914.50	0.16	-0.3	U			R04	4.0	64De15



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{176}\text{Yb}-^{174}\text{Yb}$	4000	50	3707.160	0.017	-1.5	U			R04	4.0	64De15
$^{176}\text{Ir}(\alpha)^{172}\text{Re}$	5237.3	8.	5230	40	-0.1	1	59	54 $^{172}\text{Re}$			67Si02
$^{176}\text{Pt}(\alpha)^{172}\text{Os}$	5890.1	5.	5885.1	2.1	-1.0	-					79Ha10 Z
	5881.4	4.			0.9	-			Bka		82Bo04 Z
	5887.3	3.			-0.7	-			Ora		82De11 Z
	5874.8	8.			1.3	-			Daa		96Pa01
	ave.	5885.3	2.1		-0.1	1	99	66 $^{172}\text{Os}$			average
$^{176}\text{Au}(\alpha)^{172}\text{Ir}$	6574.2	10.	6433	7	-14.1	B			Ora		75Ca06 *
	6541.5	10.			-10.8	C			GSa		84Sc.A *
	6433.4	7.				5			Anv		14An10 *
$^{176}\text{Au}^m(\alpha)^{172}\text{Ir}^m$	6436.6	10.	6433	4	-0.3	5			Ora		75Ca06 *
	6428.4	10.			0.5	5			GSa		84Sc.A *
	6433.4	6.			0.0	5			Ara		01Ko44 *
	6434.4	7.			-0.1	5			Anv		14An10 *
$^{176}\text{Hg}(\alpha)^{172}\text{Pt}$	6907.2	20.	6897	6	-0.5	o			GSa		83Sc24
	6924.7	10.			-2.8	C			GSa		84Sc.A
	6907.3	20.			-0.5	U			Daa		96Pa01
	6897.0	6.			0.0	-			Ara		99Po09
	6917.5	15.			-1.3	-			Anv		09An20
	ave.	6900	6		-0.5	1	95	72 $^{176}\text{Hg}$			average
$^{176}\text{Yb}(p,\alpha)^{173}\text{Tm}$	7628.8	4.4				2			NDm		78Ta10
$^{176}\text{Yb}(p,t)^{174}\text{Yb}$	-4216	5	-4207.642	0.015	1.7	U			Min		73Oo01
$^{176}\text{Hf}(p,t)^{174}\text{Hf}$	-6397	5	-6392.7	1.7	0.9	1	12	12 $^{174}\text{Hf}$	Min		73Oo01
$^{176}\text{Yb}(d,t)^{175}\text{Yb}$	-621	12	-609.85	0.07	0.9	U			Kop		66Bu16
$^{175}\text{Lu}(n,\gamma)^{176}\text{Lu}$	6293.2	1.2	6287.97	0.15	-4.4	B					70Wa20
	6287.96	0.15			0.1	1	100	79 $^{176}\text{Lu}$	ILn		91Kl02 Z
	6289.78	0.24			-7.5	C			Bdn		06Fi.A
$^{175}\text{Lu}(d,p)^{176}\text{Lu}$	4070	8	4063.41	0.15	-0.8	U			Tal		67St14
$^{176}\text{Lu}(d,t)^{175}\text{Lu}$	-25	15	-30.74	0.15	-0.4	U			Tal		71Mi01
$^{176}\text{Hf}(d,t)^{175}\text{Hf}$	-1925	8	-1908.7	1.8	2.0	U			Tal		73Za08
$^{176}\text{Tl}(p)^{175}\text{Hg}$	1265.2	18.				9			Jyp		04Ke06
$^{176}\text{Tm}(\beta^-)^{176}\text{Yb}$	4120	100				2					67Gu11 *
$^{176}\text{Lu}(\beta^-)^{176}\text{Hf}$	1162	25	1194.1	0.9	1.3	U					69Pr11 *
	1194.1	1.0			0.0	1	76	75 $^{176}\text{Hf}$			73Va11 *
$^{176}\text{Ta}(\beta^+)^{176}\text{Hf}$	3100	90	3210	30	1.2	U					71Be10 *
$^{176}\text{Au}(\alpha)^{172}\text{Ir}$	$E_\alpha=6260(10)$ coinc. with $E(\gamma)=168.4(0.5)$ keV										
$^{176}\text{Au}(\alpha)^{172}\text{Ir}$	$E_\alpha=6228(10)$ to $168.4(0.5)$ $\gamma$										
$^{176}\text{Au}(\alpha)^{172}\text{Ir}$	$E_\alpha=6260$ correlated with $^{172}\text{Ir}$ $E_\alpha=5510$ keV										
$^{176}\text{Au}(\alpha)^{172}\text{Ir}$	and $E_\alpha=6157(20),6138(15),6054(20),5798(20)$ to 126.3, 151.5, 236.6,500.0										
$^{176}\text{Au}^m(\alpha)^{172}\text{Ir}^m$	$E_\alpha=6286$ correlated with $^{172}\text{Ir}^m$ $E_\alpha=5828$ keV										
$^{176}\text{Au}^m(\alpha)^{172}\text{Ir}^m$	$E_\alpha=6119+E(\gamma)=175.1$ was misassigned to $^{177}\text{Au}$										
$^{176}\text{Au}^m(\alpha)^{172}\text{Ir}^m$	$E_\alpha=6115(6)$ coinc. with 175.1 $\gamma$ of reference										
$^{176}\text{Au}^m(\alpha)^{172}\text{Ir}^m$	$E_\alpha=6287(7), 6117(7), 6082(7)$ to 0, 175.2, 211.6 above $^{172}\text{Ir}^m$										
$^{176}\text{Tm}(\beta^-)^{176}\text{Yb}$	$E_{\beta^-}=2000(100), 1150(100)$ to $(3^+,4^+)$ level at 2053.34, $(3^+,4^+,5^+)$ 3052.2										
$^{176}\text{Lu}(\beta^-)^{176}\text{Hf}$	$E_{\beta^-}=565(25)$ to $6^+$ level at 596.82 keV										
$^{176}\text{Lu}(\beta^-)^{176}\text{Hf}$	$Q_{\beta^-}=1317(1)$ from $^{176}\text{Lu}^m$ at 122.845 keV										
$^{176}\text{Ta}(\beta^+)^{176}\text{Hf}$	KLM/ $\beta^+=119(50)$ to $2^-$ level at 1247.70 keV, $1^+$ level at 2994 keV										
$^{177}\text{Ta}-u$	-55559	30	-55518	4	1.4	U			GS2	1.0	05Li24
$^{177}\text{W}-u$	-53420	110	-53360	30	0.6	U			GS1	1.0	00Ra23
	-53357	30				2			GS2	1.0	05Li24
$^{177}\text{Re}-u$	-49620	104	-49670	30	-0.5	U			GS1	1.0	00Ra23
	-49672	30				2			GS2	1.0	05Li24
$^{177}\text{Os}-u$	-45020	104	-45042	16	-0.2	U			GS1	1.0	00Ra23
	-45012	30			-1.0	R			GS2	1.0	05Li24

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{177}\text{Ir}-u$	-38810	110	-38699	21	1.0	U			GS1	1.0	00Ra23
	-38699	30			0.0	2			GS2	1.0	05Li24
$^{177}\text{Pt}-u$	-31545	30	-31530	16	0.5	1	29	29 $^{177}\text{Pt}$	GS2	1.0	05Li24
$^{177}\text{Hf O}-\text{C}_{16}$	-61845.2	7.2	-61855.1	1.5	-0.9	U			TG1	1.5	11Ke03
$^{177}\text{Ir}(\alpha)^{173}\text{Re}$	5127.1	10.	5080	30	-0.9	F					67Si02 *
$^{177}\text{Pt}(\alpha)^{173}\text{Os}$	5654.6	6.	5642.9	2.7	-1.9	-					79Ha10 Z
	5640.5	3.1			0.8	-			Bka		82Bo04 Z
	ave.	5643.3	2.7		-0.2	1	99	55 $^{177}\text{Pt}$			average
$^{177}\text{Au}(\alpha)^{173}\text{Ir}$	6292.5	10.	6298	4	0.6	-			Daa		75Ca06
	6292.5	20.			0.3	U			GSa		84Sc.A
	6296.5	10.			0.2	-			Daa		96Pa01
	6298.6	6.			0.0	-			Ara		01Ko44
	6303.7	7.			-0.7	-			Anv		09An14
	ave.	6299	4		-0.1	1	99	86 $^{173}\text{Ir}$			average
$^{177}\text{Au}^m(\alpha)^{173}\text{Ir}^m$	6251.5	10.	6262	4	1.0	3			Ora		75Ca06
	6260.8	10.			0.1	3			GSa		84Sc.A *
	6259.7	9.			0.2	3			Daa		96Pa01 *
	6263.8	6.			-0.3	3			Ara		01Ko44
	6265.8	7.			-0.6	3			Anv		09An14
$^{177}\text{Hg}(\alpha)^{173}\text{Pt}$	6732.4	8.	6740	50	0.1	4					79Ha10
	6747.8	10.			-0.2	4					91Ko.A
	6729.4	9.2			0.1	4			Daa		96Pa01
	6734.5	15.			0.0	4			Anv		09An20
$^{177}\text{Tl}(\alpha)^{173}\text{Au}$	7067.0	7.				3			Ara		99Po09
$^{177}\text{Tl}^m(\alpha)^{173}\text{Au}^m$	7660.4	13.	7660	9	0.0	-			Ara		99Po09
	7645.1	13.			1.2	-			Jya		04Ke06
	ave.	7653	9		0.8	1	86	48 $^{173}\text{Au}^m$			average
$^{177}\text{Hf}(p,t)^{175}\text{Hf}$	-6071	5	-6059.8	2.0	2.2	1	16	14 $^{175}\text{Hf}$	Min		73Oo01
$^{176}\text{Yb}(n,\gamma)^{177}\text{Yb}$	5565.1	1.0	5566.40	0.22	1.3	U					72Al19 Z
	5566.40	0.22				2			Bdn		06Fi.A
$^{176}\text{Yb}(d,p)^{177}\text{Yb}$	3340	16	3341.83	0.22	0.1	U			Tal		63Ve09
	3337	12			0.4	U			Kop		66Bu16
$^{176}\text{Yb}(\alpha,t)^{177}\text{Lu}-^{174}\text{Yb}()^{175}\text{Lu}$	674.1	1.0	671.42	0.22	-2.7	U			McM		75Bu02
$^{176}\text{Lu}(n,\gamma)^{177}\text{Lu}$	7071.2	0.4	7072.89	0.16	4.2	B					71Ma45 Z
	7073.1	0.4			-0.5	-					72Mi16 Z
	7072.85	0.17			0.2	-			Bdn		06Fi.A
$^{176}\text{Lu}(d,p)^{177}\text{Lu}$	4843	10	4848.32	0.16	0.5	U			Tal		71Mi01
$^{176}\text{Lu}(n,\gamma)^{177}\text{Lu}$	ave.	7072.89	0.16	7072.89	0.16	0.0	1	99	92 $^{177}\text{Lu}$		average
$^{176}\text{Hf}(n,\gamma)^{177}\text{Hf}$	6385.8	0.8	6375.6	1.0	-12.7	C			Bdn		06Fi.A
$^{177}\text{Hf}(\gamma,n)^{176}\text{Hf}$	-6400	30	-6375.6	1.0	0.8	U			Phi		60Ge01
$^{176}\text{Hf}(d,p)^{177}\text{Hf}$	4150	7	4151.0	1.0	0.1	U			Tal		68Ri07
$^{177}\text{Hf}(d,t)^{176}\text{Hf}$	-127	11	-118.4	1.0	0.8	U			Tal		72Za04
$^{177}\text{Tl}(p)^{176}\text{Hg}$	1162.6	20.	1155	19	-0.4	o			Arp		99Po09 *
$^{177}\text{Tl}^m(p)^{176}\text{Hg}$	1969.2	10.	1962	7	-0.7	-			Arp		99Po09
	1965.2	12.			-0.2	-			Jyp		04Ke06
	ave.	1968	8		-0.7	1	90	62 $^{177}\text{Tl}^m$			average
$^{177}\text{Yb}(\beta^-)^{177}\text{Lu}$	1400	20	1397.4	1.2	-0.1	U					64Jo03
$^{177}\text{Lu}(\beta^-)^{177}\text{Hf}$	497	2	496.8	0.8	-0.1	-					55Ma12
	496.4	1.0			0.4	-					62El02 *
	ave.	496.5	0.9		0.3	1	78	70 $^{177}\text{Hf}$			average
$^{177}\text{Ta}(\beta^+)^{177}\text{Hf}$	1166	3				2					61We11
$^{177}\text{Au}^m(\text{IT})^{177}\text{Au}$	210	30	189	8	-0.7	o					01Ko44 *
$^{177}\text{Tl}^m(\text{IT})^{177}\text{Tl}$	807	18				2					99Po09
* $^{177}\text{Ir}(\alpha)^{173}\text{Re}$	F : final state uncertain: possibly to $5/2^-$ level at 214.7 keV										
* $^{177}\text{Au}^m(\alpha)^{173}\text{Ir}^m$	Followed by a $175.1(0.5) \gamma$										
*	$\gamma$ associated with $E_\alpha=6116$ keV from $^{176}\text{Au}$										
*	yet $E_\alpha=6118$ correlated with $E_\alpha=5672$ of $^{173}\text{Ir}^m$										
* $^{177}\text{Au}^m(\alpha)^{173}\text{Ir}^m$	$E_\alpha$ correlated with $^{173}\text{Ir} E_\alpha=5681(13)$ ; also with $^{181}\text{Tl} E_\alpha=6180$ keV										
*	evaluator doubts about correctness of latter remark										
* $^{177}\text{Tl}(p)^{176}\text{Hg}$	Replaced by $^{177}\text{Tl}^m(\text{IT})$										
* $^{177}\text{Lu}(\beta^-)^{177}\text{Hf}$	$E_{\beta^-}=384(2), 175(1)$ to $112.95, 321.32$ levels										
* $^{177}\text{Au}^m(\text{IT})^{177}\text{Au}$	Authors say $157.9+x$ , $x$ estimated by evaluator										
											AHW **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
*	x is better known from $^{181}\text{Tl}^m(\text{IT})^{181}\text{Tl}$ combined with $Q_\alpha$										09An14 **
$^{178}\text{W}-\text{u}$	-54152	30	-54114	16	1.3	U			GS2	1.0	05Li24
$^{178}\text{Re}-\text{u}$	-48800	110	-49010	30	-1.9	U			GS1	1.0	00Ra23
			-49011	30		2			GS2	1.0	05Li24
$^{178}\text{Os}-\text{u}$	-46790	104	-46747	15	0.4	U			GS1	1.0	00Ra23
			-46710	30	-1.2	1	24	24 $^{178}\text{Os}$	GS2	1.0	05Li24
$^{178}\text{Ir}-\text{u}$	-38950	110	-38918	21	0.3	U			GS1	1.0	00Ra23
			-38888	30	-1.0	2			GS2	1.0	05Li24
$^{178}\text{Pt}-\text{u}$	-34783	1181	-34351	11	0.1	U				2.5	91Br17
			-34300	110	-0.5	U			GS1	1.0	00Ra23
			-34333	30	-0.6	1	13	13 $^{178}\text{Pt}$	GS2	1.0	05Li24
$^{178}\text{Au}-^{133}\text{Cs}_{1.338}$	102562	11	102561	11	-0.1	1	97	97 $^{178}\text{Au}$	MA8	1.0	17Ma.A
$^{178}\text{Au}^m-^{133}\text{Cs}_{1.338}$	102764	11				2			MA8	1.0	17Ma.A
$^{178}\text{Hf } ^{35}\text{Cl}-^{176}\text{Hf } ^{37}\text{Cl}$	5236	5	5248.7	1.1	1.0	U			H23	2.5	70Wh01
	5239.5	1.3			2.8	U			H27	2.5	74Ba90
$^{178}\text{Hf O}-\text{C}_{16}$	-61364.8	7.9	-61376.9	1.5	-1.0	U			TG1	1.5	11Ke03
$^{178}\text{Pt}-^{175}\text{Ir}$	-472	1052	1500	17	0.7	U				2.5	91Br17
$^{178}\text{Pt}(\alpha)^{174}\text{Os}$	5583.3	5.	5573.0	2.2	-2.0	-					79Ha10 Z
	5569.9	3.			1.0	-			Bka		82Bo04 Z
	5568.4	13.			0.4	U			Lvn		94Wa23
	5572.4	4.			0.1	-			Ara		00Ko16 *
ave.	5573.2	2.2			-0.1	1	99	75 $^{174}\text{Os}$			average
$^{178}\text{Au}(\alpha)^{174}\text{Ir}$	6056.4	10.	6135	25	1.6	F					68Si01 *
	6117.7	20.			0.3	1	26	23 $^{174}\text{Ir}$	GSa		86Ke03
$^{178}\text{Hg}(\alpha)^{174}\text{Pt}$	6578.1	6.	6577.3	3.0	-0.1	3					79Ha10
	6576.1	9.			0.1	3			Daa		96Pa01
	6577.1	4.			0.1	3			Ara		00Ko48
	6578.1	8.			-0.1	3			Anv		09An14
$^{178}\text{Tl}(\alpha)^{174}\text{Au}$	7017.0	5.	7020	10	0.6	o			Bka		02Ro17 *
	7020.0	10.				8			Bka		13Li49 *
$^{178}\text{Pb}(\alpha)^{174}\text{Hg}$	7790.4	14.				3			Bka		01Ro.B
$^{178}\text{Pt}(\text{p},\alpha)^{175}\text{Ir}$	4420	980	6261	16	1.9	U					91Br17
$^{176}\text{Yb}(\text{t},\text{p})^{178}\text{Yb}$	3865	10				2			Phi		82Zu02
$^{176}\text{Lu}(\text{t},\text{p})^{178}\text{Lu}^m$	4482	5	4492.6	2.9	2.1	1	34	34 $^{178}\text{Lu}^m$	LAl		81Gi01
$^{178}\text{Hf}(\text{p},\text{t})^{176}\text{Hf}$	-5531	5	-5519.8	1.0	2.2	U			Min		73Oo01
$^{177}\text{Hf}(\text{n},\gamma)^{178}\text{Hf}$	7625	1	7625.94	0.18	0.9	U					69Fa01
	7624.4	1.5			1.0	U					77St10
	7626.2	0.3			-0.9	-			ILn		86Ha22 Z
	7625.80	0.22			0.6	-			Bdn		06Fi.A
$^{178}\text{Hf}(\text{d},\text{t})^{177}\text{Hf}$	-1364	9	-1368.71	0.18	-0.5	U			Tal		68Ri07
$^{177}\text{Hf}(\text{n},\gamma)^{178}\text{Hf}$	ave.	7625.94	0.18	7625.94	0.18	0.0	1	99	71 $^{178}\text{Hf}$		average
$^{178}\text{Yb}(\beta^-)^{178}\text{Lu}$	641	30	642	10	0.0	U					73Or03 *
$^{178}\text{Lu}^m(\text{IT})^{178}\text{Lu}$	120	3	123.8	2.6	1.3	1	76	66 $^{178}\text{Lu}^m$	McM		93Bu02
$^{178}\text{Lu}(\beta^-)^{178}\text{Hf}$	2046	50	2097.5	2.1	1.0	U					73Or03 *
	2117	30			-0.7	U					75Ka15 *
$^{178}\text{Ta}^m(\beta^+)^{178}\text{Hf}$	1937	15				2					61Ga05 *
$^{178}\text{W}(\epsilon)^{178}\text{Ta}^m$	91.3	2.				3					67Ni02
$^{178}\text{Re}(\beta^+)^{178}\text{W}$	4660	180	4750	30	0.5	U					70Go20 *
* $^{178}\text{Pt}(\alpha)^{174}\text{Os}$	Also $E_\alpha=5289(8)$ keV to $2^+$ 158.601 level (not used)										GAu **
* $^{178}\text{Au}(\alpha)^{174}\text{Ir}$	F : higher $E_\alpha$ branch seen in reference										86Ke03 **
* $^{178}\text{Tl}(\alpha)^{174}\text{Au}$	And a stronger $E_\alpha=6704$ ; both correlated with $^{174}\text{Au}$ $E_\alpha=6538$ keV										02Ro17 **
* $^{178}\text{Tl}(\alpha)^{174}\text{Au}$	Also 6693(10) and 6595(10) to 173.0, 273.0 levels										13Li49 **
* $^{178}\text{Yb}(\beta^-)^{178}\text{Lu}$	$E_{\beta^-}=250(30)$ to $1^+$ level at 390.8 keV										Ens097 **
* $^{178}\text{Lu}(\beta^-)^{178}\text{Hf}$	$E_{\beta^-}=2000(50)$ to ground state and 50% to $2^+$ level at 93.18 keV										Ens097 **
* $^{178}\text{Lu}(\beta^-)^{178}\text{Hf}$	$E_{\beta^-}=2050(50)$ to ground state and 50% to $2^+$ level at 93.18 keV and										Ens097 **
*	$E_{\beta^-}=770(30)$ from $^{178}\text{Lu}^m$ at 123.8(2.6) to $8^-$ level 1479.025 keV										Nub16b **
* $^{178}\text{Ta}^m(\beta^+)^{178}\text{Hf}$	$E_{\beta^+}=890(10)$ to ground state and $2^+$ level at 93.18 keV, ratio 2.7 to 1										Ens097 **
* $^{178}\text{Re}(\beta^+)^{178}\text{W}$	$E_{\beta^+}=3300(180)$ to $4^+$ level at 342.74 keV										Ens097 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{179}\text{Lu}-^{85}\text{Rb}_{2.106}$	132997	21	133104	6	5.1	C			MA8	1.0	13Ro.A
$\text{C}_{14}\text{H}_{11}-^{179}\text{Hf}$	140260.3	1.8	140249.5	1.5	-1.5	U			M23	4.0	79Ha32
$^{179}\text{W}-u$	-52964	76	-52920	16	0.6	U			GS2	1.0	05Li24 *
$^{179}\text{Re}-u$	-50010	30	-50010	26	0.0	1	78	78 $^{179}\text{Re}$	GS2	1.0	05Li24
$^{179}\text{Os}-u$	-46220	104	-46183	18	0.4	U			GS1	1.0	00Ra23
	-46176	30			-0.2	1	35	35 $^{179}\text{Os}$	GS2	1.0	05Li24
$^{179}\text{Ir}-u$	-40910	104	-40882	10	0.3	U			GS1	1.0	00Ra23
	-40852	30			-1.0	1	12	12 $^{179}\text{Ir}$	GS2	1.0	05Li24
$^{179}\text{Pt}-u$	-34710	110	-34641	9	0.6	U			GS1	1.0	00Ra23
	-34625	30			-0.5	U			GS2	1.0	05Li24
$^{179}\text{Au}-u$	-26811	31	-26826	13	-0.5	1	16	16 $^{179}\text{Au}$	GS2	1.0	05Li24
$^{179}\text{Hg}-^{208}\text{Pb}_{.861}$	1900	34	1930	29	0.9	1	74	74 $^{179}\text{Hg}$	MA6	1.0	01Sc41
$^{179}\text{Hf}^{35}\text{Cl}-^{177}\text{Hf}^{37}\text{Cl}$	5539	3	5545.63	0.22	0.9	U			H23	2.5	70Wh01
	5544.4	0.7			0.7	U			H27	2.5	74Ba90
$^{179}\text{Hf O}-\text{C}_{16}$	-59261.8	6.5	-59259.5	1.5	0.2	U			TG1	1.5	11Ke03
$^{179}\text{Pt}(\alpha)^{175}\text{Os}$	5370	10	5412	9	4.2	F					66Si08 *
	5416	10			-0.4	1	89	82 $^{175}\text{Os}$			79Ha10 *
	5382	3			10.1	F			Bka		82Bo04 *
$^{179}\text{Au}(\alpha)^{175}\text{Ir}$	5981.8	5.	5981	5	-0.1	1	97	80 $^{175}\text{Ir}$			68Si01
	5986.9	15.			-0.4	U			Jya		04Ra28 *
$^{179}\text{Hg}(\alpha)^{175}\text{Pt}$	6431.0	5.	6360	30	-1.4	-			ISa		79Ha10 Z
	6418.7	9.			-1.2	-			Daa		96Pa01
	6430.0	4.			-1.4	-			Ara		02Ko09
	ave.	6429.1	3.0		-1.4	1	35	26 $^{179}\text{Hg}$			average
$^{179}\text{Tl}(\alpha)^{175}\text{Au}$	6710.2	20.	6711	3	0.0	U			GSa		83Sc24
	6718.4	18.			-0.4	U			Daa		96Pa01
	6719.4	10.			-0.9	7			Ara		98To14
	6706.1	8.			0.5	7			Ara		11Ko.B
	6710.2	4.			0.1	7			Anv		13An10
$^{179}\text{Tl}^m(\alpha)^{175}\text{Au}^m$	7364.5	20.	7368	4	0.2	U			GSa		83Sc24
	7366.0	20.			0.1	U			Daa		96Pa01
	7378.1	10.			-1.0	o			Ara		98To14
	7372.0	5.1			-0.7	9			Anv		10An01
	7358.7	8.2			1.2	9			Ara		11Ko.B
$^{179}\text{Pb}(\alpha)^{175}\text{Hg}$	7598.3	20.				9			Anv		10An01 *
$^{179}\text{Hf}(p,t)^{177}\text{Hf}$	-5249	5	-5243.13	0.19	1.2	U			Min		73Oo01
$^{179}\text{Hf}(t,\alpha)^{178}\text{Lu}-^{178}\text{Hf}(\gamma)^{177}\text{Lu}$	-72	2	-73.7	1.9	-0.8	1	89	89 $^{178}\text{Lu}$	McM		93Bu02
$^{178}\text{Hf}(n,\gamma)^{179}\text{Hf}$	6099.02	0.10	6098.99	0.08	-0.3	-			ILn		89Ri03 Z
	6098.95	0.12			0.3	-			Bdn		06Fi.A
$^{179}\text{Hf}(\gamma,n)^{178}\text{Hf}$	-6000	70	-6098.99	0.08	-1.4	U			Phi		60Ge01
$^{178}\text{Hf}(d,p)^{179}\text{Hf}$	3877	14	3874.42	0.08	-0.2	U			Tal		63Ve09
$^{178}\text{Hf}(n,\gamma)^{179}\text{Hf}$	ave.	6098.99	0.08	6098.99	0.08	0.0	1	100	70 $^{179}\text{Hf}$		average
$^{179}\text{Lu}(\beta^-)^{179}\text{Hf}$	1350	50	1404	5	1.1	U					61Ku10
	1380	70			0.3	U					63St06
$^{179}\text{Ta}(\epsilon)^{179}\text{Hf}$	129	16	105.6	0.4	-1.5	U					61Jo15 *
	105.61	0.41			-0.1	1	99	93 $^{179}\text{Ta}$			01Hi06
$^{179}\text{Re}(\beta^+)^{179}\text{W}$	2710	50	2711	27	0.0	1	29	22 $^{179}\text{Re}$			75Me20 *
* $^{179}\text{W}-u$	$M-A=-49225(29)$ keV for mixture gs+m at 221.91 keV										
* $^{179}\text{Pt}(\alpha)^{175}\text{Os}$	F : part of double line (with $^{180}\text{Pt}$ )										
* $^{179}\text{Pt}(\alpha)^{175}\text{Os}$	$E_\alpha=5150(10)$ $5195(10)$ $5161(3)$ respectively, to $1/2^-$ level at 102.3 keV, recalibrated										
* $^{179}\text{Au}(\alpha)^{175}\text{Ir}$	$E_\alpha=5853(15)$ , $5810(15)$ to ground state, 49 keV level										
* $^{179}\text{Pb}(\alpha)^{175}\text{Hg}$	$E_\alpha=7350(20)$ to 80 keV level										
* $^{179}\text{Ta}(\epsilon)^{179}\text{Hf}$	As corrected in reference										
* $^{179}\text{Re}(\beta^+)^{179}\text{W}$	$E_{\beta^+}=950(50)$ to $3/2^+$ level at 720.18 and $5/2^+$ at 773.65 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$C_{14} H_{12} -^{180}Hf$	147356.6	4.8	147340.7	1.5	-0.8	U			M23	4.0	79Ha32	
$^{180}W-u$	-53299	30	-53286.6	1.5	0.4	U			GS2	1.0	05Li24	
$^{180}Re-u$	-49209	30	-49208	23	0.0	2			GS2	1.0	05Li24	
$^{180}Os-u$	-47650	104	-47620	18	0.3	U			GS1	1.0	00Ra23	
	-47626	30			0.2	1	34	34	$^{180}Os$	GS2	1.0	05Li24
$^{180}Ir-u$	-40800	104	-40771	23	0.3	U			GS1	1.0	00Ra23	
	-40765	30			-0.2	2			GS2	1.0	05Li24	
$^{180}Pt-u$	-36900	104	-36968	12	-0.7	U			GS1	1.0	00Ra23	
	-36918	30			-1.7	R			GS2	1.0	05Li24	
$^{180}Au-u$	-27496	30	-27510	5	-0.5	U			GS2	1.0	05Li24	
$^{180}Au-^{133}Cs_{1.353}$	100411.5	5.3	100413	5	0.4	1	94	94	$^{180}Au$	MA8	1.0	16Ma.1
$^{180}Hg-^{208}Pb_{.865}$	-1569	22	-1544	14	1.2	1	38	38	$^{180}Hg$	MA6	1.0	01Sc41
$^{180}Hf-^{35}Cl_2-^{176}Hf-^{37}Cl_2$	11036.1	3.0	11050.0	1.1	1.9	U			H27	2.5	74Ba90	
$^{180}Hf-^{35}Cl-^{178}Hf-^{37}Cl$	5797	3	5801.32	0.19	0.6	U			H23	2.5	70Wh01	
	5798.4	0.7			1.7	U			H27	2.5	74Ba90	
$^{180}W-^{180}Hf$	153.73	0.30	153.77	0.30	0.1	1	98	82	$^{180}W$	SH1	1.0	12Dr01
$^{180}Hf-^{179}Hf$	730.8	4.7	733.83	0.16	0.3	U			M24	2.5	79Ha32	
$^{180}Hf O-C_{16}$	-58524.5	6.5	-58525.7	1.5	-0.1	U			TG1	1.5	11Ke03	
$^{180}W(\alpha)^{176}Hf$	2516.4	1.6	2515.3	1.0	-0.7	1	41	23	$^{176}Hf$			04Co26
$^{180}Pt(\alpha)^{176}Os$	5257.1	10.	5240	30	-2.0	F						66Si08 *
	5279	3			-14.0	F						Bka 82Bo04 *
$^{180}Au(\alpha)^{176}Ir$	5845	30	5828	17	-0.6	-			GSa			86Ke03 *
	5857	30			-1.0	-			Lvn			93Wa03 *
	ave.	5851	21		-1.1	1	61	59	$^{176}Ir$			average
$^{180}Hg(\alpha)^{176}Pt$	6258.3	5.	6258.5	2.4	0.0	-			ISa			79Ha10 Z
	6259.5	5.			-0.2	-			Lvn			93Wa03 *
	6258.3	4.			0.0	-			Ara			00Ko48
	6259.3	5.			-0.2	-			Anv			03An27
	ave.	6258.8	2.4		-0.1	1	99	66	$^{176}Pt$			average
$^{180}Tl(\alpha)^{176}Au$	6709.4	10.				6			Ara			98To14 *
$^{180}Pb(\alpha)^{176}Hg$	7394.6	40.	7419	5	0.6	U			ORa			96To08
	7415.1	15.			0.2	2			Ara			99To11
	7419.2	10.			-0.1	2			Anv			09An20
	7419.2	7.			-0.1	2			Jya			10Ra12
$^{180}Hf(p,t)^{178}Hf$	-5011	5	-5004.95	0.17	1.2	U			Min			73Oo01
$^{180}Hf(t,\alpha)^{179}Lu-^{178}Hf(^{177}Lu)$	-669	5	-669	5	0.0	1	100	100	$^{179}Lu$	McM		92Bu12
$^{179}Hf(n,\gamma)^{180}Hf$	7387.3	0.4	7387.76	0.15	1.1	-						74Bu22 Z
	7387.8	0.6			-0.1	-						90Bo52 Z
	7387.85	0.17			-0.5	-						06Fi.A
$^{180}Hf(\gamma,n)^{179}Hf$	-7470	110	-7387.76	0.15	0.7	U			Phi			60Ge01
$^{179}Hf(d,p)^{180}Hf$	5167	7	5163.19	0.15	-0.5	U			Tal			72Za04
$^{180}Hf(d,t)^{179}Hf$	-1112	4	-1130.53	0.15	-4.6	B			Tal			68Ri07
$^{179}Hf(n,\gamma)^{180}Hf$	ave.	7387.77	0.15	7387.76	0.15	-0.1	1	99	84	$^{180}Hf$		average
$^{180}W(d,t)^{179}W$	-2155	15	-2155	15	0.0	1	94	94	$^{179}W$	Kop		72Ca01
$^{180}Lu(\beta^-)^{180}Hf$	3148	100	3100	70	-0.4	2						71Gu02 *
	3058	100			0.5	2						71Sw01 *
$^{180}Ta(\beta^-)^{180}W$	705	15	703.2	2.3	-0.1	U						51Br87
	712	15			-0.6	U						62Ga07
$^{180}Re(\beta^+)^{180}W$	3830	60	3799	21	-0.5	R						67Go22 *
	3790	40			0.2	R						67Ho12 *
* $^{180}Pt(\alpha)^{176}Os$	F : part of double line (with $^{179}Pt$ ); $E_\alpha=5140(10)$ keV										AHW **	
* $^{180}Pt(\alpha)^{176}Os$	F : part of double line (with $^{179}Pt$ )										AHW **	
* $^{180}Au(\alpha)^{176}Ir$	$E_\alpha=5685(10)$ to $40(30)$ level										93Wa03 **	
* $^{180}Au(\alpha)^{176}Ir$	$E_\alpha=5647(10,Z)$ to $80(30)$ level										93Wa03 **	
* $^{180}Hg(\alpha)^{176}Pt$	$E_\alpha=6120$ $5862$ $5689(5)$ to ground state, $2^+$ level at $264.0$ , $0^+$ at $443$ keV										Ens062 **	
* $^{180}Tl(\alpha)^{176}Au$	Highest $E_\alpha$ ; not necessarily ground state to ground state										98To14 **	
* $^{180}Lu(\beta^-)^{180}Hf$	$E_{\beta^-}=1540(100)$ $1450(100)$ respectively, to $4^+$ level at $1607.67$ keV										Ens156 **	
* $^{180}Re(\beta^+)^{180}W$	$E_{\beta^+}=1800(60)$ $1760(40)$ respectively, to $2^-$ level $1006.381$ keV										Ens156 **	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{181}\text{Lu}-u$	-48092	135				2			GS3	1.0	13Sh30
$^{181}\text{Ta O}-^{133}\text{Cs}_{1,481}$	82943.7	3.6	82939.6	1.5	-1.1	1	18	18 $^{181}\text{Ta}$	MA8	1.0	16We.A
$^{181}\text{Ta O}-^{202}\text{Tl}_{975}$	-29891.4	2.5	-29892.4	1.8	-0.4	1	53	31 $^{202}\text{Tl}$	MA8	1.0	16We.A
$^{181}\text{Re}-u$	-49915	30	-49938	13	-0.8	R			GS2	1.0	05Li24
$^{181}\text{Os}-u$	-46670	110	-46753	27	-0.8	U			GS1	1.0	00Ra23 *
	-46756	34			0.1	1	64	64 $^{181}\text{Os}$	GS2	1.0	05Li24 *
$^{181}\text{Ir}-u$	-42330	104	-42365	6	-0.3	U			GS1	1.0	00Ra23
	-42372	30			0.2	U			GS2	1.0	05Li24
$^{181}\text{Pt}-u$	-36880	104	-36910	15	-0.3	U			GS1	1.0	00Ra23
	-36900	30			-0.3	-			GS2	1.0	05Li24
	ave.	-36890	21		-0.9	1	48	48 $^{181}\text{Pt}$			average
$^{181}\text{Au}-u$	-30030	110	-29921	21	1.0	U			GS1	1.0	00Ra23
	-29920	30			0.0	R			GS2	1.0	05Li24
$^{181}\text{Hg}-^{208}\text{Pb}_{870}$	-1929	40	-1868	17	1.5	1	17	17 $^{181}\text{Hg}$	MA6	1.0	01Sc41
$^{181}\text{Tl}-^{133}\text{Cs}_{1,361}$	114936	11	114940	10	0.4	1	79	79 $^{181}\text{Tl}$	MA8	1.0	08We02
$^{181}\text{Ta }^{35}\text{Cl}-^{179}\text{Hf }^{37}\text{Cl}$	5128.6	2.1	5123.6	2.0	-1.0	1	14	7 $^{179}\text{Hf}$	H35	2.5	80Sh06
$^{181}\text{Ta }^{17}\text{O }^{35}\text{Cl}-^{180}\text{Ta}^m \text{O }^{37}\text{Cl}$	7572	21	7617.31	0.21	0.9	U			H35	2.5	80Sh06
$^{181}\text{Pt}(\alpha)^{177}\text{Os}$	5133.7	20.	5150	5	0.8	U					66Si08
	5150.1	5.				2			ORa		95Bi01
$^{181}\text{Au}(\alpha)^{177}\text{Ir}$	5750.1	5.	5751.4	2.9	0.2	3					68Si01 Z
	5751.9	5.			-0.1	3					79Ha10 Z
	5735	4			4.1	F			IRa		92Sa03 *
	5752	5			-0.1	3			ORa		95Bi01 *
$^{181}\text{Hg}(\alpha)^{177}\text{Pt}$	6288	5	6284	4	-0.7	-					79Ha10 *
	6283	10			0.1	-			GSa		86Ke03 *
	6269.3	13.			1.2	-			Daa		96Pa01 *
	ave.	6285	4		-0.2	1	99	83 $^{181}\text{Hg}$			average
$^{181}\text{Tl}(\alpha)^{177}\text{Au}$	6319.9	20.	6321	6	0.1	U					92Bo.D
	6326.1	10.			-0.5	-			Ara		98To14
	6320.9	7.			0.1	-			Anv		09An14
	ave.	6323	6		-0.2	1	97	88 $^{177}\text{Au}$			average
$^{181}\text{Tl}(\alpha)^{177}\text{Au}^m$	6120.3	20.	6132	5	0.6	2			GSa		84Sc.A *
	6132.6	10.			-0.1	2			Ara		98To14 *
	6133.1	6.4			-0.2	2			Anv		09An14 *
$^{181}\text{Pb}(\alpha)^{177}\text{Hg}$	7374.3	10.	7240	7	-13.4	F			GSa		86Ke03 *
	7203.5	15.			2.4	5			ORa		89To01
	7224.9	20.			0.7	o			Ara		96To01 *
	7250.7	10.			-1.0	5			Ara		05Ca.A *
	7252.0	15.			-0.8	5			Anv		09An20 *
$^{181}\text{Ta}(p,t)^{179}\text{Ta}$	-5738	5	-5741.8	1.9	-0.8	1	14	7 $^{179}\text{Ta}$	Min		73Oo01
$^{180}\text{Hf}(n,\gamma)^{181}\text{Hf}$	5695.2	0.6	5694.80	0.07	-0.7	U					71Al22
	5694.80	0.07				2			Prn		02Bo41
	5695.58	0.20			-3.9	C			Bdn		06Fi.A
$^{180}\text{Hf}(d,p)^{181}\text{Hf}$	3440	25	3470.23	0.07	1.2	U			Sac		66Ga06
	3475	10			-0.5	U			Tal		68Ri07
$^{181}\text{Ta}(\gamma,n)^{180}\text{Ta}$	-7713	25	-7576.8	1.3	5.4	B			Phi		60Ge01 *
	-7852	26			10.6	B			Phi		60Ge01
	-7580	5			0.6	U			McM		79Ba06
	-7579	2			1.1	2			McM		81Co17
$^{181}\text{Ta}(d,t)^{180}\text{Ta}$	-1317.7	1.8	-1319.5	1.3	-1.0	2			NDm		79Ta.B
$^{180}\text{Ta}^m(n,\gamma)^{181}\text{Ta}$	7651.8	0.5	7652.08	0.19	0.6	2			MMn		81Co17 Z
	7652.13	0.20			-0.2	2			ILn		84Fo.A Z
$^{180}\text{W}(n,\gamma)^{181}\text{W}$	6669.02	0.16				2			Bdn		15Hu07
$^{180}\text{W}(d,p)^{181}\text{W}$	4468	15	4444.45	0.16	-1.6	U			Kop		72Ca01
$^{181}\text{Hg}(\epsilon p)^{180}\text{Pt}$	6150	200	6486	19	1.7	F					72Ho19 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{181}\text{Hf}(\beta^-)^{181}\text{Ta}$	1023	8	1035.5	1.8	1.6	U					52Fa14 *
	1020	5			3.1	B					53Ba81 *
$^{181}\text{W}(\epsilon)^{181}\text{Ta}$	184	12	204.5	1.9	1.7	U					66Ra03
	190	6			2.4	U					83Se17
$^{181}\text{Os}(\beta^+)^{181}\text{Re}$	2990	200	2967	28	-0.1	U					67Go25 *
$^{181}\text{Hg}^m(\text{IT})^{181}\text{Hg}$	212	50				2					09An17 *
* $^{181}\text{Os}-\text{u}$	$M-A=-43450(100)$ keV for mixture gs+m at 49.20 keV										Nub16b **
* $^{181}\text{Os}-\text{u}$	$M-A=-43529(28)$ keV for mixture gs+m at 49.20 keV										Nub16b **
* $^{181}\text{Au}(\alpha)^{177}\text{Ir}$	$E_\alpha=5609(8)$ , $5462(4)$ to ground state and $(3/2^-)$ level at 148.00 keV										Ens035 **
*	F : all lines in $^{181}\text{Au}$ and $^{183}\text{Au}$ shifted by 16–20keV										GAU **
* $^{181}\text{Au}(\alpha)^{177}\text{Ir}$	$E_\alpha=5626(5)$ to gs; favored $5479(5)$ to $(3/2^-)$ level at 148.00 keV										Ens035 **
* $^{181}\text{Hg}(\alpha)^{177}\text{Pt}$	$E_\alpha=6147.0(10,Z)$ , $6005.0(5,Z)$ to ground state and $1/2^-$ isomer at 147.4 keV										Nub16c **
* $^{181}\text{Hg}(\alpha)^{177}\text{Pt}$	$E_\alpha=6136.6(10,Z)$ , $6005.6(10,Z)$ to ground state and $1/2^-$ isomer at 147.4 keV										Nub16c **
* $^{181}\text{Hg}(\alpha)^{177}\text{Pt}$	$E_\alpha=5986(13)$ to $1/2^-$ isomer at 147.4 keV										Nub16c **
* $^{181}\text{Tl}(\alpha)^{177}\text{Au}^m$	$E_\alpha=6566(20)$ $Q_\alpha=6956.2$ from $^{181}\text{Tl}^m$ at 835.9 to 241.5 above $^{177}\text{Au}^m$										Nub16b **
* $^{181}\text{Tl}(\alpha)^{177}\text{Au}^m$	$E_\alpha=6578(10)$ $Q_\alpha=6968.5$ from $^{181}\text{Tl}^m$ at 835.9 to 241.5 above $^{177}\text{Au}^m$										Nub16b **
* $^{181}\text{Tl}(\alpha)^{177}\text{Au}^m$	$E_\alpha=6818(15)$ , $6578(7)$ from $^{181}\text{Tl}^m$ to $^{177}\text{Au}^m$ and level 241.5 above $^{177}\text{Au}^m$										09An14 **
* $^{181}\text{Pb}(\alpha)^{177}\text{Hg}$	F : This $\alpha$ -line not found in same reaction										96To01 **
* $^{181}\text{Pb}(\alpha)^{177}\text{Hg}$	Seen in correlation with $^{177}\text{Hg}$ $E_\alpha=6580$ keV										96To01 **
* $^{181}\text{Pb}(\alpha)^{177}\text{Hg}$	$E_\alpha=7015(10)$ to level at 77.2 keV										09An20 **
* $^{181}\text{Pb}(\alpha)^{177}\text{Hg}$	$E_\alpha=7016(15)$ to level at 77.2 keV										09An20 **
* $^{181}\text{Ta}(\gamma,n)^{180}\text{Ta}$	$Q=-7640(25)$ to $^{180}\text{Ta}^m$ at 75.3(1.4) keV										Nub16b **
* $^{181}\text{Hg}(\epsilon p)^{180}\text{Pt}$	F : retracted by authors in PrvCom										AHW **
* $^{181}\text{Hf}(\beta^-)^{181}\text{Ta}$	$E_{\beta^-}=408(8)$ $405(5)$ respectively, to $^{181}\text{Ta}^n$ at 615.19 keV										Nub16b **
* $^{181}\text{Os}(\beta^+)^{181}\text{Re}$	$E_{\beta^+}=1750(200)$ from $^{181}\text{Os}^m$ at 49.20 to $^{181}\text{Re}^m$ at 262.91 keV										Nub16b **
* $^{181}\text{Hg}^m(\text{IT})^{181}\text{Hg}$	From cascade x+90.3+71.4, with x estimated 50#										09An17 **
$^{182}\text{Re}-\text{u}$	-48311	65	-48790	110	-7.3	C			GS2	1.0	03Li.A *
$^{182}\text{Os}-\text{u}$	-47883	30	-47890	23	-0.2	1	61	61 $^{182}\text{Os}$	GS2	1.0	05Li24
$^{182}\text{Ir}-\text{u}$	-41942	30	-41924	23	0.6	1	56	56 $^{182}\text{Ir}$	GS2	1.0	05Li24
$^{182}\text{Pt}-\text{u}$	-38870	104	-38828	14	0.4	U			GS1	1.0	00Ra23
	-38860	30			1.1	1	22	22 $^{182}\text{Pt}$	GS2	1.0	05Li24
$^{182}\text{Au}-\text{u}$	-30420	110	-30382	22	0.3	U			GS1	1.0	00Ra23
	-30412	30			1.0	R			GS2	1.0	05Li24
$^{182}\text{Hg}-\text{u}$	-25297	30	-25311	11	-0.5	1	12	12 $^{182}\text{Hg}$	GS2	1.0	05Li24
$^{182}\text{Hg}-^{208}\text{Pb}_{.875}$	-4893	19	-4881	11	0.6	-			MA6	1.0	01Sc41
	-4898	21			0.8	-			MA6	1.0	01Sc41
ave.	-4895	14			1.0	1	56	55 $^{182}\text{Hg}$			average
$^{182}\text{Pt}(\alpha)^{178}\text{Os}$	4928.5	30.	4951	5	0.7	U					63Gr08
	4948.9	20.			0.1	U					66Si08
	4952.0	5.			-0.2	1	97	76 $^{178}\text{Os}$	ORa		95Bi01
$^{182}\text{Au}(\alpha)^{178}\text{Ir}$	5529	10	5526	4	-0.3	3					79Ha10 *
	5525.5	5.			0.1	3			ORa		95Bi01 *
$^{182}\text{Hg}(\alpha)^{178}\text{Pt}$	5998.1	5.	5996	5	-0.5	-					79Ha10 Z
	5989.9	13.3			0.4	-			Lvn		94Wa23
ave.	5997	5			-0.3	1	95	62 $^{178}\text{Pt}$			average
$^{182}\text{Tl}(\alpha)^{178}\text{Au}$	6550.2	10.	6551	6	0.1	F			GSa		86Ke03 *
	6593.1	15.			-2.8	B					04Ra28 *
	6550.9	6.2				2			ISa		16Va01 *
$^{182}\text{Tl}(\alpha)^{178}\text{Au}^p$	6186.2	20.				3					92Bo.D
$^{182}\text{Pb}(\alpha)^{178}\text{Hg}$	7076.8	10.	7066	6	-1.1	4			GSa		86Ke03
	7074.8	15.			-0.6	4			ORa		87To09
	7050.2	10.			1.5	4			Ara		99To11
	7066.6	10.			-0.1	4			Jya		00Je09
$^{180}\text{Hf}(t,p)^{182}\text{Hf}$	3931	6				2			McM		83Bu03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{180}\text{W}(t,p)^{182}\text{W}$	6265	5	6270.8	1.6	1.2	U			LAI		76Ca10 *
$^{182}\text{W}(p,t)^{180}\text{W}$	-6261	10	-6270.8	1.6	-1.0	U			Min		73Oo01
$^{181}\text{Ta}(n,\gamma)^{182}\text{Ta}$	6063.0	0.4	6062.94	0.11	-0.2	-					71He13 Z
	6063.1	0.5			-0.3	-					77St15 Z
	6063.1	0.5			-0.3	-			MMn		81Co17 Z
	6062.95	0.2			-0.1	-			ILn		83Fo.B
	6062.89	0.14			0.4	-			Bdn		06Fi.A
$^{181}\text{Ta}(d,p)^{182}\text{Ta}$	3832	8	3838.37	0.11	0.8	U			MIT		64Er02
$^{181}\text{Ta}(n,\gamma)^{182}\text{Ta}$	ave.	6062.93	0.11	6062.94	0.11	0.1	1	100	74 $^{182}\text{Ta}$		average
$^{182}\text{W}(d,t)^{181}\text{W}$	-1809	10	-1826.3	1.6	-1.7	U			Kop		72Ca01
$^{182}\text{Hf}(\beta^-)^{182}\text{Ta}$	431	50	380	6	-1.0	U					74Wa14 *
$^{182}\text{Ta}(\beta^-)^{182}\text{W}$	1809	5	1816.1	1.4	1.4	-					64Da15 *
	1813	3			1.0	-					67Ba01 *
	ave.	1811.9	2.6		1.6	1	30	26 $^{182}\text{Ta}$			average
$^{182}\text{Re}^m(\beta^+)^{182}\text{W}$	2860	20				2					63Ba37 *
$^{182}\text{Re}^m(\text{IT})^{182}\text{Re}$	60	100				3					63Ba37
$^{182}\text{Os}(\epsilon)^{182}\text{Re}^m$	848	15	777	30	-4.7	B					70Ak02 *
$^{182}\text{Ir}(\beta^+)^{182}\text{Os}$	5700	200	5560	30	-0.7	U					72We.A
$^{182}\text{Pt}(\beta^+)^{182}\text{Ir}$	2900	200	2883	25	-0.1	U					72We.A
$^{182}\text{Au}(\beta^+)^{182}\text{Pt}$	6850	200	7868	24	5.1	C					72We.A
$^{182}\text{Hg}(\beta^+)^{182}\text{Au}$	4950	200	4724	23	-1.1	U					72We.A
$^{182}\text{Re}-u$	$M-A=-44972(29)$ keV for mixture gs+m at 60(100) keV										Nub16b **
$^{182}\text{Au}(\alpha)^{178}\text{Ir}$	$E_\alpha=5353(10)$ to $2^+$ level at 54.4(0.5) keV										Ens097 **
$^{182}\text{Au}(\alpha)^{178}\text{Ir}$	$E_\alpha=5403(5), 5352(5)$ to ground state, 54.4 level										95Bi01 **
$^{182}\text{Tl}(\alpha)^{178}\text{Au}$	F : identification from excitation function assuming 100% $\alpha$ decay										WgM118**
$^{182}\text{Tl}(\alpha)^{178}\text{Au}$	$E_\alpha=6403(15)$ in coincidence with 46 keV $\gamma$										04Ra28 **
$^{182}\text{Tl}(\alpha)^{178}\text{Au}$	$E_\alpha=6165(6)$ in coincidence with 247.2(0.5) keV $\gamma$										16Va01 **
$^{180}\text{W}(t,p)^{182}\text{W}$	$Q-Q(^{170}\text{Y}(t,p))=112(5,\text{Ca}), Q(170)=-6153(4)$ keV										AHW **
$^{182}\text{Hf}(\beta^-)^{182}\text{Ta}$	$E_{\beta^-}=970(70) 480(50)$ from $^{182}\text{Hf}^m$ to $651.215(4)^-, 1115.96(7^-)$ levels										Ens15c **
$^{182}\text{Ta}(\beta^-)^{182}\text{W}$	$E_{\beta^-}=520(5)$ to $2^-$ level at 1289.1498 keV										Ens15c **
$^{182}\text{Ta}(\beta^-)^{182}\text{W}$	$E_{\beta^-}=1713(3)$ to $2^+$ level at 100.10598 keV										Ens15c **
$^{182}\text{Re}^m(\beta^+)^{182}\text{W}$	$E_{\beta^+}=1740(20), 550(20)$ to $2^+$ level at 100.10598, $2^-$ at 1289.1498 keV										Ens15c **
$^{182}\text{Os}(\epsilon)^{182}\text{Re}^m$	pK=0.47(0.07) to $1^+$ level at 726.97 keV above $^{182}\text{Re}^m$ , recalculated Q										Ens15c **
$^{183}\text{Lu}-u$	-42637	86				2			GS3	1.0	13Sh30
$^{183}\text{W O}-\text{C}_2 \text{ } ^{35}\text{Cl}_5$	100858.0	2.7	100875.6	0.8	2.6	U			H29	2.5	77Sh04
	100873.6	0.8			1.0	1	16	15 $^{183}\text{W}$	H48	2.5	03Ba49
$^{183}\text{Re}-u$	-49151	30	-49179	9	-0.9	U			GS2	1.0	05Li24
$^{183}\text{Os}-u$	-46879	61	-46880	50	0.1	1	77	77 $^{183}\text{Os}$	GS2	1.0	05Li24 *
$^{183}\text{Ir}-u$	-43160	104	-43160	26	0.0	U			GS1	1.0	00Ra23
	-43145	30			-0.5	1	76	76 $^{183}\text{Ir}$	GS2	1.0	05Li24
$^{183}\text{Pt}-u$	-38440	107	-38403	17	0.3	U			GS1	1.0	00Ra23
	-38400	32			-0.1	1	27	27 $^{183}\text{Pt}$	GS2	1.0	05Li24 *
$^{183}\text{Au}-u$	-32440	104	-32412	10	0.3	U			GS1	1.0	00Ra23
	-32371	30			-1.4	1	11	11 $^{183}\text{Au}$	GS2	1.0	05Li24
$^{183}\text{Hg}-u$	-25537	35	-25555	8	-0.5	U			GS2	1.0	05Li24 *
$^{183}\text{Hg}-^{208}\text{Pb}_{.880}$	-5009	19	-5009	8	0.0	-			MA6	1.0	01Sc41
	-5002	19			-0.4	-			MA6	1.0	01Sc41
	ave.	-5006	13			-0.3	1	32	32 $^{183}\text{Hg}$		average
$^{183}\text{Tl}-^{133}\text{Cs}_{1.376}$	112286	11	112291	10	0.5	1	83	83 $^{183}\text{Tl}$	MA8	1.0	08We02
$^{183}\text{W O}_2-^{178}\text{Hf } ^{37}\text{Cl}$	30455.7	5.0	30442.7	1.7	-1.0	U			H35	2.5	80Sh06
$^{183}\text{W O}_2-^{180}\text{W } ^{35}\text{Cl}$	24421	9	24487.6	1.7	3.0	B			H24	2.5	73Ba40
	24509	6			-1.4	U			H28	2.5	77Sh04
$^{183}\text{W } ^{35}\text{Cl}-^{181}\text{Ta } ^{37}\text{Cl}$	5177.2	1.2	5175.3	1.5	-0.6	1	25	22 $^{181}\text{Ta}$	H35	2.5	80Sh06
$^{183}\text{W O}_2 \text{ } ^{37}\text{Cl}-^{182}\text{W } ^{35}\text{Cl}_2$	20045.6	1.8	20045.21	0.11	-0.1	U			H28	2.5	77Sh04



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{183}\text{Pt}(\alpha)^{179}\text{Os}$	4846.1	30.	4822	9	-0.8	U					63Gr08
	4835.9	20.0			-0.7	-					66Si08
	4819.6	10.2			0.2	-					95Bi01
$^{183}\text{Au}(\alpha)^{179}\text{Ir}$	ave. 4823	9			-0.1	1	93	65	$^{179}\text{Os}$		average
	5462.6	5.	5465.3	2.9	0.5	-					68Si01 Z
	5465.5	5.			0.0	-			Bka		82Bo04 Z
	5449.3	10.			1.6	F					84Br.A *
$^{183}\text{Hg}(\alpha)^{179}\text{Pt}$	5468.8	5.			-0.7	-			ORa		95Bi01
	ave. 5465.6	3.0			-0.1	1	99	88	$^{179}\text{Ir}$		average
	6043.4	6.	6039	4	-0.8	-			ORa		76To06
	6036.2	5.			0.5	-					79Ha10 Z
$^{183}\text{Tl}^m(\alpha)^{179}\text{Au}$	ave. 6039	4			-0.1	1	98	93	$^{179}\text{Pt}$		average
	6593.4	30.	6605	9	0.4	U			GSa		80Sc09 *
	6600.6	30.			0.2	U			Jya		04Ra28 *
	6609.5	10.			-0.4	1	84	67	$^{179}\text{Au}$	Anv	11Ve01 *
$^{183}\text{Pb}(\alpha)^{179}\text{Hg}$	6928	7				2			Anv		02Je09 *
	$^{183}\text{Pb}^m(\alpha)^{179}\text{Hg}$	6950.1	25.	7022	4	2.9	B		GSa		80Sc09
7029		20			-0.3	U			GSa		84Sc.A *
7026.9		10.			-0.5	2			GSa		86Ke03
6868.4		10.			15.4	B			ORa		87To09
7034		10			-1.2	2			ORa		89To01 *
7018		5			0.8	2			Anv		02Je09 *
7018		5			0.8	2			Anv		02Je09 *
$^{183}\text{W}(\text{p,t})^{181}\text{W}$	-5810	10	-5792.6	1.6	1.7	U			Min		73Oo01
$^{182}\text{Ta}(\text{n},\gamma)^{183}\text{Ta}$	6934.18	0.20				2			ILn		83Fo.B
$^{182}\text{W}(\text{n},\gamma)^{183}\text{W}$	6191.6	2.0	6190.84	0.04	-0.4	U					67Sp03 Z
	6190.1	1.5			0.5	U					70Or.A
	6190.76	0.12			0.6	-			Ltn		93Pr.A
	6190.89	0.13			-0.4	o			Bdn		06Fi.A
	6190.81	0.06			0.4	-			ILn		11Bo09
	6190.88	0.06			-0.7	-			Bdn		14Hu02
	6190.88	0.06			-0.7	-			Bdn		14Hu02
$^{183}\text{W}(\gamma,\text{n})^{182}\text{W}$	-6290	50	-6190.84	0.04	2.0	U			Phi		60Ge01
$^{182}\text{W}(\text{d,p})^{183}\text{W}$	3967	5	3966.27	0.04	-0.1	U			ANL		65Er03
	3979	10			-1.3	U			Kop		72Ca01
	3979	10			-1.3	U			Kop		72Ca01
$^{183}\text{W}(\text{d,t})^{182}\text{W}$	57	15	66.39	0.04	0.6	U			Kop		72Ca01
$^{182}\text{W}(\text{n},\gamma)^{183}\text{W}$	ave. 6190.84	0.04	6190.84	0.04	0.0	1	100	100	$^{182}\text{W}$		average
$^{182}\text{W}(\text{}^3\text{He,d})^{183}\text{Re}$	-610	40	-641	8	-0.8	U			Roc		71Lu01
$^{183}\text{Hg}(\text{ep})^{182}\text{Pt}$	5000	200	5075	15	0.4	F					72Ho19
$^{183}\text{Hf}(\beta^-)^{183}\text{Ta}$	2010	30				3					67Mo13 *
$^{183}\text{Ta}(\beta^-)^{183}\text{W}$	1068	10	1072.8	1.4	0.5	U					55Mu19 *
$^{183}\text{Re}(\epsilon)^{183}\text{W}$	556	8				2					69Ku03 *
$^{183}\text{Ir}(\beta^+)^{183}\text{Os}$	3450	100	3460	50	0.1	1	28	23	$^{183}\text{Os}$		70Be.A *
$^{183}\text{Tl}^m(\text{IT})^{183}\text{Tl}$	628.7	0.5	628.7	0.5	0.0	1	100	83	$^{183}\text{Tl}^m$		11Ve.A
* $^{183}\text{Os}-u$	$M-A=-43582(28)$ keV for mixture gs+m at 170.73 keV										Nub16b **
* $^{183}\text{Pt}-u$	$M-A=-35752(28)$ keV for mixture gs+m at 34.74 keV										Nub16c **
* $^{183}\text{Hg}-u$	Existence of isomeric state under discussion (see Nubase); not corrected										Nub16b **
* $^{183}\text{Au}(\alpha)^{179}\text{Ir}$	F : all lines in $^{181}\text{Au}$ and $^{183}\text{Au}$ shifted by 16-20keV										GAu **
* $^{183}\text{Tl}^m(\alpha)^{179}\text{Au}$	$E_\alpha=6449(15)$ , error increased since partially summed with electrons										GAu **
* $^{183}\text{Tl}^m(\alpha)^{179}\text{Au}$	$E_\alpha=6456(15)$ , error increased since partially summed with electrons										GAu **
* $^{183}\text{Tl}^m(\alpha)^{179}\text{Au}$	$E_\alpha=6400(10)$ , 6378(10) summed with $e^-$ , in coinc. with 62.4, 89.5 $\gamma$										GAu **
* $^{183}\text{Pb}(\alpha)^{179}\text{Hg}$	$E_\alpha=6775(7)$ , 6570(10) to ground state, 217 level										02Je09 **
* $^{183}\text{Pb}^m(\alpha)^{179}\text{Hg}$	$E_\alpha=6868(20)$ , 6715(20) to ground state, 171.4 isomer										02Je09 **
*	original assignment to $^{182}\text{Pb}$ changed										AHW **
* $^{183}\text{Pb}^m(\alpha)^{179}\text{Hg}$	$E_\alpha=6874(15)$ , 6712(10) to ground state, 171.4 isomer; and an 6784(15) line										Nub16b **
* $^{183}\text{Pb}^m(\alpha)^{179}\text{Hg}$	$E_\alpha=6860(11)$ , 6698(5) to ground state, 171.4 isomer										Nub16b **
* $^{183}\text{Hg}(\text{ep})^{182}\text{Pt}$	F : retracted by authors in PrvCom										AHW **
* $^{183}\text{Hf}(\beta^-)^{183}\text{Ta}$	$E_{\beta^-}=1540(30)$ to $(5/2^+)$ level at 459.062 keV, and other $E_{\beta^-}$										Ens164 **
* $^{183}\text{Ta}(\beta^-)^{183}\text{W}$	$E_{\beta^-}=615(10)$ to $7/2^-$ level at 453.0695 keV										Ens164 **
* $^{183}\text{Re}(\epsilon)^{183}\text{W}$	pK=0.40(0.07) to $7/2^-$ level at 453.0695 keV										Ens164 **
* $^{183}\text{Ir}(\beta^+)^{183}\text{Os}$	$Q_{\beta^+}=3190(100)$ mainly to $3/2^-$ level at 258.34 keV										Ens164 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{184}\text{W}-u$	-49066.76	0.99	-49066.7	0.8	0.0	1	28	28 $^{184}\text{W}$	TG1	1.5	12Sm07
$^{184}\text{Os}-u$	-47504.3	1.2	-47507.1	0.9	-1.5	1	24	24 $^{184}\text{Os}$	TG1	1.5	12Sm07
$^{184}\text{Ir}-u$	-42460	110	-42520	30	-0.6	U			GS1	1.0	00Ra23
	-42524	30				2			GS2	1.0	05Li24
$^{184}\text{Pt}-u$	-40120	104	-40080	17	0.4	U			GS1	1.0	00Ra23
	-40068	30			-0.4	1	31	31 $^{184}\text{Pt}$	GS2	1.0	05Li24
$^{184}\text{Au}-u$	-32540	104	-32548	24	-0.1	U			GS1	1.0	00Ra23
	-32557	37			0.2	R			GS2	1.0	05Li24
$^{184}\text{Hg}-u$	-28230	110	-28287	11	-0.5	U			GS1	1.0	00Ra23
	-28296	30			0.3	-			GS2	1.0	05Li24
ave.	-28279	17			-0.4	1	39	39 $^{184}\text{Hg}$			average
$^{184}\text{Hg}-^{204}\text{Pb}_{.902}$	-3986	20	-3972	11	0.7	1	29	29 $^{184}\text{Hg}$	MA6	1.0	01Sc41
$^{184}\text{Hg}-^{208}\text{Pb}_{.885}$	-7620	19	-7624	11	-0.2	1	32	32 $^{184}\text{Hg}$	MA6	1.0	01Sc41
$^{184}\text{Tl}-u$	-18115	112	-18125	11	-0.1	U			GS2	1.0	05Li24
$^{184}\text{Tl}-^{133}\text{Cs}_{1.383}$	112645.4	23.2	112635	11	-0.4	1	21	21 $^{184}\text{Tl}$	MA8	1.0	14Bo26
$^{184}\text{W O}_2-^{181}\text{Ta }^{35}\text{Cl}$	23917.5	2.8	23910.5	1.5	-1.0	U			H35	2.5	80Sh06
$^{184}\text{W }^{35}\text{Cl}-^{182}\text{W }^{37}\text{Cl}$	5675	3	5677.65	0.16	0.4	U			H22	2.5	70Mc03
	5676.3	2.2			0.2	U			H28	2.5	77Sh04
$^{184}\text{W O}_2 ^{37}\text{Cl}-^{183}\text{W }^{35}\text{Cl}_2$	18734.7	3.0	18735.19	0.17	0.1	U			H28	2.5	77Sh04
$^{184}\text{Os}-^{184}\text{W}$	1560.59	0.70	1559.7	0.7	-0.9	1	46	31 $^{184}\text{Os}$	TG1	1.5	12Sm07
$^{184}\text{Pt}(\alpha)^{180}\text{Os}$	4579.8	20.	4599	8	0.9	-					63Gr08
	4600.2	20.			-0.1	-					66Si08
	4602.2	10.			-0.4	-			ORa		95Bi01
ave.	4598	8			0.1	1	94	66 $^{180}\text{Os}$			average
$^{184}\text{Au}(\alpha)^{180}\text{Ir}$	5218.6	15.	5234	5	1.0	U			ISa		70Ha18
	5233.9	5.				3			ORa		95Bi01
$^{184}\text{Hg}(\alpha)^{180}\text{Pt}$	5658.2	15.	5662	4	0.2	2					70Ha18
	5662.3	5.1			-0.1	2			ORa		76To06
	5662.3	10.2			0.0	2			Lvn		93Wa03
$^{184}\text{Tl}(\alpha)^{180}\text{Au}$	6299.4	5.	6317	9	0.4	U			ORa		76To06
	6292.9	10.			0.5	U			GSa		80Sc09
	6315.2	10.2			0.2	1	83	79 $^{184}\text{Tl}$	ISa		16Va01
$^{184}\text{Pb}(\alpha)^{180}\text{Hg}$	6765.4	10.	6774	3	0.8	-					80Du02
	6779.6	10.			-0.6	-			GSa		80Sc09
	6773.6	10.			0.0	-			GSa		84Sc.A
	6781.7	10.2			-0.8	-			ORa		87To09
	6773.6	6.			0.1	-			Jya		98Co27
	6772.5	10.			0.1	-			Ara		99To11
	6773.6	6.			0.1	-			Anv		04An07
ave.	6774	3			0.0	1	99	70 $^{184}\text{Pb}$			average
$^{184}\text{Bi}(\alpha)^{180}\text{Tl}$	8024.8	50.				7			Anv		03An27
$^{182}\text{W}(\text{t,p})^{184}\text{W}$	5127	7	5120.15	0.14	-1.0	U			LAl		76Ca10
$^{184}\text{W}(\text{p,t})^{182}\text{W}$	-5124	5	-5120.15	0.14	0.8	U			Min		73Oo01
$^{183}\text{W}(\text{n},\gamma)^{184}\text{W}$	7411.2	0.5	7411.11	0.13	-0.2	U					74Gr11
	7411.8	0.3			-2.3	B					75Bu01
	7411.15	0.16			-0.2	o			Bdn		06Fi.A
	7411.11	0.13			0.0	1	99	72 $^{183}\text{W}$	Bdn		14Hu02
$^{183}\text{W}(\text{d,p})^{184}\text{W}$	5187	15	5186.54	0.13	0.0	U			Kop		72Ca01
$^{184}\text{W}(\text{d,t})^{183}\text{W}$	-1154	10	-1153.88	0.13	0.0	U			Kop		72Ca01
$^{184}\text{Hf}(\beta^-)^{184}\text{Ta}$	1340	30				3					73Wa18
$^{184}\text{Ta}(\beta^-)^{184}\text{W}$	2866	26				2					73Ya02
$^{184}\text{Ir}(\beta^+)^{184}\text{Os}$	5100	250	4642	28	-1.8	U					70Be.A
	4300	100			3.4	B					73Ho09
	4285	70			5.1	B					89Po09

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{184}\text{Au}(\beta^+)^{184}\text{Pt}$	6380	50	7016	27	12.7	C					84Da.A *
$^{184}\text{Hg}(\beta^+)^{184}\text{Au}$	3760	30	3970	24	7.0	C					84Da.A
* $^{184}\text{Au}-u$	$M-A=-30280(100)$ keV for mixture gs+m at 68.46 keV										Nub16b **
* $^{184}\text{Au}-u$	$M-A=-30292(28)$ keV for mixture gs+m at 68.46 keV										Nub16b **
* $^{184}\text{Tl}-u$	$M-A=-16899(102)$ keV for mixture gs+m at -50(30) keV										Nub16b **
* $^{184}\text{Tl}-^{133}\text{Cs}_{1,383}$	$D_M=112618.6(6.2)$ $\mu\text{u}$ for mixture gs+m at -50(30) keV; $M-A=-16898.5(5.8)$ keV										Nub16b **
* $^{184}\text{Au}(\alpha)^{180}\text{Ir}$	$E_{\alpha}=5172(15)$ from $^{184}\text{Au}^m$ at 68.6(0.1) keV transition to ground state in $^{180}\text{Ir}$										94Ib01 **
* $^{184}\text{Au}(\alpha)^{180}\text{Ir}$	$E_{\alpha}=5187(5)$ from $^{184}\text{Au}^m$ at 68.6(0.1) keV										95Bi01 **
* $^{184}\text{Tl}(\alpha)^{180}\text{Au}$	$E_{\alpha}=6161(10)$ to level at 17.1(3) keV										94Ib01 **
* $^{184}\text{Hf}(\beta^-)^{184}\text{Ta}$	$E_{\beta^-}=1100(30)$ to $1^-$ level at 228.4 keV, and other $E_{\beta^-}$										16Va01 **
* $^{184}\text{Ta}(\beta^-)^{184}\text{W}$	$E_{\beta^-}=1165(26)$ to $6^+$ level at 1746.03 keV, and other $E_{\beta^-}$										Ens102 **
* $^{184}\text{Ir}(\beta^+)^{184}\text{Os}$	$Q_{\beta^+}=4720(250)$ to $4^+$ level at 383.68 keV										Ens102 **
* $^{184}\text{Ir}(\beta^+)^{184}\text{Os}$	$E_{\beta^+}=2900(100)$ to $4^+$ level at 383.68 keV										Ens102 **
* $^{184}\text{Ir}(\beta^+)^{184}\text{Os}$	$E_{\beta^+}=2320(70)$ to $2^+$ level at 942.86 keV										Ens102 **
* $^{184}\text{Au}(\beta^+)^{184}\text{Pt}$	$Q_{\beta^+}=6450(50)$ from $^{184}\text{Au}^m$ at 68.6(0.1) keV										94Ib01 **
$^{185}\text{Hf}-u$	-41138	69				2			GS3	1.0	13Sh30
$^{185}\text{Os}-u$	-46037	31	-45954.0	0.9	2.7	F			GS2	1.0	03Li.A *
$^{185}\text{Ir}-u$	-43340	110	-43300	30	0.3	U			GS1	1.0	00Ra23
	-43302	30				2			GS2	1.0	05Li24
$^{185}\text{Pt}-u$	-39334	112	-39386	28	-0.5	U			GS1	1.0	00Ra23 *
	-39381	44			-0.1	1	40	$^{185}\text{Pt}$	GS2	1.0	05Li24 *
$^{185}\text{Au}-u$	-34213	115	-34201.1	2.8	0.1	o			GS1	1.0	00Ra23 *
	-34224	69			0.3	U			GS2	1.0	05Li24 *
$^{185}\text{Au}-^{133}\text{Cs}_{1,391}$	97315.2	2.8				2			MA8	1.0	16Ma.1
$^{185}\text{Hg}-u$	-28070	107	-28109	15	-0.4	U			GS1	1.0	00Ra23
	-28088	44			-0.5	1	11	$^{185}\text{Hg}$	GS2	1.0	05Li24 *
$^{185}\text{Hg}-^{208}\text{Pb}_{,889}$	-7373	29	-7353	15	0.7	1	26	$^{185}\text{Hg}$	MA6	1.0	01Sc41 *
$^{185}\text{Tl}-u$	-21354	145	-21211	22	1.0	U			GS2	1.0	05Li24 *
$^{185}\text{Re }^{16}\text{O}_2-^{182}\text{W }^{35}\text{Cl}$	25731	6	25729.2	0.7	-0.1	U			H22	2.5	70Mc03
$^{185}\text{Re }^{35}\text{Cl}-^{183}\text{W }^{37}\text{Cl}$	5695	3	5683.9	0.7	-1.5	U			H22	2.5	70Mc03
	5678.7	1.0			2.1	U			H28	2.5	77Sh04
$^{185}\text{Re}(\alpha, ^8\text{He})^{181}\text{Re}$	-26480	14	-26486	13	-0.5	2			INS		90Ka19
$^{185}\text{Pt}(\alpha)^{181}\text{Os}$	4436.6	10.2	4437	10	0.0	1	96	$^{185}\text{Pt}$	ORa		91Bi04 *
$^{185}\text{Au}(\alpha)^{181}\text{Ir}$	5180.2	5.	5180	5	0.0	3					68Si01 Z
	5182.9	15.			-0.2	U					70Ha18 Z
	5179	10			0.1	3			ORa		91Bi04 *
$^{185}\text{Hg}(\alpha)^{181}\text{Pt}$	5777	15	5773	4	-0.3	-					70Ha18 *
	5775	5			-0.4	-			ORa		76To06 *
	5761	15			0.8	-					76Gr.A *
ave.	5774	5			-0.2	1	97	$^{181}\text{Pt}$			average
$^{185}\text{Tl}^m(\alpha)^{181}\text{Au}$	6112.6	7.	6143	5	4.4	C					75Co.A Z
	6143.3	5.				4			ORa		76To06 *
	6145.6	15.			-0.2	U			GSa		80Sc09 Z
$^{185}\text{Pb}(\alpha)^{181}\text{Hg}$	6693	15	6695	5	0.1	U			GSa		80Sc09 *
	6555.0	15.			2.8	B			ORa		87To09
	6695	5				2			Anv		02An15 *
$^{185}\text{Pb}^m(\alpha)^{181}\text{Hg}^m$	6622.9	20.	6550	5	-3.7	B			Ora		75Ca06
	6679.7	20.			-6.5	B			GSa		80Sc09
	6549.8	5.				3			Anv		02An15
$^{185}\text{Bi}^m(\alpha)^{181}\text{Tl}$	8258.9	30.	8218	12	-1.3	-			Ara		01Po05 *
	8207.8	15.3			0.7	-			Anv		04An07
ave.	8218	14			0.0	1	76	$^{185}\text{Bi}^m$			average
$^{184}\text{W}(n,\gamma)^{185}\text{W}$	5753.7	0.3	5753.74	0.05	0.1	U			BNn		87Br05 Z
	5754.62	0.24			-3.7	C			Bdn		06Fi.A
	5753.74	0.05			-0.1	1	100	$^{185}\text{W}$	Bdn		14Hu02

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{184}\text{W}(\text{d,p})^{185}\text{W}$	3524	5	3529.17	0.05	1.0	U			65Er03
	3533	10			-0.4	U	ANL		72Ca01
$^{184}\text{W}({}^3\text{He,d})^{185}\text{Re}$	-98	40	-90.9	0.7	0.2	U	Roc		71Lu01
$^{185}\text{Re}(\text{d,t})^{184}\text{Re}-^{187}\text{Re}(\text{e})^{186}\text{Re}$	-310	4	-310	4	0.0	1	100	100 $^{184}\text{Re}$	76El12
$^{184}\text{Os}(\text{n},\gamma)^{185}\text{Os}$	6625.4	0.9	6624.66	0.27	-0.8	U			74Pr15
	6624.52	0.28			0.5	1	95	51 $^{185}\text{Os}$	06Fi.A
$^{185}\text{Bi}^m(\text{p})^{184}\text{Pb}$	1669	50	1607	13	-1.2	o			95Da.A *
	1606.8	16.			0.0	1	67	36 $^{185}\text{Bi}^m$	01Po05 *
	1568.6	50.			0.8	o			03An27 *
	1591.7	5.			3.0	F			04An07 *
$^{185}\text{Ta}(\beta^-)^{185}\text{W}$	2013	20	1994	14	-1.0	2			69Ku07 *
$^{185}\text{W}(\beta^-)^{185}\text{Re}$	432.6	1.0	431.2	0.7	-1.4	1	44	28 $^{185}\text{Re}$	67Wi19
$^{185}\text{Os}(\varepsilon)^{185}\text{Re}$	1012.7	1.0	1013.1	0.4	0.4	-			67Sc15 *
	1012.8	0.5			0.7	-			70Sc06 *
ave.	1012.8	0.4			0.8	1	88	49 $^{185}\text{Os}$	average
$^{185}\text{Au}(\beta^+)^{185}\text{Pt}$	4707	40	4830	26	3.1	F			86Da.A *
$^{185}\text{Tl}^m(\text{IT})^{185}\text{Tl}$	454.8	1.5				5			Ens061
* $^{185}\text{Os}-\text{u}$	F : contaminated by isomeric state AND by other nuclides								
* $^{185}\text{Pt}-\text{u}$	$M-A=-36590(100)$ keV for mixture gs+m at 103.41 keV								
* $^{185}\text{Pt}-\text{u}$	$M-A=-36631(28)$ keV for mixture gs+m at 103.41 keV								
* $^{185}\text{Au}-\text{u}$	$M-A=-31820(90)$ keV for mixture gs+m at 100#100 keV								
* $^{185}\text{Au}-\text{u}$	$M-A=-31829(28)$ keV for mixture gs+m at 100#100 keV								
* $^{185}\text{Hg}-\text{u}$	$M-A=-26112(28)$ keV for mixture gs+m at 103.7 keV								
* $^{185}\text{Hg}-^{208}\text{Pb}_{.889}$	Original error (17keV) increased by 20 due to isomer+ground state lines in trap								
* $^{185}\text{Tl}-\text{u}$	$M-A=-19664(31)$ keV for mixture gs+m at 454.8(1.5) keV								
* $^{185}\text{Pt}(\alpha)^{181}\text{Os}$	$E_\alpha=4444(10)$ assumed from $(1/2^-)$ isomer at 103.41 keV								
* $^{185}\text{Au}(\alpha)^{181}\text{Ir}$	$E_\alpha=5069(10)$ , 4826(10) to ground state, 243.3 level								
*	unhindered $E_\alpha=5069(10)$ to ground state or very low level; from coinc.								
* $^{185}\text{Hg}(\alpha)^{181}\text{Pt}$	$E_\alpha=5653.4(15,Z)$ , 5576.4(15,Z) to ground state, $3/2^-$ level at 79.41 keV								
*	and $E_\alpha=5376.4(15,Z)$ from $^{185}\text{Hg}^m$ at 103.8 to $13/2^+$ level at 380.92 keV								
* $^{185}\text{Hg}(\alpha)^{181}\text{Pt}$	$E_\alpha=5653(5)$ , 5569(5) to ground state, $3/2^-$ level at 79.41 keV								
*	and 5371(10) from $^{185}\text{Hg}^m$ at 103.8 to $13/2^+$ level at 380.92 keV								
* $^{185}\text{Hg}(\alpha)^{181}\text{Pt}$	$E_\alpha=5365(15)$ from $^{185}\text{Hg}^m$ at 103.8 to $13/2^+$ level at 380.92 keV								
* $^{185}\text{Tl}^m(\alpha)^{181}\text{Au}$	$E_\alpha=6010.2(5,Z)$ ; also an $E_\alpha=5975.2(5,Z)$ , 4 times stronger branch								
* $^{185}\text{Tl}^m(\alpha)^{181}\text{Au}$	$E_\alpha=6012.5(15,Z)$ ; also an $E_\alpha=5970.5(15,Z)$ , 4 times stronger branch								
* $^{185}\text{Pb}(\alpha)^{181}\text{Hg}$	$E_\alpha=6485(15)$ to 64 level								
* $^{185}\text{Pb}(\alpha)^{181}\text{Hg}$	$E_\alpha=6486(5)$ , 6288(5) to 64, 269 levels								
* $^{185}\text{Bi}^m(\alpha)^{181}\text{Tl}$	$E_\alpha=8030$ , by same authors, from only one event								
* $^{185}\text{Bi}^m(\text{p})^{184}\text{Pb}$	Read from graph								
* $^{185}\text{Bi}^m(\text{p})^{184}\text{Pb}$	Average by authors of $E_p=1618(11)$ , and 1585(9) in reference								
* $^{185}\text{Bi}^m(\text{p})^{184}\text{Pb}$	As read from graph								
* $^{185}\text{Bi}^m(\text{p})^{184}\text{Pb}$	F : rejected by authors: no dedicated calibration with known proton activity								
* $^{185}\text{Ta}(\beta^-)^{185}\text{W}$	$E_{\beta^-}=1770(20)$ to $7/2^-$ level at 243.62 keV								
* $^{185}\text{Os}(\varepsilon)^{185}\text{Re}$	$L/K=0.600(0.006)$ to $3/2^+$ level at 874.81 and $1/2^+$ at 880.33 keV								
* $^{185}\text{Os}(\varepsilon)^{185}\text{Re}$	$pK=0.109(0.005)$ to $3/2^+$ level at 931.06 keV, and other pK, recalculated								
* $^{185}\text{Au}(\beta^+)^{185}\text{Pt}$	F : insufficient information								
									GAU **
$^{186}\text{Hf}-\text{u}$	-39103	55				2			GS3 1.0 13Sh30
$^{186}\text{W O}-\text{C }^{13}\text{C }^{35}\text{Cl}_4 \text{ }^{37}\text{Cl}$	104592.7	3.2	104611.6	1.3	2.4	U			H29 2.5 77Sh04
$^{186}\text{Ir}-\text{u}$	-42063	30	-42053	18	0.3	2			GS2 1.0 05Li24 *
$^{186}\text{Pt}-\text{u}$	-40656	30	-40649	23	0.2	1	61	61 $^{186}\text{Pt}$	GS2 1.0 05Li24
$^{186}\text{Au}-\text{u}$	-34029	30	-34047	23	-0.6	1	56	56 $^{186}\text{Au}$	GS2 1.0 05Li24
$^{186}\text{Hg}-\text{u}$	-30660	104	-30638	13	0.2	U			GS1 1.0 00Ra23
	-30630	30			-0.3	1	17	17 $^{186}\text{Hg}$	GS2 1.0 05Li24
$^{186}\text{Hg}-^{204}\text{Pb}_{.912}$	-6065	20	-6054	13	0.6	-			MA6 1.0 01Sc41
ave.	-6058	17			0.2	1	56	56 $^{186}\text{Hg}$	average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{186}\text{Tl}-u$	-21653	218	-21349	24	1.4	o			GS1	1.0	00Ra23 *
	-21513	105			1.6	U			GS2	1.0	05Li24 *
$^{186}\text{Tl}-^{133}\text{Cs}_{1.398}$	110831	24	110829	24	-0.1	o			MA8	1.0	08We02 *
	110829	24				2			MA8	1.0	14Bo26 *
$^{186}\text{Tl}^n-^{133}\text{Cs}_{1.398}$	111254	34				2			MA8	1.0	14Bo26
$^{186}\text{W O}_2-^{183}\text{W }^{35}\text{Cl}$	25122	5	25117.3	1.3	-0.4	U			H28	2.5	77Sh04
$^{186}\text{W }^{35}\text{Cl}-^{184}\text{W }^{37}\text{Cl}$	6374	3	6382.1	1.2	1.1	U			H22	2.5	70Mc03
	6382.0	1.4			0.0	1	13	11 $^{186}\text{W}$	H28	2.5	77Sh04
$^{186}\text{Os}(\alpha)^{182}\text{W}$	2820.6	50.	2821.2	0.9	0.0	U					75Vi01
$^{186}\text{Pt}(\alpha)^{182}\text{Os}$	4323.2	20.	4320	18	-0.2	1	79	39 $^{182}\text{Os}$			63Gr08
$^{186}\text{Au}(\alpha)^{182}\text{Ir}$	4907	15	4912	14	0.3	1	87	44 $^{182}\text{Ir}$	ORa		90Ak04 *
$^{186}\text{Hg}(\alpha)^{182}\text{Pt}$	5206.2	15.	5204	10	-0.1	-					70Ha18
	5204.2	15.			0.0	-					96Ri12
	ave.	5205			-0.1	1	83	57 $^{182}\text{Pt}$			average
$^{186}\text{Tl}^m(\alpha)^{182}\text{Au}^p$	5891.9	7.	5892	5	0.0	4					75Co.A
	5891.9	7.			0.0	4			ORa		77Ij01
$^{186}\text{Pb}(\alpha)^{182}\text{Hg}$	6458.2	20.	6470	6	0.6	2			Ora		74Le02 Z
	6470.1	10.			0.0	2			GSa		80Sc09 Z
	6474.7	10.			-0.5	2			ORa		84To09 Z
	6476.5	15.			-0.4	2			ORa		97Ba25
	6459.2	15.			0.7	2			Anv		97An09
$^{186}\text{Bi}(\alpha)^{182}\text{Tl}$	7760	20	7757	12	-0.2	3			Ara		97Ba21 *
	7755	15			0.1	3			Anv		03An27 *
$^{186}\text{Bi}^m(\alpha)^{182}\text{Tl}^p$	7349.3	25.	7423	5	2.9	U			GSa		84Sc.A
	7420.9	20.			0.1	U			Ara		97Ba21
	7422.9	5.				4			Anv		03An27
$^{186}\text{Po}(\alpha)^{182}\text{Pb}$	8493	30	8501	14	0.3	5					05Hu.A
	8503.2	15.3			-0.1	5					13An13
$^{186}\text{W}(p,t)^{184}\text{W}$	-4474	5	-4464.0	1.2	2.0	U			Min		73Oo01
$^{186}\text{W}(p,t)^{184}\text{W}-^{184}\text{W}^{182}\text{W}$	660.1	1.6	656.2	1.2	-2.5	o					09Le03
	657.0	1.8			-0.5	1	42	35 $^{186}\text{W}$			09Le.A
$^{186}\text{W}(t,\alpha)^{185}\text{Ta}$	11430	20	11411	14	-1.0	R			LAl		80Lo10
$^{186}\text{W}(\gamma,n)^{185}\text{W}$	-7120	60	-7192.1	1.2	-1.2	U			Phi		60Ge01
$^{186}\text{W}(d,t)^{185}\text{W}$	-939	10	-934.8	1.2	0.4	U			Kop		72Ca01
$^{185}\text{Re}(n,\gamma)^{186}\text{Re}$	6179.8	0.8	6179.38	0.17	-0.5	-			Tal		69La11 Z
	6178.6	1.5			0.5	U					70Or.A
	6179.34	0.18			0.2	-			Bdn		06Fi.A
$^{185}\text{Re}(d,p)^{186}\text{Re}$	3939	25	3954.81	0.17	0.6	U			Tal		69La11
$^{185}\text{Re}(n,\gamma)^{186}\text{Re}$	ave.	6179.36	6179.38	0.17	0.1	1	99	72 $^{186}\text{Re}$			average
$^{186}\text{Ta}(\beta^-)^{186}\text{W}$	3901	60				2					69Mo16 *
$^{186}\text{Re}(\beta^-)^{186}\text{Os}$	1064	2	1072.9	0.8	4.4	B					56Jo05
	1071.5	1.3			1.0	-					56Po28
	1076	3			-1.0	-					64Ma36
	1064	3			3.0	B					68An11
	ave.	1072.2			0.5	1	49	28 $^{186}\text{Re}$			average
$^{186}\text{Ir}(\beta^+)^{186}\text{Os}$	3760	200	3828	17	0.3	U					62Bo22 *
	3831	20			-0.2	R					63Em02
$^{186}\text{Au}(\beta^+)^{186}\text{Pt}$	5950	200	6150	30	1.0	U					72We.A
$^{186}\text{Hg}(\beta^+)^{186}\text{Au}$	3250	200	3176	24	-0.4	U					72We.A
$^{186}\text{Tl}^n(\text{IT})^{186}\text{Tl}^m$	373.9	0.5	374.00	0.20	0.2	o			Lvn		91Va04
	374.0	0.2				3					Ens036
* $^{186}\text{Ir}-u$	$M - A = -39181(28)$ keV for mixture gs+m at 0.8 keV										Nub16b **
* $^{186}\text{Tl}-u$	$M - A = -20030(180)$ keV for mixture gs+m+n at 20(40) and 400(40) keV										Nub16b **
* $^{186}\text{Tl}-u$	$M - A = -19900(29)$ keV for mixture gs+m+n at 20(40) and 400(40) keV										Nub16b **
* $^{186}\text{Tl}-^{133}\text{Cs}_{1.398}$	$D_M = 110842.1(9.2)$ $\mu\text{u}$ for mixture gs+m at 20(40) keV; $M - A = -19874.4(8.6)$ keV										Nub16b **
* $^{186}\text{Tl}-^{133}\text{Cs}_{1.398}$	$D_M = 110840.4(8.6)$ $\mu\text{u}$ for mixture gs+m at 20(40) keV; $M - A = -19876.0(8.0)$ keV										Nub16b **
* $^{186}\text{Au}(\alpha)^{182}\text{Ir}$	$E_\alpha = 4653(15)$ to $3^-$ level at 152.3 keV										95Sa42 **
* $^{186}\text{Bi}(\alpha)^{182}\text{Tl}$	$E_\alpha = 7158(20)$ followed by $E(\gamma) = 444$ keV										03An27 **
* $^{186}\text{Bi}(\alpha)^{182}\text{Tl}$	$E_\alpha = 7152(15), 7085(15)$ followed by $E(\gamma) = 444, 520$ keV										03An27 **
* $^{186}\text{Ta}(\beta^-)^{186}\text{W}$	$E_{\beta^-} = 2240(60)$ to $(2^-, 3^-)$ level at 1661.3817 keV										Ens036 **
* $^{186}\text{Ir}(\beta^+)^{186}\text{Os}$	$E_{\beta^+} = 2600(200)$ assumed to $2^+$ level at 137.159 keV, also other $E_{\beta^+}$										Ens036 **
* $^{186}\text{Ir}(\beta^+)^{186}\text{Os}$	$E_{\beta^+} = 1940(20)$ to $6^+$ level at 868.94 keV										Ens036 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{187}\text{Ta}-u$	-39609	60				2			GS3	1.0	13Sh30
$^{187}\text{Ir}-u$	-42458	30				2			GS2	1.0	05Li24
$^{187}\text{Pt}-u$	-39500	110	-39383	26	1.1	U			GS1	1.0	00Ra23
	-39413	30			1.0	1	74	74 $^{187}\text{Pt}$	GS2	1.0	05Li24
$^{187}\text{Au}-u$	-35470	114	-35457	24	0.1	U			GS1	1.0	00Ra23
	-35441	30			-0.5	1	64	64 $^{187}\text{Au}$	GS2	1.0	05Li24
$^{187}\text{Hg}-u$	-30188	109	-30186	15	0.0	U			GS1	1.0	00Ra23
	-30155	36			-0.9	1	17	17 $^{187}\text{Hg}$	GS2	1.0	05Li24
$^{187}\text{Hg}_{-208}\text{Pb}_{.899}$	-9210	20	-9196	15	0.7	1	56	56 $^{187}\text{Hg}$	MA6	1.0	01Sc41
$^{187}\text{Hg}^m_{-208}\text{Pb}_{.899}$	-9152	19	-9132	21	1.0	o			MA6	1.0	01Sc41
$^{187}\text{Tl}-u$	-24120	107	-24095	9	0.2	U			GS1	1.0	00Ra23
	-23928	109			-1.5	U			GS2	1.0	05Li24
$^{187}\text{Tl}^m_{-133}\text{Cs}_{1.406}$	109151	24	109198	8	1.9	F			MA8	1.0	08We02
$^{187}\text{Pb}-u$	-16076	45	-16089	5	-0.3	U			GS2	1.0	05Li24
$^{187}\text{Pb}_{-133}\text{Cs}_{1.406}$	116843.5	5.9	116845	5	0.3	1	86	86 $^{187}\text{Pb}$	MA8	1.0	05We11
$^{187}\text{Pb}^m_{-133}\text{Cs}_{1.406}$	116871.6	5.6	116866	12	-1.0	o			MA8	1.0	05We11
$^{187}\text{Re O}_2_{-184}\text{W }^{35}\text{Cl}$	25797.4	3.5	25795.6	0.9	-0.2	U			H28	2.5	77Sh04
$^{187}\text{Re }^{35}\text{Cl}_{-185}\text{Re }^{37}\text{Cl}$	5737	3	5744.1	0.9	0.9	U			H22	2.5	70Mc03
	5744.2	1.2			0.0	1	10	6 $^{185}\text{Re}$	H28	2.5	77Sh04
$^{187}\text{Re}_{-187}\text{Os}$	2.676	0.036	2.6481	0.0017	-0.8	U			SH1	1.0	14Ne15
$^{187}\text{Au}(\alpha)^{183}\text{Ir}$	4792.7	20.	4751	29	-0.8	1	35	19 $^{183}\text{Ir}$			68Si01
$^{187}\text{Hg}(\alpha)^{183}\text{Pt}$	5229.9	20.	5230	14	0.0	1	49	30 $^{183}\text{Pt}$	ISa		70Ha18
$^{187}\text{Hg}^m(\alpha)^{183}\text{Pt}$	5293.4	20.	5289	16	-0.2	1	64	49 $^{187}\text{Hg}^m$	ISa		70Ha18
$^{187}\text{Tl}^m(\alpha)^{183}\text{Au}$	5643	20	5656	6	0.6	-			ORa		76To06
	5661.5	10.			-0.6	-			GSa		80Sc09
	5645.1	12.			0.9	o			Lvn		85Co06
	5661.5	10.			-0.6	-			Lvn		91Wa21
ave.	5659	7			-0.6	1	91	77 $^{183}\text{Au}$			average
$^{187}\text{Pb}(\alpha)^{183}\text{Hg}$	6393.0	10.	6393	6	0.0	-			Ora		75Ca06
	6395.0	19.			-0.1	o			GSa		80Sc09
	6398.4	10.			-0.6	-			GSa		81Mi12
ave.	6396	7			-0.4	1	77	63 $^{183}\text{Hg}$			average
$^{187}\text{Pb}^m(\alpha)^{183}\text{Hg}^m$	6213.1	20.	6208	7	-0.2	o			Ora		74Le02
	6213.1	10.			-0.5	2			Ora		75Ca06
	6223.3	10.			-1.5	o			GSa		80Sc09
	6206.0	10.2			0.2	2			GSa		81Mi12
	6202.9	15.			0.4	2			Anv		99An36
$^{187}\text{Bi}(\alpha)^{183}\text{Tl}^m$	7139.0	10.	7150	4	1.1	2			GSa		84Sc.A
	7153.3	8.			-0.4	2			ORa		99Ba45
	7158.4	10.			-0.8	o			Anv		03An27
	7147.2	8.			0.4	2			Jya		03Ke08
	7153.3	10.			-0.3	o			Anv		04An07
	7153.3	5.			-0.6	2			Anv		06An11
$^{187}\text{Bi}^m(\alpha)^{183}\text{Tl}$	7749.1	10.	7887	7	13.8	F			GSa		84Sc.A
	7890.1	15.			-0.2	2			ORa		99Ba45
	7882.9	11.			0.4	2			Jya		03Ke08
	7890.1	10.			-0.3	2			Anv		06An11
$^{187}\text{Po}(\alpha)^{183}\text{Pb}$	7978.9	15.				3			Anv		06An11
$^{187}\text{Po}^m(\alpha)^{183}\text{Pb}^m$	7889.1	20.				3			Anv		06An11
$^{186}\text{W}(n,\gamma)^{187}\text{W}$	5466.3	0.3	5466.76	0.04	1.5	U			BNn		87Br05
	5467.22	0.3			-1.5	U			Ltn		92Be17
	5466.59	0.12			1.4	o			Bdn		06Fi.A
	5466.83	0.05			-1.4	-			Prn		08Bo26
	5466.62	0.07			2.0	-			Bdn		14Hu02
$^{186}\text{W}(d,p)^{187}\text{W}$	3236	5	3242.19	0.04	1.2	U			ANL		65Er03
	3240	10			0.2	U			Kop		72Ca01
$^{186}\text{W}(n,\gamma)^{187}\text{W}$	ave.	5466.76	0.04	5466.76	0.04	0.0	1	100	55 $^{186}\text{W}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{186}\text{W}(\text{}^3\text{He,d})^{187}\text{Re}$	530	40	503.4	1.1	-0.7	U			Roc		71Lu01
$^{187}\text{Re}(\gamma,\text{n})^{186}\text{Re}$	-7180	80	-7360.7	0.9	-2.3	U			Phi		60Ge01
$^{187}\text{Re}(\text{d,t})^{186}\text{Re}$	-1055	25	-1103.5	0.9	-1.9	U			Tal		69La11
$^{186}\text{Os}(\text{n},\gamma)^{187}\text{Os}$	6291.1	1.0	6290.3	0.5	-0.8	-					74Pr15 Z
	6289.4	0.8			1.1	-			Bdn		06Fi.A
ave.	6290.1	0.6			0.4	1	70	40 $^{186}\text{Os}$			average
$^{187}\text{W}(\beta^-)^{187}\text{Re}$	1314	2	1312.5	1.1	-0.7	-					69Na03
	1310	2			1.3	-					70He14
ave.	1312.0	1.4			0.4	1	63	55 $^{187}\text{W}$			average
$^{187}\text{Re}(\beta^-)^{187}\text{Os}$	2.62	0.09	2.4667	0.0016	-1.7	U					65Br12
	2.64	0.05			-3.5	B					67Hu05
	2.667	0.020			-10.0	B					92Co23
	2.70	0.09			-2.6	U					93As02
	2.460	0.011			0.6	U					99Al20
	2.470	0.004			-0.8	-					01Ga01
	2.4661	0.0017			0.4	-					03Ar36
ave.	2.4667	0.0016			0.0	1	100	89 $^{187}\text{Re}$			average
$^{187}\text{Ir}(\beta^+)^{187}\text{Os}$	1550	200	1670	28	0.6	U					71Ma24 *
$^{187}\text{Os}(\text{}^3\text{He,t})^{187}\text{Ir}$	-1521	6	-1688	28	-27.9	B			INS		90Ka27
$^{187}\text{Au}(\beta^+)^{187}\text{Pt}$	3600	40	3657	27	1.4	1	47	26 $^{187}\text{Pt}$			83Gn01 *
$^{187}\text{Hg}^m(\text{IT})^{187}\text{Hg}$	54	21	59	16	0.2	1	60	51 $^{187}\text{Hg}^m$	MA6		01Sc41 *
$^{187}\text{Tl}^m(\text{IT})^{187}\text{Tl}$	330	5	334	3	0.7	1	45	31 $^{187}\text{Tl}$			77Sc03
$^{187}\text{Pb}^m(\text{IT})^{187}\text{Pb}$	33	13	19	10	-1.1	1	61	61 $^{187}\text{Pb}^m$	MA8		05We11
* $^{187}\text{Au}-\text{u}$	$M - A = -32980(100)$ keV for mixture gs+m at 120.33 keV										
* $^{187}\text{Hg}-\text{u}$	$M - A = -28090(100)$ keV for mixture gs+m at 59(16) keV										
* $^{187}\text{Hg}-\text{u}$	$M - A = -28060(28)$ keV for mixture gs+m at 59(16) keV										
* $^{187}\text{Hg}^m - ^{208}\text{Pb}_{.899}$	Use instead their difference between ground state and $^{187}\text{Hg}^m$ lines										
* $^{187}\text{Tl}-\text{u}$	$M - A = -22121(28)$ keV for mixture gs+m at 334(3) keV										
* $^{187}\text{Tl}^m - ^{133}\text{Cs}_{1.406}$	F : contamination from ground state not resolved										
* $^{187}\text{Pb}-\text{u}$	$M - A = -14965(41)$ keV for mixture gs+m at 19(10) keV										
* $^{187}\text{Pb} - ^{133}\text{Cs}_{1.406}$	$D_M = 116851.3(4.3)$ $\mu\text{u}$ for mixture gs+m at 19(10) keV with $R=0.62(0.02)$ ;										
*	$M - A = -14981.5(4.0)$ keV										
* $^{187}\text{Pb}^m - ^{133}\text{Cs}_{1.406}$	$D_M = 116869.5(5.5)$ for mixture gs+m at 19(10) $R=8.7(0.7)$ ; $M - A = -14964.5(5.1)$ keV										
*	used are only the equations for the $^{187}\text{Pb}$ doublet and $^{187}\text{Pb}^m(\text{IT})^{187}\text{Pb}$										
* $^{187}\text{Au}(\alpha)^{183}\text{Ir}$	Assignment uncertain										
* $^{187}\text{Hg}(\alpha)^{183}\text{Pt}$	$E_\alpha = 5035(20)$ to $3/2^-$ level at 84.73 keV										
* $^{187}\text{Hg}^m(\alpha)^{183}\text{Pt}$	$E_\alpha = 4870(20)$ to $(13/2^+)$ level at 316.9 level										
* $^{187}\text{Tl}^m(\alpha)^{183}\text{Au}$	$E_\alpha = 5510(20)$ 5528(10) 5512(12) 5528(10) respectively, to $(9/2)^-$ at 12.4(0.4)keV										
* $^{187}\text{Pb}(\alpha)^{183}\text{Hg}$	$E_\alpha = 6190(10)$ to $3/2^-$ level at 67.16 keV										
* $^{187}\text{Pb}(\alpha)^{183}\text{Hg}$	$E_\alpha = 6194(10)$ 5993(10) to $3/2^-$ levels at 67.16 and 275.33 keV										
* $^{187}\text{Bi}(\alpha)^{183}\text{Tl}^m$	Also $E_\alpha = 7612(15)$ keV to ground state										
* $^{187}\text{Bi}(\alpha)^{183}\text{Tl}^m$	Also $E_\alpha = 7605(16)$ keV to ground state										
* $^{187}\text{Bi}(\alpha)^{183}\text{Tl}^m$	Also $E_\alpha = 7612(5)$ , 7342(15) keV to ground state, 273(1) keV										
* $^{187}\text{Bi}^m(\alpha)^{183}\text{Tl}$	F : for T=700 $\mu\text{s}$ instead of Nubase=370(20) $\mu\text{s}$										
* $^{187}\text{Po}(\alpha)^{183}\text{Pb}$	$E_\alpha = 7528(15)$ to 286(1) keV level; also 1 event $E_\alpha = 7796(15)$ to ground state										
* $^{186}\text{W}(\text{n},\gamma)^{187}\text{W}$	Only statistical error 0.04 keV given; Z recalibrated										
* $^{187}\text{Ir}(\beta^+)^{187}\text{Os}$	$p^+ < 0.15(0.05)$ , resulting $Q_i(1550)$ keV										
* $^{187}\text{Au}(\beta^+)^{187}\text{Pt}$	$K/\beta^+ = 31.6(2.8)$ to $1/2^+$ level at 1341.07 keV, recalculated										
* $^{187}\text{Hg}^m(\text{IT})^{187}\text{Hg}$	Original error (7 keV) increased by 20 due to isomer+ground state lines in trap										
$^{188}\text{Ta}-\text{u}$	-36084	59				2			GS3	1.0	13Sh30
$^{188}\text{Au}-\text{u}$	-34750	104	-34752.0	2.9	0.0	U			GS1	1.0	00Ra23
	-34674	30			-2.6	B			GS2	1.0	05Li24
$^{188}\text{Au} - ^{133}\text{Cs}_{1.414}$	98938.9	2.9				2			MA8	1.0	16Ma.1
$^{188}\text{Hg}-\text{u}$	-32500	104	-32423	13	0.7	U			GS1	1.0	00Ra23
	-32428	30			0.2	1	19	19 $^{188}\text{Hg}$	GS2	1.0	05Li24
$^{188}\text{Hg} - ^{208}\text{Pb}_{.904}$	-11330	20	-11316	13	0.7	-			MA6	1.0	01Sc41
ave.	-11317	17			0.0	1	63	62 $^{188}\text{Hg}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{188}\text{Tl}-u$	-23827	110	-23980	30	-1.4	U			GS1	1.0	00Ra23 *
	-23994	38			0.4	2			GS2	1.0	05Li24 *
$^{188}\text{Pb}-u$	-19070	110	-19125	11	-0.5	U			GS1	1.0	00Ra23
	-19144	30			0.6	R			GS2	1.0	05Li24
$^{188}\text{Os } ^{35}\text{Cl}-^{186}\text{W } ^{37}\text{Cl}$	4426	3	4422.3	1.2	-0.5	U			H22	2.5	70Mc03
$^{188}\text{Pt}(\alpha)^{184}\text{Os}$	4015.7	10.	4007	5	-0.9	-					63Gr08
	4000.3	10.			0.6	-			ORa		78El11
	3990.1	15.			1.1	-					79Ha10
	ave.	4005	7		0.3	1	65	65 $^{188}\text{Pt}$			average
$^{188}\text{Hg}(\alpha)^{184}\text{Pt}$	4710.4	20.	4707	16	-0.1	1	59	40 $^{184}\text{Pt}$			79Ha10
$^{188}\text{Pb}(\alpha)^{184}\text{Hg}$	6110.3	10.	6109	3	-0.1	2			Ora		74Le02 Z
	6109.2	10.			0.0	2			Ora		77De32 Z
	6120.5	15.			-0.8	2			GSa		80Sc09 Z
	6110.5	5.			-0.3	2			ORa		81To02 Z
	6109.3	10.			0.0	2			Lvn		93Wa03 Z
	6100.0	8.			1.1	2			Jya		03Ke04
$^{188}\text{Bi}(\alpha)^{184}\text{Tl}$	7274.5	25.	7264	5	-0.4	o			GSa		80Sc09 *
	7279.7	10.			-1.6	2			GSa		84Sc.A *
	7255.2	7.			1.2	2			Lvn		97Wa05 *
	7259.3	5.			0.9	o			Anv		03An26 *
	7264.8	10.			-0.1	2			Anv		06An04 *
	7462.9	5.				5			Anv		03An26 *
$^{188}\text{Bi}^n(\alpha)^{184}\text{Tl}^m$	6968.5	20.	6965	5	-0.2	o			GSa		80Sc09
$^{188}\text{Bi}^n(\alpha)^{184}\text{Tl}^n$	6968.5	10.			-0.4	5			GSa		84Sc.A
	6963.5	6.			0.2	5			Lvn		97Wa05
	6961.3	5.			0.6	o			Anv		03An26
	6963.5	5.			0.1	5			Anv		06An04
	8087.4	25.	8082	15	-0.2	o			Anv		99An52
	8080.2	15.			0.1	o			Anv		01Va.B
$^{188}\text{Po}(\alpha)^{184}\text{Pb}$	8082.3	15.				2			Anv		03Va16
	-5802	5	-5798.1	0.5	0.8	U			Min		73Oo01
$^{187}\text{Re}(n,\gamma)^{188}\text{Re}$	-5803	4			1.2	U			McM		75Th04
	5871.77	0.3	5871.65	0.04	-0.4	U					72Sh13 Z
	5871.75	0.13			-0.8	U			Bdn		06Fi.A
	5871.65	0.04				2			Prn		10Ba48
$^{188}\text{Os}(t,\alpha)^{187}\text{Re}$	12604	10	12604.14	0.15	0.0	U			McM		76Hi08
$^{187}\text{Os}(n,\gamma)^{188}\text{Os}$	7989.6	0.3	7989.61	0.15	0.0	-					83Fe06 Z
	7989.58	0.17			0.1	-			Bdn		06Fi.A
	ave.	7989.58	0.15		0.1	1	98	57 $^{187}\text{Os}$			average
$^{188}\text{W}(\beta^-)^{188}\text{Re}$	349	3				3					64Bu10
$^{188}\text{Re}(\beta^-)^{188}\text{Os}$	2116	2	2120.42	0.15	2.2	U					56Jo05
	2111	3			3.1	B					68An11
$^{188}\text{Ir}(\beta^+)^{188}\text{Os}$	2833	10	2792	9	-4.1	B					62Wa20 *
	2781	20			0.6	-					69Ya02 *
	2827	30			-1.2	-					70Ag03 *
ave.	2795	17			-0.2	1	32	32 $^{188}\text{Ir}$			average
$^{188}\text{Pt}(\epsilon)^{188}\text{Ir}$	525	10	524	9	-0.1	1	75	68 $^{188}\text{Ir}$	ORa		78El11 *
$^{188}\text{Au}(\beta^+)^{188}\text{Pt}$	5520	30	5450	6	-2.3	U					84Da.A
$^{188}\text{Hg}(\beta^+)^{188}\text{Au}$	2040	20	2169	13	6.5	B					84Da.A
* $^{188}\text{Tl}-u$	$M - A = -22180(100)$ keV for mixture gs+m at 30(40) keV										
* $^{188}\text{Tl}-u$	$M - A = -22335(28)$ keV for mixture gs+m at 30(40) keV										
* $^{188}\text{Bi}(\alpha)^{184}\text{Tl}$	$E_\alpha = 7005(25)$ 7010(10) 6987(6) respectively, to $(3^+)$ level at 117.5(0.5) keV										
*	$E_\alpha = 7029(7)$ 3 times weaker exists too, possible mixture in older results										
* $^{188}\text{Bi}(\alpha)^{184}\text{Tl}$	$E_\alpha = 7106(5)$ , 6992(5), 6889(10) to ground state, 117.5, 216 levels										
* $^{188}\text{Bi}(\alpha)^{184}\text{Tl}$	$E_\alpha = 6995(10)$ to 117.5 level										
* $^{188}\text{Bi}^n(\alpha)^{184}\text{Tl}^m$	$E_\alpha = 7302(5)$ , 7232(10), 6995(15) to ground state, 70.5, 320 levels										
* $^{188}\text{Ir}(\beta^+)^{188}\text{Os}$	$E_{\beta^+} = 1656(10)$ 1605(20) 1650(30) respectively, to $2^+$ level at 155.021 keV										
* $^{188}\text{Pt}(\epsilon)^{188}\text{Ir}$	$pL = 0.67(0.05)$ to $1^+$ level at 478.17 keV										



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{189}\text{W}-u$	-38237	43				2			GS3	1.0	13Sh30
$\text{C}_{14}\text{H}_{21}-^{189}\text{Os}$	206188.3	6.2	206179.7	0.7	-0.3	U			M23	4.0	79Ha32
$^{189}\text{Au}-u$	-36080	140	-36052	22	0.2	U			GS1	1.0	00Ra23 *
	-36045	31			-0.2	2			GS2	1.0	05Li24
	-36058	30			0.2	2			GS2	1.0	05Li24 *
$^{189}\text{Hg}-u$	-31788	111	-31810	30	-0.2	U			GS1	1.0	00Ra23 *
	-31791	42			-0.3	1	65	65 $^{189}\text{Hg}$	GS2	1.0	05Li24 *
$^{189}\text{Hg}^m-^{208}\text{Pb}_{909}$	-10501	20	-10498	19	0.2	1	92	92 $^{189}\text{Hg}^m$	MA6	1.0	01Sc41
$^{189}\text{Tl}-u$	-26497	139	-26426	9	0.5	U			GS1	1.0	00Ra23 *
	-26313	93			-1.2	U			GS2	1.0	05Li24 *
$^{189}\text{Pb}-u$	-19206	97	-19156	15	0.5	U			GS1	1.0	00Ra23 *
	-19193	34			1.1	1	20	20 $^{189}\text{Pb}$	GS2	1.0	05Li24 *
$^{189}\text{Os } ^{35}\text{Cl}-^{187}\text{Re } ^{37}\text{Cl}$	5341	3	5343.8	0.5	0.4	U			H22	2.5	70Mc03
$^{189}\text{Pb}(\alpha)^{185}\text{Hg}$	5954.2	10.	5915	4	-3.9	B			Ora		72Ga27 *
	5943.9	10.			-2.9	B			Ora		74Le02 *
	5915	10			0.0	-					05Fr.A *
	5914.8	5.4			0.0	-			ISa		13Sa43 *
	ave.	5915	5		0.0	1	82	67 $^{189}\text{Pb}$			average
$^{189}\text{Pb}^m(\alpha)^{185}\text{Hg}$	5958	10	5955	5	-0.3	1	28	25 $^{189}\text{Pb}^m$			05Fr.A *
	5955	5			0.0	o			ISa		13Sa43 *
$^{189}\text{Bi}(\alpha)^{185}\text{Tl}$	7269.4	10.	7268.2	2.7	-0.1	6			Ora		74Le02 *
	7274.5	10.			-0.6	6			GSa		84Sc.A *
	7271.2	5.			-0.6	6			Lvn		85Co06 *
	7271.8	15.			-0.2	U			Anv		97An09 *
	7268.1	6.			0.0	6			Lvn		97Wa05
	7271.5	5.			-0.7	o			Jya		02Hu14 *
	7264.2	4.5			0.9	6			Jya		03Ke08 *
$^{189}\text{Bi}^m(\alpha)^{185}\text{Tl}$	7362.1	20.	7452	4	1.8	U			GSa		84Sc.A *
	7499.0	30.			-1.6	U			Dbb		93An19
	7458.2	40.			-0.2	U			ORa		95Ba75
	7458.2	15.			-0.4	6			Anv		97An09
	7450.0	6.			0.4	6			Lvn		97Wa05
	7453.1	6.			-0.2	6			Jya		03Ke08
$^{189}\text{Po}(\alpha)^{185}\text{Pb}$	7699.4	15.	7694	15	-0.3	o			Anv		99An52 *
	7694.3	15.				3			Anv		05Va04 *
$^{189}\text{Os}(p,t)^{187}\text{Os}$	-5431	5	-5428.6	0.5	0.5	U			Min		73Oo01
	-5432	4			0.8	U			McM		75Th04
$^{188}\text{Os}(n,\gamma)^{189}\text{Os}$	5920.8	2.	5920.8	0.4	0.0	U					76Be50
	5920.6	0.5			0.5	1	80	59 $^{188}\text{Os}$	ILn		92Br17
	5922.0	0.4			-2.9	C			Bdn		06Fi.A
$^{188}\text{Os}(d,p)^{189}\text{Os}$	3689	10	3696.3	0.4	0.7	U			Kop		75Mo29
$^{189}\text{Os}(d,t)^{188}\text{Os}$	335	15	336.4	0.4	0.1	U			Tal		75Th06
$^{189}\text{W}(\beta^-)^{189}\text{Re}$	2500	200	2360	40	-0.7	U					65Ka07
$^{189}\text{Re}(\beta^-)^{189}\text{Os}$	1000	20	1008	8	0.4	R					63Cr06
	1015	20			-0.4	R					65B106
$^{189}\text{Pt}(\beta^+)^{189}\text{Ir}$	1950	20	1980	14	1.5	1	46	30 $^{189}\text{Ir}$			71P108 *
$^{189}\text{Au}(\beta^+)^{189}\text{Pt}$	3160	300	2887	22	-0.9	U					75Un.A
$^{189}\text{Hg}(\beta^+)^{189}\text{Au}$	4200	200	3960	40	-1.2	U					75Un.A
$^{189}\text{Hg}^m(\text{IT})^{189}\text{Hg}$	100	50	80	30	-0.4	1	43	35 $^{189}\text{Hg}$	MA6		01Sc41
$^{189}\text{Tl}^m(\beta^+)^{189}\text{Hg}$	5460	200	5300	30	-0.8	U					75Un.A *
$^{189}\text{Pb}^m(\text{IT})^{189}\text{Pb}$	40	4	40	4	0.1	1	88	75 $^{189}\text{Pb}^m$	ISa		13Sa43
* $^{189}\text{Au}-u$	$M-A=-33490(100)$ keV for mixture gs+m at 247.23 keV										Nub16b **
* $^{189}\text{Au}-u$	$M-A=-33341(28)$ keV for $^{189}\text{Au}^m$ at 247.23 keV										Nub16b **
* $^{189}\text{Hg}-u$	$M-A=-29570(100)$ keV for mixture gs+m at 80(30) keV										Nub16b **
* $^{189}\text{Hg}-u$	$M-A=-29573(28)$ keV for mixture gs+m at 80(30) keV										Nub16b **
* $^{189}\text{Tl}-u$	$M-A=-24540(100)$ keV for mixture gs+m at 285(6) keV										Nub16b **
* $^{189}\text{Tl}-u$	$M-A=-24369(28)$ keV for mixture gs+m at 285(6) keV										Nub16b **
* $^{189}\text{Pb}-u$	$M-A=-17870(90)$ keV for mixture gs+m at 40(4) keV										Nub16b **
* $^{189}\text{Pb}-u$	$M-A=-17858(29)$ keV for mixture gs+m at 40(4) keV										Nub16b **
* $^{189}\text{Pb}(\alpha)^{185}\text{Hg}$	$E_\alpha=5730.1(10,Z)$ possibly from ground state, to $3/2^-$ level at 26.1 keV										Ens061 **
* $^{189}\text{Pb}(\alpha)^{185}\text{Hg}$	$E_\alpha=5720(10)$ possibly from ground state, to $3/2^-$ level at 26.1 keV										Ens061 **
* $^{189}\text{Pb}(\alpha)^{185}\text{Hg}$	$E_\alpha=5761$ to $3/2^-$ level at 26.1 and $E_\alpha=5623$ to 173.8 level										05Fr.A **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>189</sup> Pb( $\alpha$ ) <sup>185</sup> Hg	$E_\alpha=5764(3)(5)$ to $3/2^-$ level at 26.02 and $E_\alpha=5619(2)(5)$ to 173.9 level										13Sa43 **
* <sup>189</sup> Pb <sup>m</sup> ( $\alpha$ ) <sup>185</sup> Hg	$E_\alpha=5730$ to 103.8 level										05Fr.A **
* <sup>189</sup> Pb <sup>m</sup> ( $\alpha$ ) <sup>185</sup> Hg	$E_\alpha=5727(2)(5)$ to 103.7(0.4) level										13Sa43 **
* <sup>189</sup> Pb <sup>m</sup> ( $\alpha$ ) <sup>185</sup> Hg	use instead <sup>189</sup> Pb <sup>m</sup> (IT) <sup>189</sup> Pb where systematic errors cancel										GAu **
* <sup>189</sup> Bi( $\alpha$ ) <sup>185</sup> Tl	$E_\alpha=6670.1(10,Z)$ to <sup>185</sup> Tl <sup>m</sup> at 454.8(1.5) keV										Nub16b **
* <sup>189</sup> Bi( $\alpha$ ) <sup>185</sup> Tl	$E_\alpha=6675(10)$ to <sup>185</sup> Tl <sup>m</sup> at 454.8(1.5) keV										Nub16b **
* <sup>189</sup> Bi( $\alpha$ ) <sup>185</sup> Tl	$E_\alpha=7115.6(15,Z)$ and $6671.6(5,Z)$ to <sup>185</sup> Tl <sup>m</sup> at 454.8(1.5) keV										Nub16b **
* <sup>189</sup> Bi( $\alpha$ ) <sup>185</sup> Tl	$E_\alpha=7120(15)$ , $6670(15)$ to ground state and <sup>185</sup> Tl <sup>m</sup> at 454.8(1.5) keV										Nub16b **
* <sup>189</sup> Bi( $\alpha$ ) <sup>185</sup> Tl	$E_\alpha=6674(5)$ to <sup>185</sup> Tl <sup>m</sup> at 452.8(2.0) keV										77Sc03 **
* <sup>189</sup> Bi( $\alpha$ ) <sup>185</sup> Tl	$E_\alpha=6667(4)$ to <sup>185</sup> Tl <sup>m</sup> at 452.8(2.0) keV										77Sc03 **
*	and also $E_\alpha=7114(6)$ to ground state										03Ke08 **
* <sup>189</sup> Bi <sup>m</sup> ( $\alpha$ ) <sup>185</sup> Tl	Only one event; not seen in reference										93An19 **
* <sup>189</sup> Po( $\alpha$ ) <sup>185</sup> Pb	$E_\alpha=7264(15)$ to 278(1) level										99An52 **
* <sup>189</sup> Po( $\alpha$ ) <sup>185</sup> Pb	$E_\alpha=7259(15)$ to 278(1) level										05Va04 **
* <sup>189</sup> Pt( $\beta^+$ ) <sup>189</sup> Ir	$E_{\beta^+}=885(10)$ to ground state, $1/2^+$ level at 94.34 and $3/2^+$ at 176.53 keV										Ens039 **
* <sup>189</sup> Tl <sup>m</sup> ( $\beta^+$ ) <sup>189</sup> Hg	$E_{\beta^+}=4140(200)$ to several levels around 300 keV										75Un.A **
<sup>190</sup> W-u	-36917	44	-36910	40	0.1	1	94	94 <sup>190</sup> W	GS3	1.0	13Sh30
<sup>190</sup> Au-u	-35213	106	-35248	4	-0.3	U			GS2	1.0	05Li24 *
<sup>190</sup> Au- <sup>133</sup> Cs <sub>1.429</sub>	99860.9	3.7				2			MA8	1.0	16Ma.1
<sup>190</sup> Hg-u	-33670	107	-33678	17	-0.1	U			GS1	1.0	00Ra23
<sup>190</sup> Hg- <sup>208</sup> Pb <sub>.913</sub>	-12361	20	-12361	17	0.0	1	73	73 <sup>190</sup> Hg	MA6	1.0	01Sc41
<sup>190</sup> Tl <sup>m</sup> -u	-26055	107	-26076	7	-0.2	o			GS1	1.0	00Ra23 *
	-26048	30			-0.9	U			GS2	1.0	05Li24 *
<sup>190</sup> Tl <sup>m</sup> - <sup>133</sup> Cs <sub>1.429</sub>	109033.5	6.9				2			MA8	1.0	14Bo26
<sup>190</sup> Pb-u	-21940	104	-21918	13	0.2	U			GS1	1.0	00Ra23
	-21905	30			-0.4	R			GS2	1.0	05Li24
<sup>190</sup> Bi <sup>m</sup> - <sup>133</sup> Cs <sub>1.429</sub>	123800	27	123870	30	2.4	F			MA8	1.0	08We02 *
<sup>186</sup> Os- <sup>190</sup> Pt <sub>.979</sub>	-6953.13	0.86	-6953.3	0.6	-0.2	1	53	39 <sup>186</sup> Os	MS1	1.0	16Ei01
<sup>190</sup> Os- <sup>35</sup> Cl- <sup>188</sup> Os <sub>.37</sub> Cl	5557	3	5558.2	0.5	0.2	U			H22	2.5	70Mc03
<sup>190</sup> Os- <sup>190</sup> Pt	-1504.31	0.59	-1504.4	0.5	-0.1	1	62	33 <sup>190</sup> Pt	MS1	1.0	16Ei01
<sup>190</sup> Os-C <sub>14</sub> H <sub>21</sub>	-205897.8	5.8	-205880.2	0.7	0.8	U			M23	4.0	79Ha32
<sup>190</sup> Os- <sup>189</sup> Os	285.2	5.2	299.49	0.20	1.1	U			M24	2.5	79Ha32
<sup>190</sup> Pt( $\alpha$ ) <sup>186</sup> Os	3238.3	20.	3268.6	0.6	1.5	U					61Pe23
	3248.5	20.			1.0	U					63Gr08
<sup>190</sup> Pb( $\alpha$ ) <sup>186</sup> Hg	5699.8	10.	5698	5	-0.2	2			Ora		74Le02 Z
	5697.0	5.			0.1	2			ORa		81Ei03 Z
<sup>190</sup> Bi( $\alpha$ ) <sup>186</sup> Tl	6862.2	5.	6862	3	0.0	3			Lvn		91Va04 *
	6863.3	5.			-0.2	3			Anv		03An26 *
	6860.3	6.			0.3	3			Jya		13Ny01 *
<sup>190</sup> Bi <sup>m</sup> ( $\alpha$ ) <sup>186</sup> Tl <sup>m</sup>	6967.9	5.	6966.4	2.8	-0.3	4			Lvn		91Va04 *
	6969.1	5.			-0.5	4			Anv		03An26 *
	6963.1	5.			0.7	4			Jya		13Ny01 *
<sup>190</sup> Bi <sup>m</sup> ( $\alpha$ ) <sup>186</sup> Tl <sup>n</sup>	6589.0	10.	6592.4	2.8	0.3	R			Ora		74Le02
<sup>190</sup> Po( $\alpha$ ) <sup>186</sup> Pb	7643.2	20.	7693	7	2.5	C			GSa		88Qu.A
	7651.4	40.			1.0	U			ORa		96Ba35
	7691.2	10.			0.2	3			ORa		97Ba25
	7695.3	10.			-0.2	3			Anv		00An14 *
<sup>190</sup> Os(p,t) <sup>188</sup> Os	-5234	5	-5231.4	0.5	0.5	U			Min		73Oo01
	-5237	4			1.4	U			McM		75Th04
<sup>190</sup> Pt(p,t) <sup>188</sup> Pt	-7150	10	-7146	5	0.4	1	28	28 <sup>188</sup> Pt	Ors		78Ve10
<sup>190</sup> Os(t, $\alpha$ ) <sup>189</sup> Re	11796	10	11796	8	0.0	2			McM		76Hi08
<sup>189</sup> Os(n, $\gamma$ ) <sup>190</sup> Os	7791.8	1.0	7792.34	0.19	0.5	U			BNn		79Ca02 Z
	7792.31	0.19			0.2	1	97	79 <sup>189</sup> Os	Bdn		06Fi.A
<sup>190</sup> Os(d,t) <sup>189</sup> Os	-1541	10	-1535.11	0.19	0.6	U			Kop		75Mo29
	-1530	4			-1.3	U			Tal		76Be50

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{190}\text{Pt}(p,d)^{189}\text{Pt}$	-6693	11	-6684	10	0.8	1	84	84 $^{189}\text{Pt}$	Ors		80Ka19
$^{190}\text{W}(\beta^-)^{190}\text{Re}$	1270	70	1250	60	-0.2	1	82	76 $^{190}\text{Re}$			76Ha39 *
$^{190}\text{Re}(\beta^-)^{190}\text{Os}$	3090	300	3070	70	-0.1	-					55At21 *
	3190	300			-0.4	-					69Ha44 *
	3146	200			-0.4	-					64Fl02 *
ave.	3140	150			-0.5	1	24	24 $^{190}\text{Re}$			average
$^{190}\text{Ir}(\beta^+)^{190}\text{Os}$	2000	200	1954.2	1.2	-0.2	U					60Ka14 *
$^{190}\text{Au}(\beta^+)^{190}\text{Pt}$	4442	15	4473	4	2.1	U					73Jo11
	4380	55			1.7	U					74Di.A
	4380	200			0.5	U					75Un.A
$^{190}\text{Hg}(\beta^+)^{190}\text{Au}$	2105	80	1463	16	-8.0	C					74Di.A
$^{190}\text{Tl}(\beta^+)^{190}\text{Hg}$	7000	400	6999	18	0.0	U					75Un.A *
$^{190}\text{Tl}^m(\beta^+)^{190}\text{Hg}$	6975	300	7081	17	0.4	U					76Bi09 *
$^{190}\text{Bi}(\beta^+)^{190}\text{Pb}$	8700	500	9817	26	2.2	F					76Bi09 *
* $^{190}\text{Au}-u$	$M-A=-32701(28)$ keV for mixture gs+m at 200#150 keV										Nub16b **
* $^{190}\text{Tl}^m-u$	Assumed by evaluator to be the $7^+$ excited isomer										GAu **
* $^{190}\text{Bi}^m-^{133}\text{Cs}_{1.429}$	F: contamination due to ground state not resolved										08We02 **
* $^{190}\text{Bi}(\alpha)^{186}\text{Tl}$	$E_\alpha=6716(5), 6507(5), 6431(5)$ to ground state, 215.2, 293.7 levels										91Va04 **
* $^{190}\text{Bi}(\alpha)^{186}\text{Tl}$	$E_\alpha=6431(5)$ to 293.7 level										03An26 **
* $^{190}\text{Bi}(\alpha)^{186}\text{Tl}$	$E_\alpha=6428(6)$ to 293.7 level										13Ny01 **
* $^{190}\text{Bi}^m(\alpha)^{186}\text{Tl}^m$	$E_\alpha=6819(5), 6734(5), 6456(5)$ to levels 0, 89.5, 373.9 above $^{186}\text{Tl}^m$										91Va04 **
* $^{190}\text{Bi}^m(\alpha)^{186}\text{Tl}^m$	$E_\alpha=6456(5)$ to 374.0 level above $^{186}\text{Tl}^m$										03An26 **
* $^{190}\text{Bi}^m(\alpha)^{186}\text{Tl}^m$	$E_\alpha=6450(5)$ to 374.0 level above $^{186}\text{Tl}^m$										Nub16b **
* $^{190}\text{Po}(\alpha)^{186}\text{Pb}$	$E_\alpha=7545(15)$ same dataset as in reference 2000An14										97An09 **
* $^{190}\text{W}(\beta^-)^{190}\text{Re}$	$E_{\beta^-}=950(70)$ to $1^+$ level at 319.7 keV										Ens036 **
* $^{190}\text{Re}(\beta^-)^{190}\text{Os}$	$E_{\beta^-}=1700(300)$ 1800(300) respectively, to $3^-$ level at 1387.00 keV										Ens036 **
* $^{190}\text{Re}(\beta^-)^{190}\text{Os}$	$E_{\beta^-}=1600(200)$ from isomer at 204(10) to several levels around 1750 keV										Nub16b **
* $^{190}\text{Ir}(\beta^+)^{190}\text{Os}$	$p^+=6(1)\times 10^{-5}$ to $4^+$ levels at 1163.19 and 955.37 keV, level at 1872.15 keV fed										Ens036 **
* $^{190}\text{Tl}(\beta^+)^{190}\text{Hg}$	$E_{\beta^+}=5700(400)$ to ground state and $2^+$ level at 416.32 keV										Ens036 **
* $^{190}\text{Tl}^m(\beta^+)^{190}\text{Hg}$	$E_{\beta^+}=4180(300)$ to $6^+$ level at 1772.94 keV										Ens036 **
* $^{190}\text{Bi}(\beta^+)^{190}\text{Pb}$	F: $E_{\beta^+}=5700(300)$ to a level around 2000 at least										AHW **
$^{191}\text{W}-u$	-33469	45				2			GS3	1.0	13Sh30
$^{191}\text{Au}-u$	-36180	88	-36284	5	-1.2	U			GS2	1.0	05Li24 *
$^{191}\text{Au}-^{133}\text{Cs}_{1.436}$	99487.3	5.3	99487	5	0.0	1	100	100 $^{191}\text{Au}$	MA8	1.0	13Kr15
$^{191}\text{Hg}-u$	-32811	51	-32842	24	-0.6	1	22	22 $^{191}\text{Hg}$	GS2	1.0	05Li24 *
$^{191}\text{Hg}-^{208}\text{Pb}_{.918}$	-11414	29	-11408	24	0.2	1	68	68 $^{191}\text{Hg}$	MA6	1.0	01Sc41 *
$^{191}\text{Tl}-u$	-28340	130	-28216	8	1.0	U			GS1	1.0	00Ra23 *
	-28234	30			0.6	U			GS2	1.0	05Li24 *
	-28192	31			-0.8	U			GS2	1.0	05Li24 *
$^{191}\text{Pb}-u$	-21770	110	-21720	40	0.5	U			GS1	1.0	00Ra23 *
	-21719	40				2			GS2	1.0	05Li24 *
$^{191}\text{Bi}-^{133}\text{Cs}_{1.436}$	121552.1	8.6	121558	8	0.7	1	87	87 $^{191}\text{Bi}$	MA8	1.0	08We02
$^{191}\text{Pb}^m(\alpha)^{187}\text{Hg}^m$	5403.4	20.				2			Ora		74Le02
$^{191}\text{Bi}(\alpha)^{187}\text{Tl}$	6780.8	5.	6780	3	-0.1	-			Lvn		85Co06 Z
	6785.3	10.2			-0.5	-			ORa		98Bi.A
	6782.3	10.2			-0.2	-			Anv		99An36
	6783.3	7.2			-0.4	-			Jya		13Ny01
ave.	6782	4			-0.5	1	71	69 $^{187}\text{Tl}$			average
$^{191}\text{Bi}(\alpha)^{187}\text{Tl}^m$	6440.0	5.	6446.6	2.5	1.3	-					67Tr06 Z
	6455.0	10.			-0.8	U			Ora		74Le02 Z
	6445.9	5.			0.2	-			Lvn		85Co06 Z
	6447	10			0.0	U			ORa		98Bi.A
	6458.5	20.			-0.6	U			RIa		99Ta20
	6445	10			0.2	U			Anv		99An36
	6443.2	3.			1.1	o			Jya		03Ke04
	6445.2	10.			0.1	U			Jya		13Uu01
	6450.3	4.			-0.9	-			Jya		13Ny01
ave.	6446.2	2.7			0.2	1	83	72 $^{187}\text{Tl}^m$			average
$^{191}\text{Bi}^m(\alpha)^{187}\text{Tl}$	7022.8	5.	7023	3	0.0	2			Lvn		85Co06 Z
	7023.4	10.			-0.1	U			ORa		98Bi.A

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{191}\text{Bi}^m(\alpha)^{187}\text{Tl}$	7016.2	20.	7023	3	0.3	U			RIa		99Ta20
	7017.2	3.			1.7	o			Jya		03Ke04
	7031.0	15.			-0.6	U			Jya		13Uu01 *
	7022.4	4.			0.0	2			Jya		13Ny01
$^{191}\text{Po}(\alpha)^{187}\text{Pb}$	7470.8	20.	7493	5	1.1	F			GSa		93Qu03 *
	7487.1	15.			0.4	U			ORa		97Ba25
	7491.2	5.			0.4	1	94	94 $^{191}\text{Po}$	Anv		02An19 *
$^{191}\text{Po}(\alpha)^{187}\text{Pb}^m$	7493.2	15.	7474	10	-1.2	1	45	39 $^{187}\text{Pb}^m$	Anv		02An19 *
$^{191}\text{Po}^m(\alpha)^{187}\text{Pb}^m$	7535	5				2			Anv		02An19 *
$^{191}\text{At}(\alpha)^{187}\text{Bi}^m$	7713.9	11.				3			Jya		03Ke08
$^{191}\text{At}^m(\alpha)^{187}\text{Bi}$	7880.4	15.				3			Jya		03Ke08
$^{191}\text{Ir}(\text{p,t})^{189}\text{Ir}$	-5903	15	-5920	13	-1.1	1	70	70 $^{189}\text{Ir}$	McM		78Lo07
$^{190}\text{Os}(\text{n},\gamma)^{191}\text{Os}$	5758.2	2.	5758.73	0.11	0.3	U					77Be15
	5759.1	1.5			-0.2	U					77Ca19
	5758.67	0.16			0.4	-			ILn		91Bo35 Z
	5758.81	0.15			-0.5	-			Bdn		06Fi.A
	ave.	5758.74	0.11			-0.1	1	100	99 $^{191}\text{Os}$		average
$^{190}\text{Os}(\alpha,\text{t})^{191}\text{Ir}$	-14569	15	-14523.9	1.1	3.0	B			McM		71Pr13
$^{191}\text{Ir}(\text{d,t})^{190}\text{Ir}$	-1769.3	0.4				2					95Ga04 *
$^{191}\text{Os}(\beta^-)^{191}\text{Ir}$	313.3	3.	313.6	1.1	0.1	-					48Sa18 *
	314.3	2.			-0.4	-					51Ko17 *
	316.3	3.			-0.9	-					58Na15 *
	314.3	3.			-0.2	-					60Fe03 *
	318.3	3.			-1.6	-					63Pl01 *
ave.	315.1	1.2			-1.3	1	90	90 $^{191}\text{Ir}$		average	
$^{191}\text{Pt}(\epsilon)^{191}\text{Ir}$	1000	15	1011	4	0.7	U					70Sc20 *
$^{191}\text{Au}(\beta^+)^{191}\text{Pt}$	1830	50	1900	6	1.4	U					76Vi.A *
$^{191}\text{Hg}(\beta^+)^{191}\text{Au}$	3430	200	3206	23	-1.1	U					75Un.A *
	3180	70			0.4	1	11	10 $^{191}\text{Hg}$			76Vi.A *
$^{191}\text{Tl}^m(\beta^+)^{191}\text{Hg}$	5178	200	4606	23	-2.9	C					75Un.A *
* $^{191}\text{Au}-\text{u}$	$M-A=-33568(28)$ keV for mixture gs+m at 266.2 keV										Nub16b **
* $^{191}\text{Hg}-\text{u}$	$M-A=-30499(28)$ keV for mixture gs+m at 128(22) keV										Nub16b **
* $^{191}\text{Hg}-^{208}\text{Pb}_{918}$	Original error (19keV) increased by 20 due to isomer+ground state lines in trap										01Sc41 **
* $^{191}\text{Tl}-\text{u}$	$M-A=-26250(90)$ keV for mixture gs+m at 297(7) keV										Nub16b **
* $^{191}\text{Tl}-\text{u}$	$M-A=-25964(28)$ keV for $^{191}\text{Tl}^m$ at 297(7) keV										Nub16b **
* $^{191}\text{Pb}-\text{u}$	Possible isomeric contamination										00Ra23 **
* $^{191}\text{Pb}-\text{u}$	$M-A=-20226(28)$ keV for mixture gs+m at 10(50) keV										Nub16b **
* $^{191}\text{Bi}^m(\alpha)^{187}\text{Tl}$	average 6882(20) 6885(15)										13Uu01 **
* $^{191}\text{Po}(\alpha)^{187}\text{Pb}$	F : probably mainly $^{189}\text{Bi}^m$										97Ba25 **
* $^{191}\text{Po}(\alpha)^{187}\text{Pb}$	$E_\alpha=7334(10)$ , 6960(15) to ground state, 375(1) superseded by 02An19										99An10 **
* $^{191}\text{Po}^m(\alpha)^{187}\text{Pb}^m$	$E_\alpha=7376(5)$ , 6888(5) to $^{187}\text{Pb}^m$ and 494(1) above										02An19 **
* $^{191}\text{Po}^m(\alpha)^{187}\text{Pb}^m$	$E_\alpha=7378(10)$ , 6888(15) superseded by 02An19										99An10 **
* $^{191}\text{Ir}(\text{d,t})^{190}\text{Ir}$	Feeds ground state										96Ga30 **
* $^{191}\text{Os}(\beta^-)^{191}\text{Ir}$	$E_{\beta^-}=142(3)$ 143(2) 145(3) 143(3) 147(3) respectively, to $11/2^-$ level at 171.29 keV										Ens07a **
* $^{191}\text{Pt}(\epsilon)^{191}\text{Ir}$	$\text{pL}=0.73(0.12)$ to $(1/2^+, 3/2, 5/2^+)$ at 935.46 keV , no K capture										Ens07a **
* $^{191}\text{Au}(\beta^+)^{191}\text{Pt}$	$E_{\beta^+}=850(30)$ to ground state and $(5/2^-, 7/2^-)$ level at 9.547 keV; also $E_{\beta^+}=470(60)$ to $(3/2^-, 5/2^-)$ level at 277.88 and $5/2^-$ level at 293.458 keV										Ens07a **
* $^{191}\text{Hg}(\beta^+)^{191}\text{Au}$	Reassigned by evaluator to mainly ground state, partly $3/2^+$ 207.9 level										Ens07a **
* $^{191}\text{Tl}^m(\beta^+)^{191}\text{Hg}$	$E_{\beta^+}=3820(200)$ to level $(5/2^-)$ at 336.32 keV										Ens07a **
$^{192}\text{Re}-\text{u}$	-33912	76				2			GS3	1.0	13Sh30
$^{192}\text{Hg}-\text{u}$	-34440	104	-34366	17	0.7	U			GS1	1.0	00Ra23
	-34342	30			-0.8	R			GS2	1.0	05Li24
$^{192}\text{Hg}-^{208}\text{Pb}_{923}$	-12826	20	-12816	17	0.5	2			MA6	1.0	01Sc41
$^{192}\text{Tl}-\text{u}$	-27804	125	-27780	30	0.2	U			GS1	1.0	00Ra23 *
	-27775	34				2			GS2	1.0	05Li24

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{192}\text{Pb-u}$	-24280	104	-24215	14	0.6	U			GS1	1.0	00Ra23
	-24185	30			-1.0	R			GS2	1.0	05Li24
$^{192}\text{Bi-u}$	-14783	128	-14530	30	2.0	U			GS1	1.0	00Ra23 *
	-14489	60			-0.7	R			GS2	1.0	05Li24 *
$^{192}\text{Bi}^m-^{133}\text{Cs}_{1,444}$	122143.5	9.6				2			MA8	1.0	08We02
$^{192}\text{Os O}_2-^{189}\text{Os }^{35}\text{Cl}$	24301	6	24309.4	2.4	0.6	U			H22	2.5	70Mc03
$^{192}\text{Os }^{35}\text{Cl}-^{190}\text{Os }^{37}\text{Cl}$	5984	3	5983.5	2.4	-0.1	U			H22	2.5	70Mc03
$^{192}\text{Pb}(\alpha)^{188}\text{Hg}$	5221.0	5.				2			ORa		79To06 Z
$^{192}\text{Bi}(\alpha)^{188}\text{Tl}$	6376.0	5.	6377	4	0.2	3			Lvn		91Va04 *
	6377.9	5.			-0.2	3			Jya		13Ny01 *
$^{192}\text{Bi}^m(\alpha)^{188}\text{Tl}^m$	6481.4	5.	6485	3	0.7	3					67Tr06 *
	6491.6	10.			-0.7	3			Ora		74Le02 *
	6483.3	5.			0.3	3			Lvn		91Va04 *
	6494	8			-1.1	3			Jya		03Ke04 *
$^{192}\text{Po}(\alpha)^{188}\text{Pb}$	7322.8	20.	7320	3	-0.2	U					81Le23
	7319.8	7.			0.0	3			Lvn		93Wa04
	7364.6	35.			-1.3	U			RIa		95Mo14
	7349.4	30.			-1.0	U			RIa		97Pu01
	7319.8	11.			0.0	o			Jya		01Ke06
	7318.8	8.			0.1	3			Jya		03Ke04
	7319.8	4.			0.0	3			Anv		03Va16
$^{192}\text{At}(\alpha)^{188}\text{Bi}$	7695.6	25.				3			Anv		06An04
$^{192}\text{At}(\alpha)^{188}\text{Bi}^m$	7629.3	15.				4			Anv		06An04 *
$^{192}\text{At}^m(\alpha)^{188}\text{Bi}$	7695.6	25.				3			Anv		06An04
$^{192}\text{At}^m(\alpha)^{188}\text{Bi}^m$	7542.4	15.				4			Anv		06An04 *
$^{192}\text{Os}(p,t)^{190}\text{Os}$	-4835	5	-4835.3	2.2	-0.1	-			Min		73Oo01
	-4837	4			0.4	-			McM		75Th04
	ave.	-4836	3		0.3	1	51	51 $^{192}\text{Os}$			average
$^{192}\text{Pt}(p,t)^{190}\text{Pt}$	-6629	7	-6642.8	2.5	-2.0	1	13	13 $^{192}\text{Pt}$	Ors		80Ka19
$^{192}\text{Os}(t,\alpha)^{191}\text{Re}$	10993	10				2			McM		76Hi08
$^{192}\text{Os}(d,t)^{191}\text{Os}$	-1265	15	-1301.1	2.2	-2.4	U			Tal		77Be15
$^{191}\text{Ir}(n,\gamma)^{192}\text{Ir}$	6197.7	0.3	6198.12	0.11	1.4	o			ILn		87Ke.A
	6198.1	0.2			0.1	-			ILn		91Ke10
	6198.14	0.13			-0.1	-			Bdn		06Fi.A
	ave.	6198.13	0.11		-0.1	1	100	91 $^{192}\text{Ir}$			average
$^{192}\text{Pt}(p,d)^{191}\text{Pt}$	-6448	6	-6437.0	2.9	1.8	1	23	26 $^{191}\text{Pt}$	Ors		80Ka19
$^{192}\text{Pt}(p,d)^{191}\text{Pt}-^{194}\text{Pt}(t)^{193}\text{Pt}$	-307	3	-309.8	2.7	-0.9	1	81	74 $^{191}\text{Pt}$	Ors		78Be09
$^{192}\text{Ir}(\beta^+)^{192}\text{Os}$	1468	10	1046.6	2.4	-42.1	B					60An04 *
$^{192}\text{Ir}(\beta^-)^{192}\text{Pt}$	1456.7	4.	1452.9	2.3	-1.0	-					65Jo04 *
	1453.3	3.			-0.1	-					77Ra17 *
	ave.	1454.5	2.4		-0.7	1	90	87 $^{192}\text{Pt}$			average
$^{192}\text{Au}(\beta^+)^{192}\text{Pt}$	3514	20	3516	16	0.1	2					66Ny01
	3520	25			-0.1	2					74Di.A
$^{192}\text{Hg}(\beta^+)^{192}\text{Au}$	1745	30	761	22	-32.8	F					74Di.A *
$^{192}\text{Tl}(\beta^+)^{192}\text{Hg}$	6380	200	6140	40	-1.2	U					75Un.A *
* $^{192}\text{Tl-u}$	$M - A = -25830(100)$ keV for mixture gs+m at 138(45) keV										Nub16b **
* $^{192}\text{Bi-u}$	$M - A = -13700(110)$ keV for mixture gs+m at 140(30) keV										Nub16b **
* $^{192}\text{Bi-u}$	$M - A = -13426(31)$ keV for mixture gs+m at 140(30) keV										Nub16b **
* $^{192}\text{Bi}(\alpha)^{188}\text{Tl}$	$E_\alpha = 6245(5), 6060(5)$ to ground state, 184.6 level										91Va04 **
* $^{192}\text{Bi}(\alpha)^{188}\text{Tl}$	$E_\alpha = 6064(5)$ to 184.6 level										Ens024 **
* $^{192}\text{Bi}^m(\alpha)^{188}\text{Tl}^m$	$E_\alpha = 6050(5)$ to $(10^-)$ level, 302.4 above $^{188}\text{Tl}^m$										91Va04 **
* $^{192}\text{Bi}^m(\alpha)^{188}\text{Tl}^m$	$E_\alpha = 6060(10)$ to $(10^-)$ level, 302.4 above $^{188}\text{Tl}^m$										91Va04 **
* $^{192}\text{Bi}^m(\alpha)^{188}\text{Tl}^m$	$E_\alpha = 6348(5), 6253(5), 6081(10), 6052(5)$ to $^{188}\text{Tl}^m$ and to levels 103.2, 268.8, 302.4 above $^{188}\text{Tl}^m$										91Va04 **
* $^{192}\text{Bi}^m(\alpha)^{188}\text{Tl}^m$	$E_\alpha = 6062(5)$ to level 302.4 above $^{188}\text{Tl}^m$										03Ke04 **
* $^{192}\text{At}(\alpha)^{188}\text{Bi}^m$	Also $E_\alpha = 7435(15)$ keV followed by 36 keV $\gamma$										06An04 **
* $^{192}\text{At}^m(\alpha)^{188}\text{Bi}^m$	Also $E_\alpha = 7224(15), 7195(15)$ keV followed by 165 and 188 keV $\gamma$										06An04 **
* $^{192}\text{Ir}(\beta^+)^{192}\text{Os}$	$E_{\beta^+} = 240(10)$ to $2^+$ level at 205.7944 keV										Ens129 **
* $^{192}\text{Ir}(\beta^-)^{192}\text{Pt}$	$E_{\beta^-} = 672(4) 666(2)$ respectively, to $4^+$ level at 784.5759, and other $E_{\beta^-}$										Ens129 **
* $^{192}\text{Hg}(\beta^+)^{192}\text{Au}$	F: most probably due to backscattering of 2.5 MeV Au positons										AHW **
* $^{192}\text{Tl}(\beta^+)^{192}\text{Hg}$	$E_{\beta^+} = 4940(200)$ to $2^+$ level at 422.79 keV										Ens129 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{193}\text{Re}-u$	-32455	42					GS3	1.0	13Sh30
$^{193}\text{Au}-u$	-35736	96	-35862	9	-1.3	U	GS2	1.0	05Li24 *
$^{193}\text{Hg}-u$	-33288	53	-33347	17	-1.1	U	GS2	1.0	05Li24 *
$^{193}\text{Hg}-^{208}\text{Pb}_{.928}$	-11673	29	-11680	17	-0.2	1	33	33 $^{193}\text{Hg}$	01Sc41 *
$^{193}\text{Tl}-u$	-29690	158	-29498	7	1.2	U	GS1	1.0	00Ra23 *
	-29329	120			-1.4	U	GS2	1.0	05Li24 *
$^{193}\text{Tl}-^{133}\text{Cs}_{1.451}$	107691.2	7.2				2	MA8	1.0	14Bo26 *
$^{193}\text{Pb}-u$	-23865	125	-23830	50	0.3	o	GS1	1.0	00Ra23 *
	-23846	66			0.3	2	GS2	1.0	05Li24 *
$^{193}\text{Bi}-u$	-16980	110	-17053	8	-0.7	U	GS1	1.0	00Ra23
	-17025	30			-0.9	R	GS2	1.0	05Li24
$^{193}\text{Bi}-^{133}\text{Cs}_{1.451}$	120147	11	120136	8	-1.0	-	MA8	1.0	08We02
	ave. 120149	10			-1.2	1	62	62 $^{193}\text{Bi}$	average
$^{193}\text{Bi}(\alpha)^{189}\text{Tl}$	6304.5	5.	6307	5	0.4	1	92	70 $^{189}\text{Tl}$	85Co06 Z
$^{193}\text{Bi}(\alpha)^{189}\text{Tl}^m$	6017.8	5.	6021	3	0.7	2		Lvn	67Tr06 Z
	6024.6	10.			-0.3	2		Ora	74Le02 Z
	6023.7	5.			-0.5	2		Lvn	85Co06 Z
$^{193}\text{Bi}^m(\alpha)^{189}\text{Tl}$	6617.4	10.	6611	4	-0.6	-		Ora	74Le02
	6611.9	5.			-0.1	-		Lvn	85Co06 Z
	6618.4	14.			-0.5	U		Jya	05Uu02
	ave. 6613	5			-0.4	1	94	64 $^{193}\text{Bi}^m$	average
$^{193}\text{Po}(\alpha)^{189}\text{Pb}$	7128.1	20.	7094	4	-1.7	U			67Si09
	7087.1	20.			0.3	U		Ora	77De32
	7096.4	5.			-0.5	2		Lvn	93Wa04
	7093.3	30.			0.0	U		RIa	95Mo14
	7089.2	6.			0.8	2		Jya	96En02
	7096.4	10.			-0.3	2		Anv	02Va13
$^{193}\text{Po}^m(\alpha)^{189}\text{Pb}^m$	7143.3	10.	7154	3	1.0	2		Ora	77De32
	7148.4	20.			0.3	U			81Le23
	7152.5	5.			0.2	2		Lvn	93Wa04
	7139.2	30.			0.5	U		RIa	95Mo14
	7159.7	6.			-1.0	2		Jya	96En02
	7152.5	10.			0.1	2		Anv	02Va13
$^{193}\text{At}(\alpha)^{189}\text{Bi}^m$	7388.5	5.				7		Jya	03Ke08
$^{193}\text{At}^m(\alpha)^{189}\text{Bi}$	7556.9	20.	7580	5	1.1	o		Jya	95Le15
	7490	6			15.1	C		Jya	98En.A
	7580.4	5.				7		Jya	03Ke08 *
$^{193}\text{At}^m(\alpha)^{189}\text{Bi}$	7614.3	5.				7		Jya	03Ke08 *
$^{193}\text{Rn}(\alpha)^{189}\text{Po}$	8040.0	12.				4		Anv	06An36 *
$^{193}\text{Ir}(p,t)^{191}\text{Ir}$	-5490	15	-5488.32	0.23	0.1	U		McM	78Lo07
$^{192}\text{Os}(n,\gamma)^{193}\text{Os}$	5583.5	2.	5583.42	0.20	0.0	U			78Be22
	5583.40	0.20			0.1	1	100	81 $^{193}\text{Os}$	79Wa04
	5584.01	0.16			-3.7	C		Bdn	06Fi.A
$^{192}\text{Os}(\alpha,t)^{193}\text{Ir}$	-13923	15	-13870.8	2.4	3.5	B		McM	71Pr13
$^{193}\text{Ir}(t,\alpha)^{192}\text{Os}-^{191}\text{Ir}(\text{I})^{190}\text{Os}$	-661	4	-653.1	2.2	2.0	1	31	31 $^{192}\text{Os}$	LAl
$^{192}\text{Ir}(n,\gamma)^{193}\text{Ir}$	7772.0	0.2	7771.99	0.20	0.0	1	100	94 $^{193}\text{Ir}$	85Co.B Z
$^{193}\text{Ir}(\gamma,n)^{192}\text{Ir}$	-7790	50	-7771.99	0.20	0.4	U		Phi	60Ge01
$^{192}\text{Pt}(n,\gamma)^{193}\text{Pt}$	6247	3	6262.5	2.3	5.2	B			68Sa13
$^{193}\text{Os}(\beta^-)^{193}\text{Ir}$	1132	5	1141.9	2.4	2.0	1	23	19 $^{193}\text{Os}$	58Na15
$^{193}\text{Pt}(\epsilon)^{193}\text{Ir}$	56.6	0.3	56.63	0.30	0.1	1	100	96 $^{193}\text{Pt}$	83Jo04
$^{193}\text{Au}(\beta^+)^{193}\text{Pt}$	1355	20	1075	9	-14.0	B			76Di15 *
$^{193}\text{Hg}(\beta^+)^{193}\text{Au}$	2341	30	2343	14	0.1	-			58Br88 *
	2340	20			0.1	-			76Di15 *
	ave. 2340	17			0.1	1	75	67 $^{193}\text{Hg}$	average
* $^{193}\text{Au}-u$	$M - A = -33143(29)$ keV for mixture gs+m at 290.19 keV								
* $^{193}\text{Hg}-u$	$M - A = -30937(28)$ keV for mixture gs+m at 140.76 keV								
* $^{193}\text{Hg}-^{208}\text{Pb}_{.928}$	Original error (18keV) increased by 20 due to isomer+ground state lines in trap								
* $^{193}\text{Tl}-u$	$M - A = -27470(100)$ keV for mixture gs+m at 372(4) keV								
* $^{193}\text{Tl}-u$	$M - A = -27134(28)$ keV for mixture gs+m at 372(4) keV								
* $^{193}\text{Tl}-^{133}\text{Cs}_{1.451}$	$D_M = 108091.5(5.6)$ $\mu\text{u}$ for $^{193}\text{Tl}^m$ at 372(4) keV; $M - A = -27104.3(5.2)$ keV								
* $^{193}\text{Pb}-u$	$M - A = -22160(100)$ keV for mixture gs+m at 130#80 keV								

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>193</sup> Pb-u	$M - A = -22147(28)$ keV for mixture gs+m at 130#80 keV										
* <sup>193</sup> At <sup>m</sup> ( $\alpha$ ) <sup>189</sup> Bi	$E_{\alpha} = 7423(5), 7325(5)$ to ground state and 99.6(5) level										
* <sup>193</sup> At <sup>n</sup> ( $\alpha$ ) <sup>189</sup> Bi	$E_{\alpha} = 7106(5)$ to 357.6(0.5) <sup>189</sup> Bi <sup>n</sup> level										
* <sup>193</sup> Rn( $\alpha$ ) <sup>189</sup> Po	$E_{\alpha} = 7875(20), 7685(15)$ to ground state and 194 level										
* <sup>193</sup> Au( $\beta^+$ ) <sup>193</sup> Pt	$E_{\beta^+} = 153(15)$ to 3/2 <sup>-</sup> level at 187.81 keV, and other $E_{\beta^+}$										
* <sup>193</sup> Hg( $\beta^+$ ) <sup>193</sup> Au	$E_{\beta^-} = 1170(30)$ from <sup>193</sup> Hg <sup>m</sup> at 140.76 to 11/2 <sup>-</sup> level at 290.19 keV										
* <sup>193</sup> Hg( $\beta^+$ ) <sup>193</sup> Au	$E_{\beta^+} = 1287(15)$ reinterpreted by AHW as going to ground state and 1/2 <sup>+</sup> level at 38.22keV										
<sup>194</sup> Pt-u	-37315.72	0.67	-37316.5	0.5	-1.1	1	63	63 <sup>194</sup> Pt	MS1	1.0	16Ei01
<sup>194</sup> Au-u	-34768	114	-34580.9	2.3	1.6	U			GS2	1.0	05Li24 *
<sup>194</sup> Hg-u	-34527	30	-34551	3	-0.8	U			GS2	1.0	05Li24
<sup>194</sup> Hg- <sup>133</sup> Cs <sub>1,459</sub>	103394.7	3.1				2			MA8	1.0	10E111
<sup>194</sup> Hg- <sup>208</sup> Pb <sub>933</sub>	-12766	19	-12767	3	-0.1	U			MA6	1.0	01Sc41
<sup>194</sup> Tl-u	-28803	135	-28919	15	-0.9	o			GS1	1.0	00Ra23 *
	-28778	87			-1.6	U			GS2	1.0	05Li24 *
<sup>194</sup> Tl- <sup>133</sup> Cs <sub>1,459</sub>	109027	15				2			MA8	1.0	14Bo26
<sup>194</sup> Tl <sup>m</sup> - <sup>133</sup> Cs <sub>1,459</sub>	109306.4	4.1	109306	4	0.0	o			MA8	1.0	14Bo26
	109306.4	4.1				2			MA8	1.0	13St25
<sup>194</sup> Pb-u	-25980	104	-25988	19	-0.1	U			GS1	1.0	00Ra23
<sup>194</sup> Bi-u	-17162	128	-17208	7	-0.4	o			GS1	1.0	00Ra23 *
	-17178	76			-0.4	U			GS2	1.0	05Li24 *
<sup>194</sup> Bi <sup>m</sup> - <sup>133</sup> Cs <sub>1,459</sub>	120900	54				2			MA8	1.0	08We02 *
<sup>190</sup> Os- <sup>194</sup> Pt <sub>979</sub>	-5022.45	0.68	-5021.7	0.5	1.1	1	57	52 <sup>190</sup> Os	MS1	1.0	16Ei01
<sup>190</sup> Pt- <sup>194</sup> Pt <sub>979</sub>	-3516.95	0.68	-3517.3	0.5	-0.5	1	58	53 <sup>190</sup> Pt	MS1	1.0	16Ei01
<sup>194</sup> Pt- <sup>197</sup> Au <sub>985</sub>	-4396.4	3.2	-4388.0	0.6	2.6	U			CP1	1.0	05Sh52
<sup>194</sup> Au- <sup>197</sup> Au <sub>985</sub>	-1652.5	2.2				2			MA8	1.0	10E111
<sup>194</sup> Pb( $\alpha$ ) <sup>190</sup> Hg	4737.9	20.	4738	17	0.0	1	67	40 <sup>194</sup> Pb	ORa		87E109
<sup>194</sup> Bi( $\alpha$ ) <sup>190</sup> Tl	5918.3	5.				4			Lvn		91Va04 *
<sup>194</sup> Bi <sup>m</sup> ( $\alpha$ ) <sup>190</sup> Tl <sup>m</sup>	6015.7	5.				3			Lvn		91Va04 *
<sup>194</sup> Po( $\alpha$ ) <sup>190</sup> Pb	6991.5	10.	6987	3	-0.4	3					67Si09 Z
	6990.9	7.			-0.5	3					67Tr06 Z
	6984.4	5.			0.5	3			Ora		77De32 Z
	6990.0	5.			-0.6	o			Lvn		85Va03 Z
	6986.3	6.			0.1	3			Lvn		93Wa04
	6993.4	4.			-1.6	o			Jya		96En02
	6987.3	14.			0.0	3			Jya		05Uu02
<sup>194</sup> At( $\alpha$ ) <sup>190</sup> Bi	7412.5	20.	7454	11	2.1	o			Jya		95Le15 *
	7462.5	15.			-0.5	4			Anv		09An11 *
	7446.5	15.			0.5	4			Jya		13Ny01 *
<sup>194</sup> At <sup>m</sup> ( $\alpha$ ) <sup>190</sup> Bi <sup>m</sup>	7362.1	20.	7309	5	-2.6	o					80Ya.A
	7351.9	20.			-2.1	U					84Ya.A
	7341.7	20.			-1.6	o			Jya		95Le15
	7329.4	15.3			-1.3	5			Anv		09An11
	7306.9	5.1			0.4	5			Jya		13Ny01
<sup>194</sup> Rn( $\alpha$ ) <sup>190</sup> Po	7862.5	10.				4			Anv		06An36
<sup>193</sup> Ir(n, $\gamma$ ) <sup>194</sup> Ir	6067.0	0.4	6066.79	0.11	-0.5	2					82Ra.A
	6066.9	0.2			-0.6	2					98Ba85
	6066.71	0.14			0.6	2			Bdn		06Fi.A
<sup>194</sup> Pt(t, $\alpha$ ) <sup>193</sup> Ir	12286	20	12301.1	1.3	0.8	U			Tal		78Ya07
<sup>194</sup> Pt(d,t) <sup>193</sup> Pt	-2126	20	-2094.6	1.3	1.6	U			Pit		64Co11
<sup>194</sup> Pt(p,d) <sup>193</sup> Pt- <sup>196</sup> Pt( $\gamma$ ) <sup>195</sup> Pt	-445	3	-429.8	1.3	5.1	B			Ors		78Be09
<sup>194</sup> Os( $\beta^-$ ) <sup>194</sup> Ir	96.6	2.				3					64Wi07 *
<sup>194</sup> Ir( $\beta^-$ ) <sup>194</sup> Pt	2254	4	2228.4	1.3	-6.4	B					76Ra33
<sup>194</sup> Ir <sup>m</sup> ( $\beta^-$ ) <sup>194</sup> Pt	2600	70				2					68Su02 *
<sup>194</sup> Au( $\beta^+$ ) <sup>194</sup> Pt	2465	20	2548.1	2.1	4.2	B					56Th11
	2509	15			2.6	U					60Ba17
	2485	30			2.1	U					70Ag03 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{194}\text{Hg}(\epsilon)^{194}\text{Au}$	40	20	28	4	-0.6	U					81Ho18
* $^{194}\text{Au}-u$	$M-A=-32192(29)$ keV for mixture gs+m+n at 107.4 and 475.8 keV										Nub16b **
* $^{194}\text{Tl}-u$	$M-A=-26700(100)$ keV for mixture gs+m at 260(14) keV										Nub16b **
* $^{194}\text{Tl}-u$	$M-A=-26677(28)$ keV for mixture gs+m at 260(14) keV										Nub16b **
* $^{194}\text{Bi}-u$	$M-A=-15870(100)$ keV for mixture gs+m+n at 150(50) and 180(10) keV										Nub16b **
* $^{194}\text{Bi}-u$	$M-A=-15885(28)$ keV for mixture gs+m+n at 150(50) and 180(10) keV										Nub16b **
* $^{194}\text{Bi}^m-^{133}\text{Cs}_{1.459}$	Original error $16 \mu u$ increased to include possible $3^+$ and $10^-$ contam.										08We02 **
* $^{194}\text{Bi}(\alpha)^{190}\text{Tl}$	$E_\alpha=5799(5), 5645(5)$ to ground state, 151.3 level										91Va04 **
* $^{194}\text{Bi}^m(\alpha)^{190}\text{Tl}^m$	$E_\alpha=5892(5), 5781(5)$ to levels 0, 112.2 above $^{190}\text{Tl}^m$										91Va04 **
* $^{194}\text{At}(\alpha)^{190}\text{Bi}$	$E_\alpha=7140(20)$ to 121(15)										09An11 **
* $^{194}\text{At}(\alpha)^{190}\text{Bi}$	$E_\alpha=7190(15)$ to 121(15); further $E_\alpha: 7310(15), 7266(15), 7145(15)$ keV										09An11 **
* $^{194}\text{At}(\alpha)^{190}\text{Bi}$	$E_\alpha=7174(8)$ to 121(15)										13Ny01 **
* $^{194}\text{Os}(\beta^-)^{194}\text{Ir}$	$E_{\beta^-}=54.5(2.0)$ to $0^-$ level at 43.119 keV, and other $E_{\beta^-}$										Ens066 **
* $^{194}\text{Ir}^n(\beta^-)^{194}\text{Pt}$	$E_{\beta^-}<250$ to $10^+$ level at 2438.41 keV										Ens066 **
* $^{194}\text{Au}(\beta^+)^{194}\text{Pt}$	$E_{\beta^+}=1230(30)$ to $2^+$ level at 328.464 keV, and other $E^+$										Ens066 **
$^{195}\text{Os}-u$	-31682	60				2			GS3	1.0	13Sh30
$^{195}\text{Hg}-u$	-33283	62	-33294	25	-0.2	U			GS2	1.0	05Li24 *
$^{195}\text{Hg}-^{208}\text{Pb}_{.938}$	-11381	28	-11394	25	-0.5	1	79	79 $^{195}\text{Hg}$	MA6	1.0	01Sc41 *
$^{195}\text{Tl}-u$	-30320	200	-30226	12	0.5	U			GS1	1.0	00Ra23 *
	-30209	40			-0.4	-			GS2	1.0	05Li24 *
	-30264	33			1.2	-			GS2	1.0	05Li24 *
	ave.	-30242	25		0.6	1	22	22 $^{195}\text{Tl}$			average
$^{195}\text{Tl}-^{133}\text{Cs}_{1.466}$	108375	27	108382	12	0.2	-			MA8	1.0	14Bo26
	108472	79			-1.1	-			MA8	1.0	14Bo26 *
	ave.	108385	26		-0.1	1	22	22 $^{195}\text{Tl}$			average
$^{195}\text{Pb}-u$	-25423	150	-25451	19	-0.2	o			GS1	1.0	00Ra23 *
	-25461	70			0.1	-			GS2	1.0	05Li24 *
	ave.	-25457	25		0.2	1	59	59 $^{195}\text{Pb}$			average
$^{195}\text{Bi}-u$	-19320	100	-19351	6	-0.3	U			GS1	1.0	00Ra23
	-19537	128			1.5	U			GS2	1.0	05Li24 *
$^{195}\text{Bi}-^{133}\text{Cs}_{1.466}$	119258.2	6.0	119256	6	-0.3	1	89	89 $^{195}\text{Bi}$	MA8	1.0	08We02
$^{195}\text{Pt}-^{197}\text{Au}_{.990}$	-2119.9	3.2	-2110.1	0.6	3.1	B			CP1	1.0	05Sh52
$^{195}\text{Bi}(\alpha)^{191}\text{Tl}$	5832.5	5.				2			Lvn		85Co06 Z
$^{195}\text{Bi}(\alpha)^{191}\text{Tl}^m$	5542.9	10.	5535	5	-0.8	2			Ora		74Le02 Z
	5533.3	5.			0.4	2			Lvn		85Co06 Z
$^{195}\text{Bi}^m(\alpha)^{191}\text{Tl}$	6228.1	5.	6232	3	0.7	3					67Tr06 Z
	6238.4	10.			-0.6	3			Ora		74Le02 Z
	6233.7	5.			-0.4	3			Lvn		85Co06 Z
$^{195}\text{Po}(\alpha)^{191}\text{Pb}$	6763.1	8.	6749.9	2.8	-1.6	3					67Si09 Z
	6747.4	5.			0.5	3					67Tr06 Z
	6744.6	5.			1.0	3			Lvn		93Wa04
	6752.8	14.			-0.2	o			Jya		96Le09
	6744.6	10.			0.5	3			Anv		02Va13
	6755.9	6.			-1.0	3			Jya		05Uu02
$^{195}\text{Po}^m(\alpha)^{191}\text{Pb}^m$	6850.8	10.	6840.6	2.9	-1.0	3					67Si09
	6839.4	5.			0.2	3					67Tr06 Z
	6839.6	5.			0.2	3			Lvn		93Wa04
	6852.8	10.			-1.2	o			Jya		96Le09
	6839.6	10.			0.1	3			Anv		02Va13
	6840.6	6.			0.0	3			Jya		05Uu02
$^{195}\text{At}(\alpha)^{191}\text{Bi}^m$	7095.8	20.	7102	4	0.3	U			Jya		95Le15
	7105	20			-0.2	U			RIa		99Ta20
	7098.9	3.			1.0	o			Jya		03Ke04 *
	7113.2	10.			-1.1	o			Jya		13Uu01
	7101.9	4.				3			Jya		13Ny01



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{195}\text{At}^m(\alpha)^{191}\text{Bi}$	7340.9	30.	7373	4	1.1	o			Jya		83Le.A *
	7371.5	30.			0.1	o			Jya		95Le.A *
	7403	30			-1.0	U			RIa		99Ta20 *
	7372.5	4.0			0.2	o			Jya		03Ke04 *
	7373.5	4.0				2			Jya		13Ny01 *
$^{195}\text{Rn}(\alpha)^{191}\text{Po}$	7694.1	11.				2			Jya		01Ke06
$^{195}\text{Rn}^m(\alpha)^{191}\text{Po}^m$	7713.5	11.				3			Jya		01Ke06
$^{194}\text{Ir}(n,\gamma)^{195}\text{Ir}$	7231.92	0.11	7231.86	0.06	-0.5	o			ILn		87Ci.A
	7231.86	0.06				3			ILn		87Co08 Z
$^{194}\text{Pt}(n,\gamma)^{195}\text{Pt}$	6105.06	0.12	6105.10	0.12	0.3	1	99	72 $^{195}\text{Pt}$	ILn		81Ho.B Z
	6109.17	0.13			-31.3	F			Bdn		06Fi.A
$^{195}\text{Pt}(\gamma,n)^{194}\text{Pt}$	-6205	44	-6105.10	0.12	2.3	U			Phi		60Ge01
$^{194}\text{Pt}(d,p)^{195}\text{Pt}$	3908	20	3880.53	0.12	-1.4	U			Pit		64Co11
$^{195}\text{Pt}(d,t)^{194}\text{Pt}$	140	20	152.13	0.12	0.6	U			Pit		64Co11
$^{195}\text{Os}(\beta^-)^{195}\text{Ir}$	2000	500	2180	60	0.4	U					57Ba08
$^{195}\text{Ir}(\beta^-)^{195}\text{Pt}$	1116	20	1101.6	1.3	-0.7	U					73Ja10 *
$^{195}\text{Au}(\epsilon)^{195}\text{Pt}$	226.8	1.0	226.8	1.0	0.0	1	100	100 $^{195}\text{Au}$			Averag *
$^{195}\text{Hg}(\beta^+)^{195}\text{Au}$	1510	50	1554	23	0.9	1	21	21 $^{195}\text{Hg}$			71Fr03 *
$^{195}\text{Tl}(\beta^+)^{195}\text{Hg}$	3000	300	2858	26	-0.5	U					78Go15 *
$^{195}\text{Pb}(IT)^{195}\text{Pb}$	202.9	0.7	202.9	0.7	0.0	1	100	59 $^{195}\text{Pb}^m$	Oak		91Gr12
$^{195}\text{Bi}(\beta^+)^{195}\text{Pb}$	4850	550	5682	19	1.5	U			Oak		91Gr12
* $^{195}\text{Hg}-u$	$M-A=-30914(28)$ keV for mixture gs+m at 176.07 keV										Nub16b **
* $^{195}\text{Hg}-^{208}\text{Pb}_{.938}$	Corrected 40(20) keV for isomeric mixture $R=0.3(0.2)$ $E=176.07$ keV										Nub16b **
* $^{195}\text{Tl}-u$	$M-A=-28000(100)$ keV for mixture gs+m at 482.63 keV										Nub16b **
* $^{195}\text{Tl}-u$	$M-A=-27708(31)$ keV for $^{195}\text{Tl}^m$ at 482.63 keV										Nub16b **
* $^{195}\text{Tl}-^{133}\text{Cs}_{1.466}$	$D_M=108990(79)$ $\mu\text{u}$ for $^{195}\text{Tl}^m$ at 482.63 keV; $M-A=-27589(73)$ keV										Nub16b **
* $^{195}\text{Pb}-u$	$M-A=-23580(100)$ keV for mixture gs+m at 202.9 keV										Nub16b **
* $^{195}\text{Pb}-u$	$M-A=-23615(28)$ keV for mixture gs+m at 202.9 keV										Nub16b **
* $^{195}\text{Bi}-u$	$M-A=-17999(28)$ keV for mixture gs+m at 399(6) keV										Nub16b **
* $^{195}\text{At}(\alpha)^{191}\text{Bi}^m$	Correlated with $E_\alpha=6313$ of $^{191}\text{Bi}^m$										03Ke04 **
* $^{195}\text{At}^m(\alpha)^{191}\text{Bi}$	$E_\alpha=7190(30)$ to 148.7(0.5) level										03Ke04 **
*	correlated with $\alpha$ of 12 s $^{191}\text{Bi}$ ground state										95Le15 **
* $^{195}\text{At}^m(\alpha)^{191}\text{Bi}$	$E_\alpha=7105(30)$ to 148.7(0.5) level										03Ke04 **
* $^{195}\text{At}^m(\alpha)^{191}\text{Bi}$	$E_\alpha=7221(4)$ and 7075(4) to 148.7(0.5) level										03Ke04 **
* $^{195}\text{At}^m(\alpha)^{191}\text{Bi}$	$E_\alpha=7222(4)$ and 7076(5) to 148.7(0.5) level										13Ny01 **
* $^{195}\text{Ir}(\beta^-)^{195}\text{Pt}$	$E_{\beta^-}=980(30)$ to $3/2^-$ level at 98.880 keV and $5/2^-$ level at 129.772 keV, and $E_{\beta^-}=410(20)$ from $^{195}\text{Ir}^m$ at 100(5) to $9/2^-$ at 814.50, and other $E_{\beta^-}$										Ens148 **
*											Ens148 **
* $^{195}\text{Au}(\epsilon)^{195}\text{Pt}$	Average $pK=0.179(0.006)$ to $5/2^-$ level at 129.772 from the following references:										Ens148 **
*	$pK=0.195(0.015)$ to 129.78 level										65De20 **
*	$pK=0.166(0.020)$ to 129.78 level										68Ja11 **
*	$pK=0.160(0.017)$ to 129.78 level										73Go05 **
*	$pK=0.183(0.009)$ to 129.78 level										80Sa11 **
*	$pK=0.176(0.012)$ to 129.78 level										82Be.A **
* $^{195}\text{Hg}(\beta^+)^{195}\text{Au}$	Assuming 511 $\gamma$ is annihilation of $\beta^+$ to ground state and $1/2^+$ level at 61.434										Ens148 **
* $^{195}\text{Tl}(\beta^+)^{195}\text{Hg}$	$K/\beta^+=6(1)$ to ground state and $3/2^-$ level at 37.083 keV										Ens148 **
$^{196}\text{Hg}-^{208}\text{Pb}_{.942}$	-12178	20	-12173	3	0.3	U			MA6	1.0	01Sc41
$^{196}\text{Tl}-u$	-29188	126	-29519	13	-2.6	U			GS2	1.0	05Li24 *
$^{196}\text{Tl}-^{133}\text{Cs}_{1.474}$	109845	13				2			MA8	1.0	08We02 *
$^{196}\text{Pb}-^{208}\text{Pb}_{.942}$	-5228	22	-5219	8	0.4	-			MA6	1.0	01Sc41
	ave.	-5231	18			0.7	1	21	21 $^{196}\text{Pb}$		average
$^{196}\text{Pb}-u$	-27200	104	-27213	8	-0.1	U			GS1	1.0	00Ra23
	-27232	30			0.6	R			GS2	1.0	05Li24
$^{196}\text{Bi}-u$	-19309	137	-19333	26	-0.2	o			GS1	1.0	00Ra23 *
	-19325	30			-0.3	2			GS2	1.0	05Li24
	-19361	54			0.5	2			MA8	1.0	08We02 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{196}\text{Pt}-^{197}\text{Au}_{.995}$	-1781.1	3.0	-1782.6	0.6	-0.5	U			CP1	1.0	05Sh52
$^{196}\text{Bi}(\alpha)^{192}\text{Tl}^p$	5260.6	5.				3			Lvn		91Va04
$^{196}\text{Po}(\alpha)^{192}\text{Pb}$	6662.3	8.2	6658.1	2.4	-0.5	3					67Si09 Z
	6653.7	5.			0.8	3					67Tr06 Z
	6658.4	8.			0.0	3					71Ho01 Z
	6656.7	5.			0.3	o			Lvn		85Va03 Z
	6656.7	5.			0.3	3			Lvn		93Wa04
	6653.1	18.			0.3	o			Ara		95Le04
	6657.1	10.			0.1	o			Jya		96Le09
	6654.0	5.0			0.8	3			Ara		96Ta18 *
	6669.4	6.			-1.8	3			Jya		05Uu02
	6658.2	25.			0.0	U			Anv		10He25
$^{196}\text{Au}(\alpha)^{192}\text{Bi}$	7202.3	7.	7195	3	-1.0	4					67Tr06
	7187.0	25.			0.3	U			Jya		95Le15
	7200.2	30.			-0.2	U			RIa		95Mo14
	7191.0	7.			0.5	o			Jya		96En01
	7195.1	5.			0.0	o			Jya		00Sm06
	7202.3	12.			-0.6	o			Anv		05De01
	7195.1	12.			0.0	o			Jya		13Uu01 *
	7194.1	5.			0.2	4			Jya		13Ny01
	7192.1	5.			0.6	4			Anv		14Ka23
$^{196}\text{At}^m(\alpha)^{192}\text{Bi}^m$	7023.6	15.				3			Jya		96En01 *
$^{196}\text{Rn}(\alpha)^{192}\text{Po}$	7583.1	35.	7617	9	0.9	o			RIa		95Mo14
	7648.4	30.			-1.1	U			RIa		97Pu01
	7616.7	9.				4			Jya		01Ke06
$^{196}\text{Pt}(t,\alpha)^{195}\text{Ir}$	11565	20	11572.6	1.3	0.4	U			Tal		78Ya07
	11545	20			1.4	U			LAL		81Fl.A
$^{195}\text{Pt}(n,\gamma)^{196}\text{Pt}$	7921.96	0.20	7921.98	0.13	0.1	-			ILn		81Ho.B Z
	7921.92	0.17			0.3	-			Bdn		06Fi.A
$^{196}\text{Pt}(\gamma,n)^{195}\text{Pt}$	-8290	140	-7921.98	0.13	2.6	U			Phi		60Ge01
$^{195}\text{Pt}(d,p)^{196}\text{Pt}$	5712	25	5697.41	0.13	-0.6	U			Pit		64Co11
$^{196}\text{Pt}(d,t)^{195}\text{Pt}$	-1686	20	-1664.75	0.13	1.1	U			Pit		64Co11
$^{195}\text{Pt}(n,\gamma)^{196}\text{Pt}$	ave. 7921.94	0.13	7921.98	0.13	0.3	1	99	71 $^{196}\text{Pt}$			average
$^{196}\text{Os}(\beta^-)^{196}\text{Ir}$	900	40	1160	60	6.5	B					77Ha32 *
$^{196}\text{Ir}(\beta^-)^{196}\text{Pt}$	3150	60	3210	40	1.0	2					66Vo05 *
	3250	50			-0.8	2					67Mo10
$^{196}\text{Ir}^m(\beta^-)^{196}\text{Pt}$	3418	20				2					65Bi04 *
	3630	100	3418	20	-2.1	U					68Ja06 *
$^{196}\text{Au}(\beta^+)^{196}\text{Pt}$	1498	7	1505.8	3.0	1.1	1	18	18 $^{196}\text{Au}$			63Ik01 *
$^{196}\text{Au}(\epsilon)^{196}\text{Pt}$	1490	10			1.6	U					62Wa16 *
$^{196}\text{Au}(\beta^-)^{196}\text{Hg}$	685	4	687	3	0.6	1	61	31 $^{196}\text{Au}$			62Li03 *
* $^{196}\text{Tl}-u$	$M-A=-26991(28)$ keV for mixture gs+m at 394.2 keV										Nub16b **
* $^{196}\text{Tl}-^{133}\text{Cs}_{1.474}$	$Q=110268(13)$ $\mu\text{u}$ $M-A=-27103(12)$ keV for $^{196}\text{Tl}^m$ at 394.2 keV										Nub16b **
* $^{196}\text{Bi}-u$	$M-A=-17850(100)$ keV for mixture gs+n at 272(3) keV										Nub16b **
* $^{196}\text{Bi}-u$	$Q=120182(15)$ $\mu\text{u}$ for $^{196}\text{Bi}^m-^{133}\text{Cs}_{1.474}$ , $M-A(^{196}\text{Bi}^m)=-17868(14)$ keV at 167(3) keV; error increased to include possible $3^+$ and $10^-$ contam.										08We02 **
* $^{196}\text{Po}(\alpha)^{192}\text{Pb}$	Including systematic uncertainty 5 keV										96Ta18 **
* $^{196}\text{At}(\alpha)^{192}\text{Bi}$	Same group, but much less events than in 00Sm06										WgM151**
* $^{196}\text{At}^m(\alpha)^{192}\text{Bi}^m$	Correlated with $E_\alpha=7550$ of $^{200}\text{Fr}(\alpha)$										96En01 **
* $^{196}\text{Os}(\beta^-)^{196}\text{Ir}$	$E_{\beta^-}=435(20)$ to $(0,1)^+$ levels at 407.88, 522.37 keV										Ens076 **
* $^{196}\text{Ir}(\beta^-)^{196}\text{Pt}$	Original value 3170(60) recalibrated using $^{62}\text{Cu}$										AHW **
* $^{196}\text{Ir}^m(\beta^-)^{196}\text{Pt}$	$E_{\beta^-}=950(20)$ to $(10^-,11^-)$ level at 2468.0 keV										Ens076 **
* $^{196}\text{Ir}^m(\beta^-)^{196}\text{Pt}$	$E_{\beta^-}=1160(100)$ to $(10^-,11^-)$ level at 2468.0 keV										Ens076 **
* $^{196}\text{Au}(\beta^+)^{196}\text{Pt}$	$\text{KL}/\beta^+=2.0(0.4)\times 10^{+6}$ to $2^+$ level at 355.68 keV, recalculated										Ens076 **
* $^{196}\text{Au}(\epsilon)^{196}\text{Pt}$	$\text{pL}=0.64(0.06)$ to $3^-$ level at 1447.043 keV										Ens076 **
* $^{196}\text{Au}(\beta^-)^{196}\text{Hg}$	$E_{\beta^-}=259(4)$ to $2^+$ level at 425.98 keV										Ens076 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{197}\text{Hg}-u$	-32766	30	-32786	3	-0.7	U			GS2	1.0	05Li24
	-32765	30			-0.7	U			GS2	1.0	05Li24 *
$^{197}\text{Hg}-^{208}\text{Pb}_{.947}$	-10664	30	-10676	4	-0.4	U			MA6	1.0	01Sc41
$^{197}\text{Tl}-u$	-30450	30	-30426	18	0.8	R			GS2	1.0	05Li24
$^{197}\text{Pb}-u$	-26520	110	-26565	5	-0.4	U			GS1	1.0	00Ra23
	-26609	30			1.5	U			GS2	1.0	05Li24
	-26543	30			-0.7	U			GS2	1.0	05Li24 *
$^{197}\text{Pb}^m-^{133}\text{Cs}_{1.481}$	113799.6	6.0	113803	5	0.6	1	74	74 $^{197}\text{Pb}^m$	MA8	1.0	08We02
$^{197}\text{Bi}-^{208}\text{Pb}_{.947}$	982	22	976	9	-0.3	R			MA6	1.0	01Sc41
$^{197}\text{Bi}-u$	-21381	192	-21135	9	1.3	U			GS1	1.0	00Ra23 *
	-21187	31			1.7	U			GS2	1.0	05Li24
$^{197}\text{Bi}-^{133}\text{Cs}_{1.481}$	118870	26	118891	9	0.8	R			MA8	1.0	08We02 *
$^{197}\text{Po}-u$	-14434	145	-14340	50	0.6	o			GS1	1.0	00Ra23 *
	-14305	90			-0.4	R			GS2	1.0	05Li24 *
$^{197}\text{At}-^{133}\text{Cs}_{1.481}$	133186	20	133203	9	0.8	1	18	18 $^{197}\text{At}$	MA8	1.0	16Ma.1
$^{197}\text{At}^m-^{133}\text{Cs}_{1.481}$	133234	15	133251	10	1.1	1	42	42 $^{197}\text{At}^m$	MA8	1.0	16Ma.1
$^{197}\text{Au}-\text{C}_{16}$	-33432.5	7.3	-33429.9	0.6	0.2	o			TG1	1.5	09Ke.A
	-33432.9	5.4			0.4	U			TG1	1.5	10Ke09
$^{197}\text{Au}(\alpha, ^8\text{He})^{193}\text{Au}$	-26919	9	-26920	9	-0.1	1	93	93 $^{193}\text{Au}$			89Ka04
$^{197}\text{Bi}^m(\alpha)^{193}\text{Tl}$	5890.8	10.	5898	5	0.7	o			Ora		72Ga27
	5889.7	10.			0.8	3			Ora		74Le02 Z
	5899.6	5.			-0.4	3			Lvn		85Co06 Z
$^{197}\text{Po}(\alpha)^{193}\text{Pb}$	6420.7	10.	6412	3	-0.9	3					67Si09 Z
	6410.1	5.			0.3	3					67Tr06 Z
	6409.4	9.			0.2	3					71Ho01 Z
	6411.4	5.0			0.0	3			Ara		96Ta18 *
	6510.1	5.	6514.7	2.1	0.9	4					67Tr06 Z
$^{197}\text{Po}^m(\alpha)^{193}\text{Pb}^m$	6511.4	9.			0.4	U					71Ho01 Z
	6518.0	3.			-1.1	4			Bka		82Bo04 Z
	6512.4	5.0			0.4	4			Ara		96Ta18 *
	6517.6	10.			-0.3	o			Anv		02Va13
	6516.6	30.			-0.1	U			Anv		10He25
	6513.0	4.6			0.4	4			Tex		12Fo09
	7103.0	5.	7104	3	0.3	-					67Tr06 Z
$^{197}\text{At}(\alpha)^{193}\text{Bi}$	7100.5	5.			0.8	o			Jya		96En01
	7104.5	5.			0.0	o			Jya		99Sm07
	7103.5	6.			0.1	-			Jya		05Uu02
	7107.5	5.			-0.6	-			Anv		14Ka23
	ave.	7105	3			-0.1	1	98	82 $^{197}\text{At}$		average
$^{197}\text{At}^m(\alpha)^{193}\text{Bi}^m$	6846.2	10.	6844	4	-0.2	o			Lvn		86Co12
	6846.2	5.			-0.4	-			Jya		99Sm07
	6845.2	9.			-0.1	-			Jya		05Uu02
	6837.0	16.			0.5	U			Anv		14Ka23
$^{197}\text{Rn}(\alpha)^{193}\text{Po}$	ave.	6846	4		-0.4	1	94	58 $^{197}\text{At}^m$			average
	7410.8	20.	7411	7	0.0	o			RIa		95No.A
	7411.8	30.			0.0	U			RIa		95Mo14
$^{197}\text{Rn}^m(\alpha)^{193}\text{Po}^m$	7410.8	7.				3			Jya		96En02
	7523.1	30.	7509	6	-0.5	U			RIa		95Mo14
	7508.7	7.			0.1	3			Jya		96En02
	7510.7	14.			-0.1	3			Jya		05Uu02
$^{197}\text{Fr}(\alpha)^{193}\text{At}^m$	7888.4	15.				8		Anv		13Ka16	
$^{196}\text{Pt}(\text{n},\gamma)^{197}\text{Pt}$	5846.4	0.4	5846.56	0.26	0.4	-					78Ya07 Z
	5846.0	0.9			0.6	-			ILn		81Ho.B Z
	5846.6	0.5			-0.1	-			BNn		83Ca04 Z
	5846.0	0.7			0.8	-			Bdn		06Fi.A
$^{196}\text{Pt}(\text{d},\text{p})^{197}\text{Pt}$	3627	20	3621.99	0.26	-0.3	U			Pit		64Co11
	3606	20			0.8	U			Tal		78Ya07
$^{196}\text{Pt}(\text{n},\gamma)^{197}\text{Pt}$	ave.	5846.36	0.27	5846.56	0.26	0.7	1	94	65 $^{197}\text{Pt}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{197}\text{Au}(\gamma,n)^{196}\text{Au}$	-8057	22	-8072.3	2.9	-0.7	U			Phi		60Ge01	
	-8080	5			1.5	-			McM		79Ba06	
	-8072	7			0.0	-					79Be.A	
$^{197}\text{Au}(d,t)^{196}\text{Au}$	-1820	30	-1815.1	2.9	0.2	U			Pit		64Co11	
$^{197}\text{Au}(\gamma,n)^{196}\text{Au}$	ave.	-8077	4	-8072.3	2.9	1.2	1	52	52 $^{196}\text{Au}$		average	
$^{196}\text{Hg}(n,\gamma)^{197}\text{Hg}$	6785.3	1.5	6785.6	1.5	0.2	1	97	84 $^{197}\text{Hg}$	BNn		78Zg.A Z	
$^{197}\text{Ir}(\beta^-)^{197}\text{Pt}$	2000	200	2156	20	0.8	U					61Ho10	
$^{197}\text{Pt}(\beta^-)^{197}\text{Au}$	719.0	0.6	720.0	0.5	1.6	1	70	36 $^{197}\text{Au}$			71Pr03	
$^{197}\text{Hg}(\epsilon)^{197}\text{Au}$	415	20	600	3	9.2	B					65De20 *	
	610	100			-0.1	U					92Da14 *	
$^{197}\text{Tl}(\beta^+)^{197}\text{Hg}$	2220	100	2199	17	-0.2	U					61Ju05	
$^{197}\text{Pb}^m(\text{IT})^{197}\text{Pb}$	319.31	0.11	319.31	0.11	0.0	1	100	74 $^{197}\text{Pb}$			Ens053	
* $^{197}\text{Hg}-u$	$M-A=-30221(28)$ keV for $^{197}\text{Hg}^m$ at 298.93 keV										Nub16b **	
* $^{197}\text{Pb}-u$	$M-A=-24405(28)$ keV for $^{197}\text{Pb}^m$ at 319.31 keV										Nub16b **	
* $^{197}\text{Bi}-u$	$M-A=-19650(90)$ keV for mixture gs+m at 532(12) keV										Nub16b **	
* $^{197}\text{Bi}-^{133}\text{Cs}_{1.481}$	$Q=118887(12)$ $\mu\text{u}$ $M-A=-19690(11)$ keV corrected by $-16(22)$ keV due to possible contamination from $^{197}\text{Bi}^m$										08We02 **	
*											08We02 **	
* $^{197}\text{Po}-u$	$M-A=-13330(110)$ keV for mixture gs+m at 230#80 keV										Nub16b **	
* $^{197}\text{Po}-u$	$M-A=-13210(32)$ keV for mixture gs+m at 230#80 keV										Nub16b **	
* $^{197}\text{Po}(\alpha)^{193}\text{Pb}$	Also $E_\alpha=6283(5)$ keV from uncorrelated decays										96Ta18 **	
* $^{197}\text{Po}^m(\alpha)^{193}\text{Pb}^m$	Also $E_\alpha=6381(5)$ keV from uncorrelated decays										96Ta18 **	
* $^{197}\text{Hg}(\epsilon)^{197}\text{Au}$	pK=0.54(0.06) to $3/2^+$ level at 268.788 keV										Ens053 **	
* $^{197}\text{Hg}(\epsilon)^{197}\text{Au}$	pK=0.746(0.033) to 268.75 level $\rightarrow Q=574(+139-62)$ keV										Ens053 **	
$^{198}\text{Hg}-^{161}\text{Dy } ^{37}\text{Cl}$	74130	60	73927.5	1.0	-0.8	U			R04	4.0	64De15	
$^{198}\text{Hg}-^{163}\text{Dy } ^{35}\text{Cl}$	68979	37	69179.6	1.0	1.4	U			R04	4.0	64De15	
$^{198}\text{Hg}-u$	-33231.6	0.6	-33230.8	0.5	1.3	1	67	67 $^{198}\text{Hg}$	ST2	1.0	02Bf02	
$^{198}\text{Tl}-^{133}\text{Cs}_{1.489}$	111228.7	8.1				2			MA8	1.0	14Bo26 *	
$^{198}\text{Pb}-^{208}\text{Pb}_{.952}$	-5748	23	-5757	9	-0.4	-			MA6	1.0	01Sc41	
	ave.	-5739	18			-1.0	1	26	26 $^{198}\text{Pb}$		average	
$^{198}\text{Pb}-u$	-27990	104	-27985	9	0.1	U			GS1	1.0	00Ra23	
	-27951	30			-1.1	R			GS2	1.0	05Li24	
$^{198}\text{Bi}-u$	-21063	162	-20790	30	1.7	o			GS1	1.0	00Ra23 *	
	-20794	30				2			GS2	1.0	05Li24	
$^{198}\text{Bi}^n-u$	-20222	30				2			GS2	1.0	05Li24	
$^{198}\text{Po}-^{208}\text{Pb}_{.952}$	5616	24	5616	19	0.0	1	61	61 $^{198}\text{Po}$	MA6	1.0	01Sc41	
$^{198}\text{Po}-u$	-16600	104	-16611	19	-0.1	U			GS1	1.0	00Ra23	
$^{198}\text{At}-^{133}\text{Cs}_{1.489}$	133570.0	7.3	133574	6	0.5	o			MA8	1.0	13Ma.A	
	133573.7	6.3				2			MA8	1.0	13St25	
$^{198}\text{At}^m-^{133}\text{Cs}_{1.489}$	133898	39	133879	9	-0.5	U			MA8	1.0	13St25	
$^{198}\text{Hg } ^{35}\text{Cl}-^{196}\text{Hg } ^{37}\text{Cl}$	3885.91	1.66	3886	3	0.0	1	57	57 $^{196}\text{Hg}$	H33	2.5	80Ko25	
$^{198}\text{Pt}-^{197}\text{Au}_{1.005}$	1494.7	3.0	1493.8	2.2	-0.3	1	54	53 $^{198}\text{Pt}$	CPI	1.0	05Sh52	
$^{198}\text{Po}(\alpha)^{194}\text{Pb}$	6312.8	5.	6309.7	1.4	-0.6	U					67Si09 Z	
	6305.7	5.			0.8	U					67Tr06 Z	
	6301.2	8.			1.1	U					71Ho01 Z	
	6311.1	3.			-0.5	-			Bka		82Bo04 Z	
	6307.7	5.			0.4	U			Lvn		93Wa04	
	6309.7	5.0			0.0	U			Ara		96Ta18 *	
	6309.3	1.7			0.2	-			Tex		12Fo09	
	ave.	6309.7	1.4			0.0	1	100	60 $^{194}\text{Pb}$			average
	$^{198}\text{At}(\alpha)^{194}\text{Bi}$	6887.5	5.	6889.4	1.9	0.4	3					67Tr06 Z
		6904.9	7.			-2.2	U			Ora		75Ba.B Z
		6889.4	15.			0.0	U					80Ew03 Z
		6893.3	3.5			-1.1	3			Lvn		92Hu04 *
		6892.5	4.			-0.8	o			Jya		96En01
6887.4		6.			0.3	3			Jya		05Uu02	
6888.4		3.6			0.3	3			Tex		12Fo09	
6886.4	5.			0.6	3			Anv		14Ka23		
$^{198}\text{At}^m(\alpha)^{194}\text{Bi}^n$	6990.0	5.	6993.4	2.4	0.7	4					67Tr06 Z	
	6997.5	10.			-0.4	4					80Ew03 Z	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference		
$^{198}\text{At}^m(\alpha)^{194}\text{Bi}^n$	6997.6	4.	6993.4	2.4	-1.0	4			Lvn		92Hu04		
	6996.6	4.			-0.8	o			Jya		96En01		
	6991.5	6.			0.3	4			Jya		05Uu02		
	6990.5	5.			0.6	4			Anv		14Ka23		
$^{198}\text{Rn}(\alpha)^{194}\text{Po}$	7344.7	10.	7349	4	0.5	4					84Ca32		
	7353.8	5.			-0.9	4			Lvn		95Bi17		
	7344.7	6.			0.8	4			Jya		96En02		
$^{198}\text{Fr}(\alpha)^{194}\text{At}$	7869.2	20.4				5		Anv		13Ka16	*		
$^{198}\text{Fr}^m(\alpha)^{194}\text{At}^m$	7889.6	20.4				6		Anv		13Ka16			
$^{198}\text{Pt}(^{14}\text{C}, ^{16}\text{O})^{196}\text{Os}$	6130	40				2		BNL		83Bo29			
$^{198}\text{Pt}(t, \alpha)^{197}\text{Ir}$	10885	20				2		LAL		83Ci01			
$^{198}\text{Pt}(p, d)^{197}\text{Pt}$	-5332	3	-5331.0	2.1	0.3	1	47	47 $^{198}\text{Pt}$	Ors		78Be09	*	
$^{198}\text{Pt}(d, t)^{197}\text{Pt}$	-1305	20	-1298.3	2.1	0.3	U			Pit		64Co11		
	-1311	20			0.6	U			Tal		78Ya07		
$^{197}\text{Au}(n, \gamma)^{198}\text{Au}$	6512.35	0.11	6512.36	0.09	0.1	-			ILn		79Br26	Z	
	6512.32	0.16			0.2	-			Bdn		06Fi.A		
$^{197}\text{Au}(d, p)^{198}\text{Au}$	4282	30	4287.79	0.09	0.2	U			Pit		64Co11		
	4298	5			-2.0	U			MIT		67Sp09		
$^{197}\text{Au}(n, \gamma)^{198}\text{Au}$	ave.	6512.34	0.09	6512.36	0.09	0.2	1	99	63 $^{197}\text{Au}$		average		
$^{198}\text{Au}(\beta^-)^{198}\text{Hg}$	1372.3	0.7	1373.5	0.5	1.8	-					65Ke04	*	
	1372.8	1.2			0.6	-					65Pa08	*	
	ave.	1372.4	0.6		1.8	1	66	44 $^{198}\text{Au}$			average		
$^{198}\text{Tl}(\beta^+)^{198}\text{Hg}$	3460	80	3426	8	-0.4	U					61Gu02		
$^{198}\text{Bi}^n(\text{IT})^{198}\text{Bi}^m$	248.5	0.5				3			Lvn		92Hu04		
* $^{198}\text{Tl}-^{133}\text{Cs}_{1.489}$	$D_M=111812.3(8.1) \mu\text{u}$ for $^{198}\text{Tl}^m$ at 543.6(0.4) keV; $M-A=-26985.1(7.5)$ keV												
* $^{198}\text{Bi}-\text{u}$	$M-A=-19350(100)$ keV for mixture gs+m+n at 280(40) and 530(40) keV												
* $^{198}\text{Po}(\alpha)^{194}\text{Pb}$	Also $E_\alpha=6182(5)$ keV from uncorrelated decays												
* $^{198}\text{At}(\alpha)^{194}\text{Bi}$	$E_\alpha=6755(4), 6539(10), 6360(10)$ to ground state, 218, 396 levels												
* $^{198}\text{Fr}(\alpha)^{194}\text{At}$	Evaluator's interpretation of Fig. 3d, no 210 keV $\gamma$												
* $^{198}\text{Pt}(p, d)^{197}\text{Pt}$	$Q-Q(^{196}\text{Pt}(p, d))=365(3)$ keV												
* $^{198}\text{Au}(\beta^-)^{198}\text{Hg}$	$E_{\beta^-}=960.5(0.7) 961.0(1.2)$ respectively, to $2^+$ level at 411.80251 keV												
$^{199}\text{Hg}-\text{C}_2 \text{ } ^{35}\text{Cl}_5$	124023.43	0.53	124017.5	0.6	-4.5	B					H34	2.5	80Ko25
	124017.21	0.37			0.3	1	38	35 $^{199}\text{Hg}$	H48	2.5	03Ba49		
$^{199}\text{Hg}-^{183}\text{W O}$	23144.4	0.9	23141.9	0.9	-1.1	1	16	11 $^{183}\text{W}$	H48	2.5	03Ba49		
$^{199}\text{Hg}-^{162}\text{Dy } ^{37}\text{Cl}$	75661	41	75574.2	1.0	-0.5	U			R04	4.0	64De15		
$^{199}\text{Hg}-^{164}\text{Dy } ^{35}\text{Cl}$	70087	31	70247.8	1.0	1.3	U			R04	4.0	64De15		
$^{199}\text{Hg}-^{164}\text{Er } ^{35}\text{Cl}$	70310	80	70220.9	1.0	-0.3	U			R04	4.0	64De15		
$^{199}\text{Tl}-\text{u}$	-30123	30				2			GS2	1.0	05Li24		
$^{199}\text{Pb}-\text{u}$	-27028	137	-27087	11	-0.4	U			GS2	1.0	05Li24	*	
$^{199}\text{Bi}-\text{u}$	-22328	31	-22327	11	0.0	-			GS2	1.0	05Li24		
	-22263	30			-2.1	-			GS2	1.0	05Li24	*	
	ave.	-22294	22			-1.5	1	28	28 $^{199}\text{Bi}$		average		
$^{199}\text{Po}-\text{u}$	-16249	144	-16327	19	-0.5	U			GS1	1.0	00Ra23	*	
	-16327	38			0.0	R			GS2	1.0	05Li24		
	-16339	38			0.3	R			GS2	1.0	05Li24	*	
$^{199}\text{Bi}^m(\alpha)^{195}\text{Tl}$	5598.7	6.	5600	6	0.1	1	93	56 $^{195}\text{Tl}$			66Ma51		
$^{199}\text{Po}(\alpha)^{195}\text{Pb}$	6074.1	2.	6074.3	1.9	0.1	2			DbA		68Go.B	Z	
	6075.3	5.0			-0.2	2			Ara		96Ta18		
$^{199}\text{Po}^m(\alpha)^{195}\text{Pb}^m$	6190.7	5.	6183.3	1.7	-1.5	-					67Si09	Z	
	6177.6	5.1			1.1	-					67Tr06	Z	
	6182.3	3.1			0.3	-			DbA		68Go.B	Z	
	6183.5	3.			-0.1	-			Bka		82Bo04	Z	
	6183.5	5.0			0.0	-			Ara		96Ta18	*	
	6173.3	3.6			2.8	B			Tex		12Fo09		
	ave.	6183.3	1.7		0.0	1	100	59 $^{199}\text{Po}^m$			average		
$^{199}\text{At}(\alpha)^{195}\text{Bi}$	6775.1	5.	6777.3	1.2	0.4	-					67Tr06	Z	
	6781.3	3.			-1.3	-			Ora		75Ba.B	Z	
	6775.4	5.0			0.4	-			Ara		96Ta18		
	6779.4	6.			-0.4	U			Jya		05Uu02		
	6776.8	1.5			0.3	-			Tex		12Fo09		

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{199}\text{At}(\alpha)^{195}\text{Bi}$	ave.	6777.4	1.2	6777.3	1.2	-0.1	1	100	89 $^{199}\text{At}$		average
$^{199}\text{Rn}(\alpha)^{195}\text{Po}$		7133.7	15.	7132	4	-0.1	4				80Di07
		7132.7	10.			-0.1	4				82Hi14
		7138.8	10.			-0.7	4				84Ca32
		7112.2	15.			1.3	o		Jya		96Le09
		7132.6	6.			-0.1	4		Jya		05Uu02
		7121.4	10.2			1.0	4		Anv		14Ka23
$^{199}\text{Rn}^m(\alpha)^{195}\text{Po}^m$		7205.1	15.	7203	4	-0.1	4				80Di07
		7205.1	10.			-0.2	4				82Hi14
		7204.1	10.			-0.1	4				84Ca32
		7205.1	15.			-0.1	o		Jya		96Le09
		7205.1	6.			-0.3	4		Jya		05Uu02
		7194.9	10.2			0.8	4		Anv		14Ka23
$^{199}\text{Fr}(\alpha)^{195}\text{At}$		7812.3	40.	7817	10	0.1	U				99Ta20
		7821.5	11.			-0.4	4		Anv		13Ka16
		7801.1	20.			0.8	4		Jya		13Uu01
$^{199}\text{Fr}^m(\alpha)^{195}\text{At}^m$		7833.7	6.	7833	6	-0.2	3		Anv		13Ka16
		7825.6	15.3			0.5	3		Jya		13Uu01
$^{199}\text{Fr}^n(\alpha)^{195}\text{At}^p$		7968.4	20.				4		Jya		13Uu01
$^{199}\text{Hg}(p,t)^{197}\text{Hg}$		-6734	29	-6667	3	2.3	U		Pri		81Ko13
		-6658	8			-1.1	1	16	16 $^{197}\text{Hg}$	Ors	82Be21
$^{198}\text{Pt}(^{18}\text{O}, ^{17}\text{F})^{199}\text{Ir}$		-8240	41				2				95Zh10
$^{198}\text{Pt}(n,\gamma)^{199}\text{Pt}$		5602	3	5556.0	0.5	-15.3	B				68Sa13
		5556.0	0.5				2		BNn		83Ca04 Z
$^{198}\text{Pt}(d,p)^{199}\text{Pt}$		3347	20	3331.4	0.5	-0.8	U		Pit		64Co11
$^{198}\text{Au}(n,\gamma)^{199}\text{Au}$		7584.27	0.15	7584.28	0.06	0.1	o		ILn		79Br26 Z
		7584.28	0.06			0.0	1	100	80 $^{199}\text{Au}$	ILn	91Ma65
$^{198}\text{Hg}(n,\gamma)^{199}\text{Hg}$		6665.2	0.5	6663.1	0.6	-4.3	B		CRn		75Lo03
$^{199}\text{Hg}(\gamma,n)^{198}\text{Hg}$		-6590	90	-6663.1	0.6	-0.8	U		Phi		60Ge01
$^{199}\text{Pt}(\beta^-)^{199}\text{Au}$		1690	50	1705.1	2.1	0.3	U				64Jo09
$^{199}\text{Au}(\beta^-)^{199}\text{Hg}$		453.0	1.0	452.3	0.6	-0.7	1	38	20 $^{199}\text{Au}$		68Be06
$^{199}\text{Tl}(\beta^+)^{199}\text{Hg}$		1420	150	1487	28	0.4	U				75Ma05 *
$^{199}\text{Pb}(\beta^+)^{199}\text{Tl}$		2870	110	2828	30	-0.4	U				70Do.A *
$^{199}\text{Bi}^m(\text{IT})^{199}\text{Bi}$		667	5	667	3	-0.1	-				80Br23
		667	5			-0.1	-				85St02
	ave.	667	4			-0.1	1	98	64 $^{199}\text{Bi}^m$		average
$^{199}\text{At}^m(\text{IT})^{199}\text{At}$		255	20	244.0	1.0	-0.5	o		Jya		13Ja06 *
		244	1				2		Jya		14Au03
* $^{199}\text{Pb}-u$	$M - A = -24961(28)$ keV for mixture gs+m at 429.5(2.7) keV										
* $^{199}\text{Bi}-u$	$M - A = -20071(28)$ keV for $^{199}\text{Bi}^m$ at 667(3) keV										
* $^{199}\text{Po}-u$	$M - A = -14980(100)$ keV for mixture gs+m at 311.9(2.7) keV										
* $^{199}\text{Po}-u$	$M - A = -14909(35)$ keV for $^{199}\text{Po}^m$ at 311.9(2.7) keV										
* $^{199}\text{Po}^m(\alpha)^{195}\text{Pb}^m$	Also $E_\alpha = 6059(5)$ keV from uncorrelated decays										
* $^{199}\text{Tl}(\beta^+)^{199}\text{Hg}$	KL+<500(100) giving $Q_i(1620, (1/2^-, 3/2^-)$ level at 1221.17 fed. Reanalyzed										
* $^{199}\text{Pb}(\beta^+)^{199}\text{Tl}$	$p^+ = 0.04(0.01)$ to $3/2^+$ level at 366.89 keV, recalculated										
* $^{199}\text{At}^m(\text{IT})^{199}\text{At}$	Combining lines 110(20) 160(20) 240(30) as read from Fig. 8a										
$^{200}\text{Au}-u$	-29237	34	-29243	29	-0.2	1	71	71 $^{200}\text{Au}$	GS3	1.0	08Ch.A
$^{200}\text{Au}^m-u$	-28135	33	-28163	28	-0.8	1	73	73 $^{200}\text{Au}^m$	GS3	1.0	08Ch.A
$^{200}\text{Hg}-\text{C } ^{13}\text{C } ^{35}\text{Cl}_5$	120707.97	1.22	120708.6	0.6	0.2	U			H34	2.5	80Ko25
$^{200}\text{Hg}-^{165}\text{Ho } ^{35}\text{Cl}$	69116	33	69146.2	1.2	0.2	U			R04	4.0	64De15
$^{200}\text{Hg}-^{163}\text{Dy } ^{37}\text{Cl}$	73527	42	73687.5	1.0	1.0	U			R04	4.0	64De15
$^{200}\text{Pb}-u$	-28179	30	-28182	12	-0.1	R			GS2	1.0	05Li24
$^{200}\text{Bi}-u$	-21888	57	-21869	24	0.3	R			GS2	1.0	05Li24 *
$^{200}\text{Po}-u$	-18170	104	-18188	8	-0.2	U			GS1	1.0	00Ra23
	-18204	30			0.5	U			GS2	1.0	05Li24

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{200}\text{Hg}-^{208}\text{Pb}_{962}$	-9205	28	-9212.2	1.2	-0.3	U			MA6	1.0	01Sc41
$^{200}\text{Hg } ^{35}\text{Cl}-^{198}\text{Hg } ^{37}\text{Cl}$	4525	2	4507.9	0.6	-2.1	U			H17	4.0	64Mc07
$^{200}\text{Po}(\alpha)^{196}\text{Pb}$	4508.80	0.48			-0.8	1	28	17 $^{200}\text{Hg}$	H33	2.5	80Ko25
	5979.8	5.	5981.6	1.8	0.4	-					67Si09 Z
	5980.0	3.			0.5	-					67Tr06 Z
	5983.4	3.			-0.6	-					70Ra14 Z
	5981.8	5.0			0.0	-			Ara		96Ta18 *
$^{200}\text{At}(\alpha)^{196}\text{Bi}$	ave. 5981.5	1.9			0.1	1	99	79 $^{196}\text{Pb}$			average
	6594.9	5.	6596.2	1.3	0.3	3					67Tr06 Z
	6596.9	2.			-0.4	3			Ora		75Ba.B Z
	6593.1	5.			0.6	o			Lvn		87Va09
	6596.1	2.			0.0	3			Lvn		92Hu04
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}$	6593.1	5.0			0.6	3			Ara		96Ta18
	6599.1	6.			-0.5	U			Jya		05Uu02
	6708.3	5.	6709.1	2.6	0.2	3			Ora		75Ba.B Z
	6705.4	5.			0.7	o			Lvn		87Va09
	6709.5	3.			-0.1	3			Lvn		92Hu04
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^m$	6542.8	5.	6542.7	1.3	0.0	4					67Tr06 Z
	6542.9	2.			-0.1	4			Ora		75Ba.B Z
	6540.0	5.			0.5	o			Lvn		87Va09
	6542.1	2.			0.3	4			Lvn		92Hu04
	6545.1	5.0			-0.5	4			Ara		96Ta18
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^n$	6544.1	6.			-0.2	U			Jya		05Uu02
	6439.5	5.	6437.5	2.0	-0.4	4					67Tr06 *
	6438.5	5.			-0.2	4			Ora		75Ba.B *
	6433.8	5.			0.7	o			Lvn		87Va09 *
	6439.2	3.			-0.6	4			Lvn		92Hu04 *
$^{200}\text{Rn}(\alpha)^{196}\text{Po}$	6430.5	5.0			1.4	4			Ara		96Ta18 *
	6436.7	6.			0.1	4			Jya		05Uu02 *
	7020.6	10.	7043.4	2.1	2.3	U					67Va.A
	7050.3	8.			-0.9	U					71Ho01
	7040.1	10.			0.3	U			Lvn		84Co.A
$^{200}\text{Fr}(\alpha)^{196}\text{At}$	7043.5	2.5			-0.1	4			Lvn		93Wa04
	7042.1	12.			0.1	o			Ara		95Le04
	7039.0	10.			0.4	o			Jya		96Le09
	7042.1	5.1			0.2	4			Ara		96Ta18
	7044.1	6.			-0.1	4			Jya		05Uu02
$^{200}\text{Fr}(\alpha)^{196}\text{At}$	7055.4	30.			-0.4	U			Anv		10He25
	7653.4	30.	7622	4	-1.0	U			RIa		95Mo14
	7620.7	9.			0.2	5			Jya		96En01
	7625.8	12.			-0.3	o			Anv		05De01
	7620.7	15.			0.1	o			Jya		13Uu01 *
$^{200}\text{Fr}^m(\alpha)^{196}\text{At}^m$	7622.7	5.			-0.1	5			Anv		14Ka23
	7704.4	15.				4			Jya		96En01 *
$^{198}\text{Pt}(t,p)^{200}\text{Pt}$	4356	20				2					81Ci01
$^{199}\text{Hg}(n,\gamma)^{200}\text{Hg}$	8029.1	0.3	8028.52	0.11	-1.9	-			BNn		67Sc30 Z
	8029.6	0.5			-2.2	U			CRn		75Lo03 Z
	8028.51	0.18			0.1	-			ILn		79Br25 Z
	8028.37	0.17			0.9	-			Bdn		06Fi.A
	ave. 8028.53	0.11			-0.1	1	98	64 $^{200}\text{Hg}$			average
$^{200}\text{Au}(\beta^-)^{200}\text{Hg}$	2273	100	2263	27	-0.1	-					59Ro53 *
	2200	100			0.6	-					60Gi01
	2260	70			0.0	-					72He36 *
$^{200}\text{Au}^m(\beta^-)^{200}\text{Hg}$	ave. 2250	50			0.3	1	29	29 $^{200}\text{Au}$			average
	3202	50	3270	26	1.4	1	27	27 $^{200}\text{Au}^m$			72Cu07 *
$^{200}\text{Tl}(\beta^+)^{200}\text{Hg}$	2450	10	2456	6	0.6	2					57He43 *
	2459	7			-0.4	2					62Va10 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{200}\text{At}^n(\text{IT})^{200}\text{At}^m$	230.9	0.2				4			Lvn		92Hu04
$^{200}\text{Bi}-u$	$M-A=-20338(28)$ keV for mixture gs+m at 100#70 keV										Nub16b **
$^{200}\text{Po}(\alpha)^{196}\text{Pb}$	Also $E_\alpha=5863(5)$ keV from uncorrelated decays										96Ta18 **
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^n$	$E_\alpha=6536.7(5,Z)$ from $^{200}\text{At}^n$ 230.9 above $^{200}\text{At}^m$										92Hu04 **
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^n$	$E_\alpha=6535.8(5,Z)$ from $^{200}\text{At}^n$ 230.9 above $^{200}\text{At}^m$										92Hu04 **
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^n$	$E_\alpha=6301(5); 6535(5)$ from $^{200}\text{At}^n$ 230.9 above $^{200}\text{At}^m$										92Hu04 **
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^n$	$E_\alpha=6306(5); 6538(3)$ from $^{200}\text{At}^n$ 230.9 above $^{200}\text{At}^m$										92Hu04 **
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^n$	$E_\alpha=6528(5)$ from $^{200}\text{At}^n$ 230.9 above $^{200}\text{At}^m$										92Hu04 **
$^{200}\text{At}^m(\alpha)^{196}\text{Bi}^n$	$E_\alpha=6534(6)$ from $^{200}\text{At}^n$ 230.9 above $^{200}\text{At}^m$										92Hu04 **
$^{200}\text{Fr}(\alpha)^{196}\text{At}$	Same group, but much less events than preceding item										WgM151**
$^{200}\text{Fr}^m(\alpha)^{196}\text{At}^m$	Correlated with $^{196}\text{At}^m$ $E_\alpha=6880(15)$ ; two events only										96En01 **
$^{200}\text{Au}(\beta^-)^{200}\text{Hg}$	$E_{\beta^-}=2250(200)$ to ground state, and $700(100)$ to levels $1^+$ at 1570.275, $2^+$ at 1573.663, $2^+$ at 1593.423 keV										Ens077 **
$^{200}\text{Au}(\beta^-)^{200}\text{Hg}$	$E_{\beta^-}=2260(100), 670(70)$ to ground state, $2^+$ level at 1593.423 keV										Ens077 **
$^{200}\text{Au}^m(\beta^-)^{200}\text{Hg}$	$E_{\beta^-}=560(50)$ to $11^-$ level at 2641.54 keV										Ens077 **
$^{200}\text{Tl}(\beta^+)^{200}\text{Hg}$	$E_{\beta^+}=1052(10) 1069(7)$ respectively, to $2^+$ level at 367.943 keV, and other $E_{\beta^+}$										Ens077 **
$^{201}\text{Hg}-^{185}\text{Re O}$	22440	5	22430.1	1.1	-0.8	U			H48	2.5	03Ba49
$^{201}\text{Hg}-\text{C}_2^{35}\text{Cl}_4^{37}\text{Cl}$	128995.43	0.61	128989.7	0.8	-3.8	B			H34	2.5	80Ko25
$^{201}\text{Hg}-^{164}\text{Dy}^{37}\text{Cl}$	75086	42	75220.0	1.1	0.8	U			R04	4.0	64De15
$^{201}\text{Hg}-^{166}\text{Er}^{35}\text{Cl}$	71186	35	71151.3	1.5	-0.2	U			R04	4.0	64De15
$^{201}\text{Pb}-u$	-27418	198	-27130	15	1.5	U			GS2	1.0	05Li24 *
$^{201}\text{Bi}-u$	-22935	30	-22991	16	-1.9	R			GS2	1.0	05Li24
	-22995	30			0.1	R			GS2	1.0	05Li24 *
$^{201}\text{Po}-u$	-17760	190	-17736	5	0.1	U			GS1	1.0	00Ra23 *
	-17649	30			-2.9	U			GS2	1.0	05Li24
$^{201}\text{Po}^m-u$	-17305	30	-17281	5	0.8	U			GS2	1.0	05Li24
$^{201}\text{At}-u$	-11573	31	-11583	9	-0.3	U			GS2	1.0	05Li24
$^{201}\text{Hg}^{35}\text{Cl}-^{199}\text{Hg}^{37}\text{Cl}$	4981	2	4972.2	0.6	-1.1	U			H17	4.0	64Mc07
	4972.65	0.37			-0.5	1	47	$39^{201}\text{Hg}$	H33	2.5	80Ko25
	4971.8	1.0			0.1	U			H48	2.5	03Ba49
$^{201}\text{Bi}(\alpha)^{197}\text{Tl}$	4500.3	6.				5					66Ma51 *
$^{201}\text{Po}(\alpha)^{197}\text{Pb}$	5793.9	5.	5799.3	1.7	1.1	-					67Tr06 Z
	5799.4	2.			-0.1	-			DbA		68Go.B Z
	5800.4	4.			-0.3	-					70Ra14 Z
	ave.	5799.0	1.7		0.2	1	98	$71^{201}\text{Po}$			average
$^{201}\text{Po}^m(\alpha)^{197}\text{Pb}^m$	5899.0	5.1	5903.8	1.7	0.9	2					67Tr06 Z
	5904.5	2.0			-0.4	2			DbA		68Go.B Z
	5903.9	4.1			0.0	2					70Ra14 Z
$^{201}\text{At}(\alpha)^{197}\text{Bi}$	6470.7	3.	6472.8	1.6	0.7	4					67Tr06 Z
	6476.2	5.			-0.7	U					74Ho27 Z
	6474.0	2.			-0.6	4			Ora		75Ba.B Z
	6471.0	5.0			0.4	U			Ara		96Ta18
	6472.0	4.			0.2	4			Anv		05De01
$^{201}\text{Rn}(\alpha)^{197}\text{Po}$	6862.8	8.	6860.7	2.3	-0.3	U					67Va.A
	6858.8	8.			0.2	U					71Ho01
	6866.9	20.			-0.3	U			GSa		87He10
	6860.5	2.5			0.1	4			Lvn		93Wa04
	6863.8	7.			-0.4	o			Ara		95Le04
	6861.8	5.0			-0.2	4			Ara		96Ta18
$^{201}\text{Rn}^m(\alpha)^{197}\text{Po}^m$	6906.8	5.	6909.5	2.1	0.5	5					67Va17 Z
	6909.0	8.			0.1	U					71Ho01 Z
	6907.7	20.			0.1	U					87He10
	6909.9	2.5			-0.1	5			Lvn		93Wa04
	6915.9	7.			-0.9	o			Ara		95Le04
	6910.7	5.0			-0.3	5			Ara		96Ta18
	6925.1	30.			-0.5	U			Anv		10He25



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{201}\text{Fr}(\alpha)^{197}\text{At}$	7538.0	15.	7519	4	-1.3	U					80Ew03
	7510.8	7.			1.1	o			Jya		96En01
	7529.1	7.			-1.4	o			Anv		05De01
	7519.0	8.			0.0	2			Jya		05Uu02
	7519.0	5.			0.0	2			Anv		14Ka23
$^{201}\text{Fr}^m(\alpha)^{197}\text{At}^m$	7605.7	8.	7601	6	-0.6	2			Jya		05Uu02
	7596.4	8.			0.6	2			Anv		14Ka23
$^{201}\text{Ra}(\alpha)^{197}\text{Rn}$	8001.5	12.				4			Anv		14Ka23
$^{201}\text{Ra}^m(\alpha)^{197}\text{Rn}^m$	8065.8	20.				4			Jya		05Uu02
$^{201}\text{Hg}(\gamma,n)^{200}\text{Hg}$	-6210	70	-6230.6	0.6	-0.3	U			Phi		60Ge01
$^{201}\text{Pt}(\beta^-)^{201}\text{Au}$	2660	50				2					63Go06
$^{201}\text{Au}(\beta^-)^{201}\text{Hg}$	1270	100	1262	3	-0.1	U					72Pa24
$^{201}\text{Tl}(\epsilon)^{201}\text{Hg}$	470	70	482	14	0.2	U					60Gu05 *
$^{201}\text{Pb}(\beta^+)^{201}\text{Tl}$	1900	40	1910	19	0.2	1	21	11 $^{201}\text{Tl}$			79Do09 *
* $^{201}\text{Pb}-u$	$M - A = -25225(28)$ keV for mixture gs+m at 629.1 keV										Nub16b **
* $^{201}\text{Bi}-u$	$M - A = -20573(28)$ keV for $^{201}\text{Bi}^m$ at 846.35 keV										Nub16b **
* $^{201}\text{Po}-u$	$M - A = -16330(100)$ keV for mixture gs+m at 423.8(2.4) keV										Nub16b **
* $^{201}\text{Bi}(\alpha)^{197}\text{Tl}$	$E_\alpha = 5240(6)$ from $^{201}\text{Bi}^m$ at 846.35 keV										Nub16b **
* $^{201}\text{Tl}(\epsilon)^{201}\text{Hg}$	pK=0.70(0.04) to $1/2^-$ level at 167.47 keV, recalculated										Ens073 **
* $^{201}\text{Pb}(\beta^+)^{201}\text{Tl}$	p <sup>+</sup> = 10(2) × 10 <sup>-3</sup> to $3/2^+$ level at 331.16 keV										Ens073 **
$^{202}\text{Pt}-u$	-24425	34	-24361	27	1.9	o			GS3	1.0	08Ch.A
	-24361	27				2			GS3	1.0	12Ch19
$^{202}\text{Au}-u$	-26202	34	-26144	25	1.7	o			GS3	1.0	08Ch.A
	-26144	25				2			GS3	1.0	12Ch19
$^{202}\text{Hg}-\text{C }^{13}\text{C }^{35}\text{Cl}_4 \text{ }^{37}\text{Cl}$	125976.01	1.32	125975.4	0.8	-0.2	U			H34	2.5	80Ko25
$\text{C}_{16} \text{H}_{10}-^{202}\text{Hg}$	107663	40	107606.7	0.8	-0.9	U			R08	1.5	69De19
$\text{C}_{15} \text{ }^{13}\text{C } \text{H}_9 - ^{202}\text{Hg}$	103102	60	103136.5	0.8	0.4	U			R08	1.5	69De19
$^{202}\text{Hg}-^{167}\text{Er }^{35}\text{Cl}$	69740	60	69736.8	1.5	0.0	U			R04	4.0	64De15
$^{202}\text{Hg}-^{165}\text{Ho }^{37}\text{Cl}$	74470	50	74413.0	1.3	-0.3	U			R04	4.0	64De15
$^{202}\text{Tl}-^{133}\text{Cs}_{1.519}$	115727.2	3.7	115727.6	1.7	0.1	1	22	22 $^{202}\text{Tl}$	MA8	1.0	16We.A
$^{202}\text{Pb}-u$	-27823	30	-27848	4	-0.8	U			GS2	1.0	05Li24 *
$^{202}\text{Pb}-^{133}\text{Cs}_{1.519}$	115773.4	3.6	115770	4	-0.9	o			MA8	1.0	10Bo.A
	115769.2	4.4			0.2	1	86	86 $^{202}\text{Pb}$	MA8	1.0	14Bo26
$^{202}\text{Bi}-u$	-22282	30	-22267	17	0.5	1	30	30 $^{202}\text{Bi}$	GS2	1.0	05Li24
$^{202}\text{Po}-u$	-19270	104	-19261	9	0.1	U			GS1	1.0	00Ra23
	-19243	30			-0.6	U			GS2	1.0	05Li24
$^{202}\text{Hg }^{35}\text{Cl}_2 - ^{198}\text{Hg }^{37}\text{Cl}_2$	9774.87	1.06	9774.6	0.8	-0.1	U			H33	2.5	80Ko25
$^{202}\text{Hg }^{35}\text{Cl} - ^{200}\text{Hg }^{37}\text{Cl}$	5271	3	5266.8	0.6	-0.4	U			H17	4.0	64Mc07
	5266.76	0.43			0.0	1	35	28 $^{202}\text{Hg}$	H33	2.5	80Ko25
$^{202}\text{Tl}-^{203}\text{Tl}_{995}$	-372.4	2.1	-373.2	1.7	-0.4	1	63	48 $^{202}\text{Tl}$	MA8	1.0	16We.A
$^{202}\text{Po}(\alpha)^{198}\text{Pb}$	5700.9	2.	5701.0	1.7	0.0	-			DbA		68Go.B Z
	5701.6	3.			-0.2	-					70Ra14 Z
	5701.1	1.7			-0.1	1	99	74 $^{198}\text{Pb}$			average
$^{202}\text{At}(\alpha)^{198}\text{Bi}$	6355.8	3.	6353.8	1.3	-0.6	3					63Ho18 Z
	6351.7	3.			0.7	3					67Tr06 Z
	6353.2	5.			0.1	3					74Ho27 Z
	6353.9	2.			0.0	3			Ora		75Ba.B Z
	6354	5			0.0	3			Lvn		92Hu04 *
	6355.0	6.0			-0.2	3			Ara		96Ta18
$^{202}\text{At}^m(\alpha)^{198}\text{Bi}^m$	6259.9	2.	6259.0	1.3	-0.5	4					63Ho18 Z
	6256.8	3.			0.7	4					67Tr06 Z
	6257.2	5.			0.4	U					74Ho27 Z
	6259.0	2.			0.0	4			Ora		75Ba.B *
	6260.0	5.			-0.2	U			Lvn		92Hu04 *
	6257.1	6.0			0.3	U			Ara		96Ta18
$^{202}\text{Rn}(\alpha)^{198}\text{Po}$	6771.0	3.	6773.8	1.8	0.9	2					67Va17 Z
	6772.3	10.			0.2	U			GSa		87He10
	6775.3	2.5			-0.6	2			Lvn		93Wa04
	6773.4	7.			0.1	o			Ara		95Le04
	6775.4	5.0			-0.3	2			Ara		96Ta18

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{202}\text{Fr}(\alpha)^{198}\text{At}$	7397.7	15.	7386	4	-0.8	3					80Ew03 *
	7382.5	11.			0.3	3			Lvn		92Hu04 *
	7389.6	6.			-0.6	o			Jya		96En01 *
	7387.6	8.			-0.2	3			Jya		05Uu02
	7384.5	5.1			0.3	3			Anv		14Ka23
$^{202}\text{Fr}^m(\alpha)^{198}\text{At}^m$	7382.5	11.	7376	4	-0.6	5			Lvn		92Hu04 *
	7388.6	6.			-2.1	o			Jya		96En01
	7381.5	8.			-0.7	5			Jya		05Uu02
	7372.2	5.			0.7	5			Anv		14Ka23
	8019.1	60.	7880	7	-2.3	o			Jya		96Le09
$^{202}\text{Ra}(\alpha)^{198}\text{Rn}$	7896.7	20.			-0.8	5			Jya		05Uu02
	7878.3	7.1			0.3	5			Anv		14Ka23
	11567	15	11580	3	0.9	U			LAl		81Fl05
$^{202}\text{Hg}(t,\alpha)^{201}\text{Au}$	-979.9	3.1	-980	3	0.0	1	100	100 $^{201}\text{Au}$			94Gr07
$^{202}\text{Hg}(d,^3\text{He})^{201}\text{Au}-^{206}\text{Pb}(\alpha)^{205}\text{Tl}$	7754.9	0.5	7754.10	0.20	-1.6	-			BNn		75Br02 Z
$^{201}\text{Hg}(n,\gamma)^{202}\text{Hg}$	7756.4	0.5			-4.6	B			CRn		75Lo03 Z
	7753.93	0.22			0.8	-			Bdn		06Fi.A
	-7600	130	-7754.10	0.20	-1.2	U			Phi		60Ge01
$^{202}\text{Hg}(\gamma,n)^{201}\text{Hg}$	ave.	7754.09	0.20	7754.10	0.20	0.1	1	97	59 $^{201}\text{Hg}$		average
$^{201}\text{Hg}(n,\gamma)^{202}\text{Hg}$	3500	300	2992	23	-1.7	U					67Wa23
$^{202}\text{Au}(\beta^-)^{202}\text{Hg}$	2700	300			1.0	U					72Bu05
$^{202}\text{Tl}(\epsilon)^{202}\text{Hg}$	1245	25	1365.1	1.6	4.8	B					66Le06 *
$^{202}\text{Pb}(\epsilon)^{202}\text{Tl}$	55	20	40	4	-0.8	U					54Hu61
$^{202}\text{At}^n(\text{IT})^{202}\text{At}^m$	391.7	0.2				5			Lvn		92Hu04
* $^{202}\text{Pb}-u$	$M-A=-23747(28)$ keV for $^{202}\text{Pb}^m$ at 2169.85 keV										
* $^{202}\text{At}(\alpha)^{198}\text{Bi}$	$E_\alpha=6228(5), 6070(10), 5929(10)$ to ground state, 164, 303 levels										
* $^{202}\text{At}^m(\alpha)^{198}\text{Bi}^m$	Assignment to $^{202}\text{At}^m$ in reference; Z recalibrated										
* $^{202}\text{At}^m(\alpha)^{198}\text{Bi}^m$	$E_\alpha=6135(5)$ ; and $6277(5)$ from $^{202}\text{At}^n(\alpha)^{198}\text{Bi}^n$ , with										
* $^{202}\text{At}^n(\text{IT})^{202}\text{At}^m=391.7(0.5)$ and $^{198}\text{Bi}^n(\text{IT})^{198}\text{Bi}^m=248.5(0.5)$ keV											
* $^{202}\text{Fr}(\alpha)^{198}\text{At}$	$E_\alpha=7251(10)$ has a doublet structure										
* $^{202}\text{Fr}(\alpha)^{198}\text{At}$	$E_\alpha=7237(8)$ , is a doublet										
* $^{202}\text{Fr}(\alpha)^{198}\text{At}$	$^{202}\text{Fr}$ $E_\alpha$ 's in correlation with At daughters										
* $^{202}\text{Fr}^m(\alpha)^{198}\text{At}^m$	$E_\alpha=7237(8)$ , is a doublet										
* $^{202}\text{Tl}(\epsilon)^{202}\text{Hg}$	$pK=0.305(0.020)$ to $2^+$ level at 959.94 keV										
$\text{C}_{16}\text{H}_{11}-^{203}\text{Tl}$	113735	43	113731.3	1.3	-0.1	U			R08	1.5	69De19
$\text{C}_{15}\text{H}_{13}\text{H}_{10}-^{203}\text{Tl}$	109216	95	109261.1	1.3	0.3	U			R08	1.5	69De19
$\text{C}_{14}\text{N}_2\text{H}_7-^{203}\text{Tl}$	88540	48	88579.2	1.3	0.5	U			R08	1.5	69De19
$^{203}\text{Tl}-^{166}\text{Er}$ $^{37}\text{Cl}$	76190	48	76142.4	1.8	-0.7	U			R08	1.5	69De19
$^{203}\text{Tl}-^{168}\text{Er}$ $^{35}\text{Cl}$	71069	36	71115.1	1.8	0.9	U			R08	1.5	69De19
$^{203}\text{Tl}-^{133}\text{Cs}_{1.526}$	116622.5	3.8	116624.3	1.3	0.5	U			MA8	1.0	16We.A
$^{203}\text{Pb}-u$	-26594	30	-26609	7	-0.5	U			GS2	1.0	05Li24
$^{203}\text{Po}-u$	-18581	30	-18584	9	-0.1	U			GS2	1.0	05Li24
$^{203}\text{At}-u$	-13042	30	-13057	11	-0.5	1	14	14 $^{203}\text{At}$	GS2	1.0	05Li24
$^{203}\text{Fr}-^{133}\text{Cs}_{1.526}$	145205	17	145221	7	1.0	1	15	15 $^{203}\text{Fr}$	MA8	1.0	08We02
$^{203}\text{At}-^{208}\text{Pb}_{.976}$	9690	25	9731	11	1.6	1	21	21 $^{203}\text{At}$	MA6	1.0	01Sc41
$^{203}\text{Rn}^m-^{208}\text{Pb}_{.976}$	16579	30	16571	19	-0.3	1	41	41 $^{203}\text{Rn}^m$	SH1	1.0	13Dr04
$^{203}\text{Tl}$ $^{35}\text{Cl}-^{201}\text{Hg}$ $^{37}\text{Cl}$	4997	3	4991.1	1.2	-0.5	U			H17	4.0	64Mc07
	4995.23	1.49			-1.1	1	10	8 $^{203}\text{Tl}$	H36	2.5	85De40
$^{202}\text{Hg}$ $\text{H}-^{203}\text{Tl}$	6154	34	6124.6	1.2	-0.6	U			R08	1.5	69De19
$^{203}\text{Tl}-^{167}\text{Er}$ $^{35}\text{Cl}$	71436	36	71437.2	1.8	0.0	U			R08	1.5	69De19
$^{167}\text{Er}$ $^{37}\text{Cl}-^{203}\text{Tl}$	-74404	33	-74387.3	1.8	0.3	U			R08	1.5	69De19
$^{169}\text{Tm}$ $^{35}\text{Cl}-^{203}\text{Tl}$	-69257	29	-69273.0	1.5	-0.4	U			R08	1.5	69De19
$^{203}\text{Tl}-^{202}\text{Hg}$	1722	20	1700.4	1.2	-0.7	U			R08	1.5	69De19
$^{203}\text{Tl}-^{201}\text{Hg}$	1999	29	2041.0	1.2	1.0	U			R08	1.5	69De19

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{203}\text{Po}(\alpha)^{199}\text{Pb}$	5496	5				5			Db		68Go.B *
$^{203}\text{At}(\alpha)^{199}\text{Bi}$	6210.3	1.	6210.1	0.8	-0.2	-					63Ho18 Z
	6208.7	3.			0.5	-					67Tr06 Z
	6209.4	2.			0.3	-			Db		68Go.B Z
	6211.7	3.			-0.5	-			Ora		75Ba.B
	6210.6	5.0			-0.1	U			Ara		96Ta18
	ave.	6210.1	0.8		0.0	1	100	61 $^{203}\text{At}$			average
$^{203}\text{Rn}(\alpha)^{199}\text{Po}$	6628.6	5.	6629.9	2.1	0.3	3					67Va17 Z
	6630.2	2.5			-0.1	3			Lvn		93Wa04
	6630	10			0.0	U			Jya		95Uu01
	6629.8	5.0			0.0	3			Ara		96Ta18
$^{203}\text{Rn}^m(\alpha)^{199}\text{Po}^m$	6679.5	3.	6680.5	1.6	0.3	-					67Va17 Z
	6681.9	10.			-0.1	U			GSa		87He10
	6680.9	2.5			-0.2	-			Lvn		93Wa04
	6683.9	7.			-0.5	o			Ara		95Le04
	6679.8	3.			0.2	-			Jya		96Le09
	6682.9	5.0			-0.5	-			Ara		96Ta18
	ave.	6680.5	1.6		0.0	1	100	59 $^{203}\text{Rn}^m$			average
$^{203}\text{Fr}(\alpha)^{199}\text{At}$	7275.6	5.	7275	4	-0.1	-					67Va20 Z
	7281.7	10.			-0.7	-					80Ew03 Z
	7263.4	25.			0.5	o			Jya		94Le05
	7273.6	6.			0.2	-			Jya		05Uu02
	ave.	7276	4		-0.2	1	95	85 $^{203}\text{Fr}$			average
$^{203}\text{Fr}^m(\alpha)^{199}\text{At}^m$	7391.9	5.				3			Jya		13Ja06
$^{203}\text{Ra}(\alpha)^{199}\text{Rn}$	7729.6	20.	7736	6	0.3	o			Jya		96Le09
	7741.8	8.			-0.7	5			Jya		05Uu02
	7727.6	10.			0.9	5			Anv		14Ka23
$^{203}\text{Ra}^m(\alpha)^{199}\text{Rn}^m$	7768.4	20.	7763	6	-0.3	o			Jya		96Le09
	7765.3	8.			-0.3	5			Jya		05Uu02
	7760.2	8.2			0.3	5			Anv		14Ka23
$^{203}\text{Tl}(p,t)^{201}\text{Tl}$	-6240	15	-6241	14	-0.1	1	89	89 $^{201}\text{Tl}$	Yal		71Ki01
$^{202}\text{Hg}(d,p)^{203}\text{Hg}-^{204}\text{Hg}(\gamma)^{205}\text{Hg}$	325	5	326	4	0.3	1	52	48 $^{205}\text{Hg}$	Pit		72Mo12
$^{203}\text{Tl}(p,d)^{202}\text{Tl}$	-5630	20	-5627.9	1.6	0.1	U			Yal		71Ki01
$^{203}\text{Au}(\beta^-)^{203}\text{Hg}$	2040	60	2126	3	1.4	U					94We02
$^{203}\text{Hg}(\beta^-)^{203}\text{Tl}$	489.2	2.	492.1	1.2	1.5	-					54Th17 *
	493.2	2.			-0.5	-					55Ma40 *
	493.2	3.			-0.4	-					58Ni28 *
	ave.	491.6	1.3		0.4	1	92	85 $^{203}\text{Hg}$			average
$^{203}\text{Pb}(\epsilon)^{203}\text{Tl}$	940	50	975	6	0.7	U					55Ha.A *
	980	20			-0.3	1	10	10 $^{203}\text{Pb}$			65Le07 *
$^{203}\text{Bi}(\beta^+)^{203}\text{Pb}$	3260	50	3262	14	0.0	U					58No30 *
$^{203}\text{At}(\beta^+)^{203}\text{Po}$	5060	200	5148	14	0.4	U					87Se04
* $^{203}\text{Po}(\alpha)^{199}\text{Pb}$	$E_\alpha=5383.8(3,Z)$ to $4(4)$ level (this is level $x<9.3$ in Ensdf)										
* $^{203}\text{Hg}(\beta^-)^{203}\text{Tl}$	$E_{\beta^-}=210(2)$ $214(2)$ $214(3)$ respectively, to $3/2^+$ level at 279.1958 keV										
* $^{203}\text{Pb}(\epsilon)^{203}\text{Tl}$	$pK=0.36(0.07)$ $0.71(0.01)$ respectively, to $5/2^+$ level at 680.5164 keV										
* $^{203}\text{Bi}(\beta^+)^{203}\text{Pb}$	$E_{\beta^+}=1350(50)$ , $740(50)$ to levels around 840, 1550 keV										
$^{204}\text{Hg}-^{13}\text{C}$ $^{35}\text{Cl}_3$ $^{37}\text{Cl}_2$	131776.05	1.25	131775.9	0.6	0.0	U			H34	2.5	80Ko25
$^{204}\text{Hg}-^{169}\text{Tm}$ $^{35}\text{Cl}$	70420	100	70423.0	1.0	0.0	U			R04	4.0	64De15
$^{204}\text{Hg}-^{167}\text{Er}$ $^{37}\text{Cl}$	75430	60	75537.3	1.4	0.4	U			R04	4.0	64De15
$^{204}\text{Hg}-u$	-26505.8	0.6	-26506.0	0.5	-0.3	1	79	79 $^{204}\text{Hg}$	ST2	1.0	02Bf02
$^{204}\text{Po}-u$	-19689	30	-19690	12	0.0	R			GS2	1.0	05Li24
$^{204}\text{At}-u$	-12748	30	-12749	24	0.0	-			GS2	1.0	05Li24
	ave.	-12752	27		0.1	1	81	81 $^{204}\text{At}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{204}\text{Rn}-^{133}\text{Cs}_{1,534}$	136394	60	136480	8	1.4	U			SH1	1.0	13Dr04
$^{204}\text{Hg}-^{35}\text{Cl}_2-^{200}\text{Hg}-^{37}\text{Cl}_2$	11066.85	0.55	11067.3	0.7	0.3	1	24	12	$^{200}\text{Hg}$	2.5	80Ko25
$^{204}\text{Pb}-^{208}\text{Pb}_{,981}$	-4047	21	-4052.11	0.18	-0.2	U			MA6	1.0	01Sc41
$^{204}\text{Rn}-^{208}\text{Pb}_{,981}$	14358.1	9.4	14348	8	-1.1	-			SH1	1.0	13Dr04
	14303	25			1.8	-			SH1	1.0	13Dr04
	ave.	14351	9		-0.4	1	81	81	$^{204}\text{Rn}$		average
$^{204}\text{Hg}-^{35}\text{Cl}-^{202}\text{Hg}-^{37}\text{Cl}$	5807	2	5800.6	0.8	-0.8	U			H17	4.0	64Mc07
	5800.67	0.53			-0.1	1	35	26	$^{202}\text{Hg}$	2.5	80Ko25
$^{204}\text{Hg}-^{203}\text{Tl}$	1161	25	1150.0	1.3	-0.3	U			R08	1.5	69De19
$^{204}\text{Pb}(\alpha)^{200}\text{Hg}$	2650	100	1968.5	1.1	-6.8	B					58Ri23
$^{204}\text{Pb}(\alpha, ^8\text{He})^{200}\text{Pb}$	-28043	13	-28044	11	0.0	2			INS		90Ka10
$^{204}\text{Po}(\alpha)^{200}\text{Pb}$	5484.6	1.5	5484.9	1.4	0.2	3			DbA		69Go23 *
	5486.3	3.			-0.5	3					70Ra14
$^{204}\text{At}(\alpha)^{200}\text{Bi}$	6069.9	3.	6070.4	1.2	0.2	2					63Ho18 Z
	6066.2	3.			1.4	2					67Tr06 Z
	6071.3	3.			-0.3	2			Ora		75Ba.B
	6071.3	2.0			-0.4	2					79Sc.A
	6072.0	3.			-0.5	2			DbA		81Va27 Z
$^{204}\text{Rn}(\alpha)^{200}\text{Po}$	6544.3	3.	6546.7	1.8	0.8	-					67Va17 Z
	6547.5	2.5			-0.3	-			Lvn		93Wa04
	6537.4	7.			1.3	o			Ara		95Le04
	6548.6	5.0			-0.4	-			Ara		96Ta18
	ave.	6546.5	1.8		0.1	1	99	80	$^{200}\text{Po}$		average
$^{204}\text{Fr}(\alpha)^{200}\text{At}$	7170.4	5.	7170.3	2.4	0.0	4					67Va20 Z
	7169.4	5.			0.2	4					74Ho27 Z
	7170.6	5.			-0.1	4			Lvn		92Hu04 *
	7179.0	6.			-1.4	o			Jya		94Le05
	7167.8	7.			0.3	4			Ara		95Le04
	7173.9	6.			-0.6	o			Jya		05Uu02
	7171.9	5.			-0.3	4			Jya		13Ja06
$^{204}\text{Fr}^m(\alpha)^{200}\text{At}$	7218.8	8.	7221	4	0.2	o			Lvn		92Hu04
$^{204}\text{Fr}^m(\alpha)^{200}\text{At}^m$	7108.2	5.	7107.6	2.0	-0.1	4					74Ho27 Z
	7105.5	3.			0.7	4			Bka		82Bo04 Z
	7108.4	5.			-0.2	4			Lvn		92Hu04 *
	7115.6	7.			-1.1	o			Jya		94Le05 *
	7114.7	7.			-1.0	4			Ara		95Le04
	7117.7	6.			-1.7	o			Jya		05Uu02 *
	7108.7	5.			-0.2	4			Jya		13Ja06
$^{204}\text{Fr}^n(\alpha)^{200}\text{At}^n$	7157.6	6.1	7152.8	2.1	-0.8	o			Jya		05Uu02
	7153.5	5.			-0.1	o			Jya		13Ja06
$^{204}\text{Ra}(\alpha)^{200}\text{Rn}$	7638.1	12.	7637	7	-0.1	5			Ara		95Le04
	7638.1	25.			-0.1	o			Jya		95Le15
	7634.0	10.			0.3	o			Jya		96Le09
	7636.1	8.			0.1	5			Jya		05Uu02
	7638.1	25.			-0.1	U			Anv		10He25
$^{204}\text{Pb}(p,t)^{202}\text{Pb}$	-6835	10	-6830	4	0.5	1	15	14	$^{202}\text{Pb}$		Yal
$^{204}\text{Hg}(t,\alpha)^{203}\text{Au}$	10962	15	10978	3	1.1	U			LAl		81FI05
$^{204}\text{Hg}(d, ^3\text{He})^{203}\text{Au}-^{206}\text{Pb}(\gamma)^{205}\text{Tl}$	-1582.0	3.0	-1582.0	3.0	0.0	1	100	100	$^{203}\text{Au}$		94Gr07
$^{204}\text{Hg}(d,t)^{203}\text{Hg}$	-1242	5	-1235.0	1.6	1.4	1	11	10	$^{203}\text{Hg}$		Ald
$^{203}\text{Tl}(n,\gamma)^{204}\text{Tl}$	6656.0	0.3	6656.08	0.29	0.3	1	94	65	$^{203}\text{Tl}$		MMn
	6654.88	0.14			8.6	C			Bdn		06Fi.A
$^{204}\text{Pb}(p,d)^{203}\text{Pb}$	-6165	10	-6170	6	-0.5	-			Yal		71Ki01
$^{204}\text{Pb}(d,t)^{203}\text{Pb}$	-2160	20	-2137	6	1.1	-			Ald		67Bj01
$^{204}\text{Pb}(p,d)^{203}\text{Pb}$	ave.	-6171	9	-6170	6	0.0	1	52	$^{203}\text{Pb}$		average
$^{204}\text{Au}(\beta^-)^{204}\text{Hg}$	4500	300	4040#	200#	-1.5	F					67Wa23 *
$^{204}\text{Tl}(\epsilon)^{204}\text{Hg}$	314	20	344.0	1.2	1.5	U					64Ch17
	332	20			0.6	U					66Kl02
	385	20			-2.0	U					73La17
$^{204}\text{Tl}(\beta^-)^{204}\text{Pb}$	764.24	0.31	763.75	0.18	-1.6	-					67Pa08
	763.47	0.22			1.3	-					68Wo02
	ave.	763.73	0.18		0.1	1	97	68	$^{204}\text{Tl}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{204}\text{At}(\beta^+)^{204}\text{Po}$	6220	160	6466	25	1.5	U					86Ve.B *
$^{204}\text{Fr}^n(\text{IT})^{204}\text{Fr}^m$	276.1	0.5				5					95Bi.A
$^{204}\text{Po}(\alpha)^{200}\text{Pb}$	Printing error in reference: $^{204}\text{Po}$ not $^{206}\text{Po}$ ; Z recalibrated										
$^{204}\text{Fr}(\alpha)^{200}\text{At}$	$E_\alpha=7031(5)$ , $6916(8)$ to ground state, 113 level										
$^{204}\text{Fr}^m(\alpha)^{200}\text{At}^m$	$E_\alpha=6969(5)$ ; and $7013(5)$ from $^{204}\text{Fr}^n$ 276.1 above $^{204}\text{Fr}^m$ to $^{200}\text{At}^n$										
*	230.9 above $^{200}\text{At}^m$										
$^{204}\text{Fr}^m(\alpha)^{200}\text{At}^m$	$E_\alpha=7020(7)$ from $^{204}\text{Fr}^n$ 276.1 above $\text{Fr}^m$ to $^{200}\text{At}^n$ 230.9 above $^{200}\text{At}^m$										
$^{204}\text{Fr}^m(\alpha)^{200}\text{At}^m$	$E_\alpha=6976(6)$ ; and $7017(6)$ from $^{204}\text{Fr}^n$ 276.1 above $^{204}\text{Fr}^m$ to $^{200}\text{At}^n$										
$^{204}\text{Au}(\beta^-)^{204}\text{Hg}$	F : reported 4 s activity does not exist										
$^{204}\text{At}(\beta^+)^{204}\text{Po}$	$E_{\beta^+}=2950(160)$ to $8^+$ level at 2248.17 keV										
$\text{C}_{16}\text{H}_{13}-^{205}\text{Tl}$	127345	29	127298.2	1.3	-1.1	U			R08	1.5	69De19
$\text{C}_{14}\text{N}_2\text{H}_9-^{205}\text{Tl}$	102091	36	102146.1	1.3	1.0	U			R08	1.5	69De19
$^{205}\text{Tl}-^{168}\text{Er}^{37}\text{Cl}$	76198	44	76148.5	1.8	-0.8	U			R08	1.5	69De19
$^{205}\text{Tl}-^{170}\text{Er}^{35}\text{Cl}$	70034	23	70103.9	2.1	2.0	U			R08	1.5	69De19
$^{205}\text{Tl}-^{133}\text{Cs}_{1.541}$	120129	11	120125.8	1.3	-0.3	U			MA8	1.0	08We02
$^{205}\text{Bi}-u$	-22559	30	-22614	5	-1.8	U			GS2	1.0	05Li24
$^{205}\text{Po}-u$	-18773	30	-18810	11	-1.2	-			GS2	1.0	05Li24
	ave.	-18790	25		-0.8	1	19	19 $^{205}\text{Po}$			average
$^{205}\text{Rn}-^{133}\text{Cs}_{1.541}$	137456	50	137422	5	-0.7	U			SH1	1.0	13Dr04
		137458	29		-1.3	U			SH1	1.0	13Dr04
$^{205}\text{Fr}-^{133}\text{Cs}_{1.541}$	144293.8	9.7	144292	8	-0.1	2			MA8	1.0	08We02
$^{205}\text{Rn}-^{208}\text{Pb}_{986}$	14748	11	14744	6	-0.3	-			SH1	1.0	13Dr04
		14772	20		-1.4	-			SH1	1.0	13Dr04
	ave.	14754	10		-1.0	1	33	32 $^{205}\text{Rn}$			average
$^{205}\text{Tl}^{35}\text{Cl}-^{203}\text{Tl}^{37}\text{Cl}$	5040	4	5033.3	0.6	-0.4	U			H17	4.0	64Mc07
		5031.43	1.07		0.7	-			H36	2.5	85De40
		5032.88	1.01		0.3	-			H42	1.5	93Si05
	ave.	5032.5	1.3		0.6	1	19	15 $^{205}\text{Tl}$			average
$^{205}\text{Po}-^{205}\text{Fr}$	-18450	2050	-17404	14	0.5	U			RI1	1.0	16Sc.A
$^{205}\text{At}-^{205}\text{Fr}$	-12420	450	-12520	18	-0.2	U			RI1	1.0	16Sc.A
$^{205}\text{Rn}-^{205}\text{Fr}$	-6640	330	-6871	10	-0.7	U			RI1	1.0	16Sc.A
$^{205}\text{Tl}-^{167}\text{Er}^{37}\text{Cl}$	76426	47	76470.5	1.8	0.6	U			R08	1.5	69De19
$^{205}\text{Tl}-^{169}\text{Tm}^{35}\text{Cl}$	71355	25	71356.2	1.6	0.0	U			R08	1.5	69De19
$^{169}\text{Tm}^{37}\text{Cl}-^{205}\text{Tl}$	-74316	32	-74306.3	1.6	0.2	U			R08	1.5	69De19
$^{205}\text{Tl}-^{204}\text{Hg}$	938	27	933.2	1.4	-0.1	U			R08	1.5	69De19
$^{205}\text{Tl}-^{203}\text{Tl}$	2092	20	2083.2	0.6	-0.3	U			R08	1.5	69De19
$^{205}\text{Po}(\alpha)^{201}\text{Pb}$	5324.1	10.	5325	10	0.1	1	95	90 $^{201}\text{Pb}$			67Ti04
$^{205}\text{At}(\alpha)^{201}\text{Bi}$	6016.3	4.	6019.6	1.7	0.8	4					63Ho18 Z
	6020.5	2.			-0.5	4			DbA		68Go.B Z
	6018.9	5.			0.1	4					74Ho27 Z
$^{205}\text{Rn}(\alpha)^{201}\text{Po}$	6386.6	3.	6386.5	1.8	0.0	-					67Va17 Z
	6386.6	6.			0.0	-					71Ho01 Z
	6385.7	2.5			0.3	-			Lvn		93Wa04
	ave.	6386.1	1.9		0.2	1	97	68 $^{205}\text{Rn}$			average
$^{205}\text{Fr}(\alpha)^{201}\text{At}$	7056.5	5.	7054.7	2.4	-0.3	3					67Va20 Z
	7052.2	5.			0.5	3					74Ho27 Z
	7057.3	5.			-0.5	3			ORa		81Ri04 Z
	7052.9	7.			0.3	3			Ara		95Le04
	7053.9	5.			0.2	3			Anv		05De01
$^{205}\text{Ra}(\alpha)^{201}\text{Rn}$	7506.7	20.	7490	50	-0.4	F			GSa		87He10 *
	7496.6	25.			-0.2	o			Jya		95Le15
	7486.4	20.				5			Jya		96Le09
$^{205}\text{Ra}^m(\alpha)^{201}\text{Rn}^m$	7501.7	10.	7505	9	0.3	6			Ara		95Le04
	7522.1	25.			-0.7	o			Jya		95Le15
	7517.0	20.			-0.6	6			Jya		96Le09
	7526.1	30.			-0.7	U			Anv		10He25

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{205}\text{Ac}(\alpha)^{201}\text{Fr}$	8093.2	30.6				3			Lza		14Zh03
$^{204}\text{Hg}(\text{d,p})^{205}\text{Hg}$	3443	5	3444	4	0.3	1	52	$^{205}\text{Hg}$	Ald		70An14
$^{205}\text{Tl}(\gamma,\text{n})^{204}\text{Tl}$	-7515	29	-7546.0	0.5	-1.1	U			Phi		60Ge01
	-7548	3			0.7	U			McM		79Ba06
$^{205}\text{Tl}(\text{d,t})^{204}\text{Tl}$	-1288.7	0.6	-1288.8	0.5	-0.2	1	64	$^{205}\text{Tl}$	Mun		90Li40
$^{204}\text{Pb}(\text{n},\gamma)^{205}\text{Pb}$	6731.53	0.15	6731.66	0.11	0.9	-			ILn		83Hu13 Z
	6731.80	0.16			-0.9	-			Bdn		06Fi.A
$^{204}\text{Pb}(\text{d,p})^{205}\text{Pb}$	4516	20	4507.10	0.11	-0.4	U			Ald		67Bj01
$^{204}\text{Pb}(\text{n},\gamma)^{205}\text{Pb}$	ave.	6731.66	0.11	6731.66	0.11	0.1	1	99	$^{204}\text{Pb}$		average
$^{205}\text{Hg}(\beta^-)^{205}\text{Tl}$	1620	200	1533	4	-0.4	U					40Kr08
	1750	200			-1.1	U					51Ly10
$^{205}\text{Pb}(\varepsilon)^{205}\text{Tl}$	41.4	1.1	50.6	0.5	8.4	B					78Pe08
$^{205}\text{Bi}(\beta^+)^{205}\text{Pb}$	2701.4	10.	2706	5	0.4	-					62Bo25 *
	2715.4	10.			-1.0	-					62Pe08 *
	ave.	2708	7		-0.4	1	52	51 $^{205}\text{Bi}$			average
$^{205}\text{Po}(\beta^+)^{205}\text{Bi}$	3390	150	3543	11	1.0	U					69Ho37 *
$^{205}\text{Ra}(\alpha)^{201}\text{Rn}$	F : possibly mixed with $^{205}\text{Ra}^m(\alpha)^{201}\text{Rn}^m$										
$^{205}\text{Bi}(\beta^+)^{205}\text{Pb}$	$E_{\beta^+}=976(10) 990(10)$ respectively, to $7/2^-$ level at 703.427 keV										
$^{205}\text{Po}(\beta^+)^{205}\text{Bi}$	$p^+=3(1)\times 10^{-3}$ to $7/2^-$ level at 849.84 and $7/2^-$ at 1001.22 keV										
											87He10 **
											Ens044 **
											Ens044 **
$^{206}\text{Bi}-\text{u}$	-21429	30	-21501	8	-2.4	U			GS2	1.0	05Li24
$^{206}\text{Po}-\text{u}$	-19471	30	-19526	4	-1.8	U			GS2	1.0	05Li24
$^{206}\text{At}-\text{u}$	-13305	30	-13344	16	-1.3	1	29	29 $^{206}\text{At}$	GS2	1.0	05Li24
$^{206}\text{Rn}-^{133}\text{Cs}_{1,549}$	136641	15	136650	9	0.6	1	38	38 $^{206}\text{Rn}$	SH1	1.0	13Dr04
$^{206}\text{Pb }^{35}\text{Cl}_2-^{202}\text{Hg }^{37}\text{Cl}_2$	9722.09	0.57	9721.8	1.1	-0.2	1	62	54 $^{206}\text{Pb}$	H36	2.5	85De40
$^{206}\text{Pb }^{35}\text{Cl}-^{204}\text{Hg }^{37}\text{Cl}$	3929	4	3921.2	1.3	-0.5	U			H17	4.0	64Mc07
$^{206}\text{Pb }^{35}\text{Cl}-^{204}\text{Pb }^{37}\text{Cl}$	4378	3	4371.81	0.15	-0.5	U			H17	4.0	64Mc07
	4370.72	1.17			0.4	U			H36	2.5	85De40
	4371.29	0.81			0.4	U			H42	1.5	93Si05
$^{206}\text{Rn}-^{208}\text{Pb}_{990}$	13307	15	13310	9	0.2	1	38	37 $^{206}\text{Rn}$	SH1	1.0	13Dr04
$^{206}\text{At}-^{205}\text{Fr}_{1,005}$	-12110	2260	-11931	18	0.1	U			RI1	1.0	16Sc.A
$^{206}\text{Rn}-^{205}\text{Fr}_{1,005}$	-7800	620	-8391	13	-1.0	U			RI1	1.0	16Sc.A
$^{206}\text{Fr}-^{205}\text{Fr}_{1,005}$	130	140	80	30	-0.4	U			RI1	1.0	16Sc.A *
$^{206}\text{Po}(\alpha)^{202}\text{Pb}$	5327.4	4.	5327.0	1.3	-0.1	2					67Ti04 Z
	5327.4	1.5			-0.3	2			DbA		69Go23 *
	5325.1	3.			0.6	2					70Ra14 Z
$^{206}\text{At}(\alpha)^{202}\text{Bi}$	5884.4	3.6	5887	5	0.7	o			DbA		68Go.B *
	5886.4	5.			0.1	1	98	70 $^{202}\text{Bi}$	DbA		81Va27 *
$^{206}\text{Rn}(\alpha)^{202}\text{Po}$	6381.8	3.	6383.7	1.6	0.6	-					67Va17 Z
	6384.6	3.			-0.3	-			DbA		71Go35 Z
	6384.8	2.5			-0.4	-			Lvn		93Wa04
	ave.	6383.9	1.6		-0.1	1	99	75 $^{202}\text{Po}$			average
$^{206}\text{Fr}(\alpha)^{202}\text{At}$	6925.9	7.	6923	4	-0.4	4					67Va20 *
	6918.9	7.			0.6	4					74Ho27 *
	6924.0	7.			-0.1	4			ORa		81Ri04 *
	6924.8	7.			-0.2	4			Lvn		92Hu04 *
$^{206}\text{Fr}^n(\alpha)^{202}\text{At}^n$	7068.8	5.	7068	4	-0.2	6			ORa		81Ri04 Z
	7067.1	5.			0.2	6			Lvn		92Hu04 *
$^{206}\text{Ra}(\alpha)^{202}\text{Rn}$	7416.3	5.	7415	4	-0.2	3					67Va22 Z
	7414.3	10.			0.1	3			GSa		87He10
	7412.2	10.			0.3	o			Jya		95Le15
	7406	15			0.6	o			Jya		95Uu01
	7412.2	10.			0.3	3			Jya		96Le09
$^{206}\text{Ac}(\alpha)^{202}\text{Fr}$	7944.6	30.	7960	50	0.3	4			Jya		98Es02
	7972.0	30.			-0.3	4			Lza		14Zh03

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{206}\text{Ac}^m(\alpha)^{202}\text{Fr}^m$	7903.8	30.				6			Jya		98Es02
$^{204}\text{Pb}(t,p)^{206}\text{Pb}$	6322	20	6336.53	0.12	0.7	U			Ald		67Ha.A
$^{204}\text{Pb}(\alpha,d)^{206}\text{Bi}$	-15798.	11.5	-15792	8	0.5	R			Pit		76Da20
$^{205}\text{Tl}(n,\gamma)^{206}\text{Tl}$	6503.7	0.4	6503.8	0.4	0.3	1	93	84 $^{206}\text{Tl}$	MMn		74Co21 Z
	6502.87	0.27			3.5	C			Bdn		06Fi.A
$^{205}\text{Tl}(d,p)^{206}\text{Tl}$	4276	5	4279.2	0.4	0.6	U			ANL		65Er02
$^{205}\text{Tl}(^3\text{He},d)^{206}\text{Pb}$	1761.7	1.4	1760.2	0.5	-1.1	1	13	12 $^{205}\text{Tl}$	Mun		90Li40
$^{205}\text{Pb}(n,\gamma)^{206}\text{Pb}$	8086.66	0.06	8086.66	0.06	0.0	1	100	69 $^{205}\text{Pb}$			96Ra16 Z
$^{206}\text{Pb}(\gamma,n)^{205}\text{Pb}$	-8090	70	-8086.66	0.06	0.0	U			Phi		60Ge01
	-8087	3			0.1	U			McM		79Ba06
$^{206}\text{Pb}(d,t)^{205}\text{Pb}$	-1830	100	-1829.43	0.06	0.0	U			MIT		53Ha66
	-1831.2	0.5			3.5	B			Mun		90Li40
$^{206}\text{Hg}(\beta^-)^{206}\text{Tl}$	1240	62	1308	20	1.1	U					68Wo09 *
$^{206}\text{Tl}(\beta^-)^{206}\text{Pb}$	1534	5	1532.2	0.6	-0.4	U					71Pe23
	1527	4			1.3	U					72Wi18
$^{206}\text{Bi}(\beta^+)^{206}\text{Pb}$	3683	33	3757	8	2.3	U					62Pe08 *
$^{206}\text{Bi}(\epsilon)^{206}\text{Pb}$	3753	10			0.4	2					74Go20 *
$^{206}\text{At}(\beta^+)^{206}\text{Po}$	5687	150	5759	16	0.5	U					77Li16 *
$^{206}\text{Fr}^n(\text{IT})^{206}\text{Fr}^m$	531	2				7			ORa		81Ri04
$^{206}\text{Fr}^x(\text{IT})^{206}\text{Fr}$	100	100				5					AHW *
$^{*206}\text{Fr}-^{205}\text{Fr}_{1.005}$	$D_M=230(110) \mu\text{u}$ for mixture gs+m at 190(40) keV; $M-A=-1100(110)$ keV										
$^{*206}\text{Po}(\alpha)^{202}\text{Pb}$	Printing error in reference: $^{206}\text{Po}$ not $^{211}\text{Po}$ ; Z recalibrated										
$^{*206}\text{At}(\alpha)^{202}\text{Bi}$	$E_\alpha=5702.8(2,Z)$ to $(5)^+$ level at 68(3) keV										
$^{*206}\text{At}(\alpha)^{202}\text{Bi}$	$E_\alpha=5773.8(5,Z)$ , 5702.8(5,Z) to ground state and $(5)^+$ level at 68(3) keV										
$^{*206}\text{Fr}(\alpha)^{202}\text{At}$	$E_\alpha=6793.1(5,Z)$ ; correction -2 for being a doublet										
$^{*206}\text{Fr}(\alpha)^{202}\text{At}$	$E_\alpha=6786.3(5,Z)$ ; correction -2 for being a doublet										
$^{*206}\text{Fr}(\alpha)^{202}\text{At}$	$E_\alpha=6791.3(5,Z)$ ; correction -2 for being a doublet										
$^{*206}\text{Fr}(\alpha)^{202}\text{At}$	$E_\alpha=6792(5)$ ; correction -2 for being a doublet										
$^{*206}\text{Fr}^n(\alpha)^{202}\text{At}^n$	$E_\alpha=6930(5)$ and 6792(7) combined with $E(\gamma)$ 's 531, 391.7 keV										
$^{*206}\text{Hg}(\beta^-)^{206}\text{Tl}$	$E_{\beta^-}=935(62)$ to $1^-$ level at 304.896 keV										
$^{*206}\text{Bi}(\beta^+)^{206}\text{Pb}$	$E_{\beta^+}=977(33)$ to $4^+$ level at 1683.99 keV										
$^{*206}\text{Bi}(\epsilon)^{206}\text{Pb}$	LK=0.509(0.015) to $5^-$ level at 3562.92 keV, original error 22, recalculated										
$^{*206}\text{At}(\beta^+)^{206}\text{Po}$	$E_{\beta^+}=3092(150)$ to $6^+$ level at 1573.38 keV										
$^{*206}\text{Fr}^x(\text{IT})^{206}\text{Fr}$	Assuming a 0.15(0.20)% isomeric mixture										
	AHW **										
$^{207}\text{Hg}-u$	-17721	33	-17700	30	0.6	o			GS3	1.0	08Ch.A
	-17700	32				2			GS3	1.0	12Ch19
$^{207}\text{Rn}-^{133}\text{Cs}_{1.556}$	137794	28	137847	9	1.9	U			SH1	1.0	13Dr04
$^{207}\text{Fr}-^{133}\text{Cs}_{1.556}$	144062	20	144063	19	0.1	1	88	88 $^{207}\text{Fr}$	MA8	1.0	14Bo26
$^{207}\text{Pb }^{35}\text{Cl}-^{205}\text{Tl }^{37}\text{Cl}$	4413	4	4419.6	0.6	0.4	U			H17	4.0	64Mc07
	4415.60	2.40			0.7	U			H36	2.5	85De40
	4417.32	1.40			1.1	U			H42	1.5	93Si05
$^{206}\text{Pb} H-^{207}\text{Pb}$	6394.2	1.1	6393.42	0.10	-0.3	U			C4	2.5	71Ke02
$^{206}\text{Fr}^x-^{207}\text{Fr}_{.498} \ ^{205}\text{Fr}_{.502}$	930	90	930	100	0.0	U			P24	2.5	82Au01
$^{207}\text{Po}(\alpha)^{203}\text{Pb}$	5216.0	2.5	5215.9	2.5	0.0	1	96	59 $^{207}\text{Po}$	DbA		70Af.A
$^{207}\text{At}(\alpha)^{203}\text{Bi}$	5872.5	3.				3			DbA		69Go23 Z
$^{207}\text{Rn}(\alpha)^{203}\text{Po}$	6256.3	3.	6251.2	1.6	-1.6	4					67Va20 Z
	6247.3	3.			1.3	4			DbA		71Go35 Z
	6250.4	2.5			0.3	4			Lvn		93Wa04
$^{207}\text{Fr}(\alpha)^{203}\text{At}$	6907.8	5.	6893	20	-0.3	-					67Va20 Z
	6895.8	5.			-0.1	-					74Ho27 Z
	6900.9	5.			-0.2	-			ORa		81Ri04 Z
	ave.	2.9			-0.2	1	16	12 $^{207}\text{Fr}$			average
$^{207}\text{Ra}(\alpha)^{203}\text{Rn}$	7273.8	5.	7270	50	0.0	4					67Va22 Z
	7268.7	10.			0.1	4			GSa		87He10
	7276.9	12.2			-0.1	4			Jya		95Uu01
$^{207}\text{Ra}^m(\alpha)^{203}\text{Rn}^m$	7464.5	10.2	7468	8	0.3	2			GSa		87He10
	7474.7	15.			-0.4	o			Jya		95Le15
	7475.7	15.			-0.5	2			Jya		96Le09

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{207}\text{Ac}(\alpha)^{203}\text{Fr}$	7864.3	25.	7840	50	-0.4	o			Jya		94Le05
	7844.9	25.				2			Jya		98Es02
$^{205}\text{Tl}(t,p)^{207}\text{Tl}$	4880	15	4874	5	-0.4	1	13	13 $^{207}\text{Tl}$	Ald		69Ha11
$^{207}\text{Pb}(t,\alpha)^{206}\text{Tl}$	12321	25	12326.2	0.6	0.2	U			Ald		67Ha.A
$^{206}\text{Pb}(n,\gamma)^{207}\text{Pb}$	6737.85	0.15	6737.78	0.10	-0.5	-			MMn		81Ke11 Z
	6737.72	0.18			0.3	-			ILn		83Hu13 Z
	6737.74	0.17			0.2	-			Bdn		06Fi.A
$^{207}\text{Pb}(\gamma,n)^{206}\text{Pb}$	-6742	3	-6737.78	0.10	1.4	U			McM		79Ba06
$^{206}\text{Pb}(d,p)^{207}\text{Pb}$	4480	30	4513.21	0.10	1.1	U			MIT		53Ha66
	4510	20			0.2	U					58Mc64
	4526	30			-0.4	U			Pit		64Co11
$^{206}\text{Pb}(n,\gamma)^{207}\text{Pb}$	ave.	6737.78	0.10	6737.78	0.10	0.0	1	100	87 $^{207}\text{Pb}$		average
$^{207}\text{Hg}(\beta^-)^{207}\text{Tl}$	4815	150	4550	30	-1.8	U					81Jo.B *
$^{207}\text{Tl}(\beta^-)^{207}\text{Pb}$	1431	8	1418	5	-1.7	1	46	45 $^{207}\text{Tl}$			67Da10
$^{207}\text{Bi}(\epsilon)^{207}\text{Pb}$	2392	10	2397.4	2.1	0.5	U					Averag *
$^{207}\text{Po}(\beta^+)^{207}\text{Bi}$	2907	10	2909	7	0.2	1	44	41 $^{207}\text{Po}$			58Ar56 *
$^{207}\text{Rn}(\beta^+)^{207}\text{At}$	4617	70	4593	15	-0.3	U					75Ze.A *
$^{207}\text{Hg}(\beta^-)^{207}\text{Tl}$	$E_{\beta^-}=1800(150)$ , 14%, 32%, 16%, 7% to $(7/2^-, 9/2)$ level at 2911.83 keV										
	$(9/2)^+$ at 2985.23 and 3104.43, $(7/2^-, 9/2, 11/2)$ at 3143.1 keV										
*	Average $pL=0.61(0.05)$ to $7/2^-$ level at 2339.921 keV from two references:										
*	$pL=0.663(0.014)$										
*	$pL=0.56(0.04)$ ; original error 0.08 is $2\sigma$										
$^{207}\text{Po}(\beta^+)^{207}\text{Bi}$	$E_{\beta^+}=893(10)$ to $7/2^-$ level at 992.43 keV, and other $E_{\beta^+}$										
$^{207}\text{Rn}(\beta^+)^{207}\text{At}$	$E_{\beta^+}=3250(70)$ to $7/2^-$ level at 344.55 keV										
											81Jo.B **
											Ens112 **
*	Average $pL=0.61(0.05)$ to $7/2^-$ level at 2339.921 keV from two references:										
*	$pL=0.663(0.014)$										
*	$pL=0.56(0.04)$ ; original error 0.08 is $2\sigma$										
$^{207}\text{Po}(\beta^+)^{207}\text{Bi}$	$E_{\beta^+}=893(10)$ to $7/2^-$ level at 992.43 keV, and other $E_{\beta^+}$										
$^{207}\text{Rn}(\beta^+)^{207}\text{At}$	$E_{\beta^+}=3250(70)$ to $7/2^-$ level at 344.55 keV										
											Ens112 **
$^{208}\text{Hg}-u$	-14241	33	-14240	30	0.0	o			GS3	1.0	08Ch.A
	-14241	33				2			GS3	1.0	09Ch08
$^{208}\text{Pb}-^{133}\text{Cs}_{1.564}$	124532.0	5.6	124525.1	1.2	-1.2	U			MA8	1.0	08We02
	124524.3	5.5			0.1	U			MA8	1.0	14Bo26
$^{208}\text{Po}-u$	-18710	31	-18754.4	1.9	-1.4	U			GS2	1.0	05Li24
$^{208}\text{Fr}-^{133}\text{Cs}_{1.564}$	144984	20	145011	13	1.4	o			MA8	1.0	12Bo.A
	145030	16			-1.2	o			MA8	1.0	12Bo.A
	145012	13			-0.1	1	95	95 $^{208}\text{Fr}$	MA8	1.0	14Bo26
$^{208}\text{Pb } ^{35}\text{Cl}-^{206}\text{Pb } ^{37}\text{Cl}$	5136	2	5136.90	0.14	0.1	U			H17	4.0	64Mc07
	5136.23	1.08			0.2	U			H36	2.5	85De40
	5136.93	0.41			0.0	U			H42	1.5	93Si05
$^{207}\text{Fr}-^{208}\text{Fr}_{498} \text{ } ^{206}\text{Fr}_{502}$	-890	60	-940	60	-0.4	U			P24	2.5	82Au01
$^{208}\text{Po}(\alpha)^{204}\text{Pb}$	5216.3	2.	5215.4	1.3	-0.5	2			DbA		69Go23 Z
	5214.0	3.			0.5	2					70Ra14 Z
	5215.1	2.			0.1	2					89Ma05
$^{208}\text{At}(\alpha)^{204}\text{Bi}$	5750.6	3.	5751.1	2.2	0.2	3			DbA		69Go23 Z
	5751.6	3.			-0.2	3			DbA		81Va27 Z
$^{208}\text{Rn}(\alpha)^{204}\text{Po}$	6269.3	4.	6260.7	1.7	-2.1	4					55Mo69 Z
	6260.0	3.			0.2	4			DbA		71Go35 Z
	6257.5	5.			0.6	4					74Ho27
	6258.7	2.5			0.8	4			Lvn		93Wa04
$^{208}\text{Fr}(\alpha)^{204}\text{At}$	6778.3	5.	6785	24	0.1	-					67Va20 Z
	6767.7	5.			0.3	-					74Ho27 Z
	6767.7	5.			0.3	-			ORa		81Ri04 Z
	6739.8	51.0			0.9	-			GSa		15De22
	ave.	6771.1	2.9		0.3	1	23	19 $^{204}\text{At}$			average
$^{208}\text{Ra}(\alpha)^{204}\text{Rn}$	7273.1	5.				2					67Va22 Z
$^{208}\text{Ac}(\alpha)^{204}\text{Fr}$	7720.8	15.	7720	50	0.0	5			Jya		94Le05
	7769.7	40.			-1.0	5			JAA		96Ik01
	7707.5	21.4			0.3	5			Lza		14Ya19
$^{208}\text{Ac}^m(\alpha)^{204}\text{Fr}^m$	7892.1	20.	7899	14	0.3	6			Dbb		94An01
	7910.4	20.			-0.6	6			Jya		94Le05
	7871.7	50.			0.5	6			JAA		96Ik01



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{208}\text{Th}(\alpha)^{204}\text{Ra}$	8202.0	30.				6			Anv		10He25
$^{206}\text{Pb}(t,p)^{208}\text{Pb}$	5622	30	5623.85	0.11	0.1	U			Ald		67Ha.A
$^{207}\text{Pb}(n,\gamma)^{208}\text{Pb}$	7367.95	0.15	7367.87	0.05	-0.5	-			MMn		81Ke11 Z
	7367.96	0.10			-0.9	-					81Su.A Z
	7367.81	0.11			0.5	-			ILn		83Hu13 Z
	7367.774	0.098			1.0	-					98Be19 Z
	7367.92	0.16			-0.3	-			Bdn		06Fi.A
$^{208}\text{Pb}(\gamma,n)^{207}\text{Pb}$	-7370	3	-7367.87	0.05	0.7	U			McM		79Ba06
$^{208}\text{Pb}(d,t)^{207}\text{Pb}$	-1114	25	-1110.64	0.05	0.1	U			Pit		64Co11
$^{207}\text{Pb}(n,\gamma)^{208}\text{Pb}$	ave.	7367.87	0.05	7367.87	0.05	0.0	1	100	87 $^{208}\text{Pb}$		average
$^{208}\text{Tl}(\beta^-)^{208}\text{Pb}$	4989.7	7.	4998.5	1.7	1.3	U					48Ma29 *
	4997.7	10.			0.1	U					54El24 *
$^{208}\text{Bi}(\epsilon)^{208}\text{Pb}$	2810	4	2878.4	2.0	17.1	B					59Mi19 **
$^{208}\text{Tl}(\beta^-)^{208}\text{Pb}$	$E_{\beta^-} = 1792(7) 1800(10)$ respectively, to $5^-$ level at 3197.711 keV										Ens077 **
$^{208}\text{Bi}(\epsilon)^{208}\text{Pb}$	$pK=0.24(0.01)$ to $3^-$ level at 2614.522 keV, recalculated										Ens077 **
$^{209}\text{Bi}-^{133}\text{Cs}_{1.571}$	128937.6	4.7	128933.5	1.5	-0.9	U			MA8	1.0	08We02
$^{209}\text{Fr}-^{226}\text{Ra}_{925}$	-27584	36	-27550	16	1.0	2			MA3	1.0	92Bo28
$^{209}\text{Bi } ^{35}\text{Cl}-^{207}\text{Pb } ^{37}\text{Cl}$	7444	3	7451.9	0.8	0.7	U			H17	4.0	64Mc07
	7454.13	1.51			-0.6	U			H36	2.5	85De40
$^{208}\text{Fr}-^{209}\text{Fr}_{498} \ ^{207}\text{Fr}_{502}$	720	60	639	16	-0.5	U			P24	2.5	82Au01
$^{209}\text{Bi}(\alpha)^{205}\text{Tl}$	3137.0	2.2	3137.3	0.8	0.1	1	12	10 $^{209}\text{Bi}$			03De11
$^{209}\text{Po}(\alpha)^{205}\text{Pb}$	4974	5	4979.2	1.4	1.0	2					66Ha29 *
	4980.0	2.			-0.4	2			DbA		69Go23 *
	4979.3	2.			0.0	2					89Ma05 *
$^{209}\text{At}(\alpha)^{205}\text{Bi}$	5757.2	2.	5757.0	2.0	-0.1	1	96	49 $^{205}\text{Bi}$	DbA		69Go23 Z
$^{209}\text{Rn}(\alpha)^{205}\text{Po}$	6157.5	3.	6155.4	2.0	-0.7	-			DbA		71Go35 Z
	6154.2	2.5			0.5	-			Lvn		93Wa04
	ave.	6155.5	2.0			-0.1	1	99	75 $^{205}\text{Po}$		average
$^{209}\text{Fr}(\alpha)^{205}\text{At}$	6777.7	5.	6777	4	0.0	3					67Va20 Z
	6777.3	5.			0.0	3					74Ho27 Z
$^{209}\text{Ra}(\alpha)^{205}\text{Rn}$	7147.0	5.	7143.1	2.7	-0.8	2					67Va22 Z
	7141	5			0.4	2			GSa		03He06 *
	7142.0	4.			0.3	2					08Ha12
	7733.3	15.	7730	50	-0.1	3					68Va04
	7738.4	20.			-0.2	3			Dbb		94An01
$^{209}\text{Ac}(\alpha)^{205}\text{Fr}$	7729.2	15.			0.0	3			Jya		94Le05
	7728.2	40.			0.0	U			JAA		96Ik01
	7725.1	10.			0.1	3			GSa		00He17
	7723.0	23.			0.1	3			Lza		14Ya19
	8238.0	50.	8273	23	0.7	7			JAA		96Ik01 *
	8281.8	25.			-0.3	7			Anv		10He25 *
	2814	12	2823.4	1.3	0.8	U			Ald		68Bj03
$^{209}\text{Bi}(p,t)^{207}\text{Bi}$	-5864.8	2.0	-5864.9	2.0	0.0	1	98	97 $^{207}\text{Bi}$	MSU		76Be.B *
$^{208}\text{Pb}(d,p)^{209}\text{Pb}$	1705	15	1712.8	1.3	0.5	U			MIT		64Sp12
	1700	10			1.3	U					67Mu16
	1718	4			-1.3	1	11	11 $^{209}\text{Pb}$	Pit		72Ko03 *
	1715	10			-0.2	U			Yal		74Ko20
	16003	25	16014.8	0.8	0.5	U			Ald		68Bj01
$^{209}\text{Bi}(\gamma,n)^{208}\text{Bi}$	-7432	10	-7459.8	1.9	-2.8	U			Phi		60Ge01
$^{209}\text{Bi}(d,t)^{208}\text{Bi}$	-7460	2			0.1	2			McM		79Ba06
	-1216	30	-1202.5	1.9	0.4	U			Pit		64Co11
	-1201	5			-0.3	2			ANL		64Er06
$^{209}\text{Pb}(\beta^-)^{209}\text{Bi}$	644.6	1.2	644.0	1.1	-0.5	1	91	87 $^{209}\text{Pb}$			72Be44
$^{209}\text{Rn}(\beta^+)^{209}\text{At}$	3928	40	3942	11	0.3	R					74Vy01 *
$^{209}\text{Po}(\alpha)^{205}\text{Pb}$	$E_{\alpha}=4876.8(5.Z) 4882.8(2.Z)$ respectively, to (20% ground state + 80% $^{205}\text{Pb}^m$ at 2.329 keV)										Ens044 **
$^{209}\text{Po}(\alpha)^{205}\text{Pb}$	$E_{\alpha}=4882.6(2.0) 4622(5)$ to (ground state + 80% $^{205}\text{Pb}^m$ at 2.329), $3/2^-$ at 262.8keV										Ens044 **
$^{209}\text{Ra}(\alpha)^{205}\text{Rn}$	$E_{\alpha}=7003(10)$ to ground state, $6625(5)$ to $387.0$ level										03He06 **
$^{209}\text{Th}^m(\alpha)^{205}\text{Ra}^m$	the decay is from $^{209}\text{Th}$ isomer, following $Ea_1 - Ea_2 - Ea_3 - Ea_4$ correlations										FGK141 **
$^{209}\text{Bi}(p,t)^{207}\text{Bi}$	$Q - Q(^{208}\text{Pb}(p,t)) = -241(2, \text{Be}), Q(\text{Pb}) = -5623.82(0.20)$ keV										AHW **
$^{208}\text{Pb}(d,p)^{209}\text{Pb}$	$Q - Q(^{209}\text{Bi}(d,p)) = -662(4), Q(\text{Bi}) = 2380.01(0.14)$ keV										AHW **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
* <sup>209</sup> Rn( $\beta^+$ ) <sup>209</sup> At	$E_{\beta^+}=2160(40)$ to $7/2^-$ level at 745.81 keV										Ens156 **
<sup>210</sup> Fr- <sup>226</sup> Ra <sub>929</sub>	-27198	24	-27183	16	0.6	1	46	46 <sup>210</sup> Fr	MA3	1.0	92Bo28
<sup>209</sup> Fr- <sup>210</sup> Fr <sub>498</sub> <sup>208</sup> Fr <sub>502</sub>	-770	50	-771	17	0.0	U			P24	2.5	82Au01
<sup>210</sup> Pb( $\alpha$ ) <sup>206</sup> Hg	3792.4	20.				2					62Ka27
<sup>210</sup> Bi( $\alpha$ ) <sup>206</sup> Tl	5042.8	2.	5036.5	0.8	-3.2	B			Orm		60Wa14 *
	5037.3	1.1			-0.8	1	50	33 <sup>210</sup> Bi			76Tu.A *
<sup>210</sup> Po( $\alpha$ ) <sup>206</sup> Pb	5407.53	0.07	5407.53	0.07	0.0	1	100	98 <sup>210</sup> Po			73Go39 Z
	5407.7	2.			-0.1	U					89Ma05
<sup>210</sup> At( $\alpha$ ) <sup>206</sup> Bi	5630.9	1.5	5631.2	1.0	0.2	3			DbA		69Go23 *
	5631.4	1.3			-0.2	3			DbA		81Va27 *
<sup>210</sup> Rn( $\alpha$ ) <sup>206</sup> Po	6162.1	3.	6159.0	2.2	-1.0	3					55Mo69 Z
	6155.9	3.			1.0	3			DbA		71Go35 Z
<sup>210</sup> Fr( $\alpha$ ) <sup>206</sup> At	6699.9	5.	6672	5	-5.7	B					67Va20 *
	6672.3	5.			-0.1	1	97	54 <sup>210</sup> Fr	GSa		05Ku06
<sup>210</sup> Ra( $\alpha$ ) <sup>206</sup> Rn	7156.6	5.	7151	3	-1.1	2					67Va22 Z
	7147	5			0.8	2			GSa		03He06 *
	7146.4	5.			0.5	2			JyA		07Le14
<sup>210</sup> Ac( $\alpha$ ) <sup>206</sup> Fr	7607.2	8.	7610	50	0.0	5					68Va04
	7607.2	10.			0.0	5			GSa		00He17
<sup>210</sup> Th( $\alpha$ ) <sup>206</sup> Ra	8052.7	17.	8069	6	0.9	4			JyA		95Uu01
	7962.0	50.			2.1	F			JAA		96Ik01 *
	8071.0	6.			-0.3	4			Anv		10He25
<sup>208</sup> Pb(t,p) <sup>210</sup> Pb	628	12	640.7	0.9	1.1	U			Ald		68Bj03
<sup>209</sup> Bi(n, $\gamma$ ) <sup>210</sup> Bi	4604.5	0.3	4604.63	0.08	0.4	-					71Mo03
	4604.68	0.14			-0.3	-			MMn		83Ts01 Z
	4604.63	0.10			0.0	-			Bdn		06Fi.A
<sup>209</sup> Bi(d,p) <sup>210</sup> Bi	2369	10	2380.07	0.08	1.1	U			MIT		64Sp12
<sup>209</sup> Bi(n, $\gamma$ ) <sup>210</sup> Bi	ave.	4604.64	0.08	4604.63	0.08	0.0	1	100	86 <sup>209</sup> Bi		average
<sup>210</sup> Tl( $\beta^-$ ) <sup>210</sup> Pb	5500	100	5482	12	-0.2	U					64We06 *
<sup>210</sup> Pb( $\beta^-$ ) <sup>210</sup> Bi	63.5	0.5	63.5	0.5	0.0	1	100	97 <sup>210</sup> Pb			67Ha03 *
<sup>210</sup> Bi( $\beta^-$ ) <sup>210</sup> Po	1160.5	1.5	1161.2	0.8	0.4	-					62Da03
	1161.5	1.5			-0.2	-					67Hs01
	ave.	1161.0	1.1		0.1	1	52	50 <sup>210</sup> Bi			average
<sup>210</sup> At( $\epsilon$ ) <sup>210</sup> Po	3870	30	3981	8	3.7	B					63Sc15 *
* <sup>210</sup> Bi( $\alpha$ ) <sup>206</sup> Tl	$E_{\alpha}=4685.3(2,Z)$ , $4648.3(2,Z)$ to $2^-$ level at 265.832, $1^-$ at 304.896 keV										Ens085 **
* <sup>210</sup> Bi( $\alpha$ ) <sup>206</sup> Tl	$E_{\alpha}=4946(1)$ , $4909(1)$ from <sup>210</sup> Bi <sup>m</sup> at 271.31 keV										Nub16b **
*	to $2^-$ level at 265.832, $1^-$ level at 304.896 keV										Ens085 **
* <sup>210</sup> At( $\alpha$ ) <sup>206</sup> Bi	$E_{\alpha}=5523.8$ , $5464.8$ , $5441.8(1.5,Z)$ to ground state, $4^+$ at 59.897, $5^+$ at 82.818										Ens085 **
* <sup>210</sup> At( $\alpha$ ) <sup>206</sup> Bi	$E_{\alpha}=5524.1$ , $5465.3$ , $5442.8(1.3,Z)$ to ground state, $4^+$ at 59.897, $5^+$ at 82.818										Ens085 **
* <sup>210</sup> Fr( $\alpha$ ) <sup>206</sup> At	$E_{\alpha}=6572.0(5,Z)$ $6542(5,Z)$ to ground state and level at 31.05 keV										Ens085 **
* <sup>210</sup> Ra( $\alpha$ ) <sup>206</sup> Rn	$E_{\alpha}=7003(10)$ to ground state, $6447(5)$ to 574.9 level										03He06 **
* <sup>210</sup> Th( $\alpha$ ) <sup>206</sup> Ra	F : Low energy; may be escape										96Ik01 **
* <sup>210</sup> Tl( $\beta^-$ ) <sup>210</sup> Pb	$E_{\beta^-}=1870(100)$ to $3625(19)$ level, and other $E_{\beta^-}$										Ens148 **
* <sup>210</sup> Pb( $\beta^-$ ) <sup>210</sup> Bi	$E_{\beta^-}=17.0(0.5)$ to $0^-$ level at 46.5390 keV										Ens148 **
* <sup>210</sup> At( $\epsilon$ ) <sup>210</sup> Po	pK=0.46(0.10) to $(6)^-$ level at 3727.34 keV										Ens148 **
<sup>211</sup> Tl-u	-6525	45				2			GS3	1.0	08Ch.A
<sup>211</sup> Fr- <sup>133</sup> Cs <sub>1.586</sub>	145517	15	145508	13	-0.6	1	74	74 <sup>211</sup> Fr	MA8	1.0	09Ko35
<sup>211</sup> Fr- <sup>226</sup> Ra <sub>934</sub>	-28200	25	-28176	13	1.0	1	27	26 <sup>211</sup> Fr	MA3	1.0	92Bo28
<sup>211</sup> Ra- <sup>133</sup> Cs <sub>1.586</sub>	150846.4	8.5				2			MA8	1.0	09Ko35
<sup>211</sup> Po <sup>m</sup> - <sup>211</sup> Po	1580	129	1570	5	0.0	U				2.5	15Di03
<sup>207</sup> Fr- <sup>211</sup> Fr <sub>327</sub> <sup>205</sup> Fr <sub>673</sub>	-930	100	-609	19	1.3	U			P24	2.5	82Au01
<sup>208</sup> Fr- <sup>211</sup> Fr <sub>394</sub> <sup>206</sup> Fr <sub>606</sub>	-260	50	-340	60	-0.7	U			P24	2.5	82Au01

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{210}\text{Fr}-^{211}\text{Fr}_{.498}$ $^{209}\text{Fr}_{.502}$	580	50	621	18	0.3	U			P24	2.5	82Au01
$^{211}\text{Bi}(\alpha)^{207}\text{Tl}$	6749.5	0.7	6750.4	0.5	1.2	-					61Ry02 Z
	6751.1	0.6			-1.2	-					71Gr17 Z
	6750.4	0.5			-0.1	1	100	57 $^{211}\text{Bi}$			average
$^{211}\text{Po}(\alpha)^{207}\text{Pb}$	7594.5	0.5				2			Orm		62Wa18 Z
	7594.3	3.	7594.6	0.5	0.1	U			DbA		69Go23 Z
	7600.6	2.			-3.0	B					85La17 Z
	7586.0	15.3			0.6	U					15Di03
$^{211}\text{Po}^m(\alpha)^{207}\text{Pb}$	9057.1	5.1				2			Bka		82Bo04
	9049.0	15.	9057	5	0.5	U					89Ku08 *
	9043.0	15.			0.9	U					15Di03 *
$^{211}\text{At}(\alpha)^{207}\text{Bi}$	5979.4	2.	5982.4	1.3	1.5	2			DbA		69Go23 Z
	5981.6	3.			0.3	2			Bka		82Bo04 *
	5985.9	2.			-1.7	2					85La17 Z
$^{211}\text{Rn}(\alpha)^{207}\text{Po}$	5967.9	2.	5965.5	1.4	-1.2	2					55Mo69 Z
	5963.1	2.			1.2	2			DbA		71Go35 Z
$^{211}\text{Fr}(\alpha)^{207}\text{At}$	6660.2	5.1	6662	3	0.4	2					67Va20 Z
	6663.5	4.			-0.3	2			GSa		05Ku06
$^{211}\text{Ra}(\alpha)^{207}\text{Rn}$	7045.3	5.	7042	3	-0.7	3					67Va22 Z
	7040	5			0.4	3			GSa		03He06 *
	7039.7	6.			0.4	3			Jya		07Le14
$^{211}\text{Ac}(\alpha)^{207}\text{Fr}$	7624.8	8.	7620	50	-0.1	2					68Va04
	7616.7	10.			0.1	2			GSa		00He17
$^{211}\text{Th}(\alpha)^{207}\text{Ra}$	7942.9	14.	7940	50	0.0	5			Jya		95Uu01
	7930.6	30.6			0.2	5			Lza		15Ya13
$^{211}\text{Pb}(\beta^-)^{211}\text{Bi}$	1378	8	1366	5	-1.5	1	47	43 $^{211}\text{Bi}$			65Co06
* $^{211}\text{Po}^m(\alpha)^{207}\text{Pb}$	$E_\alpha=7275(15)$ to $13/2^+$ level at 1633.356 keV										Ens112 **
* $^{211}\text{Po}^m(\alpha)^{207}\text{Pb}$	$E_\alpha=7269(15)$ to $13/2^+$ level at 1633.356 keV										Ens112 **
* $^{211}\text{At}(\alpha)^{207}\text{Bi}$	Recalibrated as in reference										91Ry01 **
* $^{211}\text{Ra}(\alpha)^{207}\text{Rn}$	Average of $E_\alpha=6907(5)$ and several branches to known levels										03He06 **
$^{212}\text{Bi}^n-u$	-7127	32				2			GS3	1.0	08Ch.A
$^{212}\text{Fr}-^{133}\text{Cs}_{1.594}$	146938	10	146935	9	-0.3	1	89	89 $^{212}\text{Fr}$	MA8	1.0	09Ko35
$^{212}\text{Fr}-^{226}\text{Ra}_{.938}$	-27631	28	-27608	10	0.8	1	12	11 $^{212}\text{Fr}$	MA3	1.0	92Bo28
$^{209}\text{Fr}-^{212}\text{Fr}_{.563}$ $^{205}\text{Fr}_{.437}$	-1270	70	-1218	16	0.3	U			P24	2.5	82Au01
$^{206}\text{Fr}^x-^{212}\text{Fr}_{.139}$ $^{205}\text{Fr}_{.861}$	340	130	470	100	0.4	U			P24	2.5	82Au01
$^{207}\text{Fr}-^{212}\text{Fr}_{.163}$ $^{206}\text{Fr}_{.837}$	-1150	70	-1320	90	-0.9	U			P24	2.5	82Au01
$^{212}\text{Bi}(\alpha)^{208}\text{Tl}$	6207.12	0.04	6207.262	0.028	3.5	B			BIP		61Ry02 Z
	6207.09	0.08			2.1	o			BIP		69Gr28 *
	6207.262	0.028				2			BIP		72Go.A *
$^{212}\text{Bi}^m(\alpha)^{208}\text{Tl}$	6458.1	30.				3					78Ba44
$^{212}\text{Po}(\alpha)^{208}\text{Pb}$	8953.6	0.8	8954.20	0.11	0.7	U					61Ry02 Z
	8953.85	0.31			1.1	-					71De52 Z
	8953.3	0.6			1.5	U					71Gr17 Z
	8954.25	0.12			-0.5	-					74Hu15 Z
	8954.20	0.11			0.0	1	100	91 $^{212}\text{Po}$			average
$^{212}\text{Po}^m(\alpha)^{208}\text{Pb}$	11874.6	20.	11877	4	0.1	2					62Pe15
	11859.3	15.			1.2	o					75Fr.B
	11884.8	10.2			-0.7	2					76Fr.A
	11875.6	5.1			0.3	2			GSa		12Ho12
$^{212}\text{At}(\alpha)^{208}\text{Bi}$	7829.0	9.	7817.1	0.6	-1.3	U					70Re02 *
	7817.0	0.6				3					76Fr.A *
	7828.0	10.			-1.1	U					96Li37 *
$^{212}\text{At}^m(\alpha)^{208}\text{Bi}$	8049.3	10.	8040.0	0.6	-0.9	U					68Va18
	8054.3	9.			-1.6	U					70Re02 *
	8040.00	0.61				3					76Fr.A *
	8051.2	10.			-1.1	U					96Li37 *
$^{212}\text{Rn}(\alpha)^{208}\text{Po}$	6392.3	5.	6385.1	2.6	-1.4	3					55Mo69 Z
	6382.5	3.			0.9	3			DbA		71Go35 Z
$^{212}\text{Fr}(\alpha)^{208}\text{At}$	6531.3	3.	6529.0	1.6	-0.7	2					66Va.A Z
	6528.0	3.			0.3	2			DbA		81Va27

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{212}\text{Fr}(\alpha)^{208}\text{At}$	6527.5	3.	6529.0	1.6	0.5	2			Bka		82Bo04 *
	6529.5	4.			-0.1	2			GSa		05Ku06
$^{212}\text{Ra}(\alpha)^{208}\text{Rn}$	7030.0	5.	7031.7	1.7	0.3	5					67Va22 Z
	7034.0	5.			-0.4	5					74Ho27 Z
	7032.2	2.			-0.3	5			Bka		82Bo04 Z
	7028	5			0.7	5			GSa		03He06 *
	7040	24			-0.3	U			Lza		14Ya19
$^{212}\text{Ac}(\alpha)^{208}\text{Fr}$	7521.2	8.	7520	50	-0.1	2					68Va04
	7515.1	10.			0.1	2			GSa		00He17
	7514.0	18.			0.1	2			Lza		14Ya19
	7952.3	10.	7958	5	0.6	3					80Ve01
$^{212}\text{Th}(\alpha)^{208}\text{Ra}$	7959.5	5.			-0.3	3			Anv		10He25
	7980.8	20.4			-1.1	U			GSa		15De22
	8429.4	30.	8420	50	-0.3	6			JAA		97Mi03
$^{212}\text{Pa}(\alpha)^{208}\text{Ac}$	8408.9	20.			0.1	6			Lza		14Ya19
$^{210}\text{Pb}(\text{t,p})^{212}\text{Pb}$	515	25	481.2	2.1	-1.4	U					71El05
$^{212}\text{Pb}(\beta^-)^{212}\text{Bi}$	569.3	2.5	569.1	1.8	-0.1	-					48Ma30 *
	576.6	5.			-1.5	-					58Se71 *
$^{212}\text{Bi}(\beta^-)^{212}\text{Po}$	ave.	570.8	2.2		-0.7	1	67	34 $^{212}\text{Bi}$			average
	2256	3	2251.5	1.7	-1.5	-					48Fe09
	2250.5	2.5			0.4	-					48Ma30
ave.	2252.8	1.9			-0.6	1	75	66 $^{212}\text{Bi}$			average
* $^{212}\text{Bi}(\alpha)^{208}\text{Tl}$	$E_\alpha=6089.86(0.08,Z)$ , 6050.57(0.07,Z) to ground state, $4^+$ level at 39.858 keV										
* $^{212}\text{Bi}(\alpha)^{208}\text{Tl}$	$E_\alpha=6089.883(0.037,Z)$ , 6050.837(0.028,Z) to ground state, $4^+$ level at 39.858 keV										
* $^{212}\text{At}(\alpha)^{208}\text{Bi}$	Original $E_\alpha=7679(8)$ ; calibration $^{211}\text{Po}$ 7448(1), now 7450.3(0.5) keV										
* $^{212}\text{At}(\alpha)^{208}\text{Bi}$	Original $E_\alpha=7669.0(0.2)$ ; calibration $^{211}\text{Po}$ 7450(2), now 7450.3(0.5)										
* $^{212}\text{At}(\alpha)^{208}\text{Bi}$	$E_\alpha=7679(10)$ to ground state, 7618(10) to 63.3 level										
*	error estimated by the evaluators										
* $^{212}\text{At}^m(\alpha)^{208}\text{Bi}$	Original $E_\alpha=7900(8)$ ; calibration $^{211}\text{Po}$ 7448(1), now 7450.3(0.5) keV										
* $^{212}\text{At}^m(\alpha)^{208}\text{Bi}$	Original $E_\alpha=7887.7(0.2)$ ; calibration $^{211}\text{Po}$ 7450(2), now 7450.3(0.5)										
* $^{212}\text{At}^m(\alpha)^{208}\text{Bi}$	$E_\alpha=7897(10)$ to ground state, 7837(10) to 63.3 level										
*	error estimated by the evaluators										
* $^{212}\text{Fr}(\alpha)^{208}\text{At}$	$E_\alpha=6341(3)$ (recalibrated as in reference) to 63.70 level										
* $^{212}\text{Ra}(\alpha)^{208}\text{Rn}$	$E_\alpha=6898(5)$ to ground state, 6269(5) to 635.1 level										
* $^{212}\text{Pb}(\beta^-)^{212}\text{Bi}$	$E_{\beta^-}=330.7(2.5)$ 338(5) respectively to $0^-$ level at 238.632 keV										
$^{213}\text{Tl}-u$	1893	65	1915	29	0.3	o			GS3	1.0	10Ch19
	1915	29				2			GS3	1.0	12Ch19
$^{213}\text{Fr}-^{133}\text{Cs}_{1.602}$	147649.1	7.4	147652	5	0.4	1	55	55 $^{213}\text{Fr}$	MA8	1.0	09Ko35
$^{213}\text{Ra}-^{133}\text{Cs}_{1.602}$	151833	12	151837	11	0.3	1	77	77 $^{213}\text{Ra}$	SH1	1.0	13Dr04
$^{207}\text{Fr}-^{213}\text{Fr}_{.324}$ $^{204}\text{Fr}_{.676}$	-2540	330	-2104	24	0.5	U			P24	2.5	82Au01 *
$^{208}\text{Fr}-^{213}\text{Fr}_{.279}$ $^{206}\text{Fr}_{.721}$	-700	60	-850	80	-1.0	U			P24	2.5	82Au01
$^{209}\text{Fr}-^{213}\text{Fr}_{.327}$ $^{207}\text{Fr}_{.673}$	-670	60	-694	19	-0.2	U			P24	2.5	82Au01
$^{209}\text{Fr}-^{213}\text{Fr}_{.196}$ $^{208}\text{Fr}_{.804}$	-980	60	-930	17	0.3	U			P24	2.5	82Au01
$^{211}\text{Fr}-^{213}\text{Fr}_{.330}$ $^{210}\text{Fr}_{.670}$	-830	60	-735	16	0.6	U			P24	2.5	82Au01
$^{212}\text{Fr}-^{213}\text{Fr}_{.498}$ $^{211}\text{Fr}_{.502}$	270	50	332	11	0.5	U			P24	2.5	82Au01
$^{213}\text{Bi}(\alpha)^{209}\text{Tl}$	5982.6	6.	5988	3	0.9	2					64Gr11
	5990.7	4.			-0.6	2			Gea		13Ma13
$^{213}\text{Po}(\alpha)^{209}\text{Pb}$	8537.1	5.	8536.1	2.6	-0.2	-					64Va20 Z
	8536.5	3.			-0.1	-			Bka		82Bo04 Z
	ave.	8536.7	2.6		-0.2	1	95	93 $^{213}\text{Po}$			average
$^{213}\text{At}(\alpha)^{209}\text{Bi}$	9254.2	12.	9254	5	0.0	2					70Bo13
	9254.2	5.			0.0	2			Lvn		87De.A
$^{213}\text{Rn}(\alpha)^{209}\text{Po}$	8245.1	8.	8245.2	2.9	0.0	3					67Va20
	8240.0	10.			0.5	U					70Va13
	8242	10			0.3	U			GSa		00He17 *
	8245.2	3.1			0.0	3			Jya		01Ku07
	8218.6	44.			0.6	U					05Li17
$^{213}\text{Fr}(\alpha)^{209}\text{At}$	6904.0	5.	6904.9	1.2	0.2	-					67Va20 Z
	6908.0	5.			-0.6	-					74Ho27 Z
	6904.6	2.			0.1	-			Bka		82Bo04 Z

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{213}\text{Fr}(\alpha)^{209}\text{At}$	6904.9	1.7	6904.9	1.2	0.0	-			GSa		05Ku06
	6880	20			1.2	U			GSa		15De22
$^{213}\text{Ra}(\alpha)^{209}\text{Rn}$	ave. 6904.9	1.2			-0.1	1	99	53 $^{209}\text{At}$			average
	6860.3	5.	6861.7	2.3	0.3	-					67Va22 *
	6862.4	5.			-0.1	-					76Ra37 *
	ave. 6862.2	3.			-0.2	-			GSa		06Ku26 *
$^{213}\text{Ra}^m(\alpha)^{209}\text{Rn}$	6861.8	2.3			-0.1	1	99	76 $^{209}\text{Rn}$			average
	8630.3	5.1	8630	4	-0.1	2					76Ra37
$^{213}\text{Ac}(\alpha)^{209}\text{Fr}$	8629.3	5.			0.1	2			GSa		06Ku26 *
	7505.2	8.	7499	4	-0.7	3					68Va04 *
$^{213}\text{Th}(\alpha)^{209}\text{Ra}$	7497.0	10.			0.2	o			GSa		00He17
	7497.0	5.			0.4	3			GSa		02He.A
	7837.4	10.	7837	7	-0.1	3					68Va18 *
$^{213}\text{Pa}(\alpha)^{209}\text{Ac}$	7836.5	10.			0.0	3					80Ve01
	8393.9	15.				4			GSa		00He17
$^{213}\text{Bi}(\beta^-)^{213}\text{Po}$	1430	10	1422	5	-0.8	1	30	23 $^{213}\text{Bi}$			68Va17 *
$^{207}\text{Fr}-^{213}\text{Fr}_{324}$ $^{204}\text{Fr}$ .	$D_M=-2470(330)$ keV for $^{204}\text{Fr}$ mixture gs+m+n at 50(4), 326(4) keV										
$^{213}\text{Rn}(\alpha)^{209}\text{Po}$	$E_\alpha=8088(10)$ , 7550(15) to ground state, 540.3 level										
$^{213}\text{Ra}(\alpha)^{209}\text{Rn}$	$E_\alpha=6730.7$ , 6623.7, 6520.7(3,Z) to ground state, $1/2^-$ at 110.25, $3/2^-$ at 214.93										
$^{213}\text{Ra}(\alpha)^{209}\text{Rn}$	$E_\alpha=6731.9$ , 6624.9, 6523.9(5,Z) to ground state, $1/2^-$ at 110.25, $3/2^-$ at 214.93										
$^{213}\text{Ra}(\alpha)^{209}\text{Rn}$	$E_\alpha=6733(3)$ , 6625(3) to ground state and $1/2^-$ level at 110.25 keV										
$^{213}\text{Ra}^m(\alpha)^{209}\text{Rn}$	$E_\alpha=8467$ (5) 8358(10) to ground state and $1/2^-$ level at 110.25 keV										
$^{213}\text{Ac}(\alpha)^{209}\text{Fr}$	Original $Q$ increased by 2 keV, as in reference										
$^{213}\text{Th}(\alpha)^{209}\text{Ra}$	Original $Q$ decreased by 0.5 keV, as in reference										
$^{213}\text{Bi}(\beta^-)^{213}\text{Po}$	$E_{\beta^-}=1420(10)$ 1018(15) to ground state and $7/2^+$ level at 440.45 keV										
$^{214}\text{Ra}-^{133}\text{Cs}_{1.609}$	152235	22	152227	6	-0.3	U			MA8	1.0	08We02
$^{214}\text{Bi}(\alpha)^{210}\text{Tl}$	5621.3	3.0				2					91Ry01 *
$^{214}\text{Po}(\alpha)^{210}\text{Pb}$	7833.54	0.06	7833.54	0.06	0.0	1	100	97 $^{214}\text{Po}$			71Gr17 Z
$^{214}\text{At}(\alpha)^{210}\text{Bi}$	8987.2	4.				2			Bka		82Bo04 Z
$^{214}\text{At}^m(\alpha)^{210}\text{Bi}$	9046.4	8.				2					82Ew01
$^{214}\text{At}^n(\alpha)^{210}\text{Bi}$	9220.8	5.				2					82Ew01 *
$^{214}\text{Rn}(\alpha)^{210}\text{Po}$	9212.6	20.	9208	9	-0.2	2					70To07
	9207.5	10.			0.1	2					70Va13
$^{214}\text{Fr}(\alpha)^{210}\text{At}$	8585.5	8.	8589	4	0.4	4					68Va18 *
	8590.9	5.			-0.5	4					70To18 *
	8583.8	10.			0.5	4			Dbb		89An.A
	8590.8	20.			-0.1	U			GSa		90Ni05
$^{214}\text{Fr}^m(\alpha)^{210}\text{At}$	8578.7	48.			0.2	U					05Li17
	8711.7	8.	8710	3	-0.2	4					68Va04 Z
	8711.7	5.			-0.3	4					70To18 *
$^{214}\text{Ra}(\alpha)^{210}\text{Rn}$	8708.1	5.			0.4	4			GSa		05Ku06
	7271.7	5.	7272.6	2.6	0.2	4					67Va22 Z
	7275.6	5.			-0.6	4					74Ho27 Z
	7273.2	10.			-0.1	4			GSa		00He17 *
$^{214}\text{Ra}(\alpha)^{210}\text{Rn}^m$	7271.2	4.			0.3	4			GSa		06Ku26 *
	5563.9	30.				5			GSa		06Ku26 *
$^{214}\text{Ac}(\alpha)^{210}\text{Fr}$	7351.7	5.	7352.1	2.5	0.1	2					68Va04 Z
	7347.6	10.			0.4	2			Dbb		89An13
	7347.6	10.			0.5	o			GSa		00He17 *
	7349.6	5.			0.5	o			GSa		02He.A
	7352.7	3.			-0.2	2			GSa		04Ku24 *
$^{214}\text{Th}(\alpha)^{210}\text{Ra}$	7828.6	10.	7827	5	-0.1	3					68Va18
	7823.5	10.			0.4	3					80Ve01
	7828.6	8.			-0.2	3			Jya		07Le14

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{214}\text{Pa}(\alpha)^{210}\text{Ac}$	8270.9	15.				6			GSa		00He17
$^{214}\text{Pb}(\beta^-)^{214}\text{Bi}$	1024	20	1018	11	-0.3	1	32	31 $^{214}\text{Bi}$			52Be78 *
$^{214}\text{Bi}(\beta^-)^{214}\text{Po}$	3260	30	3269	11	0.3	-					56Da06
	3275	15			-0.4	-					60Lu07
ave.	3272	13			-0.2	1	69	69 $^{214}\text{Bi}$			average
* $^{214}\text{Bi}(\alpha)^{210}\text{Tl}$	$E_\alpha=5516(3)$ recommended in place of the following $E_\alpha$ :										
*	$E_\alpha=5510.5(1.0)$ keV										
*	$E_\alpha=5515.8(3.0)$ keV										
* $^{214}\text{At}^m(\alpha)^{210}\text{Bi}$	$E_\alpha=8782(5)$ to $9^-$ level at 271.31 keV										
* $^{214}\text{Fr}(\alpha)^{210}\text{At}$	$E_\alpha=8425.5, 8352.5(8,Z)$ to ground state, $(4)^+$ level at 72.65 keV										
* $^{214}\text{Fr}(\alpha)^{210}\text{At}$	$E_\alpha=8428.3, 8360.3(5,Z)$ to ground state, $(4)^+$ level at 72.65 keV										
* $^{214}\text{Fr}^m(\alpha)^{210}\text{At}$	$E_\alpha=8546.8, 8477.8(5,Z)$ to ground state, $(4)^+$ level at 72.65 keV										
* $^{214}\text{Ra}(\alpha)^{210}\text{Rn}$	$E_\alpha=7137(10), 6505(15)$ to ground state, 641.9 level										
* $^{214}\text{Ra}(\alpha)^{210}\text{Rn}$	Also $E_\alpha=8950(30)$ keV $Q_\alpha=9120.9$ keV from $^{214}\text{Ra}^n$ at 1865.2 keV										
* $^{214}\text{Ra}(\alpha)^{210}\text{Rn}^m$	$E_\alpha=7290(30)$ $Q_\alpha=7429.1$ from $^{214}\text{Ra}^n$ at 1865.2 keV										
* $^{214}\text{Ac}(\alpha)^{210}\text{Fr}$	$E_\alpha=7210(10), 7080(15)$ to ground state, 138.6 level										
* $^{214}\text{Ac}(\alpha)^{210}\text{Fr}$	Also $E_\alpha=7081(4)$ keV to 139.0(1) level										
* $^{214}\text{Pb}(\beta^-)^{214}\text{Bi}$	$E_{\beta^-}=670(20)$ to $(0^-, 1^-)$ level at 351.9324 keV, and another branch										
$^{215}\text{Bi}-^{133}\text{Cs}_{1.617}$	154654	16	154633	6	-1.3	1	14	14 $^{215}\text{Bi}$	MA8	1.0	08We02
$^{215}\text{Po}(\alpha)^{211}\text{Pb}$	7526.45	0.8	7526.3	0.8	-0.2	1	99	96 $^{211}\text{Pb}$			71Gr17 Z
$^{215}\text{At}(\alpha)^{211}\text{Bi}$	8178.5	4.				2			Bka		82Bo04 Z
$^{215}\text{Rn}(\alpha)^{211}\text{Po}$	8834.7	20.	8839	8	0.2	3			ORa		69Ha32
	8839.8	8.			-0.1	3					70Va13
$^{215}\text{Fr}(\alpha)^{211}\text{At}$	9543.0	15.	9540	7	-0.2	3					70Bo13
	9532.7	10.			0.8	3					74No02
	9546.9	10.			-0.6	3					84De16
$^{215}\text{Ra}(\alpha)^{211}\text{Rn}$	8862.7	5.	8864	3	0.3	3					68Va18 Z
	8865.5	5.			-0.2	3					70To18 Z
	8865.3	10.			-0.1	3			GSa		00He17
	8865.3	46.			0.0	U					05Li17
$^{215}\text{Ac}(\alpha)^{211}\text{Fr}$	7748.4	5.	7746	3	-0.5	2					68Va04 Z
	7746	10			0.0	o			GSa		00He17 *
	7740.3	5.			1.1	o			GSa		02He.A
	7744.4	4.			0.4	2			GSa		04Ku24
$^{215}\text{Th}(\alpha)^{211}\text{Ra}$	7664.9	8.	7665	4	0.0	3					68Va18
	7667.0	10.			-0.2	o			GSa		89He03
	7664	15			0.0	o			GSa		00He17 *
	7665	5			-0.1	3			GSa		05Ku31 *
	7662.8	10.			0.2	3			Jya		07Le14 *
$^{215}\text{Pa}(\alpha)^{211}\text{Ac}$	8238.6	15.	8240	50	0.0	3					79Sc09
	8244.7	15.			-0.2	o			GSa		00He17
	8233.5	20.4			0.1	3			GSa		15De22
$^{215}\text{U}(\alpha)^{211}\text{Th}$	8588.0	30.6				6			Lza		15Ya13
* $^{215}\text{Ac}(\alpha)^{211}\text{Fr}$	$E_\alpha=7602(10)$ $7026(15)$ $6960(15)$ to ground state, $11/2^-$ at 583.28, $13/2^-$ at 652.62										
* $^{215}\text{Th}(\alpha)^{211}\text{Ra}$	$E_\alpha=7520(15), 7387(15), 7336(15)$ to ground state, 133.6, 192.4 levels										
* $^{215}\text{Th}(\alpha)^{211}\text{Ra}$	$E_\alpha=7523(5), 7392(4), 7335(5), 7236(7)$ to ground state, 133.9, 194.5, 295.1 levels										
* $^{215}\text{Th}(\alpha)^{211}\text{Ra}$	Also $E_\alpha=7399(20)$ keV to 133.9 level										
$^{216}\text{Bi}-^{133}\text{Cs}_{1.624}$	159852	12				2			MA8	1.0	08We02
$^{216}\text{Po}(\alpha)^{212}\text{Pb}$	6906.44	0.5	6906.4	0.5	-0.2	1	98	67 $^{212}\text{Pb}$			71Gr17 Z
$^{216}\text{At}(\alpha)^{212}\text{Bi}$	7949.7	3.				2			Bka		82Bo04 Z
$^{216}\text{At}^m(\alpha)^{212}\text{Bi}$	8110.5	10.				2					71Br13
$^{216}\text{Rn}(\alpha)^{212}\text{Po}$	8199.2	10.	8197	6	-0.2	2					61Ru06
	8201.2	10.			-0.4	2					70Va13
	8192.0	10.2			0.5	2					71Br13
$^{216}\text{Fr}(\alpha)^{212}\text{At}$	9175.3	12.	9174	3	-0.1	4					70Bo13
	9174.1	5.			0.0	4					96Li37 *
	9174.3	5.			0.0	4			GSa		07Ku30

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{216}\text{Fr}^m(\alpha)^{212}\text{At}^m$	9170.2	5.				4			GSa		07Ku30
$^{216}\text{Ra}(\alpha)^{212}\text{Rn}$	9525.8	8.				4					73No09
$^{216}\text{Ac}(\alpha)^{212}\text{Fr}$	9243.3	8.	9235	6	-1.0	2					70To18 Z
	9223.1	10.			1.2	2		GSa			00He17
	9241.4	50.9			-0.1	U					05Li17
$^{216}\text{Ac}^m(\alpha)^{212}\text{Fr}$	9280.0	5.	9279	4	-0.2	2					70To18 Z
	9284	10			-0.5	o		GSa			00He17 *
	9278.2	5.			0.2	o		GSa			02He.A
	9277.2	7.			0.3	2		GSa			04Ku24 *
$^{216}\text{Th}(\alpha)^{212}\text{Ra}$	8070.7	8.	8072	4	0.2	6					68Va18
	8071	10			0.1	o		GSa			00He17 *
	8073	5			-0.1	6		GSa			05Ku31 *
	8069.7	44.			0.1	U					05Li17
	8070.7	23.			0.1	U		Lza			14Ya19
$^{216}\text{Th}^m(\alpha)^{212}\text{Ra}$	10099.4	20.	10116	8	0.8	6					83Hi08
	10107.4	40.			0.2	U			Dbb		93An07
	10120.8	15.			-0.3	6		GSa			00He17
	10117.5	10.			-0.2	6					05Ku31 *
$^{216}\text{Pa}(\alpha)^{212}\text{Ac}$	8013.7	20.	8097	15	4.2	B					79Sc09
	8110.5	50.			-0.3	U		JAA			98Ik01
	8097	15				3		GSa			00He17 *
$^{216}\text{U}(\alpha)^{212}\text{Th}$	8542.5	30.6	8531	26	-0.4	4		Lza			15Ma37
	8497.6	50.9			0.6	4		GSa			15De22
$^{216}\text{U}^m(\alpha)^{212}\text{Th}$	10782.0	30.6				4		Lza			15Ma37
$^{*216}\text{Fr}(\alpha)^{212}\text{At}$	$E_\alpha=9004(5)$ ; and $E_\alpha=8933(8)$ from 133.3 level to 205.6 keV										96Li37 **
$^{*216}\text{Ac}^m(\alpha)^{212}\text{Fr}$	$E_\alpha=9110(10)$ , 9026(15), 8586(15) to ground state, 82.4, 542.2 levels										00He17 **
$^{*216}\text{Ac}^m(\alpha)^{212}\text{Fr}$	Also $E_\alpha=9029(7)$ keV to 82.6(1) level										04Ku24 **
$^{*216}\text{Th}(\alpha)^{212}\text{Ra}$	$E_\alpha=7923(10)$ , 7302(15) to ground state, 618.3 level										00He17 **
$^{*216}\text{Th}(\alpha)^{212}\text{Ra}$	$E_\alpha=7923(5)$ , 7304(4) to ground state, 629.3(1) level										05Ku31 **
$^{*216}\text{Th}^m(\alpha)^{212}\text{Ra}$	$E_\alpha=9930(10)$ , 9312(12) to ground state, 629.3(1) level										05Ku31 **
$^{*216}\text{Pa}(\alpha)^{212}\text{Ac}$	$E_\alpha=7948(15)$ , 7815(15) to ground state, 133.6 level										00He17 **
$^{217}\text{Bi}-u$	9420	32	9372	19	-1.5	o			GS3	1.0	08Ch.A
	9372	19				2			GS3	1.0	12Ch19
$^{217}\text{Po}(\alpha)^{213}\text{Pb}$	6660.3	4.	6662.1	2.4	0.4	4			DbA		77Vy02 Z
	6660.0	4.			0.5	4			Orm		97Li23
	6666.1	4.1			-1.0	4			Anv		03Ku25
$^{217}\text{At}(\alpha)^{213}\text{Bi}$	7200.3	3.	7201.4	1.2	0.4	-					60Vo05 Z
	7200.3	2.			0.5	-			Orm		62Wa28 Z
	7204.6	5.			-0.6	-					64Va20 Z
	7193.1	5.			1.6	-			DbA		77Vy02 Z
	7204.0	2.			-1.3	-			Bka		82Bo04
	ave.	7201.5	1.2		-0.1	1	99	77 $^{213}\text{Bi}$			average
$^{217}\text{Rn}(\alpha)^{213}\text{Po}$	7887.5	4.	7887.2	2.9	-0.1	2					61Ru06 Z
	7886.9	4.			0.1	2			Bka		82Bo04 Z
$^{217}\text{Fr}(\alpha)^{213}\text{At}$	8471.5	8.	8469	4	-0.3	3					70Bo13
	8468.4	5.			0.2	3			Lvn		87De.A
$^{217}\text{Ra}(\alpha)^{213}\text{Rn}$	9159.1	8.	9161	6	0.2	4					70To07
	9163.2	10.			-0.2	4					70Va13
$^{217}\text{Ac}(\alpha)^{213}\text{Fr}$	9831.6	10.				2					73No09
$^{217}\text{Ac}^m(\alpha)^{213}\text{Fr}$	11843.8	17.				2					85De14
$^{217}\text{Th}(\alpha)^{213}\text{Ra}$	9424.1	10.	9435	4	1.1	2					68Va18
	9424.1	20.			0.6	U					73Ha32
	9421.1	15.			0.9	U					00Ni02
	9442	15			-0.4	o			GSa		00He17 *
	9435.6	5.			-0.1	2			GSa		02He29 *
	9443.5	9.			-0.9	2			GSa		05Ku31
	9424.1	47.			0.2	U					05Li17
$^{217}\text{Pa}(\alpha)^{213}\text{Ac}$	8486.7	10.	8489	4	0.2	4					68Va18
	8489.8	15.			-0.1	U					79Sc09
	8486.7	50.			0.0	U			JAA		98Ik01

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{217}\text{Pa}(\alpha)^{213}\text{Ac}$	8490.8	15.	8489	4	-0.1	U			GSa		00He17
	8489.3	5.			-0.1	4			GSa		02He29 *
$^{217}\text{Pa}^m(\alpha)^{213}\text{Ac}$	10351	20	10349	5	-0.1	U					79Sc09
	10330.8	50.			0.4	U			JAA		98Ik01
	10346.1	15.			0.2	o			GSa		00He17
	10349.1	5.				4			GSa		02He29 *
$^{217}\text{U}(\alpha)^{213}\text{Th}^p$	8155.6	20.	8170	50	0.3	5					00Ma65
	8175.0	14.3			-0.1	5			Jya		05Le42 *
$^{*217}\text{Th}(\alpha)^{213}\text{Ra}$	$E_\alpha=9268(15), 8731(15), 8459(15)$ to ground state, 546.35, 822.7 levels										
$^{*217}\text{Th}(\alpha)^{213}\text{Ra}$	$E_\alpha=9261(5), 8725(5), 8455(5)$ to ground state, 546.35, 822.7 levels										
$^{*217}\text{Pa}(\alpha)^{213}\text{Ac}$	$E_\alpha=8337(5), 7873(5), 7728(5), 7710(5)$ to ground state, 466.1, 612.5, 634.3 levels										
$^{*217}\text{Pa}^m(\alpha)^{213}\text{Ac}$	Average of 5 $E_\alpha$ 's to known levels										
$^{*217}\text{U}(\alpha)^{213}\text{Th}^p$	Only one event. Not reported in later publication 07Le14										
$^{218}\text{Bi}-u$	14178	34	14188	29	0.3	o			GS3	1.0	08Ch.A
	14188	29				2			GS3	1.0	12Ch19
$^{218}\text{Po}(\alpha)^{214}\text{Pb}$	6114.76	0.09	6114.75	0.09	0.0	1	100	99 $^{214}\text{Pb}$			71Gr17 Z
$^{218}\text{At}(\alpha)^{214}\text{Bi}$	6874	3				2			Orm		58Wa.A *
$^{218}\text{Rn}(\alpha)^{214}\text{Po}$	7265.0	5.	7262.5	1.9	-0.5	-					56As38 Z
	7262.4	2.			0.0	-			Bka		82Bo04 Z
	ave.	7262.8	1.9		-0.2	1	96	93 $^{218}\text{Rn}$			average
$^{218}\text{Fr}(\alpha)^{214}\text{At}$	8014.0	2.				3		Bka		82Bo04 Z	
$^{218}\text{Fr}^m(\alpha)^{214}\text{At}$	8099.9	5.	8100	4	0.1	3					82Ew01 Z
	8100.9	5.			-0.1	3					99Sh03
$^{218}\text{Ra}(\alpha)^{214}\text{Rn}$	8549.1	8.	8546	6	-0.4	3					70To07
	8541.0	10.			0.5	3					70Va13
$^{218}\text{Ac}(\alpha)^{214}\text{Fr}$	9377.4	15.				5					70Bo13
$^{218}\text{Th}(\alpha)^{214}\text{Ra}$	9861.3	20.4	9849	9	-0.6	5					73Ha32
	9846.1	10.			0.3	5					73No09
	9851.0	81.5			0.0	U			GSa		15Kh09
	9794.1	20.	9815	10	1.0	F					79Sc09 *
$^{218}\text{U}(\alpha)^{214}\text{Th}$	9815	10				3			GSa		00He17 *
	8786.6	25.	8775	9	-0.5	4			Dbb		92An04
$^{218}\text{U}^m(\alpha)^{214}\text{Th}$	8773.2	9.			0.2	4			Jya		07Le14
	10878.1	17.	10884	15	0.3	4			Jya		07Le14
	10901.4	30.6			-0.6	4			Lza		15Ma37
$^{*218}\text{At}(\alpha)^{214}\text{Bi}$	$E_\alpha=6696.3(3.0, Z)$ to $(2)^-$ level at 53.2282 keV										
$^{*218}\text{Pa}(\alpha)^{214}\text{Ac}$	$E_\alpha=9614(20)$ ; F : probably piled-up with $e^-$										
$^{*218}\text{Pa}(\alpha)^{214}\text{Ac}$	$E_\alpha=9544(10)$ to 91.8 level										
$^{219}\text{Po}-u$	13601	32	13614	17	0.4	o			GS3	1.0	08Ch.A
	13614	17				2			GS3	1.0	12Ch19
$^{219}\text{At}-^{133}\text{Cs}_{1.647}$	166879.4	8.4	166881	3	0.2	1	17	17 $^{219}\text{At}$	MA8	1.0	16Ma.1
$^{219}\text{Po}(\alpha)^{215}\text{Pb}$	5914.2	5.				3			ISa		15Fi07
$^{219}\text{At}(\alpha)^{215}\text{Bi}$	6390.9	50.	6342	5	-1.0	U					53Hy83
	6344.0	5.			-0.4	1	90	86 $^{215}\text{Bi}$	ISa		15Fi07
$^{219}\text{Rn}(\alpha)^{215}\text{Po}$	6946.21	0.3	6946.2	0.3	-0.1	1	100	96 $^{215}\text{Po}$			71Gr17 Z
$^{219}\text{Fr}(\alpha)^{215}\text{At}$	7448.7	2.0	7448.6	1.8	-0.1	3			Orm		68Ba73 Z
	7448.2	4.			0.1	3			Bka		82Bo04 Z
$^{219}\text{Ra}(\alpha)^{215}\text{Rn}$	8139.0	20.	8138	3	-0.1	U			ORa		69Ha32
	8128.7	10.			0.9	U					70Va13
	8128.7	20.			0.5	U			Dbb		89An13
	8138.0	3.				4					94Sh02
$^{219}\text{Ac}(\alpha)^{215}\text{Fr}$	8826.5	10.				4					70Bo13
$^{219}\text{Th}(\alpha)^{215}\text{Ra}$	9514.1	20.	9510	50	0.0	4					73Ha32
	9503.9	50.9			0.2	4			GSa		15Kh09



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{219}\text{Pa}(\alpha)^{215}\text{Ac}$	10084.6	50.				3					87Fa.A
$^{219}\text{U}(\alpha)^{215}\text{Th}$	9860.4	40.	9940	50	1.6	4			Dbb		93An07
	9956.2	18.			-0.3	4			Jya		07Le14
$^{219}\text{Np}(\alpha)^{215}\text{Pa}$	9167.8	203.7				4			GSa		15De22 *
* $^{219}\text{Np}(\alpha)^{215}\text{Pa}$	$E_\alpha > 9000$ keV										15De22 **
$^{220}\text{Po}-u$	16420	32	16386	19	-1.1	o			GS3	1.0	08Ch.A
	16386	19				2			GS3	1.0	12Ch19
$^{220}\text{At}-u$	15427	32	15433	15	0.2	o			GS3	1.0	08Ch.A
	15433	15				2			GS3	1.0	12Ch19
$^{220}\text{Rn}-^{133}\text{Cs}_{1,654}$	167777	11	167775.0	1.9	-0.2	U			MA8	1.0	09Ne03
$^{210}\text{Fr}-^{220}\text{Fr}_{159}$	-2930	60	-2917	18	0.1	U			P24	2.5	82Au01
$^{211}\text{Fr}-^{220}\text{Fr}_{240}$	-4850	70	-4867	15	-0.1	U			P24	2.5	82Au01
$^{212}\text{Fr}-^{220}\text{Fr}_{321}$	-5450	60	-5392	12	0.4	U			P24	2.5	82Au01
$^{212}\text{Fr}-^{220}\text{Fr}_{263}$	-3730	60	-3754	14	-0.2	U			P24	2.5	82Au01
$^{213}\text{Fr}-^{220}\text{Fr}_{352}$	-5170	50	-5148	11	0.2	U			P24	2.5	82Au01
$^{212}\text{Fr}-^{220}\text{Fr}_{193}$	-3160	60	-3039	15	0.8	U			P24	2.5	82Au01
$^{220}\text{At}(\alpha)^{216}\text{Bi}^m$	6053.3	6.				3					89Bu09
$^{220}\text{Rn}(\alpha)^{216}\text{Po}$	6404.75	0.10	6404.74	0.10	0.0	1	100	69 $^{216}\text{Po}$			71Gr17 Z
$^{220}\text{Fr}(\alpha)^{216}\text{At}$	6799.0	2.	6800.7	1.9	0.9	3			Orm		68Ba73 *
	6811.6	5.			-2.2	3					74Ho27 *
$^{220}\text{Ra}(\alpha)^{216}\text{Rn}$	7593.3	10.	7592	6	-0.1	3					61Ru06
	7595.3	10.			-0.3	3					70Va13
	7598.4	20.4			-0.3	3			Dbb		90An19
	7587.2	10.			0.5	3			GSa		00He17
$^{220}\text{Ac}(\alpha)^{216}\text{Fr}$	8347.1	10.	8348	4	0.1	5					70Bo13
	8348	5			0.0	5					97Sh09 *
$^{220}\text{Th}(\alpha)^{216}\text{Ra}$	8953.1	20.				5					73Ha32
$^{220}\text{Pa}(\alpha)^{216}\text{Ac}$	9829.1	50.	9650#	50#	-3.6	D					87Fa.A *
* $^{220}\text{Fr}(\alpha)^{216}\text{At}$	$E_\alpha=6675.2, 6631.0, 6570.2(2,Z)$ to ground state, $(2)^-$ at 44.59, $(0)^-$ at 105.89										Ens075 **
* $^{220}\text{Fr}(\alpha)^{216}\text{At}$	$E_\alpha=6687.5, 6642.5, 6583.5(2,Z)$ to ground state, $(2)^-$ at 44.59, $(0)^-$ at 105.89										Ens075 **
* $^{220}\text{Ac}(\alpha)^{216}\text{Fr}$	$E_\alpha=7792, 7855$ to levels at 409.3, 349.3 keV										Ens075 **
* $^{220}\text{Pa}(\alpha)^{216}\text{Ac}$	Trends from Mass Surface TMS suggest $^{220}\text{Pa}$ 180 more bound										GAU **
$^{221}\text{Po}-u$	21238	62	21228	21	-0.2	o			GS3	1.0	10Ch19
	21228	21				2			GS3	1.0	12Ch19
$^{221}\text{At}-u$	18028	32	18017	15	-0.3	o			GS3	1.0	08Ch.A
	18017	15				2			GS3	1.0	12Ch19
$^{221}\text{Fr}-^{226}\text{Ra}_{978}$	-10590	34	-10596	5	-0.2	U			MA3	1.0	92Bo28
$^{211}\text{Fr}-^{221}\text{Fr}_{159}$	-3080	60	-3081	17	0.0	U			P24	2.5	82Au01
$^{221}\text{Rn}(\alpha)^{217}\text{Po}$	6161.8	5.8	6163	3	0.2	3			Dbb		77Vy02 *
	6163.5	5.4			-0.1	3			Orm		97Li23 *
	6163.5	5.4			-0.1	3			Orm		04Li28 *
$^{221}\text{Fr}(\alpha)^{217}\text{At}$	6457.3	2.0	6457.7	1.4	0.2	-			Orm		62Wa28 *
	6458.5	2.0			-0.4	-			Orm		68Le07 *
	ave.	6457.9	1.4		-0.1	1	99	78 $^{217}\text{At}$			average
$^{221}\text{Ra}(\alpha)^{217}\text{Rn}$	6883.7	5.	6880.4	2.0	-0.7	3					61Ru06 *
	6881.3	3.			-0.3	3					95Ch74 *
	6878.3	3.			0.7	3					97Li12 *
$^{221}\text{Ac}(\alpha)^{217}\text{Fr}$	7786.2	10.	7780	50	-0.1	4					70Bo13
	7782.1	5.			0.0	4			Lvn		87De.A
	7791.3	15.			-0.2	4			Dbb		92An.A
$^{221}\text{Th}(\alpha)^{217}\text{Ra}$	8628.5	5.	8626	4	-0.5	5					70To07 Z
	8626.0	10.			0.0	5					70Va13 Z
	8626.4	10.			-0.1	5			Dbb		90An19
	8614.2	10.			1.1	5			GSa		00He17
	8596.9	66.			0.4	U					05Li17

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{221}\text{Pa}(\alpha)^{217}\text{Ac}$	9247.7	30.				3					89Mi17
$^{221}\text{U}(\alpha)^{217}\text{Th}$	9889.3	50.9				3			GSa		15Kh09
* $^{221}\text{Rn}(\alpha)^{217}\text{Po}$	$E_\alpha=6035.3(3,Z)$ to a tentative $(11/2^+)$ level at 15(5)keV above $(9/2^+)$ ground state										
* $^{221}\text{Rn}(\alpha)^{217}\text{Po}$	$E_\alpha=6037(2)$ to a tentative $(11/2^+)$ level at 15(5) keV above $(9/2^+)$ ground state										
* $^{221}\text{Rn}(\alpha)^{217}\text{Po}$	$E_\alpha=6037(2)$ to a tentative $(11/2^+)$ level at 15(5) keV above $(9/2^+)$ ground state										
* $^{221}\text{Fr}(\alpha)^{217}\text{At}$	$E_\alpha=6341.1(2,Z), 6125.1(3,Z)$ to ground state, $5/2^-$ level at 218.12 keV										
* $^{221}\text{Fr}(\alpha)^{217}\text{At}$	$E_\alpha=6341.3(2,Z), 6127.2(3,Z)$ to ground state, $5/2^-$ level at 218.12 keV										
* $^{221}\text{Ra}(\alpha)^{217}\text{Rn}$	$E_\alpha=6761.2\ 6668.2\ 6613.2\ 6591.2(5,Z)$ to ground state, levels at 88.9 149.18 174.3keV										
* $^{221}\text{Ra}(\alpha)^{217}\text{Rn}$	$E_\alpha=6610(3,Z)$ to 149.2 level										
* $^{221}\text{Ra}(\alpha)^{217}\text{Rn}$	$E_\alpha=6754, 6662, 6607(\dots)$ to ground state, 93.02, 149.2 level										
$^{222}\text{Po}-u$	24133	72	24140	40	0.1	o			GS3	1.0	10Ch19
	24140	43				2			GS3	1.0	12Ch19
$^{222}\text{At}-u$	22459	32	22494	17	1.1	o			GS3	1.0	08Ch.A
	22494	17				2			GS3	1.0	12Ch19
$^{222}\text{Fr}-^{133}\text{Cs}_{1.669}$	175383.3	8.0				2			MA8	1.0	14Kr09
$^{222}\text{Fr}-^{226}\text{Ra}_{.982}$	-7410	25	-7368	8	1.7	U			MA3	1.0	92Bo28
$^{213}\text{Fr}-^{222}\text{Fr}_{.240}\ ^{210}\text{Fr}_{.761}$	-4810	60	-4947	13	-0.9	U			P24	2.5	82Au01
$^{213}\text{Fr}-^{222}\text{Fr}_{.096}\ ^{212}\text{Fr}_{.904}$	-1940	60	-1947	9	0.0	U			P24	2.5	82Au01
$^{221}\text{Fr}-^{222}\text{Fr}_{.498}\ ^{220}\text{Fr}_{.502}$	-610	90	-643	6	-0.1	U			P34	2.5	86Au02
$^{222}\text{Rn}(\alpha)^{218}\text{Po}$	5590.39	0.3	5590.4	0.3	0.0	1	100	99 $^{218}\text{Po}$			71Gr17 Z
$^{222}\text{Ra}(\alpha)^{218}\text{Rn}$	6680.0	5.	6678	4	-0.4	1	69	62 $^{222}\text{Ra}$			56As38 Z
$^{222}\text{Ac}(\alpha)^{218}\text{Fr}$	7137.5	2.				4			Bka		82Bo04 Z
$^{222}\text{Ac}^m(\alpha)^{218}\text{Fr}^p$	7140.3	20.				5					72Es03 *
$^{222}\text{Th}(\alpha)^{218}\text{Ra}$	8127.7	10.	8127	5	-0.1	4					70To07
	8130.7	8.				4					70Va13
	8126.7	15.				4			Dbb		92An.A
	8120.6	10.				4			GSa		00He17
	8116.4	48.				U					05Li17
$^{222}\text{Pa}(\alpha)^{218}\text{Ac}^m$	8697.0	30.	8736	13	1.3	7					70Bo13
	8745.5	15.				7					95Ho.C *
$^{222}\text{U}(\alpha)^{218}\text{Th}$	9481.1	50.9				6			GSa		15Kh09
* $^{222}\text{Ac}^m(\alpha)^{218}\text{Fr}^p$	$E_\alpha=7011.4(20,Z)$ not to ground state										
* $^{222}\text{Pa}(\alpha)^{218}\text{Ac}^m$	$E_\alpha=8210(15)$ to $^{218}\text{Ac}^n$ at 384.49 keV above $^{218}\text{Ac}^m$										
$^{223}\text{At}-u$	25172	32	25151	15	-0.7	o			GS3	1.0	08Ch.A
	25151	15				2			GS3	1.0	12Ch19
$^{223}\text{Rn}-^{133}\text{Cs}_{1.677}$	180453	11	180446	8	-0.6	1	58	58 $^{223}\text{Rn}$	MA8	1.0	09Ne03
$^{223}\text{Rn}-u$	21899	32	21889	8	-0.3	o			GS3	1.0	08Ch.A
	21880	13				1	42	42 $^{223}\text{Rn}$	GS3	1.0	12Ch19
$^{213}\text{Fr}-^{223}\text{Fr}_{.087}\ ^{212}\text{Fr}_{.913}$	-1900	60	-1942	9	-0.3	U			P24	2.5	82Au01
$^{222}\text{Fr}-^{223}\text{Fr}_{.498}\ ^{221}\text{Fr}_{.502}$	790	100	558	8	-0.9	U			P34	2.5	86Au02
$^{223}\text{Fr}(\alpha)^{219}\text{At}$	5431.6	80.	5561.4	2.8	1.6	U					55Ad10
	5562	3				1	85	79 $^{219}\text{At}$			01Li44
$^{223}\text{Ra}(\alpha)^{219}\text{Rn}$	5978.9	0.3	5978.99	0.21	0.3	-			Orm		62Wa18 *
	5979.1	0.3				-			BIP		71Gr17 *
	ave. 5979.00	0.21				1	100	96 $^{219}\text{Rn}$			average
$^{223}\text{Ac}(\alpha)^{219}\text{Fr}$	6783.2	1.0				4			Orm		69Le.A *
$^{223}\text{Th}(\alpha)^{219}\text{Ra}$	7602	23	7567	4	-1.5	U			ORa		69Ha32 *
	7589	14				U					70Va13 *
	7570	25				U					84Mi.A
	7568	10				5					87El02 *
	7567.4	10.				5			Dbb		90An19 *
	7566.1	5.				5					92Li09 *
$^{223}\text{Pa}(\alpha)^{219}\text{Ac}$	8345.0	10.	8330	50	-0.4	5					70Bo13
	8339.9	10.2				o			Dbb		89An.A
	8350.0	15.				U			Dbb		90An19
	8339.9	15.				U			GSa		95Ho.C
	8321.6	5.				5			Jya		99Ho28

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{223}\text{U}(\alpha)^{219}\text{Th}$	8940.7	40.7				5			Dbb		91An10
$^{223}\text{Fr}(\beta^-)^{223}\text{Ra}$	1170	10	1149.1	0.8	-2.1	U					75We23 *
* $^{223}\text{Ra}(\alpha)^{219}\text{Rn}$	$E_{\alpha}=5747.0(0.4,Z)$ , $5715.7(0.3,Z)$ , $5606.7(0.3,Z)$ keV										
*	to $11/2^+$ level at 126.77, $7/2^+$ at 158.64, $3/2^+$ at 269.48 keV										
* $^{223}\text{Ra}(\alpha)^{219}\text{Rn}$	$E_{\alpha}=5747.0(0.40,Z)$ , $5716.23(0.29,Z)$ , $5606.73(0.30,Z)$ keV										
*	to $11/2^+$ level at 126.77, $7/2^+$ at 158.64, $3/2^+$ at 269.48 keV										
* $^{223}\text{Ac}(\alpha)^{219}\text{Fr}$	$E_{\alpha}=6661.6, 6646.7, 6563.7(1.0,Z)$ to ground state, $5/2^+$ at 15.0, $7/2^-$ at 98.58										
* $^{223}\text{Th}(\alpha)^{219}\text{Ra}$	$E_{\alpha}=7330(20)$ to mixture of excited states at 138(10) keV										
* $^{223}\text{Th}(\alpha)^{219}\text{Ra}$	$E_{\alpha}=7317(10)$ to mixture of excited states at 138(10) keV										
* $^{223}\text{Th}(\alpha)^{219}\text{Ra}$	$E_{\alpha}=7324(10)$ to 113.8, 7285(10) 55% to 140.0, 26% to 152.0 level										
* $^{223}\text{Th}(\alpha)^{219}\text{Ra}$	$E_{\alpha}=7290(10)$ 55% to 140.0, 26% to 152.0 level										
* $^{223}\text{Th}(\alpha)^{219}\text{Ra}$	$E_{\alpha}=7318(5), 7293(5), 7281(5)$ to 113.8, 140.0, 152.0 levels										
* $^{223}\text{Fr}(\beta^-)^{223}\text{Ra}$	$E_{\beta^-}=1120(10)$ to $3/2^-$ level at 50.128 keV										
$^{224}\text{At}-u$	29744	63	29749	24	0.1	o			GS3	1.0	10Ch19
	29749	24				2			GS3	1.0	12Ch19
$^{224}\text{Rn}-^{133}\text{Cs}_{1.684}$	183304	16	183315	11	0.7	1	43	43 $^{224}\text{Rn}$	MA8	1.0	09Ne03
$^{224}\text{Rn}-u$	24073	32	24096	11	0.7	o			GS3	1.0	08Ch.A
	24104	14			-0.6	1	57	57 $^{224}\text{Rn}$	GS3	1.0	12Ch19
$^{224}\text{Fr}-u$	23399	32	23348	12	-1.6	o			GS3	1.0	08Ch.A
	23398	14			-3.6	B			GS3	1.0	12Ch19
$^{224}\text{Fr}-^{133}\text{Cs}_{1.684}$	182567	12				2			MA8	1.0	14Kr09
$^{224}\text{Ra}-^{133}\text{Cs}_{1.684}$	179430	30	179429.4	1.9	0.0	U			MA8	1.0	14Bo26
$^{223}\text{Fr}-^{224}\text{Fr}_{.747}$ $^{220}\text{Fr}_{.253}$	-620	70	-769	9	-0.9	U			P34	2.5	86Au02
$^{222}\text{Fr}-^{224}\text{Fr}_{.496}$ $^{220}\text{Fr}_{.505}$	10	70	-260#	50#	-1.5	U			P24	2.5	82Au01
$^{223}\text{Fr}-^{224}\text{Fr}_{.747}$ $^{220}\text{Fr}_{.253}$	-410	70	-840#	80#	-2.5	B			P24	2.5	82Au01
$^{223}\text{Fr}-^{224}\text{Fr}_{.664}$ $^{221}\text{Fr}_{.336}$	780	110	-520	8	-4.7	F			P34	2.5	86Au02 *
$^{223}\text{Fr}-^{224}\text{Fr}_{.664}$ $^{221}\text{Fr}_{.336}$	-110	70	-590#	70#	-2.7	B			P24	2.5	82Au01
$^{224}\text{Ra}(\alpha)^{220}\text{Rn}$	5788.93	0.15	5788.92	0.15	-0.1	1	100	69 $^{220}\text{Rn}$			71Gr17 Z
$^{224}\text{Ac}(\alpha)^{220}\text{Fr}$	6326.9	0.7				4			Orm		69Le.A *
$^{224}\text{Th}(\alpha)^{220}\text{Ra}$	7304.7	10.	7299	6	-0.6	4					61Ru06
	7304.7	10.			-0.6	4					70Va13
	7300.7	20.			-0.1	U			Dbb		89An13
	7286.4	10.			1.2	4			GSa		00He17
$^{224}\text{Pa}(\alpha)^{220}\text{Ac}$	7695.2	10.	7694	4	-0.2	6					70Bo13 *
	7692.6	10.			0.1	F			Dbb		90An19 *
	7680	15			0.9	U			GSa		95Ho.C
	7693.3	5.			0.1	6					96Li05 *
$^{224}\text{U}(\alpha)^{220}\text{Th}$	8624.3	15.	8628	7	0.3	6			Dbb		91An10
	8612.1	20.			0.8	6			ORa		92To02
	8631.9	8.1			-0.5	6			ORm		14Lo10 *
$^{224}\text{Fr}(\beta^-)^{224}\text{Ra}$	2830	50	2923	11	1.9	U					75We23 *
* $^{223}\text{Fr}-^{224}\text{Fr}_{.664}$ $^{221}\text{Fr}$ .	F : rejection based on line-shape analysis										
* $^{224}\text{Ac}(\alpha)^{220}\text{Fr}$	$E_{\alpha}=6213.8, 6207.0, 6141.7, 6059.8(0.7,Z)$ keV										
*	to ground state, $3^+$ at 6.92, $3^+$ at 72.99, $2^-$ at 156.82 keV										
* $^{224}\text{Pa}(\alpha)^{220}\text{Ac}$	$E_{\alpha}=7490(10)$ to $5^-$ level at 68.71 keV										
* $^{224}\text{Pa}(\alpha)^{220}\text{Ac}$	F : intensities in contradiction with reference										
* $^{224}\text{Pa}(\alpha)^{220}\text{Ac}$	$E_{\alpha}=7488(5), 7375(5)$ to $(5^-)$ level at 68.71 keV and 184.21 level										
* $^{224}\text{U}(\alpha)^{220}\text{Th}$	$E_{\alpha}=8479(8), 8095(11)$ to ground state, 386.7(1.8) $2^+$ level										
* $^{224}\text{Fr}(\beta^-)^{224}\text{Ra}$	$E_{\beta^-}=1780(50)$ to $1^-$ level at 1053.041 keV, and other $E_{\beta^-}$										
$^{225}\text{Rn}-^{133}\text{Cs}_{1.692}$	188484	23	188461	12	-1.0	1	27	27 $^{225}\text{Rn}$	MA8	1.0	09Ne03
$^{225}\text{Rn}-u$	28498	32	28486	12	-0.4	o			GS3	1.0	08Ch.A
	28477	14			0.6	1	73	73 $^{225}\text{Rn}$	GS3	1.0	12Ch19

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{225}\text{Fr}-u$	25574	14	25572	13	-0.1	1	84	84 $^{225}\text{Fr}$	GS3 1.0 12Ch19
$^{221}\text{Fr}-^{225}\text{Fr}_{.655} \ ^{213}\text{Fr}_{.346}$	-1110	60	-1096	9	0.1	U			P24 2.5 82Au01
$^{224}\text{Fr}^x-^{225}\text{Fr}_{.747} \ ^{221}\text{Fr}_{.253}$	50	80	700#	100#	3.2	B			P24 2.5 82Au01
$^{224}\text{Fr}^x-^{225}\text{Fr}_{.498} \ ^{223}\text{Fr}_{.502}$	190	80	760#	100#	2.8	B			P24 2.5 82Au01
$^{225}\text{Ra}(\alpha)^{221}\text{Rn}$	5096.7	5.1				2			00Li37
$^{225}\text{Ac}(\alpha)^{221}\text{Fr}$	5936.1	2.	5935.1	1.4	-0.5	-			Orm 67Ba51 Z
	5934.5	2.			0.3	-			67Dz02 Z
	ave. 5935.3	1.4			-0.1	1	99	79 $^{221}\text{Fr}$	average
$^{225}\text{Th}(\alpha)^{221}\text{Ra}$	6920.7	3.	6921.4	2.1	0.2	4			61Ru06 *
	6922.1	3.			-0.2	4			87Li.A *
$^{225}\text{Pa}(\alpha)^{221}\text{Ac}$	7381.5	20.	7390	50	0.2	U			ORa 68Ha14
	7376.4	10.			0.3	F			70Bo13 *
	7392.7	5.1				5			Lvn 87De.A
	7383.5	19.			0.2	U			00Sa52
$^{225}\text{U}(\alpha)^{221}\text{Th}$	8012.7	20.	8015	7	0.1	o			Dbb 89An13
	8022.9	20.			-0.4	6			GSa 89He13
	8021.9	15.			-0.5	6			ORa 92To02
	8012.7	20.4			0.1	6			Dbb 94Ye08
	8010	10			0.5	6			GSa 00He17 *
$^{225}\text{Np}(\alpha)^{221}\text{Pa}$	8786.5	20.				4			Dbb 94Ye08
$^{225}\text{Fr}(\beta^-)^{225}\text{Ra}$	1820	30	1828	12	0.3	1	16	16 $^{225}\text{Fr}$	75We23 *
$^{225}\text{Ra}(\beta^-)^{225}\text{Ac}$	360	10	356	5	-0.4	1	25	20 $^{225}\text{Ac}$	55Ma.A *
	360	30			-0.1	U			55Pe24 *
* $^{225}\text{Th}(\alpha)^{221}\text{Ra}$	$E_\alpha=6800.2, 6746.2, 6503.2, 6480.2, 6443.2(3,Z)$ keV								
*	to ground state, $7/2^+ 53.14, 7/2^+ 299.16, 3/2^+ 321.39, 5/2^+ 359.02$ levels								
* $^{225}\text{Th}(\alpha)^{221}\text{Ra}$	$E_\alpha=6799.3, 6745.3, 6504.3, 6483.3, 6447.3(3,Z)$ keV								
*	to ground state, $7/2^+ 53.14, 7/2^+ 299.16, 3/2^+ 321.39, 5/2^+ 359.02$ levels								
* $^{225}\text{Pa}(\alpha)^{221}\text{Ac}$	F : average of two branches								
* $^{225}\text{U}(\alpha)^{221}\text{Th}$	$E_\alpha=7868(15), 7621(15)$ to ground state, 250.9 level								
* $^{225}\text{Fr}(\beta^-)^{225}\text{Ra}$	$E_{\beta^-}=1640(10). \ 28\%$ to $3/2^-$ level at 225.2								
* $^{225}\text{Ra}(\beta^-)^{225}\text{Ac}$	$E_{\beta^-}=320(10) \ 320(30)$ respectively, to $3/2^+$ level at 40.09 keV								
$^{226}\text{Rn}-^{133}\text{Cs}_{1.699}$	191490	17	191499	11	0.5	1	44	44 $^{226}\text{Rn}$	MA8 1.0 09Ne03
$^{226}\text{Rn}-u$	30864	32	30861	11	-0.1	o			GS3 1.0 08Ch.A
	30868	15			-0.4	1	56	56 $^{226}\text{Rn}$	GS3 1.0 12Ch19
$^{226}\text{Fr}-u$	29565	32	29545	7	-0.6	o			GS3 1.0 08Ch.A
	29566	13			-1.7	1	26	26 $^{226}\text{Fr}$	GS3 1.0 12Ch19
$^{226}\text{Fr}-^{133}\text{Cs}_{1.699}$	190173.9	7.8	190182	7	1.0	1	74	74 $^{226}\text{Fr}$	MA8 1.0 14Kr09
$^{133}\text{Cs}-^{226}\text{Ra}_{.588}$	-109487	9	-109488.2	1.2	-0.1	U			MA3 1.0 92Bo28
	-109499	13			0.8	U			MA4 1.0 99Am05
$^{223}\text{Fr}-^{226}\text{Fr}_{.493} \ ^{220}\text{Fr}_{.507}$	-800	80	-1007	4	-1.0	U			P24 2.5 82Au01
$^{225}\text{Fr}-^{226}\text{Fr}_{.796} \ ^{221}\text{Fr}_{.204}$	-570	100	-794	13	-0.9	U			P24 2.5 82Au01
$^{225}\text{Fr}-^{226}\text{Fr}_{.498} \ ^{224}\text{Fr}_{.502}$	-260	90	-850#	50#	-2.6	B			P24 2.5 82Au01
$^{226}\text{Ra}(\alpha)^{222}\text{Rn}$	4870.70	0.25	4870.70	0.25	0.0	1	100	99 $^{222}\text{Rn}$	71Gr17 Z
$^{226}\text{Ac}(\alpha)^{222}\text{Fr}$	5496.1	5.	5506	8	0.2	U			DbA 75Va.A Z
$^{226}\text{Th}(\alpha)^{222}\text{Ra}$	6448.5	3.0	6452.5	1.0	1.3	U			56As38 *
	6454.8	3.6			-0.6	U			75Va.A *
	6452.6	1.0			-0.1	1	99	61 $^{226}\text{Th}$	12Ma30
$^{226}\text{Pa}(\alpha)^{222}\text{Ac}$	6986.9	10.				5			64Mc21
$^{226}\text{U}(\alpha)^{222}\text{Th}$	7747.4	30.	7701	4	-1.6	U			73Vi10 *
	7706.6	15.			-0.4	5			Dbb 90An22
	7701.6	5.			-0.1	5			Jya 99Gr28
	7691.4	10.			0.9	o			GSa 00He17
	7696.5	10.			0.4	5			GSa 01Ca.B
	7696.4	20.4			0.2	5			Rla 16Ka13
$^{226}\text{Np}(\alpha)^{222}\text{Pa}$	8189.2	20.4	8200	50	0.2	8			GSa 90Ni05
	8205.5	20.			-0.2	8			Dbb 94Ye08

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{226}\text{Ra}(\text{p,t})^{224}\text{Ra}$	-2816	15	-2818.9	1.9	-0.2	U			ANL		74Fr01	
$^{226}\text{Ra}(\text{d,t})^{225}\text{Ra}$	-146	10	-139.4	2.9	0.7	U					83Ny01	
$^{226}\text{Fr}(\beta^-)^{226}\text{Ra}$	3804	330	3853	7	0.1	U					75We23 *	
	3704	100			1.5	U					87Ve.A *	
$^{226}\text{Ac}(\beta^-)^{226}\text{Th}$	1115	7	1112	5	-0.5	-					68Va17 *	
	ave.	1115	6		-0.5	1	52	39	$^{226}\text{Th}$		average	
* $^{226}\text{Th}(\alpha)^{222}\text{Ra}$	$E_\alpha=6334.6(3,Z)$ , $6224.6(3,Z)$ to ground state, $2^+$ level at 111.12 keV										Ens11b **	
* $^{226}\text{Th}(\alpha)^{222}\text{Ra}$	$E_\alpha=6337.1(1.0,Z)$ , $6233.6(1.0,Z)$ to ground state, $2^+$ level at 111.12 keV										Ens11b **	
* $^{226}\text{U}(\alpha)^{222}\text{Th}$	$E_\alpha=7430(30)$ to $2^+$ level at 183.3(0.3) keV										94Ye08 **	
* $^{226}\text{Fr}(\beta^-)^{226}\text{Ra}$	$E_{\beta^-}=3550(330)$ $3450(100)$ respectively, to $1^-$ level at 253.73 keV										Ens964 **	
* $^{226}\text{Ac}(\beta^-)^{226}\text{Th}$	$E_{\beta^-}=885(7)$ to $1^-$ level at 230.37 keV										Ens964 **	
$^{227}\text{Rn}-^{133}\text{Cs}_{1.707}$	196686	19	196698	15	0.6	1	63	63	$^{227}\text{Rn}$	MA8	1.0	09Ne03
$^{227}\text{Rn}-\text{u}$	35288	33	35304	15	0.5	o			GS3	1.0	08Ch.A	
	35325	25			-0.8	1	37	37	$^{227}\text{Rn}$	GS3	1.0	12Ch19
$^{227}\text{Fr}-\text{u}$	31868	32	31865	6	-0.1	o			GS3	1.0	08Ch.A	
	31869	14			-0.3	1	20	20	$^{227}\text{Fr}$	GS3	1.0	12Ch19
$^{227}\text{Fr}-^{133}\text{Cs}_{1.707}$	193258.0	7.1	193259	6	0.1	1	80	80	$^{227}\text{Fr}$	MA8	1.0	14Kr09
$^{225}\text{Fr}-^{227}\text{Fr}_{.708}$ $^{220}\text{Fr}_{.292}$	-410	130	-547	13	-0.4	U			P24	2.5	82Au01	
$^{224}\text{Fr}^x-^{227}\text{Fr}_{.493}$ $^{221}\text{Fr}_{.507}$	-220	80	480#	100#	3.5	B			P24	2.5	82Au01	
$^{227}\text{Ac}(\alpha)^{223}\text{Fr}$	5043.0	2.0	5042.27	0.14	-0.4	U					66Ba19 Z	
	5042.27	0.14			0.0	1	100	94	$^{223}\text{Fr}$			86Ry04 Z
$^{227}\text{Th}(\alpha)^{223}\text{Ra}$	6146.60	0.10	6146.60	0.10	0.0	1	100	96	$^{223}\text{Ra}$	BIP		71Gr17 *
$^{227}\text{Pa}(\alpha)^{223}\text{Ac}$	6581.5	3.	6580.4	2.1	-0.4	5					63Su.A *	
	6579.3	3.			0.4	5					90Sh15 *	
$^{227}\text{U}(\alpha)^{223}\text{Th}$	7230	30	7235	3	0.2	U			ORa			69Ha32 *
	7206	16			1.8	U						91Ho05
	7234.7	3.1				6			GSa			15Ka24
$^{227}\text{Np}(\alpha)^{223}\text{Pa}$	7818.0	10.	7816	14	0.0	o			Dbb			90An19
	7815.0	20.			0.1	6			GSa			90Ni05
	7818.0	20.			-0.1	6			Dbb			94Ye08
$^{226}\text{Ra}(\text{n},\gamma)^{227}\text{Ra}$	4561.43	0.27				2			ILn			81Vo03 Z
$^{227}\text{Fr}(\beta^-)^{227}\text{Ra}$	2476	100	2505	6	0.3	U						75We23 *
$^{227}\text{Ra}(\beta^-)^{227}\text{Ac}$	1345	20	1328.1	2.3	-0.8	U						53Bu63 *
	1335	15			-0.5	U						71Lo15 *
$^{227}\text{Ac}(\beta^-)^{227}\text{Th}$	45.5	1.0	44.8	0.8	-0.7	-						55Be20
	43.5	1.5			0.8	-						59No41
	ave.	44.9	0.8		-0.2	1	99	96	$^{227}\text{Th}$			average
* $^{227}\text{Th}(\alpha)^{223}\text{Ra}$	$E_\alpha=6038.01(0.15,Z)$ , $5977.72(0.10,Z)$ , $5756.89(0.15,Z)$ keV to ground state, $7/2^+$ at 61.424, $1/2^+$ at 286.182 keV										71Gr17 **	
*											Ens01a **	
* $^{227}\text{Pa}(\alpha)^{223}\text{Ac}$	$E_\alpha=6465.8(3,Z)$ , $6423.8(3,Z)$ , $6415.8(3,Z)$ , $6401.7(3,Z)$ , $6356.7(3,Z)$ to ground state, $7/2^-$ at 42.4, $5/2^-$ at 50.7, $5/2^+$ at 64.62, $7/2^+$ at 110.06										63Su.A **	
*											Ens01a **	
* $^{227}\text{Pa}(\alpha)^{223}\text{Ac}$	$E_\alpha=6463$ , $6421$ , $6355$ keV (all errors 3 keV, estimated by evaluator) to ground state, $7/2^-$ at 42.4, $5/2^-$ at 50.7, $7/2^+$ at 110.06 keV										90Sh15 **	
*											Ens01a **	
* $^{227}\text{U}(\alpha)^{223}\text{Th}$	$E_\alpha=6860(30)$ to $3/2^+$ level at 247(1) keV										Ens01a **	
* $^{227}\text{Fr}(\beta^-)^{227}\text{Ra}$	$E_{\beta^-}=1800(100)$ to $1/2^-$ level at 675.863 keV										Ens162 **	
* $^{227}\text{Ra}(\beta^-)^{227}\text{Ac}$	$E_{\beta^-}=1310(20)$ $1300(15)$ respectively, to $3/2^+$ level at 27.369 and $5/2^+$ at 46.354										Ens162 **	
$^{228}\text{Rn}-^{133}\text{Cs}_{1.714}$	199897	24	199891	19	-0.3	1	63	63	$^{228}\text{Rn}$	MA8	1.0	09Ne03
$^{228}\text{Rn}-\text{u}$	37856	33	37835	19	-0.6	o			GS3	1.0	08Ch.A	
	37825	31			0.3	1	37	37	$^{228}\text{Rn}$	GS3	1.0	12Ch19
$^{228}\text{Fr}-\text{u}$	35833	34	35839	7	0.2	o			MA8	1.0	11Kr.A	
	35852	32			-0.4	o			GS3	1.0	08Ch.A	
	35821	16			1.2	1	20	20	$^{228}\text{Fr}$	GS3	1.0	12Ch19

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{228}\text{Fr}-^{133}\text{Cs}_{1.714}$	197899.5	8.1	197895	7	-0.6	1	80	80 $^{228}\text{Fr}$	MA8	1.0	14Kr09
$^{224}\text{Fr}^* - ^{228}\text{Fr}_{.491} \ ^{220}\text{Fr}_{.509}$	-540	320	-390#	100#	0.2	U			P24	2.5	82Au01
$^{228}\text{Th}(\alpha)^{224}\text{Ra}$	5520.17	0.22	5520.15	0.22	-0.1	1	100	69 $^{224}\text{Ra}$			71Gr17 Z
$^{228}\text{Pa}(\alpha)^{224}\text{Ac}$	6266.7	3.	6264.5	1.5	-0.7	5					58Hi.A *
	6264.7	3.			-0.1	5					93Sh07 *
	6263.5	2.			0.5	5					94Ah03 *
$^{228}\text{U}(\alpha)^{224}\text{Th}$	6803.6	10.				5					61Ru06
$^{228}\text{Np}(\alpha)^{224}\text{Pa}$	7308.5	36.				7			JAA		03Ni10
$^{228}\text{Pu}(\alpha)^{224}\text{U}$	7949.7	20.	7940	18	-0.5	7			DBb		94An02
	7911.0	35.			0.8	7			JAA		03Ni10
$^{228}\text{Ra}(\beta^-)^{228}\text{Ac}$	46.7	2.	45.5	0.6	-0.6	3					61To10 *
	45.5	0.9			0.0	3					72He.A *
	45.3	1.0			0.2	3					95So11 *
$^{228}\text{Ac}(\beta^-)^{228}\text{Th}$	2240	20	2123.7	2.6	-5.8	B					53Ky19 *
	2158	20			-1.7	U					57Bj56 *
$^{228}\text{Pa}(\epsilon)^{228}\text{Th}$	2109	15	2153	4	2.9	B					73Ku09 *
$^{228}\text{Pa}(\alpha)^{224}\text{Ac}$	$E_{\alpha}=6119.2(3,Z), 6106.2(3,Z), 6079.2(3,Z)$ to 37.2, 51.9, 78.4 levels										
$^{228}\text{Pa}(\alpha)^{224}\text{Ac}$	$E_{\alpha}=6118(3)$ to 37.2 level										
$^{228}\text{Pa}(\alpha)^{224}\text{Ac}$	$E_{\alpha}=6117(2)$ to 37.1 level										
$^{228}\text{Ra}(\beta^-)^{228}\text{Ac}$	$E_{\beta^-}=40(2) 39(1)$ respectively, to $1^-$ level at 6.28 keV, and other $E_{\beta^-}$										
$^{228}\text{Ra}(\beta^-)^{228}\text{Ac}$	$E_{\beta^-}=39.0(1.0)$ to $1^+$ level at 6.28 keV										
$^{228}\text{Ac}(\beta^-)^{228}\text{Th}$	$E_{\beta^-}=2180(20)$ to $2^+$ level at 57.773 keV, and other $E_{\beta^-}$										
$^{228}\text{Ac}(\beta^-)^{228}\text{Th}$	$E_{\beta^-}=2100(20), 1760, 1180$ to $2^+$ at 57.773, $3^-$ at 396.094, $2^+$ at 968.984										
$^{228}\text{Pa}(\epsilon)^{228}\text{Th}$	$pK=0.33(0.08)$ to $3^+$ level at 1944.904 keV, recalculated										
$^{229}\text{Rn}-^{133}\text{Cs}_{1.722}$	205069	14				2			MA8	1.0	09Ne03
$^{229}\text{Fr}-^{133}\text{Cs}_{1.722}$	201262	40	201103	5	-4.0	B			MA8	1.0	08We02 *
	201104.5	6.4			-0.2	1	70	70 $^{229}\text{Fr}$	MA8	1.0	14Kr09
$^{229}\text{Fr}-u$	38343	32	38291	5	-1.6	o			GS3	1.0	08Ch.A
	38298	15			-0.4	1	13	13 $^{229}\text{Fr}$	GS3	1.0	12Ch19
$^{229}\text{Fr}-^{238}\text{U}_{.962}$	-10576	13	-10566	6	0.8	1	18	17 $^{229}\text{Fr}$	MA8	1.0	14Kr09
$^{229}\text{Ra}-^{133}\text{Cs}_{1.722}$	197782	21	197768	17	-0.6	2			MA8	1.0	08We02
	197746	27			0.8	2			MA8	1.0	05He26
$^{229}\text{Ac}-u$	32947	13				2			GS3	1.0	12Ch19
$^{229}\text{Th}(\alpha)^{225}\text{Ra}$	5167.4	1.2	5167.6	1.0	0.1	-			Kum		71Bb10 *
	5168.2	2.			-0.3	-					87He28 Z
	ave.	5167.6	1.0		-0.1	1	99	95 $^{225}\text{Ra}$			average
$^{229}\text{Pa}(\alpha)^{225}\text{Ac}$	5835.6	5.	5835	4	-0.2	1	73	60 $^{225}\text{Ac}$			63Su.A *
$^{229}\text{U}(\alpha)^{225}\text{Th}$	6475.5	3.				5					61Ru06 Z
$^{229}\text{Np}(\alpha)^{225}\text{Pa}$	7012.7	20.	7010	50	0.0	6			ORa		68Ha14
	7015.8	23.			0.0	6					00Sa52
$^{229}\text{Pu}(\alpha)^{225}\text{U}$	7592.9	30.	7590	50	0.0	7			DBb		94An02
	7598.0	10.			-0.1	o			GSa		01Ca.B
	7589.8	20.			0.0	7			GSa		10Kh06
$^{229}\text{Am}(\alpha)^{225}\text{Np}$	8137.4	20.4				5			GSa		15De22 *
$^{229}\text{Ra}(\beta^-)^{229}\text{Ac}$	1760	40	1872	20	2.8	B					75We23
$^{229}\text{Ac}(\beta^-)^{229}\text{Th}$	1140	150	1104	12	-0.2	U					73Ch24
	1090	50			0.3	U					75We23
$^{229}\text{Fr}-^{133}\text{Cs}_{1.722}$	Could be influenced by $^{229}\text{Rn}$ contaminant										
$^{229}\text{Th}(\alpha)^{225}\text{Ra}$	$E_{\alpha}=4978.3(1.2,Z), 4967.3(1.2,Z), 4845.1(1.2,Z)$ keV										
*	to 100.60, 111.60, 236.25 levels										
$^{229}\text{Th}(\alpha)^{225}\text{Ra}$	$E_{\alpha}=4979.3(2,Z), 4968.3(2,Z), 4845.1(2,Z)$ keV										
*	to $9/2^+$ level at 100.50, $7/2^+$ at 111.60, $5/2^+$ at 236.25 keV										
*	calibrated with 71BaB2 value for 4845 level										
$^{229}\text{Pa}(\alpha)^{225}\text{Ac}$	$E_{\alpha}=5670.2, 5630.2, 5615.2, 5580.2, 5536.2$ (all 3,Z) keV to										
*	$5/2^+ 64.70, 7/2^+ 105.06, 5/2^- 120.80, 5/2^+ 155.65, 7/2^+ 199.85$										
$^{229}\text{Am}(\alpha)^{225}\text{Np}$	$E_{\alpha}=7990(20)$ and $E_{\alpha}=8000(20)$										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{230}\text{Fr}-^{133}\text{Cs}_{1.729}$	205878	32	205864	7	-0.4	o			MA8	1.0	05He26
	205860.1	7.5			0.6	1	88	88 $^{230}\text{Fr}$	MA8	1.0	14Kr09
$^{230}\text{Fr}-u$	42401	34	42391	7	-0.3	o			GS3	1.0	08Ch.A
	42421	20			-1.5	1	12	12 $^{230}\text{Fr}$	GS3	1.0	12Ch19
$^{230}\text{Ra}-^{133}\text{Cs}_{1.729}$	200530	13	200528	11	-0.1	2			MA8	1.0	08We02
	200524	21			0.2	2			MA8	1.0	05He26
$^{230}\text{Ac}-u$	36328	32	36327	17	0.0	o			GS3	1.0	08Ch.A
	36327	17				2			GS3	1.0	12Ch19
$^{230}\text{Ra}-^{226}\text{Ra}_{1.018}$	11225	35	11189	11	-1.0	U			MA3	1.0	92Bo28
$^{230}\text{Th}(\alpha)^{226}\text{Ra}$	4770.1	1.5	4769.9	1.5	-0.2	1	99	98 $^{226}\text{Ra}$	Orm		66Ba14 Z
$^{230}\text{Pa}(\alpha)^{226}\text{Ac}$	5439.5	0.7	5439.4	0.7	-0.1	1	99	87 $^{226}\text{Ac}$	Orm		66Ba14 Z
$^{230}\text{U}(\alpha)^{226}\text{Th}$	5992.8	0.7	5992.5	0.5	-0.5	2			Orm		66Ba14 Z
	5992.1	0.7			0.5	2			Gea		12Ma30
$^{230}\text{Np}(\alpha)^{226}\text{Pa}$	6778.1	20.				6		ORa		68Ha14	
$^{230}\text{Pu}(\alpha)^{226}\text{U}$	7175.0	15.	7181	7	0.4	6			Dbb		90An22
	7180.1	17.			0.0	6			Jya		99Gr28
	7182.2	10.			-0.1	6			GSa		01Ca.B
	7185.1	20.4			-0.2	6			RIa		16Ka13
$^{230}\text{Th}(\text{p,t})^{228}\text{Th}$	-3550	15	-3569.2	1.1	-1.3	U			ANL		74Fr01
$^{230}\text{Th}(\text{p,t})^{228}\text{Th}-^{232}\text{Th}(\text{O})^{230}\text{Th}$	-493.5	1.0	-492.6	0.5	0.9	o					91Gr13
	-492.5	0.5			-0.2	1	98	69 $^{228}\text{Th}$			94Le22
$^{230}\text{Th}(\text{p,t})^{228}\text{Th}-^{184}\text{W}(\text{O})^{182}\text{W}$	1564.0	1.6	1550.9	1.2	-8.2	B					09Le03
	1564.0	1.8			-7.3	C					09Le.A
$^{230}\text{Th}(\text{d,t})^{229}\text{Th}$	-541	6	-537.1	2.2	0.7	-					90Bu17
	-525	6			-2.0	-			ANL		67Er02 *
	ave.	-533	4		-1.0	1	27	26 $^{229}\text{Th}$			average
$^{230}\text{Ra}(\beta^-)^{230}\text{Ac}$	710	300	678	19	-0.1	U					80Gi04 *
$^{230}\text{Ac}(\beta^-)^{230}\text{Th}$	2700	100	2976	16	2.8	U					80Gi04 *
$^{230}\text{Pa}(\epsilon)^{230}\text{Th}$	1310.3	3.	1311.0	2.8	0.2	1	89	88 $^{230}\text{Pa}$			70Lo02 *
$^{230}\text{Pa}(\beta^-)^{230}\text{U}$	561	15	559	5	-0.2	R					70Lo02
$^{230}\text{Th}(\text{d,t})^{229}\text{Th}$	$Q = -525(6)$ to $^{229}\text{Th}^m$ at 0.0035(0.0010) keV										
$^{230}\text{Ra}(\beta^-)^{230}\text{Ac}$	$E_{\beta^-} = 500(200)$ to level at 211.78 keV										
$^{230}\text{Ac}(\beta^-)^{230}\text{Th}$	$E_{\beta^-} = 1400(100)$ to $0^+$ level at 1297.14 keV										
$^{230}\text{Pa}(\epsilon)^{230}\text{Th}$	$pK = 0.42(0.01)$ to $3^-$ level at 1127.789, recalculated										
$^{231}\text{Fr}-u$	45191	39	45175	8	-0.4	o			GS3	1.0	08Ch.A
	45158	27			0.6	U			GS3	1.0	12Ch19
$^{231}\text{Fr}-^{133}\text{Cs}_{1.737}$	209405.3	8.3				2			MA8	1.0	14Kr09
$^{231}\text{Ra}-^{133}\text{Cs}_{1.737}$	205267	21	205257	12	-0.5	1	34	34 $^{231}\text{Ra}$	MA8	1.0	05He26
	41052	32	41027	12	-0.8	o			GS3	1.0	08Ch.A
$^{231}\text{Ra}-u$	41022	15			0.3	1	66	66 $^{231}\text{Ra}$	GS3	1.0	12Ch19
	38404	32	38393	14	-0.3	o			GS3	1.0	08Ch.A
$^{231}\text{Ac}-u$	38393	14				2			GS3	1.0	12Ch19
	5150.2	1.5	5149.9	0.8	-0.2	o			Orm		66Ba14
$^{231}\text{Pa}(\alpha)^{227}\text{Ac}$	5146.9	1.0			3.0	B			Kum		68Ba25 *
	5150.7	1.5			-0.5	-			Orm		69Le.A *
	5149.8	1.0			0.1	-			Kum		76Ba99 *
	ave.	5150.1	0.8			-0.2	1	98	91 $^{227}\text{Ac}$		average
$^{231}\text{U}(\alpha)^{227}\text{Th}$	5551.3	50.	5576.3	1.7	0.5	U					53Cr.A
	5576.9	3.			-0.2	2					94Li12 *
	5576	2			0.1	2					97Mu08
$^{231}\text{Np}(\alpha)^{227}\text{Pa}$	6368.4	8.				6				73Ja06	
$^{231}\text{Pu}(\alpha)^{227}\text{U}$	6838.6	20.				7				99La14	
$^{231}\text{Pa}(\text{p,t})^{229}\text{Pa}$	-4133	3	-4133.3	2.8	-0.1	1	90	87 $^{229}\text{Pa}$	Mun		91Gr13 *
$^{230}\text{Th}(\text{n},\gamma)^{231}\text{Th}$	5118.00	0.20	5118.02	0.20	0.1	1	98	73 $^{231}\text{Th}$	ILn		87Wh01 Z

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{230}\text{Th}(d,p)^{231}\text{Th}$	2907	7	2893.45	0.20	-1.9	U			ANL		67Er02
$^{231}\text{Ac}(\beta^-)^{231}\text{Th}$	2100	100	1947	13	-1.5	U					60Ta19
$^{231}\text{Th}(\beta^-)^{231}\text{Pa}$	389.2	2.	391.5	1.5	1.1	1	53	47 $^{231}\text{Pa}$			75Ho14 *
* $^{231}\text{Pa}(\alpha)^{227}\text{Ac}$	$E_\alpha=5057.6(1.0,Z)$ , $4985.9(1.0,Z)$ , $4950.4(1.0,Z)$ to ground state, $7/2^-$ at 74.149 keV, and $9/2^+$ at 109.992 keV										Ens129 **
* $^{231}\text{Pa}(\alpha)^{227}\text{Ac}$	$E_\alpha=5015.9(1.5,Z)$ , $4735.9(1.5,Z)$ to $5/2^+$ at 46.354, $3/2^-$ at 330.040 keV										Ens129 **
* $^{231}\text{Pa}(\alpha)^{227}\text{Ac}$	$E_\alpha=4736.2(1.0,Z)$ to 330.040 level										Ens129 **
* $^{231}\text{U}(\alpha)^{227}\text{Th}$	$E_\alpha=5471(3)$ , $5456(3)$ , $5404(3)$ to 9.3, 24.4, 77.7 levels										94Li12 **
* $^{231}\text{Pa}(p,t)^{229}\text{Pa}$	$Q=-4145(3)$ to 11.6 level										98Le15 **
* $^{231}\text{Th}(\beta^-)^{231}\text{Pa}$	$E_{\beta^-}=305(2)$ to $5/2^+$ level at 84.2148 keV										Ens136 **
$^{232}\text{Fr}-^{133}\text{Cs}_{1.744}$	214353	15				2			MA8	1.0	14Kr09
$^{232}\text{Ra}-^{133}\text{Cs}_{1.744}$	208368	13	208367	10	-0.1	1	57	57 $^{232}\text{Ra}$	MA8	1.0	05He26
$^{232}\text{Ra}-u$	43518	32	43475	10	-1.3	o			GS3	1.0	08Ch.A
	43474	15			0.1	1	43	43 $^{232}\text{Ra}$	GS3	1.0	12Ch19
$^{232}\text{Ac}-u$	42052	32	42034	14	-0.6	o			GS3	1.0	08Ch.A
	42034	14				2			GS3	1.0	12Ch19
$\text{C}_{18}\text{H}_{16}-^{232}\text{Th}$	87142.4	2.	87146.8	1.5	0.9	U			M20	2.5	73Br06
$\text{C}_{24}\text{H}_{16}-^{232}\text{Th}$ $^{37}\text{Cl}$ $^{35}\text{Cl}$	152393.4	1.8	152391.5	1.5	-0.4	1	12	12 $^{232}\text{Th}$	M20	2.5	73Br06
$^{232}\text{Th}(\alpha)^{228}\text{Ra}$	4082.5	5.	4081.6	1.4	-0.2	U					57Ha08 Z
	4084.6	5.			-0.6	U					61Ko11 Z
	4083.5	5.			-0.4	U					62Ko12 Z
	4081.6	1.4				2					89Sa01 *
$^{232}\text{U}(\alpha)^{228}\text{Th}$	5413.63	0.09	5413.63	0.09	0.0	1	100	99 $^{232}\text{U}$	BIP		72Go33 *
$^{232}\text{Pu}(\alpha)^{228}\text{U}$	6716.0	10.				6					73Ja06
$^{232}\text{Th}(p,t)^{230}\text{Th}$	-3070	15	-3076.6	1.1	-0.4	U			ANL		74Fr01
$^{232}\text{Th}(p,t)^{230}\text{Th}-^{184}\text{W}()$ $^{182}\text{W}$	2056.4	1.6	2043.5	1.1	-8.1	B					09Le03
	2056.5	1.8			-7.2	B					09Le.A
$^{232}\text{Th}(d,t)^{231}\text{Th}$	-174	6	-183.2	1.1	-1.5	U			ANL		67Er02
	-187	10			0.4	U			MIT		72Gr19
$^{232}\text{Ac}(\beta^-)^{232}\text{Th}$	3700	100	3708	13	0.1	U					90Be.B
$^{232}\text{Pa}(\beta^-)^{232}\text{U}$	1344	20	1337	7	-0.3	2					63Bj01 *
	1336	8			0.1	2					71Ka42 *
* $^{232}\text{Th}(\alpha)^{228}\text{Ra}$	$E_\alpha=4012.3(1.4)$ , $3947.2(2.0)$ to ground state, $2^+$ level at 63.823 keV										Ens143 **
* $^{232}\text{U}(\alpha)^{228}\text{Th}$	$E_\alpha=5320.12(0.14,Z)$ , $5263.36(0.09,Z)$ to ground state, $2^+$ level at 57.773 level										Ens143 **
* $^{232}\text{Pa}(\beta^-)^{232}\text{U}$	$E_{\beta^-}=1295(20)$ to $2^+$ level at 47.573 keV, and other $E_{\beta^-}$										Ens06a **
* $^{232}\text{Pa}(\beta^-)^{232}\text{U}$	$E_{\beta^-}=314(8)$ to $2^-$ level at 1016.85 keV, and other $E_{\beta^-}$										Ens06a **
$^{233}\text{Fr}-^{133}\text{Cs}_{1.752}$	218166	21				2			MA8	1.0	14Kr09
$^{233}\text{Ra}-u$	47602	32	47595	9	-0.2	o			GS3	1.0	08Ch.A
	47582	17			0.7	1	30	30 $^{233}\text{Ra}$	GS3	1.0	12Ch19
$^{233}\text{Ra}-^{133}\text{Cs}_{1.752}$	213248	11	213243	9	-0.5	1	70	70 $^{233}\text{Ra}$	MA8	1.0	14Kr09
$^{233}\text{Ac}-u$	44363	32	44346	14	-0.5	o			GS3	1.0	08Ch.A
	44346	14				2			GS3	1.0	12Ch19
$^{233}\text{U}(\alpha)^{229}\text{Th}$	4908.4	1.2	4908.7	1.2	0.2	1	93	70 $^{229}\text{Th}$	Kum		68Ba25 Z
$^{233}\text{Np}(\alpha)^{229}\text{Pa}$	5626.7	50.9				2					50Ma14
$^{233}\text{Pu}(\alpha)^{229}\text{U}$	6416.3	20.				6					57Th10
$^{233}\text{Am}(\alpha)^{229}\text{Np}^p$	6898.6	17.3				8					00Sa52
$^{233}\text{Cm}(\alpha)^{229}\text{Pu}$	7468.5	10.	7470	50	0.1	o			GSa		01Ca.B
	7473.5	20.				8			GSa		10Kh06
$^{233}\text{Bk}(\alpha)^{229}\text{Am}^p$	7905.9	20.3				7			GSa		15De22
$^{232}\text{Th}(n,\gamma)^{233}\text{Th}$	4786.69	0.25	4786.39	0.09	-1.2	-					74Ke13 Z
	4786.34	0.10			0.5	-			Bdn		06Fi.A
$^{232}\text{Th}(d,p)^{233}\text{Th}$	2555	10	2561.82	0.09	0.7	U			MIT		72Gr19
	2567	7			-0.7	U			ANL		72Vo08
$^{232}\text{Th}(n,\gamma)^{233}\text{Th}$	ave.	4786.39	0.09	4786.39	0.09	0.0	1	100	92 $^{233}\text{Th}$		average



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{233}\text{Th}(\beta^-)^{233}\text{Pa}$	1245	3	1242.2	1.1	-0.9	1	14	8	$^{233}\text{Th}$		57Fr.A *
$^{233}\text{Pa}(\beta^-)^{233}\text{U}$	568	4	570.3	2.0	0.6	-					54Br37 *
	568	5			0.5	-					55On05 *
	566	5			0.9	-					63Bi03 *
ave.	567.4	2.6			1.1	1	56	51	$^{233}\text{U}$		average
$^{*233}\text{Th}(\beta^-)^{233}\text{Pa}$	PrvCom to reference										
$^{*233}\text{Pa}(\beta^-)^{233}\text{U}$	$E_{\beta^-}=568(5)$ , 256(4) to ground state, $3/2^+$ level at 311.904 keV										
$^{*233}\text{Pa}(\beta^-)^{233}\text{U}$	$E_{\beta^-}=568(5)$ , 257(5) to ground state, $3/2^+$ level at 311.904 keV										
$^{*233}\text{Pa}(\beta^-)^{233}\text{U}$	$E_{\beta^-}=254(5)$ to $3/2^+$ level at 311.904 keV										
	Ens057 **										
	Ens057 **										
	Ens057 **										
$^{234}\text{Ra}-u$	50358	33	50382	9	0.7	o			GS3	1.0	08Ch.A
	50342	33			1.2	U			GS3	1.0	12Ch19
$^{234}\text{Ra}-^{133}\text{Cs}_{1.759}$	216692.1	9.0				2			MA8	1.0	14Kr09
$^{234}\text{Ac}-u$	48137	32	48139	15	0.1	o			GS3	1.0	08Ch.A
	48139	15				2			GS3	1.0	12Ch19
$^{234}\text{U}(\alpha)^{230}\text{Th}$	4857.4	1.0	4857.5	0.7	0.1	-					55Go.A Z
	4860.4	2.			-1.4	-			Kum		67Ba43 Z
ave.	4858.0	0.9			-0.5	1	55	39	$^{230}\text{Th}$		average
$^{234}\text{Pu}(\alpha)^{230}\text{U}$	6310.1	5.				3					60Ho.A *
$^{234}\text{Am}(\alpha)^{230}\text{Np}$	6572.6	20.	6800#	150#	11.4	F					90Ha02 *
$^{234}\text{Cm}(\alpha)^{230}\text{Pu}$	7365.2	10.	7365	9	0.0	7			GSa		01Ca.B
	7366.1	20.3			0.0	7			RIa		16Ka13
$^{234}\text{Bk}(\alpha)^{230}\text{Am}$	8087	50	8100	50	0.2	o			RIa		02Mo.B *
	8098.6	20.3				10			RIa		16Ka13
$^{232}\text{Th}(t,p)^{234}\text{Th}$	2487	20	2494.6	2.6	0.4	U			LAl		69Br11
$^{234}\text{U}(p,t)^{232}\text{U}$	-4099	15	-4125.3	1.6	-1.8	U			ANL		74Fr01
$^{234}\text{U}(p,t)^{232}\text{U}-^{184}\text{W}(l)^{182}\text{W}$	1007.6	1.6	994.8	1.6	-8.0	B					09Le03
	1007.6	1.8			-7.1	C					09Le.A
$^{233}\text{U}(d,p)^{234}\text{U}$	4656	15	4620.9	2.0	-2.3	U			Kop		68Bj05
$^{234}\text{U}(d,t)^{233}\text{U}$	-579	6	-588.2	2.0	-1.5	1	11	11	$^{233}\text{U}$	ANL	67Er02
$^{234}\text{Th}(\beta^-)^{234}\text{Pa}^m$	192	2	195.1	1.0	1.5	3					55De40 *
	193	2			1.0	3					63Bj02 *
	198.	1.5			-1.9	3					73Go40 *
$^{234}\text{Pa}^m(IT)^{234}\text{Pa}$	79	3				4					Nub16b
$^{234}\text{Pa}(\beta^-)^{234}\text{U}$	2230	40	2194	4	-0.9	U					62Bj01
$^{234}\text{Pa}^m(\beta^-)^{234}\text{U}$	2290	20	2272.9	2.6	-0.9	U					63Bj02
$^{234}\text{Np}(\beta^+)^{234}\text{U}$	1812	10	1810	8	-0.2	2					67Ha04 *
	1805	15			0.3	2					67Wa09 *
$^{*234}\text{Pu}(\alpha)^{230}\text{U}$	With correction similar to reference										
$^{*234}\text{Am}(\alpha)^{230}\text{Np}$	F : not believed to be measured in this work, replaced by estimate										
$^{*234}\text{Bk}(\alpha)^{230}\text{Am}$	$E_{\alpha}=7850(50)$ to 100 keV excited state										
$^{*234}\text{Th}(\beta^-)^{234}\text{Pa}^m$	$E_{\beta^-}=100(2)$ 100(2) 104.0(1.5) respectively, to $(1^-)$ 92.38 above $^{234}\text{Pa}^m$ , and other $E_{\beta^-}$										
$^{*234}\text{Np}(\beta^+)^{234}\text{U}$	$E_{\beta^+}=790(10)$ pK=0.48(0.03) respectively, to $1^+$ at 1570.69 and $1^+$ at 1601.8 keV										
	Ens074 **										
	Ens074 **										
$^{235}\text{Ac}-u$	50872	32	50840	15	-1.0	o			GS3	1.0	08Ch.A
	50840	15				2			GS3	1.0	12Ch19
$^{235}\text{Th}-u$	47252	32	47255	14	0.1	o			GS3	1.0	08Ch.A
	47255	14				2			GS3	1.0	12Ch19
$^{235}\text{Pa}-u$	45421	32	45399	15	-0.7	o			GS3	1.0	08Ch.A
	45399	15				2			GS3	1.0	12Ch19
$^{235}\text{U}-^{206}\text{Pb C}_2 \text{H}_5$	30341.0	10.	30337.9	1.6	-0.1	U			C4	2.5	71Ke02
$^{235}\text{U}-\text{C}_{18} \text{H}_{18}$	-96932.8	3.8	-96922.4	1.2	1.1	U			M20	2.5	73Br06
$\text{C}_{18} \text{H}_{20}-^{235}\text{U}$	112584.2	4.8	112572.5	1.2	-1.0	U			M20	2.5	73Br06
$^{235}\text{U}(\alpha)^{231}\text{Th}$	4678	2	4678.0	0.7	0.0	-			Kum		60Ba44 *
	4681	3			-1.0	-					60Vo07 *
	4675.5	3.0			0.8	-					64Sc27 *
	4677	3			0.3	-					66Ga03 *
ave.	4677.9	1.3			0.1	1	28	21	$^{231}\text{Th}$		average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{235}\text{Np}(\alpha)^{231}\text{Pa}$	5197.2	2.0	5193.8	1.5	-1.7	1	54	42 $^{231}\text{Pa}$	Bka		73Br12 *
$^{235}\text{Pu}(\alpha)^{231}\text{U}$	5951.5	20.				3					57Th10
$^{235}\text{Am}(\alpha)^{231}\text{Np}$	6559	100	6576	13	0.2	o			JAa		99Sa.D *
	6576	15			0.0	o			JAa		04Sa05 *
	6576	13				7			JAa		04As12 *
$^{233}\text{U}(\text{t,p})^{235}\text{U}$	3668	10	3661.2	2.0	-0.7	U					67Ri.A *
$^{234}\text{U}(\text{n},\gamma)^{235}\text{U}$	5297.1	0.5	5297.50	0.23	0.8	-					72Ri08 Z
	5297.4	0.3			0.3	-					77Ko15 Z
$^{234}\text{U}(\text{d,p})^{235}\text{U}$	3075	7	3072.93	0.23	-0.3	U			ANL		70Br01
$^{235}\text{U}(\text{d,t})^{234}\text{U}$	935	15	959.73	0.23	1.6	U			Kop		68Bj05
$^{234}\text{U}(\text{n},\gamma)^{235}\text{U}$	ave.	5297.32	0.26	5297.50	0.23	0.7	1	81	63 $^{234}\text{U}$		average
$^{235}\text{Th}(\beta^-)^{235}\text{Pa}$	1470	80	1729	19	3.2	B					89Yu01
$^{235}\text{Pa}(\beta^-)^{235}\text{U}$	1410	50	1370	14	-0.8	U					68Tr07
$^{235}\text{Np}(\epsilon)^{235}\text{U}$	123.5	2.	124.3	0.9	0.4	-					58Gi05
	123.6	1.			0.7	-					72Mc25
	ave.	123.6	0.9		0.8	1	91	88 $^{235}\text{Np}$			average
* $^{235}\text{U}(\alpha)^{231}\text{Th}$	$E_\alpha=4596\ 4398\ 4372(\text{all } 2,Z)$ to ground state, $(7/2^-)$ at 205.310, $9/2^-$ at 236.906										
* $^{235}\text{U}(\alpha)^{231}\text{Th}$	$E_\alpha=4598.6(3,Z)$ , $4402.6(3,Z)$ to ground state, $(7/2^-)$ at 205.310 keV										
* $^{235}\text{U}(\alpha)^{231}\text{Th}$	$E_\alpha=4595\ 4394\ 4364(\text{all } 3,Z)$ to ground state, $(7/2^-)$ at 205.310, $9/2^-$ at 236.906										
* $^{235}\text{U}(\alpha)^{231}\text{Th}$	$E_\alpha=4595.3, 4397.3, 4365.3(\text{all } 3,Z)$ to ground state, $(7/2^-)$ level at 205.310 keV and $9/2^-$ at 236.906 keV										
* $^{235}\text{Np}(\alpha)^{231}\text{Pa}$	$E_\alpha=5105.2(3), 5097.2(3), 5050.8(2,Z), 5024.8(2,Z), 4924.8(2,Z)$ to ground state, $1/2^-$ at 9.206, $7/2^-$ at 58.5699, $5/2^+$ at 84.2148, $5/2^+$ at 183.4962										
* $^{235}\text{Am}(\alpha)^{231}\text{Np}$	$E_\alpha=6440(100)$ to level below 15 keV										
* $^{235}\text{Am}(\alpha)^{231}\text{Np}$	$E_\alpha=6457(14)$ to level below 15 keV										
* $^{235}\text{Am}(\alpha)^{231}\text{Np}$	$E_\alpha=6457(12)$ to level below 15 keV										
* $^{233}\text{U}(\text{t,p})^{235}\text{U}$	$Q=3335(10,\text{Ri})$ to $5/2^+$ level at 332.845 keV										
$^{236}\text{Ac}-\text{u}$	55037	73	54990	40	-0.7	o			GS3	1.0	10Ch19
	54988	41				2			GS3	1.0	12Ch19
$^{236}\text{Th}-\text{u}$	49665	32	49657	15	-0.3	o			GS3	1.0	08Ch.A
	49657	15				2			GS3	1.0	12Ch19
$^{236}\text{Pa}-\text{u}$	48666	32	48668	15	0.1	o			GS3	1.0	08Ch.A
	48668	15				2			GS3	1.0	12Ch19
$^{236}\text{U}(\alpha)^{232}\text{Th}$	4572.2	3.	4572.9	0.9	0.2	o					60Ko04 Z
	4570.0	3.1			1.0	-					61Ko11 Z
	4573.1	1.0			-0.2	-			Kum		78Ba.C
	ave.	4572.8	1.0		0.1	1	82	83 $^{232}\text{Th}$			average
$^{236}\text{Pu}(\alpha)^{232}\text{U}$	5867.15	0.08	5867.15	0.08	0.0	1	100	99 $^{236}\text{Pu}$			84Ry02 Z
$^{236}\text{Am}(\alpha)^{232}\text{Np}$	6256.2	40.				3			JAa		04Sa05
$^{236}\text{Cm}(\alpha)^{232}\text{Pu}$	7074.1	20.	7067	5	-0.4	U			GSa		10Kh06
	7066.9	5.				7			JAa		10As.A
$^{236}\text{U}(\text{p,t})^{234}\text{U}$	-3330	15	-3361.2	0.3	-2.1	U			ANL		74Fr01
$^{235}\text{U}(\text{n},\gamma)^{236}\text{U}$	6545	2	6545.52	0.26	0.3	U					70Ka22
	6545.1	0.5			0.8	-					74Ju.B Z
	6545.4	0.5			0.2	-					75We.A Z
$^{236}\text{U}(\text{d,t})^{235}\text{U}$	-281	6	-288.29	0.26	-1.2	U			ANL		70Br01
$^{235}\text{U}(\text{n},\gamma)^{236}\text{U}$	ave.	6545.3	0.4	6545.52	0.26	0.8	1	54	30 $^{235}\text{U}$		average
$^{236}\text{Pa}(\beta^-)^{236}\text{U}$	3350	100	2889	14	-4.6	B					63Wo04
	2900	200			-0.1	U					68Tr07 *
$^{236}\text{Np}^m(\text{IT})^{236}\text{Np}$	60	50				3					Ens06a
$^{236}\text{Np}^m(\beta^-)^{236}\text{Pu}$	525	10	537	6	1.2	2					56Gr11 *
	544	8			-0.9	2					69Le05 *
* $^{236}\text{Pa}(\beta^-)^{236}\text{U}$	$E_{\beta^-}=2000(200)$ to $1^-$ level at 687.59 keV, and other $E_{\beta^-}$ , reinterpreted										
* $^{236}\text{Np}^m(\beta^-)^{236}\text{Pu}$	$E_{\beta^-}=518(10)\ 537(8)$ respectively, to ground state and $2^+$ level at 44.63 keV										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{237}\text{Th}-u$	53690	32	53629	17	-1.9	o			GS3	1.0	08Ch.A
	53629	17				2			GS3	1.0	12Ch19
$^{237}\text{Pa}-u$	51038	32	51023	14	-0.5	o			GS3	1.0	08Ch.A
	51023	14				2			GS3	1.0	12Ch19
$^{237}\text{Np}(\alpha)^{233}\text{Pa}$	4959.9	3.	4957.3	0.7	-0.9	U					61Ba44 *
	4956.7	1.5			0.4	-			Kum		68Ba25 *
	4959.9	3.			-0.9	U					69Va06 *
	4956.9	0.9			0.4	-			Gea		00Si02 *
	ave.	4956.8	0.8			0.6	1	91	90 $^{233}\text{Pa}$		average
$^{237}\text{Pu}(\alpha)^{233}\text{U}$	5753.3	20.	5747.6	2.3	-0.3	U					57Th10
	5747	5			0.1	1	20	15 $^{233}\text{U}$	DbA		93Dm02
$^{237}\text{Am}(\alpha)^{233}\text{Np}^p$	6146.2	5.				4					75Ah05 Z
$^{237}\text{Cm}(\alpha)^{233}\text{Pu}$	6774.5	10.	6770	50	-0.1	o			JAa		02As08
	6770.4	10.				7			JAa		06As03
$^{237}\text{Cf}(\alpha)^{233}\text{Cm}$	8220	20				9			GSa		10Kh06
$^{235}\text{U}(\text{t,p})^{237}\text{U}$	3206	20	3189.5	0.5	-0.8	U			Ald		64Mi.A *
	3178	20			0.6	U			LAI		69Br11
$^{237}\text{Np}(\text{p,t})^{235}\text{Np}$	-3816	15	-3832.3	0.9	-1.1	U			ANL		74Fr01
$^{236}\text{U}(\text{n},\gamma)^{237}\text{U}$	5125.9	0.5	5125.8	0.5	-0.3	1	85	84 $^{237}\text{U}$	BNn		79Vo05 Z
$^{236}\text{U}(\text{d,p})^{237}\text{U}$	2898	8	2901.2	0.5	0.4	U			ANL		67Er02
$^{237}\text{Pa}(\beta^-)^{237}\text{U}$	2250	100	2137	13	-1.1	U					74Ka05
$^{237}\text{U}(\beta^-)^{237}\text{Np}$	520	5	518.5	0.5	-0.3	U					53Wa05 *
	524	5			-1.1	U					56Ba39 *
	523	5			-0.9	U					57Ra04 *
	222	8	220.1	1.3	-0.2	U					58Ho02 *
$^{237}\text{Pu}(\epsilon)^{237}\text{Np}$	207	18			0.7	U					59Gi54 *
	$E_\alpha=4876.7$ 4774.2 4769.1(3,Z) to ground state, $7/2^+$ at 103.635, $9/2^+$ 109.07 keV										Ens057 **
* $^{237}\text{Np}(\alpha)^{233}\text{Pa}$	$E_\alpha=4787.9$ (1.5,Z) to $5/2^+$ level at 86.48 keV										Ens057 **
* $^{237}\text{Np}(\alpha)^{233}\text{Pa}$	$E_\alpha=4791.0$ (3,Z), 4774.0(3,Z), 4770.0(3,Z) keV										69Va06 **
* $^{237}\text{Np}(\alpha)^{233}\text{Pa}$	to $5/2^+$ at 86.468, $7/2^+$ at 103.635, $9/2^+$ at 109.07 keV										Ens057 **
*											Ens057 **
* $^{237}\text{Np}(\alpha)^{233}\text{Pa}$	$E_\alpha=4788.0$ (0.9) keV to $5/2^+$ at 86.468 keV										00Si02 **
* $^{235}\text{U}(\text{t,p})^{237}\text{U}$	$Q=2980$ (20) to $7/2^+$ level at 426.15 keV										Ens068 **
* $^{237}\text{U}(\beta^-)^{237}\text{Np}$	$E_{\beta^-}=245$ (5), 249(5), 248(5) respectively, to 53% $3/2^-$ level at 267.556, and										Ens068 **
*	43% $1/2^-$ level at 281.356 keV										Ens068 **
* $^{237}\text{Pu}(\epsilon)^{237}\text{Np}$	LK=2.8(0.8) capture to $5/2^-$ level at 59.541 keV, recalculated										Ens068 **
* $^{237}\text{Pu}(\epsilon)^{237}\text{Np}$	pK=0.38(0.06) to ground state, $7/2^+$ level at 33.196, $5/2^-$ at 59.541 keV										Ens068 **
$^{238}\text{Pa}-u$	54648	32	54637	17	-0.3	o			GS3	1.0	08Ch.A
	54637	17				2			GS3	1.0	12Ch19
$^{238}\text{U}-^{206}\text{Pb}$ $^{32}\text{S}$	104253.9	10.	104250.7	1.9	-0.1	U			C4	2.5	71Ke02
$\text{C}_{18} \text{H}_{22}-^{238}\text{U}$	121366.0	2.4	121363.7	1.6	-0.4	U			M20	2.5	73Br06
$\text{C}_{24} \text{H}_{20}-^{238}\text{U}$ $^{35}\text{Cl}_2$	168010.8	1.4	168008.3	1.6	-0.7	1	21	21 $^{238}\text{U}$	M20	2.5	73Br06
$^{238}\text{U}-^{235}\text{U}$	6858.6	10.	6858.8	1.3	0.0	U			C4	2.5	71Ke02
$^{238}\text{U}(\alpha)^{234}\text{Th}$	4271.5	5.	4269.9	2.1	-0.3	2					57Ha08 Z
	4265.1	5.			0.9	2					60Vo07 Z
	4272.9	5.			-0.6	2					61Ko11 Z
	4269.9	3.			0.0	2			Gea		14Po02
$^{238}\text{Pu}(\alpha)^{234}\text{U}$	5593.20	0.2	5593.27	0.19	0.4	1	90	69 $^{238}\text{Pu}$			71Gr17 Z
$^{238}\text{Am}(\alpha)^{234}\text{Np}$	6041.7	30.				3					72Ah04
$^{238}\text{Cm}(\alpha)^{234}\text{Pu}$	6611.5	50.	6670	10	1.2	U					48St.A *
	6632.0	50.			0.8	U					52Hi.A
	6672.3	10.			-0.2	o			JAa		02As08
	6670.3	10.				4			JAa		06As03
$^{238}\text{U}(\text{n},\alpha)^{235}\text{Th}$	8700	50	8936	13	4.7	B					81Wa11
$^{236}\text{U}(\text{t,p})^{238}\text{U}$	2900	20	2797.7	1.2	-5.1	F			Ald		64Mi.A *
	2782	10			1.6	U			ANL		67Er02
	2780	20			0.9	U			LAI		69Br11

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{238}\text{U}(\text{p,t})^{236}\text{U}$	-2765	15	-2797.7	1.2	-2.2	U			ANL		74Fr01
$^{238}\text{U}(\text{p,d})^{237}\text{U}$	-3951	20	-3929.2	1.3	1.1	U			Ald		64Mi.A
$^{238}\text{U}(\text{d,t})^{237}\text{U}$	116	6	103.5	1.3	-2.1	U			ANL		67Er02
$^{237}\text{Np}(\text{n},\gamma)^{238}\text{Np}$	5488.32	0.20				2			BNn		79Io01 Z
$^{238}\text{Pu}(\text{d,t})^{237}\text{Pu}$	-746	10	-742.6	1.3	0.3	U			Kop		73Gr26
$^{238}\text{Pa}(\beta^-)^{238}\text{U}$	3600	300	3586	16	0.0	U					68Tr07 *
	3460	60			2.1	U					85Ba57 *
$^{238}\text{Np}(\beta^-)^{238}\text{Pu}$	1295	10	1291.4	0.5	-0.4	U					55Ra27 *
	1300	15			-0.6	U					56Ba95 *
* $^{238}\text{Cm}(\alpha)^{234}\text{Pu}$	PrvCom to reference										
* $^{236}\text{U}(\text{t,p})^{238}\text{U}$	F : authors not satisfied with target material										
* $^{238}\text{Pa}(\beta^-)^{238}\text{U}$	$E_{\beta^-}=1700(300)$ to $(3^-)$ level at 1992.2 keV, and other $E_{\beta^-}$ , reinterpreted										
* $^{238}\text{Pa}(\beta^-)^{238}\text{U}$	Reports result from thesis										
* $^{238}\text{Np}(\beta^-)^{238}\text{Pu}$	$E_{\beta^-}=270(10)$ 280(10) respectively, to $2^+$ level at 1028.537 keV, and other $E_{\beta^-}$										
$^{239}\text{Pu}(\alpha)^{235}\text{U}$	5244.60	0.25	5244.52	0.21	-0.3	1	68	41 $^{235}\text{U}$			79Ry.A *
$^{239}\text{Am}(\alpha)^{235}\text{Np}$	5924.6	2.0	5922.4	1.4	-1.1	2			Bka		71Go01 *
	5920.2	2.0			1.1	2					75Ah05 *
$^{239}\text{Cm}(\alpha)^{235}\text{Pu}$	6539.7	140.				4			JAA		02Sh.C *
$^{239}\text{Cf}(\alpha)^{235}\text{Cm}^p$	7760.1	25.				10			GSa		81Mu12 *
$^{238}\text{U}(\text{n},\gamma)^{239}\text{U}$	4806.55	0.30	4806.38	0.17	-0.6	2			ANL		72Bo46 Z
	4806.30	0.21			0.4	2			ILn		79Br25 Z
$^{238}\text{U}(\text{d,p})^{239}\text{U}$	2588	20	2581.82	0.17	-0.3	U			Ald		64Mi.A
	2579	7			0.4	U			Tal		66Sh16
	2585	6			-0.5	U			ANL		67Er02
$^{238}\text{Pu}(\text{n},\gamma)^{239}\text{Pu}$	5646.7	0.5	5646.2	0.3	-0.9	1	38	31 $^{238}\text{Pu}$			75Ma.A Z
$^{238}\text{Pu}(\text{d,p})^{239}\text{Pu}$	3432	10	3421.7	0.3	-1.0	U			Kop		73Gr26
$^{239}\text{Pu}(\text{d,t})^{238}\text{Pu}$	604	10	611.0	0.3	0.7	U			ANL		73Fr01
$^{239}\text{U}(\beta^-)^{239}\text{Np}$	1290	20	1261.7	1.5	-1.4	U					64B111 *
$^{239}\text{Np}(\beta^-)^{239}\text{Pu}$	722.5	1.0	722.8	0.9	0.3	1	87	67 $^{239}\text{Np}$			59Co63 *
* $^{239}\text{Pu}(\alpha)^{235}\text{U}$	$E_{\alpha}=5156.59(0.25,Z)$ to $1/2^+$ level at 0.0760 keV										
* $^{239}\text{Am}(\alpha)^{235}\text{Np}$	$E_{\alpha}=5824.6(4,Z)$ 5775.6(2,Z) 5733.6(2,Z); ground state, $(5/2)^-$ 49.10, $(7/2)^-$ 91.6										
* $^{239}\text{Am}(\alpha)^{235}\text{Np}$	$E_{\alpha}=5772.7(2,Z)$ to $(5/2)^-$ level at 49.10 keV										
* $^{239}\text{Cm}(\alpha)^{235}\text{Pu}$	Private communication to reference										
* $^{239}\text{U}(\beta^-)^{239}\text{Np}$	$E_{\beta^-}=1211(20)$ to $5/2^-$ level at 74.6640 keV, and other $E_{\beta^-}$										
* $^{239}\text{Np}(\beta^-)^{239}\text{Pu}$	$E_{\beta^-}=437(1)$ to $5/2^+$ level at 285.460 keV, and other $E_{\beta^-}$										
$^{240}\text{Pu}(\alpha)^{236}\text{U}$	5255.88	0.15	5255.82	0.14	-0.4	1	90	77 $^{236}\text{U}$			72Go33 Z
$^{240}\text{Am}(\alpha)^{236}\text{Np}^p$	5468.9	1.0				3					70Go42 Z
$^{240}\text{Cm}(\alpha)^{236}\text{Pu}$	6397.8	0.6	6397.8	0.6	0.0	1	100	99 $^{240}\text{Cm}$	Kum		71Bb10 *
$^{240}\text{Cf}(\alpha)^{236}\text{Cm}$	7718.9	10.	7711	4	-0.8	8					70Si19
	7713.8	20.			-0.1	U			GSa		10Kh06
	7709.7	4.1			0.3	8			JAA		10As.A
$^{238}\text{U}(\text{t,p})^{240}\text{U}$	2242	20	2253.1	2.9	0.6	U			Ald		64Mi.A
	2253	20			0.0	U			LAI		69Br11
$^{240}\text{Pu}(\text{p,t})^{238}\text{Pu}$	-3692	15	-3698.7	0.4	-0.4	U			ANL		74Fr01
$^{239}\text{Pu}(\text{n},\gamma)^{240}\text{Pu}$	6534.1	1.0	6534.22	0.23	0.1	-					70Ch.A
	6534.3	0.4			-0.2	-					74Ju.B Z
	6534.2	0.4			0.0	-					75We.A Z
$^{239}\text{Pu}(\text{d,p})^{240}\text{Pu}$	4300	10	4309.65	0.23	1.0	U			ANL		73Fr01
$^{239}\text{Pu}(\text{n},\gamma)^{240}\text{Pu}$	ave.	6534.24	6534.22	0.23	-0.1	1	72	46 $^{239}\text{Pu}$			average
$^{240}\text{U}(\beta^-)^{240}\text{Np}^m$	386	20	381	13	-0.2	1	43	42 $^{240}\text{Np}^m$			53Kn23 *
$^{240}\text{Np}^m(\text{IT})^{240}\text{Np}$	20	15	18	14	-0.1	1	83	68 $^{240}\text{Np}$			81Hs02 *
$^{240}\text{Np}(\beta^-)^{240}\text{Pu}$	2199	30	2191	17	-0.3	1	32	32 $^{240}\text{Np}$			51Or.A *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{240}\text{Np}^m(\beta^-)^{240}\text{Pu}$	2210	20	2209	13	-0.1	1	43	43 $^{240}\text{Np}^m$			59Bu20 *
$^{240}\text{Am}(\epsilon)^{240}\text{Pu}$	1395	35	1385	14	-0.3	R					72Ah07 *
* $^{240}\text{Cm}(\alpha)^{236}\text{Pu}$	$E_\alpha=6290.5, 6247.7(0.6,Z)$ to ground state, $2^+$ level at 44.63 keV										
* $^{240}\text{U}(\beta^-)^{240}\text{Np}^m$	$E_{\beta^-}=360(20)$ to $^{240}\text{Np}^m$ , and $1^+$ level at 44.17 keV above										
* $^{240}\text{Np}^m(\text{IT})^{240}\text{Np}$	From fraction IT=0.0012(0.0001)										
* $^{240}\text{Np}(\beta^-)^{240}\text{Pu}$	$E_{\beta^-}=890(30)$ to $5^-$ level at 1308.74 keV										
* $^{240}\text{Np}^m(\beta^-)^{240}\text{Pu}$	$E_{\beta^-}=2180(20)$ to ground state and $2^+$ level at 42.824 keV, and other $E_{\beta^-}$										
* $^{240}\text{Am}(\epsilon)^{240}\text{Pu}$	pK=0.635(0.020) to $3^+$ level at 1030.55 keV, recalculated										
$^{241}\text{Am O-C}_{22}$	51744.8	1.9	51742.0	1.2	-1.0	1	18	18 $^{241}\text{Am}$	TG1	1.5	14Ei01
$^{241}\text{Pu}(\alpha)^{237}\text{U}$	5139.6	3.	5140.1	0.5	0.2	U					68Ah01 *
	5139.3	1.2			0.6	1	16	16 $^{237}\text{U}$	Kum		68Ba25 *
$^{241}\text{Am}(\alpha)^{237}\text{Np}$	5637.81	0.12	5637.82	0.12	0.1	1	100	99 $^{237}\text{Np}$			71Gr17 *
$^{241}\text{Cm}(\alpha)^{237}\text{Pu}$	6182.8	2.0	6185.2	0.6	1.2	U			Kum		67Ba42 *
	6185.2	0.6			0.0	-			Kum		71Bb10 *
	6185.0	2.0			0.1	-					75Ah05 *
ave.	6185.2	0.6			0.0	1	99	94 $^{237}\text{Pu}$			average
$^{241}\text{Cf}(\alpha)^{237}\text{Cm}^p$	7459.0	5.	7455	3	-0.9	9					70Si19
	7451.9	4.1			0.7	9			JAa		10As.A
$^{241}\text{Es}(\alpha)^{237}\text{Bk}$	8064.1	30.	8250	20	6.2	C			GSa		85Hi.A *
	8250.2	20.				10			GSa		96Ni09
$^{239}\text{Pu}(t,p)^{241}\text{Pu}$	3242	20	3293.95	0.23	2.6	U			LAI		69Br11
$^{240}\text{Pu}(n,\gamma)^{241}\text{Pu}$	5241.3	0.7	5241.522	0.030	0.3	U					75Ma.A
	5241.52	0.03			0.1	1	100	61 $^{240}\text{Pu}$	ILn		98Wh01 Z
$^{240}\text{Pu}(d,p)^{241}\text{Pu}$	3018	6	3016.956	0.030	-0.2	U			ANL		67Er02
$^{241}\text{Am}(d,t)^{240}\text{Am}$	-388	15	-390	14	-0.1	2			Kop		76Gr19
$^{241}\text{Np}(\beta^-)^{241}\text{Pu}$	1360	100	1310	70	-0.5	2					59Va32
	1250	100			0.6	2					66Qa02
$^{241}\text{Pu}(\beta^-)^{241}\text{Am}$	20.8	0.2	20.78	0.17	-0.1	-					56Sh31
	20.7	0.3			0.3	-					99Dr13
	20.78	0.20			0.0	o					99Ya.A
	21.6	0.5			-1.6	U					10Lo14 *
ave.	20.77	0.17			0.1	1	99	81 $^{241}\text{Am}$			average
$^{241}\text{Cm}(\epsilon)^{241}\text{Am}$	767.5	1.2	767.4	1.2	-0.1	1	95	94 $^{241}\text{Cm}$			89Su.A *
* $^{241}\text{Pu}(\alpha)^{237}\text{U}$	$E_\alpha=4896.6(3,Z), 4853.6(3,Z)$ to $5/2^+$ at 159.962, $11/2^+$ at 204.06 keV										
* $^{241}\text{Pu}(\alpha)^{237}\text{U}$	$E_\alpha=4896.3(1.2,Z), 4853.3(1.2,Z)$ to $5/2^+$ at 159.962, $11/2^+$ at 204.06 keV										
* $^{241}\text{Am}(\alpha)^{237}\text{Np}$	$E_\alpha=5485.56(0.12,Z), 5442.80(0.13,Z)$ to $5/2^-$ at 59.54, $7/2^-$ at 102.96										
* $^{241}\text{Cm}(\alpha)^{237}\text{Pu}$	$E_\alpha=6080.6(2,Z), 5926.6(2,Z)$ to ground state, $3/2^+$ level at 155.456 keV										
* $^{241}\text{Cm}(\alpha)^{237}\text{Pu}$	$E_\alpha=5939.0(0.6,Z), 5884.7(0.6,Z)$ to $1/2^+$ at 145.543, $5/2^+$ at 201.179 keV										
* $^{241}\text{Cm}(\alpha)^{237}\text{Pu}$	$E_\alpha=5938.7(2,Z), 5884.7(2,Z)$ to $1/2^+$ at 145.543, $5/2^+$ at 201.179 keV										
* $^{241}\text{Es}(\alpha)^{237}\text{Bk}$	C : new data from same group (next item) is more reliable										
* $^{241}\text{Pu}(\beta^-)^{241}\text{Am}$	No quoted uncertainty, estimated by evaluator										
* $^{241}\text{Cm}(\epsilon)^{241}\text{Am}$	$Q(\epsilon)=5.5(1.2)$ to $3/2^-$ level at 636.861 keV										
$^{242}\text{Pu}(\alpha)^{238}\text{U}$	4987.3	2.0	4984.2	1.0	-1.5	-					53As.A *
	4989.5	3.0			-1.8	U					56Ko67 *
	4982.9	1.2			1.1	-			Kum		68Ba25 *
ave.	4984.1	1.0			0.2	1	92	78 $^{238}\text{U}$			average
$^{242}\text{Am}(\alpha)^{238}\text{Np}$	5587.5	0.8	5588.50	0.25	1.3	U			Kum		79Ba67 *
	5589.9	0.8			-1.7	U					90Ho02 *
$^{242}\text{Cm}(\alpha)^{238}\text{Pu}$	6215.63	0.08	6215.63	0.08	0.0	1	100	100 $^{242}\text{Cm}$			71Gr17 Z
$^{242}\text{Cf}(\alpha)^{238}\text{Cm}$	7516.9	4.				5					70Si19 Z
$^{242}\text{Es}(\alpha)^{238}\text{Bk}$	8160.2	20.				9			GSa		10An08
$^{240}\text{Pu}(t,p)^{242}\text{Pu}$	3043	20	3069.3	0.7	1.3	U			LAI		69Br11

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{242}\text{Pu}(p,t)^{240}\text{Pu}$	-3045	15	-3069.3	0.7	-1.6	U			ANL		74Fr01
$^{241}\text{Pu}(n,\gamma)^{242}\text{Pu}$	6309.5	0.7	6309.6	0.7	0.1	1	95	81 $^{242}\text{Pu}$			72Ma.A
$^{242}\text{Pu}(d,t)^{241}\text{Pu}$	-49	7	-52.3	0.7	-0.5	U			ANL		67Er02
$^{241}\text{Am}(n,\gamma)^{242}\text{Am}$	5541.5	1.5	5537.64	0.10	-2.6	U					75Ij.A
	5537.64	0.1				2			ILn		88Sa18 Z
$^{241}\text{Am}(d,p)^{242}\text{Am}$	3308	15	3313.07	0.10	0.3	U			Kop		76Gr19
$^{242}\text{Np}(\beta^-)^{242}\text{Pu}$	2700	200				2					79Ha26
$^{242}\text{Am}(\beta^-)^{242}\text{Cm}$	651	5	664.3	0.4	2.7	U					50Ok52 *
	667	5			-0.5	U					55Ba.A
* $^{242}\text{Pu}(\alpha)^{238}\text{U}$	$E_\alpha=4904.6, 4860.6(2,Z)$ to ground state, $2^+$ level at 44.916 keV										Ens157 **
* $^{242}\text{Pu}(\alpha)^{238}\text{U}$	$E_\alpha=4905.2(3,Z), 4863.2(3,Z)$ to ground state, $2^+$ level at 44.916 keV										Ens157 **
* $^{242}\text{Pu}(\alpha)^{238}\text{U}$	$E_\alpha=4900.4(1.2,Z), 4856.1(1.2,Z)$ to ground state, $2^+$ level at 44.916 keV										Ens157 **
* $^{242}\text{Am}(\alpha)^{238}\text{Np}$	$E_\alpha=5206.6(0.5,Z), 5141.4(0.5,Z)$ from $^{242}\text{Am}^m$ to $5^-$ at 342.439, and $6^-$ at 407.59 keV; error increased due to conflict with next item										Ens157 **
*											GAu **
* $^{242}\text{Am}(\alpha)^{238}\text{Np}$	$E_\alpha=5208.3(0.8,Z), 5144.3(0.9,Z)$ from $^{242}\text{Am}^m$ to $5^-$ $6^-$ levels (see above)										Ens157 **
* $^{242}\text{Am}(\beta^-)^{242}\text{Cm}$	$E_{\beta^-}=628(5)$ to ground state and $2^+$ level at 42.13 keV										Ens026 **
$^{243}\text{Am O}-\text{C}_{22}$	56295.8	1.5	56294.6	1.5	-0.6	1	44	44 $^{243}\text{Am}$	TG1	1.5	14Ei01
$^{243}\text{Am}(\alpha)^{239}\text{Np}$	5438.8	1.0	5439.1	0.9	0.3	1	87	54 $^{243}\text{Am}$	Kum		68Ba25 *
$^{243}\text{Cm}(\alpha)^{239}\text{Pu}$	6165.4	3.0	6168.8	1.0	1.1	U					57As.A *
	6165.7	3.0			1.0	U					63Dz07 *
	6165.4	3.0			1.1	o			Kum		66Ba07 *
	6168.8	1.0				2					69Ba57 *
$^{243}\text{Bk}(\alpha)^{239}\text{Am}$	6874.4	4.				3			Bka		66Ah.A Z
$^{243}\text{Cf}(\alpha)^{239}\text{Cm}^p$	7178	10				6					67Fi04 *
$^{243}\text{Es}(\alpha)^{239}\text{Bk}$	8072.1	10.				9			RIa		89Ha27
$^{243}\text{Es}(\alpha)^{239}\text{Bk}^p$	8022.3	20.	8030.9	2.9	0.4	U					73Es02
	8031.4	3.			-0.2	10			RIa		89Ha27
	8027.3	20.			0.2	o			GSa		93Ho.A
	8025.4	10.			0.6	10			GSa		10An08
$^{243}\text{Fm}(\alpha)^{239}\text{Cf}$	8689.3	25.4	8690	50	0.1	o			GSa		81Mu12
	8693.4	20.3				11			GSa		08Kh10
$^{243}\text{Am}(p,t)^{241}\text{Am}$	-3407	15	-3420.2	1.2	-0.9	U			ANL		74Fr01
$^{242}\text{Pu}(n,\gamma)^{243}\text{Pu}$	5034.2	3.	5033.6	2.4	-0.2	1	63	59 $^{243}\text{Pu}$			76Ca25
$^{242}\text{Pu}(d,p)^{243}\text{Pu}$	2807	8	2809.1	2.4	0.3	U			ANL		67Er02
$^{243}\text{Am}(d,t)^{242}\text{Am}$	-111	15	-107.1	1.2	0.3	U			Kop		76Gr19
$^{243}\text{Pu}(\beta^-)^{243}\text{Am}$	578	10	579.6	2.6	0.2	-					69Ho10
	580	10			0.0	-					77Dr07
ave.	579	7			0.1	1	14	11 $^{243}\text{Pu}$			average
* $^{243}\text{Am}(\alpha)^{239}\text{Np}$	$E_\alpha=5275.2(1.0,Z), 5233.3(1.0,Z)$ to $5/2^-$ at 74.6640, $7/2^-$ at 117.715										Ens14b **
* $^{243}\text{Cm}(\alpha)^{239}\text{Pu}$	$E_\alpha=6063.7, 5989.7, 5782.7, 5738.7(3,Z)$ to ground state, $7/2^+$ level at 75.705, $5/2^+$ at 285.460, and $7/2^+$ at 330.124 keV										57As.A **
*											Ens14b **
* $^{243}\text{Cm}(\alpha)^{239}\text{Pu}$	$E_\alpha=5990.5, 5783.5, 5738.5(3,Z)$ to $7/2^+$ level at 75.705, $5/2^+$ at 285.46, and $7/2^+$ at 330.124 keV										Ens14b **
*											Ens14b **
* $^{243}\text{Cm}(\alpha)^{239}\text{Pu}$	$E_\alpha=6067.4, 5992.4(2,Z)$ to ground state, $7/2^+$ at 75.705 keV										Ens14b **
* $^{243}\text{Cm}(\alpha)^{239}\text{Pu}$	$E_\alpha=5785.7(1.0,Z), 5742.8(1.0,Z)$ to $5/2^+$ at 285.46, $7/2^+$ at 330.124keV										Ens14b **
* $^{243}\text{Cf}(\alpha)^{239}\text{Cm}^p$	Unhindered $E_\alpha=7060(10)$ ; there is a weaker $E_\alpha=7170(10)$ keV										AHW **
$^{244}\text{Pu O}-\text{C}_{22}$	59119.3	1.9	59119.0	2.5	-0.1	1	78	78 $^{244}\text{Pu}$	TG1	1.5	14Ei01
$^{244}\text{Pu}(\alpha)^{240}\text{U}$	4665.6	1.0	4665.6	1.0	0.0	1	100	99 $^{240}\text{U}$			69Be06 Z
$^{244}\text{Cm}(\alpha)^{240}\text{Pu}$	5901.60	0.03				2			BIP		71Gr17 *
$^{244}\text{Bk}(\alpha)^{240}\text{Am}$	6778.8	4.				3					66Ah.B *
$^{244}\text{Cf}(\alpha)^{240}\text{Cm}$	7327.1	2.	7329.0	1.8	0.9	-					67Fi04 Z
	7336.4	4.			-1.8	-					67Si08 Z
	7330.4	20.			-0.1	U			GSa		08Kh10
ave.	7329.0	1.8			0.0	1	99	98 $^{244}\text{Cf}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{244}\text{Es}(\alpha)^{240}\text{Bk}^p$	7696.4	20.				4					73Es02
$^{242}\text{Pu}(\text{t,p})^{244}\text{Pu}$	2576	20	2571.7	2.5	-0.2	U			LAI		69Br11
$^{244}\text{Pu}(\text{p,t})^{242}\text{Pu}$	-2560	15	-2571.7	2.5	-0.8	U			ANL		72Ma15
$^{244}\text{Pu}(\text{t},\alpha)^{243}\text{Np}^p$	12405	10				2					79FI02
$^{244}\text{Pu}(\text{d,t})^{243}\text{Pu}$	234	5	237.3	2.9	0.7	1	34	19 $^{243}\text{Pu}$	ANL		76Ca25
$^{243}\text{Am}(\text{n},\gamma)^{244}\text{Am}^m$	5277.90	0.07				2			ILn		84Vo07 Z
$^{244}\text{Cm}(\text{d,t})^{243}\text{Cm}$	-530	7	-544.2	1.0	-2.0	U			ANL		67Er02
$^{244}\text{Am}^m(\text{IT})^{244}\text{Am}$	85.0	1.0	89.3	1.6	4.3	F					84Ho02 *
$^{244}\text{Am}(\beta^-)^{244}\text{Cm}$	1427.3	1.0				3					62Va08 *
* $^{244}\text{Cm}(\alpha)^{240}\text{Pu}$	$E_\alpha=5804.77(0.05,Z)$ , $5762.65(0.03,Z)$ to ground state, $2^+$ level at 42.824 keV										
* $^{244}\text{Bk}(\alpha)^{240}\text{Am}$	$E_\alpha=6667.5(4,Z)$ , $6625.5(3,Z)$ to ground state, $2^+$ level at 42.824 keV										
* $^{244}\text{Am}^m(\text{IT})^{244}\text{Am}$	F : value in Fig. 1 only, no source no error										
* $^{244}\text{Am}(\beta^-)^{244}\text{Cm}$	$E_{\beta^-}=387(1)$ to $6^+$ level at 1040.188 keV; also $E_{\beta^-}=1498(10)$ from										
*	$^{244}\text{Am}^m$ at 89.3(1.6) to ground state and $2^+$ at 42.965 keV, not used										
$^{245}\text{Cm}(\alpha)^{241}\text{Pu}$	5621.9	0.5	5624.5	0.5	5.2	B					75Ba65 *
	5624.8	0.5			-0.6	1	96	68 $^{245}\text{Cm}$	Ara		16Ko.B *
$^{245}\text{Bk}(\alpha)^{241}\text{Am}$	6454.7	4.	6454.5	1.4	0.0	2					74Po08 *
	6454.5	1.5			0.0	2			Kum		75Ba25 *
$^{245}\text{Cf}(\alpha)^{241}\text{Cm}$	7257.5	2.0	7258.5	1.8	0.5	-					67Fi04 *
	7265	5			-1.3	-					96Ma72 *
	7260.8	11.			-0.2	U			GSa		04He28
	ave.	7258.5	1.9		0.0	1	98	97 $^{245}\text{Cf}$			average
$^{245}\text{Es}(\alpha)^{241}\text{Bk}$	7858.5	20.	7909	3	1.0	U					73Es01
	7884.0	20.			0.5	U			GSa		85He22
	7909.4	3.				3			RIa		89Ha27
$^{245}\text{Es}(\alpha)^{241}\text{Bk}^p$	7827.9	30.	7858.5	1.0	1.0	U					67Mi06
	7858.5	1.				4			RIa		89Ha27
$^{245}\text{Fm}(\alpha)^{241}\text{Cf}^p$	8285.5	20.				11					67Nu01
$^{245}\text{Md}(\alpha)^{241}\text{Es}^p$	8824.3	20.				12			GSa		96Ni09 *
$^{244}\text{Pu}(\text{d,p})^{245}\text{Pu}$	2469	15	2475	13	0.4	2			ANL		75Er.A *
$^{244}\text{Cm}(\text{d,p})^{245}\text{Cm}$	3297	7	3294.1	0.5	-0.4	U			ANL		67Er02
$^{245}\text{Pu}(\beta^-)^{245}\text{Am}$	1257	30	1278	14	0.7	R					68Da02 *
$^{245}\text{Am}(\beta^-)^{245}\text{Cm}$	905	5	895.9	1.5	-1.8	U					55Br02
$^{245}\text{Es}^p(\text{IT})^{245}\text{Es}$	283	15				4					Nub16b
* $^{245}\text{Cm}(\alpha)^{241}\text{Pu}$	$E_\alpha=5529.0$ , $5488.5$ , $5436.1(0.5,Z)$ , $5303.6$ , $5234.4(1.2,Z)$ keV										
* $^{245}\text{Cm}(\alpha)^{241}\text{Pu}$	to ground state, $7/2^+$ 41.9722, $9/2^+$ 95.7795, $9/2^+$ 231.935, $11/2^+$ 301.172 levels										
*	$E_\alpha=5361.8(1.2,Z)$ to $7/2^+$ 175.05 level in error in 75Ba65; excluded										
* $^{245}\text{Cm}(\alpha)^{241}\text{Pu}$	$E_\alpha=5532.7$ , $5491.3(0.5)$ , $5436.6(2.0)$ , $5360.6$ , $5305.2$ , $5624.5(0.5)$ keV										
*	to ground state, 41.97, 95.78, 175.05, 231.94, 301.17 levels										
* $^{245}\text{Bk}(\alpha)^{241}\text{Am}$	$E_\alpha=6349.0$ , $6309.0$ , $6146.0$ , $5886.0$ (all 4,Z)										
*	to ground state, $7/2^-$ at 41.176, $5/2^+$ at 205.883, $3/2^-$ at 471.810 keV										
* $^{245}\text{Bk}(\alpha)^{241}\text{Am}$	$E_\alpha=6347.8$ , $6307.8$ , $6146.8$ , $5885.8$ recalibrated as in reference										
*	to ground state, $7/2^-$ at 41.176, $5/2^+$ at 205.883, $3/2^-$ at 471.810 keV										
* $^{245}\text{Cf}(\alpha)^{241}\text{Cm}$	$E_\alpha=7136.8(2.0,Z)$ , $7083.8(2.0,Z)$ to gs+5.6 and 56.1 level										
* $^{245}\text{Cf}(\alpha)^{241}\text{Cm}$	$E_\alpha=7145(5)$ , $7090(5)$ to gs+5.6 and 56.1 level										
* $^{245}\text{Md}(\alpha)^{241}\text{Es}^p$	Second $E_\alpha$ 8635(20) keV										
* $^{244}\text{Pu}(\text{d,p})^{245}\text{Pu}$	$Q=2252(15)$ to 217 level (estimated energy for $15/2^-$ level)										
* $^{245}\text{Pu}(\beta^-)^{245}\text{Am}$	$E_{\beta^-}=1210(40)$ , $930(30)$ to $(9/2^+)$ level at 47.07, $7/2^+$ at 327.428 keV										
$^{246}\text{Cm}(\alpha)^{242}\text{Pu}$	5475.2	4.	5475.1	0.9	0.0	U					63Dz07 Z
	5474.9	2.			0.1	-			Kum		66Ba07 *
	5475.2	1.			-0.1	-					84Sh31 *
	ave.	5475.1	0.9		0.0	1	99	98 $^{246}\text{Cm}$			average

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{246}\text{Cf}(\alpha)^{242}\text{Cm}$	6871.0	1.0	6861.6	1.0	-9.4	B					63Fr04 *
	6861.6	1.			0.0	1	100	99 $^{246}\text{Cf}$	Kum		77Ba69 *
$^{246}\text{Es}(\alpha)^{242}\text{Bk}^p$	7451.2	30.	7492	4	1.4	U					67Mi06
	7481.9	30.			0.3	U					73Es01
	7492.0	4.				4			RIa		89Ha27
$^{246}\text{Fm}(\alpha)^{242}\text{Cf}$	8371.4	20.	8377	8	0.3	6					66Ak01
	8376.5	20.			0.0	6			Bka		67Nu01
	8386.7	20.			-0.5	o			GSa		96Ni09
	8378.4	10.			-0.2	6			GSa		10An08
$^{246}\text{Md}(\alpha)^{242}\text{Es}$	8884.7	20.	8890	40	0.1	o			GSa		96Ni09 *
	8888.8	40.				10			GSa		10An08
$^{246}\text{Md}^m(\alpha)^{242}\text{Es}$	8944.5	50.				10			GSa		10An08 *
$^{244}\text{Pu}(t,p)^{246}\text{Pu}$	2085	20	2072	15	-0.6	1	56	55 $^{246}\text{Pu}$	LAI		79Br19
$^{246}\text{Cm}(d,t)^{245}\text{Cm}$	-196	6	-201.7	1.2	-1.0	U			ANL		67Er02
$^{246}\text{Pu}(\beta^-)^{246}\text{Am}^m$	374	10	371	9	-0.3	1	89	45 $^{246}\text{Pu}$			56Ho23 *
$^{246}\text{Am}^m(\beta^-)^{246}\text{Cm}$	2300	100	2407	15	1.1	U					55En16 *
	2420	20			-0.6	1	56	56 $^{246}\text{Am}^m$			56Sm85 *
$^{246}\text{Bk}(\epsilon)^{246}\text{Cm}$	1350	60				2					89Sc.A
$^{246}\text{Cm}(\alpha)^{242}\text{Pu}$	$E_{\alpha}=5385.3(2,Z)$ , $5342.3(2,Z)$ to ground state, $2^+$ level at 44.54 keV										
$^{246}\text{Cm}(\alpha)^{242}\text{Pu}$	$E_{\alpha}=5385.6(1,Z)$ , $5342.6(1,Z)$ to ground state, $2^+$ level at 44.54 keV										
$^{246}\text{Cf}(\alpha)^{242}\text{Cm}$	$E_{\alpha}=6757.4(1.0,Z)$ , $6718.4(0.7,Z)$ to ground state, $2^+$ level at 42.13 keV										
$^{246}\text{Cf}(\alpha)^{242}\text{Cm}$	$E_{\alpha}=6750.0(1.0,Z)$ , $6708.2(1.0,Z)$ to ground state, $2^+$ level at 42.13 keV										
$^{246}\text{Md}(\alpha)^{242}\text{Es}$	Also a lower $E_{\alpha}=8530(30)$ keV										
$^{246}\text{Md}^m(\alpha)^{242}\text{Es}$	$E_{\alpha}=8178(10)$ to level at 531+x; x estimated to be 100#50 keV										
$^{246}\text{Pu}(\beta^-)^{246}\text{Am}^m$	$E_{\beta^-}=150(10)$ to $1^+$ level at 223.74 keV above $^{246}\text{Am}^m$										
$^{246}\text{Am}^m(\beta^-)^{246}\text{Cm}$	$E_{\beta^-}=1222(100)$ $1350(20)$ respectively, to $1^-$ level at 1078.845, $2^-$ level at 1104.854										
$^{247}\text{Cm}(\alpha)^{243}\text{Pu}$	5354.6	4.	5354	3	-0.2	1	71	60 $^{247}\text{Cm}$			71Fi01 *
$^{247}\text{Bk}(\alpha)^{243}\text{Am}$	5889.6	5.				2					69Fr01 *
$^{247}\text{Cf}(\alpha)^{243}\text{Cm}^p$	6399.6	5.				5					84Ah02 Z
$^{247}\text{Es}(\alpha)^{243}\text{Bk}^p$	7450.7	30.	7443.8	1.0	-0.2	U					67Mi06
	7430.5	30.			0.4	U					73Es01
	7443.8	1.				5			RIa		89Ha27
$^{247}\text{Fm}(\alpha)^{243}\text{Cf}$	8060.8	50.	8258	10	3.9	B					67Fi15
	8213	18			2.5	o			GSa		89He03 *
	8287.3	20.			-1.5	o			GSa		04He28 *
	8268.1	10.			-1.0	o			GSa		06He27 *
$^{247}\text{Fm}^m(\alpha)^{243}\text{Cf}$	8314.9	30.	8307	5	-0.3	U					67Fi15 *
	8260.0	30.			1.5	o			GSa		97He29 *
	8304.8	11.			0.2	o			GSa		04He28
	8306.8	5.				7			GSa		06He27
$^{247}\text{Md}(\alpha)^{243}\text{Es}$	8776.6	25.	8764	10	-0.5	o			GSa		81Mu12 *
	8772.5	20.			-0.4	o			GSa		93Ho.A *
	8770.5	10.			-0.6	o			GSa		05He27 *
	8764.4	10.				10			GSa		10An08 *
$^{247}\text{Md}^m(\alpha)^{243}\text{Es}$	9027.9	40.				10			GSa		10An08 *
$^{246}\text{Cm}(d,p)^{247}\text{Cm}$	2931	8	2931	4	-0.1	1	22	21 $^{247}\text{Cm}$	ANL		67Er02
$^{247}\text{Cf}(\epsilon)^{247}\text{Bk}$	646	6	614	16	-5.3	C					56Ch.A *
$^{247}\text{Cm}(\alpha)^{243}\text{Pu}$	$E_{\alpha}=5267.3(4,Z)$ $5212.3(4,Z)$ $4870.3(4,Z)$ to ground state, $9/2^+$ 58.13, $9/2^-$ 402.6										
$^{247}\text{Bk}(\alpha)^{243}\text{Am}$	$E_{\alpha}=5794$ , $5710$ , $5688(5,Z)$ to ground state, $5/2^+$ level at 84.00, $7/2^+$ at 109.22 keV										
$^{247}\text{Fm}(\alpha)^{243}\text{Cf}$	$E_{\alpha}=8060(15)$ summed with $e^-$										
$^{247}\text{Fm}(\alpha)^{243}\text{Cf}$	$E_{\alpha}=7840(20)$ to 318 level										
$^{247}\text{Fm}(\alpha)^{243}\text{Cf}$	$E_{\alpha}=7824(10)$ to 315 level										
$^{247}\text{Fm}^m(\alpha)^{243}\text{Cf}$	Only one event										
$^{247}\text{Fm}^m(\alpha)^{243}\text{Cf}$	Not observed in later work on $^{251}\text{No}$ decay										
$^{247}\text{Md}(\alpha)^{243}\text{Es}$	$E_{\alpha}=8428(25)$ to 209.6 level										
$^{247}\text{Md}(\alpha)^{243}\text{Es}$	$E_{\alpha}=8424(20)$ to 209.6 level										
$^{247}\text{Md}(\alpha)^{243}\text{Es}$	$E_{\alpha}=8422(10)$ to 209.6 level										
$^{247}\text{Md}(\alpha)^{243}\text{Es}$	$E_{\alpha}=8616(20)$ , $8416(10)$ to ground state, 209.6 level										
$^{247}\text{Md}^m(\alpha)^{243}\text{Es}$	$E_{\alpha}=8783(40)$ to $1/2^-$ level at 100 keV										
$^{247}\text{Cf}(\epsilon)^{247}\text{Bk}$	LMK=10(3) assuming first-forbidden to $(5/2^-)$ level at 447.80 keV, yields 646.8										



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
*	contradicts LMK=75 allowed to $(5/2)^+$ level at 334.92 keV, yields 550;										WgM10a**
*	both conflict with 613 from Shiptrap data to $^{255}\text{No}$ plus $\alpha$ chain										WgM10a**
$^{248}\text{Cm}(\alpha)^{244}\text{Pu}$	5161.81	0.25	5161.81	0.25	0.0	1	100	95 $^{248}\text{Cm}$	Kum		77Ba69 Z
$^{248}\text{Cf}(\alpha)^{244}\text{Cm}$	6364.8	5.1	6361	5	-0.7	o					78Gr10
	6361.2	5.				3					84Ah02 *
$^{248}\text{Es}(\alpha)^{244}\text{Bk}$	7165.8	20.	7160#	50#	-0.3	F					84Li.A *
$^{248}\text{Es}(\alpha)^{244}\text{Bk}^p$	6982.8	15.	7020	5	2.5	U					56Ch67
	6982.8	10.			3.8	B					70Ah01
	7020.4	5.				5			RIa		89Ha27
$^{248}\text{Fm}(\alpha)^{244}\text{Cf}$	8009.4	30.	7995	8	-0.5	-					66Ak01
	7999.3	20.			-0.2	-					67Nu01
	8002.3	15.			-0.5	o			GSa		85He.A
	7992.2	11.			0.2	-			GSa		04He28
	ave.	9			-0.1	1	79	77 $^{248}\text{Fm}$			average
$^{248}\text{Md}(\alpha)^{244}\text{Es}^p$	8497.3	30.				6					73Es01
$^{248}\text{Cm}(p,t)^{246}\text{Cm}$	-2894	15	-2885.1	2.7	0.6	U			ANL		74Fr01
$^{248}\text{Cm}(d,t)^{247}\text{Cm}$	49	8	46	4	-0.4	1	25	19 $^{247}\text{Cm}$	ANL		67Er02
$^{248}\text{Bk}^m(\beta^-)^{248}\text{Cf}$	870	20				4					78Gr10
* $^{248}\text{Cf}(\alpha)^{244}\text{Cm}$	$E_\alpha=6257.8(5,Z)$ , $6216.8(5,Z)$ to ground state, $2^+$ level at 42.965 keV										Ens036 **
* $^{248}\text{Es}(\alpha)^{244}\text{Bk}$	F : this line is not observed in more recent works										AHW **
$^{249}\text{Cf} \text{ O}-\text{C}_{22}$	69760.0	1.4	69765.1	1.3	2.4	1	37	37 $^{249}\text{Cf}$	TG1	1.5	14Ei01
$^{249}\text{Bk}(\alpha)^{245}\text{Am}$	5520.4	2.0	5521.0	1.4	0.3	3			Bka		66Ah.A *
	5526.1	1.0			-5.1	B			Kum		71Bb10 *
	5521.6	2.0			-0.3	3			Ara		13Ah03 *
$^{249}\text{Cf}(\alpha)^{245}\text{Cm}$	6296.0	0.7	6293.3	0.5	-3.9	B			Kum		71Bb10 *
	6293.6	0.5			-0.6	1	96	63 $^{249}\text{Cf}$	Ara		15Ah03 *
$^{249}\text{Es}(\alpha)^{245}\text{Bk}^p$	6881.3	5.	6886.0	1.9	0.9	4					70Ah01 Z
	6886.8	2.			-0.4	4			RIa		89Ha27
$^{249}\text{Fm}(\alpha)^{245}\text{Cf}$	7718.3	20.	7709	6	-0.5	-					73Es01 *
	7705.1	23.			0.2	o			GSa		85He06 *
	7705.0	14.			0.3	-			GSa		04He28 *
	7710.2	8.1			-0.1	-			Orm		11Lo06 *
	ave.	7			-0.1	1	80	77 $^{249}\text{Fm}$			average
$^{249}\text{Md}(\alpha)^{245}\text{Es}^p$	8161.3	20.	8158	9	-0.2	5					73Es01
	8157.3	20.			0.0	o			GSa		85He22
	8165	20			-0.3	o			GSa		01He35 *
	8157.3	10.			0.1	o			GSa		05He27
	8157.3	10.			0.1	5			GSa		09He20
$^{249}\text{Md}^m(\alpha)^{245}\text{Es}^q$	8212.2	20.				7			GSa		01He35
$^{248}\text{Cm}(n,\gamma)^{249}\text{Cm}$	4713.37	0.25				2			ILn		82Ho07 Z
$^{248}\text{Cm}(d,p)^{249}\text{Cm}$	2488	6	2488.80	0.25	0.1	U			ANL		67Er02
$^{249}\text{Cm}(\beta^-)^{249}\text{Bk}$	870	100	904.3	2.6	0.3	U					58Ea06 *
	885	15			1.3	U			ANB		05Ah03 *
$^{249}\text{Bk}(\beta^-)^{249}\text{Cf}$	125	2	123.6	0.4	-0.7	U					59Va02
	123	2			0.3	U					74G110
	123.6	0.4				2					14Ch47
* $^{249}\text{Bk}(\alpha)^{245}\text{Am}$	$E_\alpha=5431.8$ , $5412.8$ , $5384.8$ (all $2,Z$ ) to ground state, $7/2^+$ 19.20, $9/2^+$ 47.07 keV										Ens112 **
* $^{249}\text{Bk}(\alpha)^{245}\text{Am}$	$E_\alpha=5437.1(1.0,Z)$ to gs. Energies of higher branches										71Bb10 **
*	rather different from reference, calibrated with same ground state $\alpha$										75Ba27 **
* $^{249}\text{Bk}(\alpha)^{245}\text{Am}$	$E_\alpha=5433(2)$ , $5414(2)$ , $5386(2)$ to ground state, $7/2^+$ 19.20, $9/2^+$ 47.07 keV										Ens112 **
* $^{249}\text{Cf}(\alpha)^{245}\text{Cm}$	$E_\alpha=6193.8(0.7,Z)$ , $5813.3(1.0,Z)$ to ground state, $9/2^-$ level at 388.181 keV										Ens112 **
* $^{249}\text{Cf}(\alpha)^{245}\text{Cm}$	$E_\alpha=6192.4(0.5)$ , $5810.5(0.5)$ to ground state, $9/2^-$ level at 388.181 keV										Ens112 **
* $^{249}\text{Fm}(\alpha)^{245}\text{Cf}$	$E_\alpha=7540(20)$ to corresponding $7/2^+$ [624] level at 55(10) keV										04He28 **
* $^{249}\text{Fm}(\alpha)^{245}\text{Cf}$	$E_\alpha=7527(23)$ to corresponding $7/2^+$ [624] level at 55(10) keV										04He28 **
* $^{249}\text{Fm}(\alpha)^{245}\text{Cf}$	Also $E_\alpha=7530(10)$ keV to $7/2^+$ [624] level at 57(4) keV										11Lo06 **
* $^{249}\text{Fm}(\alpha)^{245}\text{Cf}$	$E_\alpha=7530(7)$ to $7/2^+$ [624] level at 57(4) keV										11Lo06 **
* $^{249}\text{Md}(\alpha)^{245}\text{Es}^p$	$E_\alpha=8022(20)$ partly summed with conversion electrons										01He35 **
* $^{249}\text{Cm}(\beta^-)^{249}\text{Bk}$	$E_{\beta^-}=860(100)$ $876(15)$ respectively, to $3/2^-$ level at 8.777 keV										Ens118 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$\nu_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{250}\text{Cf}(\alpha)^{246}\text{Cm}$	6129.1	0.6	6128.51	0.19	-1.0	2			Kum		71Bb10 *
	6128.44	0.2			0.3	2				86Ry04 Z	
$^{250}\text{Fm}(\alpha)^{246}\text{Cf}$	7550.9	50.	7557	8	0.1	U					57Am47
	7540.9	30.5			0.5	-					66Ak01
	7561.1	30.			-0.1	-					73Es01
	7560.1	15.			-0.2	-			ORb		77Be36
	7556.1	35.6			0.0	o			GSa		81Mu06
	7555.0	12.			0.2	-			GSa		04He28
	7544.8	35.			0.3	U			Bka		06Fo02
	ave.	7556	9			0.1	1	80	80 $^{250}\text{Fm}$		
$^{250}\text{Md}(\alpha)^{246}\text{Es}^p$	7947.4	30.	7955	24	0.2	6					73Es01
	7964.7	20.			-0.5	o			GSa		85He22
	7967.7	40.			-0.3	6			GSa		08An16
$^{248}\text{Cm}(\text{t,p})^{250}\text{Cm}$	2064	10				2					73Ba72
$^{250}\text{Bk}(\beta^-)^{250}\text{Cf}$	1760	15	1780	3	1.3	U					59Va02 *
$^{250}\text{Cf}(\alpha)^{246}\text{Cm}$	$E_\alpha=6030.6(0.6,Z)$ , $5988.9(0.6,Z)$ to ground state, $2^+$ level at 42.852 keV										Ens118 **
$^{250}\text{Bk}(\beta^-)^{250}\text{Cf}$	$E_{\beta^-}=725(15)$ to $2^+$ level at 1031.852 keV and $3^+$ level at 1071.37 keV										Ens01c **
$^{251}\text{Cf}(\alpha)^{247}\text{Cm}$	6177.2	1.0	6177.0	0.9	-0.2	2			Kum		71Bb10 *
	6176	2			0.5	2				03Ah07 *	
$^{251}\text{Es}(\alpha)^{247}\text{Bk}$	6593.5	5.	6598	3	0.9	o					70Ah01 *
	6597.8	3.				3					79Ah03 *
$^{251}\text{Fm}(\alpha)^{247}\text{Cf}$	7425.1	2.0				4					73Ah02 *
$^{251}\text{Md}(\alpha)^{247}\text{Es}$	7965.5	20.	7963	4	-0.1	U					73Es01 *
	7955.2	10.			0.8	4			GSa		05He27 *
	7965.5	5.			-0.4	4			Jya		06Ch52 *
$^{251}\text{No}(\alpha)^{247}\text{Fm}$	8739.5	20.	8752	4	0.6	U					Bka
	8732.4	15.			1.3	o					GSa
	8762.9	20.			-0.6	o					GSa
	8760.9	20.			-0.9	o					GSa
	8747.7	11.			0.4	o					GSa
	8751.8	4.				10					GSa
$^{251}\text{No}^m(\alpha)^{247}\text{Fm}^m$	8619.6	30.	8809	4	6.3	F					GSa
	8805.5	13.			0.2	o					GSa
	8808.6	4.				8					GSa
$^{251}\text{Cm}(\beta^-)^{251}\text{Bk}$	1420	20				4					78Lo13
$^{251}\text{Bk}(\beta^-)^{251}\text{Cf}$	1093	10				3					84Li05 *
$^{251}\text{No}^m(\text{IT})^{251}\text{No}$	106	6				9			GSa		06He27
$^{251}\text{Cf}(\alpha)^{247}\text{Cm}$	$E_\alpha=5680.1(1.0,Z)$ to $1/2^+$ level at 404.90 keV										Ens153 **
$^{251}\text{Cf}(\alpha)^{247}\text{Cm}$	$E_\alpha=6078(2)$ , $5679(2)$ to ground state, $1/2^+$ level at 404.90 keV, and others										Ens153 **
$^{251}\text{Es}(\alpha)^{247}\text{Bk}$	$E_\alpha=6488.5(5,Z)$ , $6458.5(5,Z)$ to ground state, $(5/2^-)$ level at 29.88 keV										Ens153 **
$^{251}\text{Es}(\alpha)^{247}\text{Bk}$	$E_\alpha=6492.8(3,Z)$ , $6462.8(3,Z)$ to ground state, $(5/2^-)$ level at 29.88 keV										Ens153 **
$^{251}\text{Fm}(\alpha)^{247}\text{Cf}$	$E_\alpha=7305.7(3,Z)$ , $6833.7(2,Z)$ to ground state and $(9/2^-)$ level at 480.40 keV										Ens153 **
$^{251}\text{Md}(\alpha)^{247}\text{Es}$	$E_\alpha=7550(20)$ $7540(10)$ $7550(1)$ respectively, to $7/2^-$ level at 293.7 keV										06Ch52 **
$^{251}\text{Md}(\alpha)^{247}\text{Es}$	Original error 1 keV in third reference increased for calibration										GAu **
$^{251}\text{No}^m(\alpha)^{247}\text{Fm}^m$	F : not observed in later work on $^{251}\text{No}$ decay										01He35 **
$^{251}\text{Bk}(\beta^-)^{251}\text{Cf}$	$E_{\beta^-}=915(10)$ to $3/2^+$ level at 177.602 keV										Ens139 **
$^{252}\text{No}-^{133}\text{Cs}_{1,895}$	268111	34	268135	10	0.7	o			SH1	1.0	10Dw01
	268133	18			0.1	1	31	31 $^{252}\text{No}$		SH1	1.0
$^{252}\text{Cf}(\alpha)^{248}\text{Cm}$	6216.9	0.5	6216.95	0.04	0.1	U			Kum		71Bb10 Z
	6216.95	0.04				2					
$^{252}\text{Es}(\alpha)^{248}\text{Bk}^p$	6739.5	3.				4					73Fi06 *
$^{252}\text{Fm}(\alpha)^{248}\text{Cf}$	7152.7	2.				4					84Ah02 *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{252}\text{No}(\alpha)^{248}\text{Fm}$	8545.9	20.	8549	5	0.1	U			Bka		67Gh01	
	8545.9	30.			0.1	U			DbA		67Mi03	
	8551.0	6.			-0.4	-					77Be09	
	8542.8	15.			0.4	o			GSa		85He.A	
	8538.7	13.			0.7	-			GSa		04He28	
	ave.	8549	6			0.0	1	92	69 $^{252}\text{No}$			average
$^{252}\text{Lr}(\alpha)^{248}\text{Md}$	9163.8	20.	9164	17	0.0	7			GSa		01He35	
	9165.8	30.			0.0	7			Bka		08Ne01 *	
$^{252}\text{Es}(\epsilon)^{252}\text{Cf}$	1260	50				3					73Fi06 *	
* $^{252}\text{Es}(\alpha)^{248}\text{Bk}^p$	$E_\alpha=6632.1(3,Z)$ , $6522.1(3,Z)$ to 0, $7^+$ level at 70.65 keV above $^{248}\text{Bk}^p$										Ens14b **	
* $^{252}\text{Fm}(\alpha)^{248}\text{Cf}$	$E_\alpha=7038.9(2,Z)$ , $6998.1(2,Z)$ to ground state, $2^+$ level at 41.53 keV										Ens14b **	
* $^{252}\text{Lr}(\alpha)^{248}\text{Md}$	Other $E_\alpha=9610(20)$ unexplained, and 8990, 8820 keV										08Ne01 **	
* $^{252}\text{Es}(\epsilon)^{252}\text{Cf}$	$pK=0.45(0.10)$ to $3^+$ level at 969.8 keV, recalculated for non-unique first forbidden or allowed transition; unique first forbidden would give 1440(100)										Ens061 **	
*											AHW **	
$^{253}\text{No}-^{133}\text{Cs}_{1,902}$	270390	13	270393	7	0.2	1	33	33 $^{253}\text{No}$	SH1	1.0	10Dw01	
$^{253}\text{Cf}(\alpha)^{249}\text{Cm}$	6127.3	5.	6126	4	-0.3	3					66Rg01 *	
	6124.6	5.			0.3	3					68Be21 *	
$^{253}\text{Es}(\alpha)^{249}\text{Bk}$	6739.24	0.05				3					71Gr17 Z	
$^{253}\text{Fm}(\alpha)^{249}\text{Cf}$	7199	3	7198.0	2.7	-0.3	2					67Ah02 *	
	7194.2	6.			0.6	2			Orm		11Lo06 *	
	7567.5	15.	7573	8	0.4	o			GSa		05He27 *	
$^{253}\text{Md}(\alpha)^{249}\text{Es}$	7574.0	10.			-0.1	5			Orm		11Lo06 *	
	7571	15			0.1	5			GSa		12He09 *	
	8419	20	8415	4	-0.2	U			Bka		67Gh01 *	
	8419	30			-0.1	U			DbA		67Mi03 *	
$^{253}\text{No}(\alpha)^{249}\text{Fm}$	8430	20			-0.8	o			GSa		85He.A *	
	8420	10			-0.5	o			GSa		01He.A *	
	8412.5	11.			0.2	o			GSa		04He28 *	
	8411.5	5.			0.6	o			Orm		06Lo12 *	
	8415.6	5.0			-0.2	-			Orm		11Lo06 *	
	8412.4	11.			0.2	-			GSa		12He09 *	
	ave.	8415	5			-0.1	1	90	67 $^{253}\text{No}$			average
	8941.6	20.	8918	20	-1.2	o			GSa		85He22	
$^{253}\text{Lr}(\alpha)^{249}\text{Md}$	8935.6	10.			-1.7	o			GSa		01He35	
	8927.4	15.			-0.6	o			GSa		09He20	
	8918.3	20.				6			GSa		10He11	
	8862.4	20.	8850	20	-0.6	o			GSa		85He22	
$^{253}\text{Lr}^m(\alpha)^{249}\text{Md}^m$	8862.4	10.			-1.2	o			GSa		01He35	
	8859.4	15.			-0.6	o			GSa		09He20	
	8850.2	20.				7			GSa		10He11	
$^{253}\text{Cf}(\beta^-)^{253}\text{Es}$	270	50	291	4	0.4	U					59Gh.A	
$^{253}\text{Md}^p(\text{IT})^{253}\text{Md}$	60	30				6					Ens139	
* $^{253}\text{Cf}(\alpha)^{249}\text{Cm}$	$E_\alpha=5981(5,Z)$ to $7/2^+$ level at 48.76 keV										Ens118 **	
* $^{253}\text{Cf}(\alpha)^{249}\text{Cm}$	$E_\alpha=5978.4(5,Z)$ , $5920.4(5,Z)$ to $7/2^+$ at 48.76, $7/2^+$ at 110.173 keV										Ens118 **	
* $^{253}\text{Fm}(\alpha)^{249}\text{Cf}$	$E_\alpha=7083.2(4,Z)$ , $6943.2(3,Z)$ , $6846.2(3,Z)$ , $6673.2(3,Z)$ keV to ground state and levels $5/2^+$ at 144.98, $9/2^+$ at 243.13, $1/2^+$ at 416.8 keV										Ens118 **	
*											Ens118 **	
* $^{253}\text{Fm}(\alpha)^{249}\text{Cf}$	$E_\alpha=6670(6)$ to 416.8 level										11Lo06 **	
* $^{253}\text{Md}(\alpha)^{249}\text{Es}$	$E_\alpha=7100(15)$ to 353.2(0.4) level										05He27 **	
* $^{253}\text{Md}(\alpha)^{249}\text{Es}$	$E_\alpha=7105(10)$ to 354.6(0.6) level										11Lo06 **	
* $^{253}\text{Md}(\alpha)^{249}\text{Es}$	$E_\alpha=7103(15)$ to 353.2(0.4) level										12He09 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8010(20)$ to 279.7 level										04He28 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8010(30)$ to 279.7 level										04He28 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8021(20)$ to 279.7 level										04He28 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8011(10)$ to 279.7 level										04He28 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8004(11)$ to 279.7(5) level										04He28 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8003(5)$ to 279.7(5) level; and 8280(10) to ground state										04He28 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8007(4)$ to 279.8(0.2) level; also $E_\alpha=7615(30)$ to 669(3) and										11Lo06 **	
*	$E_\alpha=8080(10)$ to 209.5(0.5) levels										11Lo06 **	
* $^{253}\text{No}(\alpha)^{249}\text{Fm}$	$E_\alpha=8004(10)$ to 279.5(5) level										12He09 **	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{254}\text{No}-^{133}\text{Cs}_{1,910}$	271552	15	271541	10	-0.7	o			SH1	1.0	10Dw01
	271544	16			-0.2	1	42	42 $^{254}\text{No}$	SH1	1.0	10Mi.A
$^{254}\text{Cf}(\alpha)^{250}\text{Cm}$	5926.9	5.				3					68Be21 Z
$^{254}\text{Es}(\alpha)^{250}\text{Bk}$	6615.7	1.5				6			Kum		72Bb24 *
$^{254}\text{Es}(\alpha)^{250}\text{Bk}^n$	6531.6	1.5				7			Kum		72Bb24
$^{254}\text{Es}^m(\alpha)^{250}\text{Bk}$	6699.9	2.0				5					73Ah04 *
$^{254}\text{Fm}(\alpha)^{250}\text{Cf}$	7306.8	5.	7307.5	1.9	0.1	3			Bka		64As01 Z
	7307.6	2.			-0.1	3					84Ah02 *
$^{254}\text{No}(\alpha)^{250}\text{Fm}$	8229.8	20.	8226	8	-0.2	o			Bka		67Gh01
	8240.0	30.			-0.5	U			DbA		67Mi03
	8215.6	20.			0.5	o			GSa		85He22
	8177.0	30.			1.6	U			Bka		06Fo02
	8225.7	10.			0.0	-			GSa		10He10
	8222.7	20.3			0.2	-			RIa		15KaZX
	ave.	8225	9		0.1	1	78	58 $^{254}\text{No}$			average
$^{254}\text{Lr}(\alpha)^{250}\text{Md}$	8804.7	20.	8816	12	0.6	o			GSa		85He22 *
	8804.7	20.			0.6	7			Lza		01Ga20 *
	8820.6	25.7			-0.2	7			Bka		06Fo02 *
	8825.2	20.			-0.4	7			GSa		08An16 *
$^{254}\text{Es}^m(\beta^-)^{254}\text{Fm}$	1172	2				4					62Un01 *
* $^{254}\text{Es}(\alpha)^{250}\text{Bk}$	$E_\alpha=6415.4(1.5,Z)$ to $5^-$ level at 97.49 keV										Ens01c **
* $^{254}\text{Es}^m(\alpha)^{250}\text{Bk}$	$E_\alpha=6558.9(2,Z), 6383.9(2,Z)$ to $4^+$ at 35.59, $2^+$ at 211.82 keV										Ens01c **
* $^{254}\text{Fm}(\alpha)^{250}\text{Cf}$	$E_\alpha=7192.3(2,Z), 7150.3(2,Z)$ to ground state, $5^+$ level at 42.721 keV										Ens01c **
* $^{254}\text{Lr}(\alpha)^{250}\text{Md}$	$E_\alpha=8460(20)$ to 209.1 level										08An16 **
* $^{254}\text{Lr}(\alpha)^{250}\text{Md}$	$E_\alpha=8460(20)$ to 209.1 level										08An16 **
* $^{254}\text{Lr}(\alpha)^{250}\text{Md}$	$E_\alpha=8437(50)$ to 209.1 and $E_\alpha=8394(30)$ to 306.2 keV										08An16 **
* $^{254}\text{Lr}(\alpha)^{250}\text{Md}$	$E_\alpha=8480(20)$ to 209.1 and $E_\alpha=8385(20)$ to 306.2 keV										08An16 **
* $^{254}\text{Es}^m(\beta^-)^{254}\text{Fm}$	$E_{\beta^-}=1127(2)$ to $2^+$ level at 45.000 keV										Ens05b **
$^{255}\text{No}-^{133}\text{Cs}_{1,917}$	274440	16				2			SH1	1.0	10Mi.A
$^{255}\text{Lr}-^{133}\text{Cs}_{1,917}$	277811	19				2			SH1	1.0	10Mi.A *
$^{255}\text{Es}(\alpha)^{251}\text{Bk}$	6439.3	3.0	6436.3	1.3	-1.0	4					66Rg01 *
	6435.6	1.5			0.5	4			Kum		71Bb10 *
$^{255}\text{Fm}(\alpha)^{251}\text{Cf}$	7237.0	4.	7239.7	1.8	0.7	3					64As01 *
	7240.4	2.			-0.3	3					75Ah01 *
$^{255}\text{Md}(\alpha)^{251}\text{Es}$	7901.8	5.	7905.9	2.6	0.8	4					70Fi12 *
	7910.7	5.			-1.0	4					71Ho16 *
	7905.4	4.			0.1	4			Ara		00Ah02 *
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	7891.2	15.			1.0	U			GSa		05He27 *
	8451.1	6.	8428	3	-3.8	B			ORb		71Di03 *
	8426.4	20.			0.1	o			GSa		98Ho13 *
	8391.9	35.			1.0	U			RIa		04Mo40 *
	8449.3	20.			-1.0	U			Bka		04Fo08 *
	8426.4	10.			0.2	U			GSa		06He20 *
	8403.8	60.			0.4	U			Bka		06Gr24 *
	8428.4	3.				3			JAa		11As03 *
$^{255}\text{Lr}(\alpha)^{251}\text{Md}$	8563.6	18.	8556	7	-0.4	3			ORb		76Be.A *
	8534.1	30.			0.7	U			Lza		01Ga20 *
	8554.4	10.			0.1	3			Jya		06Ch52
	8554.4	10.			0.1	3			Orm		08Ha31 *
$^{255}\text{Lr}(\alpha)^{251}\text{Md}^p$	8503.6	18.	8503	4	0.0	3			ORb		76Be.A *
	8442.7	50.			1.2	F			Bka		95Gh04 *
	8493.5	30.			0.3	U			Lza		01Ga20 *
	8498.6	5.			0.8	3			Jya		06Ch52
	8506.7	11.			-0.3	3			GSa		08An16 *
	8509.7	10.2			-0.7	3			Orm		08Ha31
	8509.7	10.2			-0.7	3			RIa		11As.A *
$^{255}\text{Lr}^m(\alpha)^{251}\text{Md}$	8592.0	5.	8597	4	0.9	4			Jya		06Ch52 *
	8602.2	11.			-0.5	4			GSa		08An16
	8603.1	10.2			-0.6	4			Orm		08Ha31 *
	8604.2	10.2			-0.7	4			RIa		11As.A *

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value	Adjusted value	$v_i$	Dg	Signf.	Main infl.	Lab	F	Reference
$^{255}\text{Rf}(\alpha)^{251}\text{No}$	9042 20	9055 4	0.7	10			Bka		69Gh01 *
	9053 15		0.2	o			GSa		85He06 *
	9064 20		-0.4	o			GSa		97He29 *
	9062 10		-0.7	o			GSa		01He35 *
	9056 4		-0.1	10			GSa		06He27 *
$^{255}\text{Rf}^m(\text{IT})^{255}\text{Rf}$	135 20	150 22	0.7	R					15An05 *
$^{255}\text{Lr}-^{133}\text{Cs}_{1,917}$	$D_M=277822(17) \mu\text{u}$ for mixture 71% ground state + 29% $^{255}\text{Lr}^m$ at 41(8); $M - A=89958(16)$ keV								
$^{255}\text{Es}(\alpha)^{251}\text{Bk}$	$E_\alpha=6303(3,Z) 6299.3(1.5,Z)$ respectively, to $(7/2^+)$ level at 35.5 keV								
$^{255}\text{Fm}(\alpha)^{251}\text{Cf}$	$E_\alpha=7121.5, 7018.5(4,Z)$ to ground state, $7/2^+$ level at 106.309 keV								
$^{255}\text{Fm}(\alpha)^{251}\text{Cf}$	$E_\alpha=7126.8, 7021.8(2,Z)$ to ground state, $7/2^+$ level at 106.309 keV								
$^{255}\text{Md}(\alpha)^{251}\text{Es}$	$E_\alpha=7323.5(5,Z) 7332.3(5,Z) 7327(4) 7313(15)$ respectively, to $7/2^-$ at 461.5 keV								
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	$E_\alpha=8312(9), 8121(6)$ to ground state and 199.9 level								
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	$E_\alpha=8296(20), 8092(20)$ to ground state and 199.9 level								
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	$E_\alpha=8060$ to 199.9 level; also $E_\alpha=7800$ keV								
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	$E_\alpha=8341, 8092$ to ground state and 199.9 level; also $E_\alpha=7873$ keV								
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	$E_\alpha=8290(20), 8095(10)$ to ground state and 199.9 level								
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	$E_\alpha=8150, 8000$ to 199.9 level								
$^{255}\text{No}(\alpha)^{251}\text{Fm}$	$E_\alpha=8100(3)$ to $^{251}\text{Fm}^m$ at 200.09(0.11) keV								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}$	This is the faint $\alpha$ from long-lived isomer to the $7/2^-$ ground state								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}$	Line is mixed with $^{254}\text{Lr}$ 's $\alpha$								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}$	As interpreted from Fig. 1								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}^p$	This is the most intense $\alpha$ from long-lived isomer to $1/2^-$								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}^p$	F : one event in a questionable $^{267}\text{Ds}$ decay chain								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}^p$	No $\gamma$ observed in coincidence								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}^p$	Original $E_\alpha=8371(10)$ corrected for recoil to $E_\alpha=8476(10)$								
$^{255}\text{Lr}(\alpha)^{251}\text{Md}^p$	Uncertainty estimated by evaluator								
$^{255}\text{Lr}^m(\alpha)^{251}\text{Md}$	Original error 2 keV increased for calibration								
$^{255}\text{Lr}^m(\alpha)^{251}\text{Md}$	Original $E_\alpha=8463(10)$ corrected for recoil to $E_\alpha=8468(10)$								
$^{255}\text{Lr}^m(\alpha)^{251}\text{Md}$	Uncertainty estimated by evaluator								
$^{255}\text{Rf}(\alpha)^{251}\text{No}$	$E_\alpha=8700(20)$ to 203 level								
$^{255}\text{Rf}(\alpha)^{251}\text{No}$	$E_\alpha=8766(15), 8715(15)$ to 142, 203 levels,								
$^{255}\text{Rf}(\alpha)^{251}\text{No}$	$E_\alpha=8905(20), 8739(20)$ to ground state, 203 level								
$^{255}\text{Rf}(\alpha)^{251}\text{No}$	$E_\alpha=8722(10)$ to 203(3) level								
$^{255}\text{Rf}(\alpha)^{251}\text{No}$	$E_\alpha=8716(4)$ to 203.6(0.2) level								
$^{255}\text{Rf}^m(\text{IT})^{255}\text{Rf}$	From 105 keV EC from K shell, error estim. by eval.								
$^{256}\text{Lr}-^{133}\text{Cs}_{1,925}$	280499 89			2			SH1	1.0	10Mi.A
$^{256}\text{Fm}(\alpha)^{252}\text{Cf}$	7027.3 5.			3					68Ho13 Z
$^{256}\text{Md}^m(\alpha)^{252}\text{Es}$	7834.6 20.	7900 50	1.2	B					71Ho16 *
	7896.6 16.			4					93Mo18 *
$^{256}\text{No}(\alpha)^{252}\text{Fm}$	7798.0 8.		2.0	B					00Ah02
	8553.9 30.	8582 5	0.9	U					67Fl05 *
	8553.9 20.		1.4	U			Bka		67Gh01 *
	8578.3 12.		0.3	5					81Be03
$^{256}\text{Lr}(\alpha)^{252}\text{Md}^p$	8582.3 6.		-0.1	5					90Ho03
	8787.6 20.	8771 11	-0.8	3					71Es01
	8761.1 25.		0.4	o			ORb		76Be.A
	8777.4 20.		-0.3	3			ORb		76Di.A
	8767.2 35.		0.1	3			RIa		04Mo26
$^{256}\text{Rf}(\alpha)^{252}\text{No}$	8749.9 20.		1.0	3			Bka		04Fo08
	8952.1 23.	8926 15	-1.1	o			GSa		85He06
	8929.8 20.		-0.2	o			GSa		97He29
	8925.7 15.2			2			GSa		10St14
$^{256}\text{Db}(\alpha)^{252}\text{Lr}$	9336.2 20.			8			Bka		08Ne01
$^{256}\text{Db}(\alpha)^{252}\text{Lr}^p$	9157.4 20.	9169 14	0.6	9			GSa		01He35
	9179.7 20.		-0.6	9			Bka		08Ne01 *
$^{256}\text{Md}^m(\alpha)^{252}\text{Es}$	Also $E_\alpha=7210(5,Z)$ keV to 520(20) level								
$^{256}\text{Md}^m(\alpha)^{252}\text{Es}$	Very weak line; more precise $E_\alpha$ to excited levels								
*	$\alpha$ summed with electrons								
$^{256}\text{No}(\alpha)^{252}\text{Fm}$	Probably mixture of two branches								
$^{256}\text{Db}(\alpha)^{252}\text{Lr}^p$	5 events $E_\alpha=9030 9060 9020 9040 9030$ keV								

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{257}\text{Fm}(\alpha)^{253}\text{Cf}$	6862.7	2.	6863.5	1.4	0.4	4			Bka		67As02 *
	6864.4	2.			-0.4	4				82Ah01 *	
$^{257}\text{Md}(\alpha)^{253}\text{Es}$	7549.3	5.	7557.6	1.0	1.7	U					70Fi12 *
	7557.6	1.				4					93Mo18 *
$^{257}\text{No}(\alpha)^{253}\text{Fm}$	8474.1	30.	8477	6	0.1	U					70Es02 *
	8480	30			-0.1	U		GSa			96Ho13 *
	8476.6	6.				3		JAA			05As05 *
$^{257}\text{Lr}(\alpha)^{253}\text{Md}^p$	9020.6	20.	9008	9	-0.6	7					71Es01
	9001.3	12.			0.5	o		ORb			76Be.A
	9001.3	12.			0.5	7		ORb			77Be36
	9015.5	15.2			-0.5	o		GSa			97He29
	9030.8	50.			-0.5	U		Lza			04Ga29
	9010.4	15.			-0.2	7		GSa			10St14
	9079.8	15.	9083	8	0.2	2		ORb			73Be33 *
$^{257}\text{Rf}(\alpha)^{253}\text{No}$	9083.7	15.			-0.1	o		GSa			85He06
	9044.0	15.			2.6	B		GSa			97He29
	9084.1	20.			-0.1	o		GSa			07St12 *
	9084.1	10.			-0.1	2		GSa			10St14 *
	9106.2	100.			-0.2	U		Ara			09Qi04 *
	9142.5	20.	9156	7	0.7	2		Bka			69Gh01
	9158.8	15.			-0.2	o		ORb			73Be33
$^{257}\text{Rf}^m(\alpha)^{253}\text{No}$	9155.8	8.			0.0	2		ORb			90Be.A
	9163.9	15.			-0.5	2		GSa			97He29
	9144.0	100.			0.1	U		Ara			09Qi04 *
	9112.1	20.	9207	20	4.7	B		GSa			85He22
	9218	10			-1.1	o		GSa			01He35 *
$^{257}\text{Db}(\alpha)^{253}\text{Lr}$	9209.2	20.			-0.1	o		GSa			09He20 *
	9206.6	20.				7		GSa			10He11
	9305.1	20.	9313	20	0.4	o		GSa			85He22
	9308.2	10.			0.5	o		GSa			01He35
$^{257}\text{Db}^m(\alpha)^{253}\text{Lr}^m$	9300.0	20.3			0.7	o		GSa			09He20
	9313.2	20.3				8		GSa			10He11
	1082	4				3					10Be16
$^{257}\text{Rf}^m(\text{IT})^{257}\text{Rf}^m$											Ens139 **
* $^{257}\text{Fm}(\alpha)^{253}\text{Cf}$	$E_\alpha=6518.5(2,Z)$ to $(9/2^+)$ level at 241.01 keV										Ens139 **
* $^{257}\text{Fm}(\alpha)^{253}\text{Cf}$	$E_\alpha=6756.5(3,Z), 6520.5(2,Z)$ to ground state, $(9/2^+)$ level at 241.01 keV										Ens139 **
* $^{257}\text{Md}(\alpha)^{253}\text{Es}$	$E_\alpha=7066(5,Z)$ to 371.4 level										93Mo18 **
* $^{257}\text{Md}(\alpha)^{253}\text{Es}$	$E_\alpha=7440(2), 7074(1)$ to ground state, 371.4 level										93Mo18 **
* $^{257}\text{No}(\alpha)^{253}\text{Fm}$	$E_\alpha=8320(30)$ to 22.3 keV										05As05 **
* $^{257}\text{No}(\alpha)^{253}\text{Fm}$	$E_\alpha=8340(20)$ ; one event only; may be summing with $e^-$										AHW **
* $^{257}\text{No}(\alpha)^{253}\text{Fm}$	$E_\alpha=8222(6)$ to 124.1(1) level; also $E_\alpha=8323(7)$ to 22.3 keV										05As05 **
* $^{257}\text{Rf}(\alpha)^{253}\text{No}$	$E_\alpha=8778(15)$ to 166.7 level										07St12 **
* $^{257}\text{Rf}(\alpha)^{253}\text{No}$	$E_\alpha=8778(20)$ and $8495(20)$ to 166.7 and 455 level										07St12 **
* $^{257}\text{Rf}(\alpha)^{253}\text{No}$	$E_\alpha=8778(10)$ to 166.7; and $8950(15)$ to ground state is sum with conversion $e^-$										10St14 **
* $^{257}\text{Rf}(\alpha)^{253}\text{No}$	$E_\alpha=8800(100)$ to 166.7 level										09Qi04 **
* $^{257}\text{Rf}^m(\alpha)^{253}\text{No}$	$E_\alpha=9000(100)$ and $8950(100)$ to ground state and 54(14) level										09Qi04 **
* $^{257}\text{Db}(\alpha)^{253}\text{Lr}$	$E_\alpha=9074(10)$ partly summed with conversion $e^-$										01He35 **
* $^{257}\text{Db}(\alpha)^{253}\text{Lr}$	$E_\alpha=8965(20)$ coinc. $E(\gamma)=102.2$ ; $E_\alpha=9066(20)$ summed with conversion $e^-$										09He20 **
$^{258}\text{Md}(\alpha)^{254}\text{Es}$	7266.8	5.	7271.3	1.9	0.9	7					70Fi12 *
	7272	2			-0.4	7					93Mo18 *
$^{258}\text{Lr}(\alpha)^{254}\text{Md}$	8870	50	8904	19	0.7	5			ORb		76Be.A *
	8910	20			-0.3	5				88Gr30 *	
$^{258}\text{Rf}(\alpha)^{254}\text{No}$	9192.8	30.5				2					08Ga08
$^{258}\text{Db}(\alpha)^{254}\text{Lr}$	9531.0	50.	9500	50	-0.6	o			GSa		97Ho14
	9500.6	15.				8				09He20	

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{258}\text{Db}(\alpha)^{254}\text{Lr}^p$	9445.7	15.	9437	14	-0.6	o		GSa			85He22
	9446.8	20.			-0.5	9		Lza			01Ga20
	9426.4	20.3			0.5	9		GSa			09He20
$^{258}\text{Db}^m(\alpha)^{254}\text{Lr}^q$	9341.1	10.				10		GSa			09He20
* $^{258}\text{Md}(\alpha)^{254}\text{Es}$	$E_\alpha=6713(5)$ to 447.9 level										93Mo18 **
* $^{258}\text{Md}(\alpha)^{254}\text{Es}$	$E_\alpha=6763(4)$ , 6718(2) to 403.8, 447.9 levels										93Mo18 **
* $^{258}\text{Lr}(\alpha)^{254}\text{Md}$	$E_\alpha=8648(10)$ in coincidence with X(L) not X(K) $\rightarrow E(\gamma)=90(50)$ keV										AHW **
* $^{258}\text{Lr}(\alpha)^{254}\text{Md}$	$E_\alpha=8752$ observed as sum of $\alpha$ 's and conversion electrons										AHW **
* $^{258}\text{Lr}(\alpha)^{254}\text{Md}$	Mass assignment confirmed										92Gr02 **
$^{259}\text{No}(\alpha)^{255}\text{Fm}$	7849.2	15.	7854	5	0.3	U					73Si40 *
	7869.6	15.			-1.0	U					93Mo18 *
	7854	5				4		JAA			13As02 *
$^{259}\text{Lr}(\alpha)^{255}\text{Md}^p$	8582.8	20.	8574	9	-0.4	6					71Es01
	8571.6	10.			0.2	6					92Ha22
	8577.7	29.			-0.1	U		Bka			92Kr01
$^{259}\text{Rf}(\alpha)^{255}\text{No}^p$	8999.2	20.	9030	11	1.5	o		Bka			69Gh01
	9030	20			0.0	4					81Be03 *
	9034.7	20.			-0.2	4		GSa			98Ho13
	9026.6	35.			0.1	4		RIa			04Mo40
	9026.6	20.3			0.2	4		Bka			04Fo08
	9017	60			0.2	U		Bka			06Gr24
	8940.4	11.			8.2	F		GSa			10Ni14 *
	8968.8	50.			1.2	U					12Zh04
$^{259}\text{Db}(\alpha)^{255}\text{Lr}$	9618.8	20.				3		Lza			01Ga20
$^{259}\text{Sg}(\alpha)^{255}\text{Rf}$	9771.2	30.5	9765	8	-0.1	o		GSa			85Mu11
	9807.7	23.			-0.9	U		Bka			09Fo02 *
	9784	50			-0.4	o		GSa			09He20 *
	9765.0	8.1				11		GSa			15An05
$^{259}\text{Sg}^m(\alpha)^{255}\text{Rf}$	9852.4	20.3				11		GSa			15An05
$^{259}\text{Sg}^m(\alpha)^{255}\text{Rf}^m$	9700.0	8.1	9702	8	0.3	12		GSa			15An05
* $^{259}\text{No}(\alpha)^{255}\text{Fm}$	suffer summing effect; highest $E_\alpha$ seen 7685(10); extra unc. added										WgM147**
* $^{259}\text{No}(\alpha)^{255}\text{Fm}$	suffer summing effect; highest $E_\alpha$ seen 7689(4); extra unc. added										WgM147**
* $^{259}\text{No}(\alpha)^{255}\text{Fm}$	$E_\alpha=7505(5)$ to 231.1 level										13As02 **
* $^{259}\text{Rf}(\alpha)^{255}\text{No}^p$	$E_\alpha=8870(20)$ ; partly summed $E_\alpha=8770(20)$ and $e^-$										AHW **
* $^{259}\text{Rf}(\alpha)^{255}\text{No}^p$	F : lifetime 107 ms is much shorter than $T=2.63$ s in Nubase										Nu16b **
* $^{259}\text{Sg}(\alpha)^{255}\text{Rf}$	One event only, resolution 23 keV; also a wide group at lower 9593(46)keV										09Fo02 **
* $^{259}\text{Sg}(\alpha)^{255}\text{Rf}$	One event with $E_\alpha$ 9050 in coincidence with 593 $\gamma$ ; also groups at										09He20 **
*	$E_\alpha=9607(10)$ , 9550(10) keV										09He20 **
$^{260}\text{Lr}(\alpha)^{256}\text{Md}^p$	8155.6	20.3				6					71Es01
$^{260}\text{Db}(\alpha)^{256}\text{Lr}$	9191.5	30.	9500#	40#	10.3	F		RIa			04Mo26 *
	9516.5	30.			-0.5	F		RIa			04Mo26 *
	9563.2	20.			-3.1	F		Bka			04Fo08 *
$^{260}\text{Db}(\alpha)^{256}\text{Lr}^p$	9283.1	20.	9271	13	-0.6	4		Bka			70Gh02
	9262.8	17.			0.5	4		ORB			77Be36
	9316.5	60.			-0.8	U		GSa			95Ho04 *
	9285.1	60.			-0.2	U		GSa			02Ho11 *
	9181.3	60.			1.5	U		Lza			04Ga29 *
	9310.4	60.			-0.7	U		RIa			04Mo26 *
$^{260}\text{Sg}(\alpha)^{256}\text{Rf}$	9923.0	30.	9901	10	-0.7	o		GSa			85Mu11
	9900.6	10.				3		GSa			09He20
$^{260}\text{Bh}(\alpha)^{256}\text{Db}$	10400.3	30.5				9					08Ne01 *
* $^{260}\text{Db}(\alpha)^{256}\text{Lr}$	Highest energy event; other two $E_\alpha=8810$ and 8500 keV										04Mo26 **
* $^{260}\text{Db}(\alpha)^{256}\text{Lr}$	F : not observed in experiments with greater statistics										77Be36 **
* $^{260}\text{Db}(\alpha)^{256}\text{Lr}^p$	Two events $E_\alpha=9200$ and 9146; error estimated by evaluator										FGK126 **
* $^{260}\text{Db}(\alpha)^{256}\text{Lr}^p$	Two events $E_\alpha=9156$ and 9129; error estimated by evaluator										FGK126 **
* $^{260}\text{Db}(\alpha)^{256}\text{Lr}^p$	Eight events out of 14, $E_\alpha=9170$ 9050 9340 9400 9010 9100 9140 9130 keV										04Mo26 **
* $^{260}\text{Db}(\alpha)^{256}\text{Lr}^p$	Two longer-lived $\alpha$ -escapes are assigned to the daughter										FGK126 **
* $^{260}\text{Bh}(\alpha)^{256}\text{Db}$	Other events $E_\alpha=10170$ , 10170 and 10190; 10080 and 10030										08Ne01 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{261}\text{Rf}(\alpha)^{257}\text{No}$	8652.9	20.	8650	50	-0.1	o			GSa		96Ho13	
	8652.9	20.			-0.1	o		GSa	02Ho11			
	8659	52			-0.3	o		GSa		03Tu05	*	
	8642.6	50.			0.1	4		GSa		08Dv02		
	8652.9	50.			-0.1	4		RIm		11Ha13	*	
	8642.6	60.			0.1	4		RIm		12Ha05	*	
$^{261}\text{Rf}^m(\alpha)^{257}\text{No}^p$	8409.1	20.	8415	15	0.3	6			Bka		70Gh01	
	8388.8	30.			0.9	o		GSa	98Tu01	*		
	8429.4	30.5			-0.5	6		Db		00La34		
	8470.0	50.			-1.1	U				08Ga08	*	
	8419.3	50.			-0.1	6		GSa		08Dv02		
	8409.1	50.			0.1	o		RIm		11Ha13		
$^{261}\text{Db}(\alpha)^{257}\text{Lr}^p$	8409.1	40.6			0.1	6		RIm		13Mu08		
	9069.2	20.	9068	14	-0.1	9			Bka		71Gh01	
	9069.2	40.			0.0	9		Lza	04Ga29			
	9066.2	20.			0.1	9		GSa		10St14		
$^{261}\text{Sg}(\alpha)^{257}\text{Rf}$	9709.0	30.	9714	15	0.2	o			GSa		85Mu11	
	9713.1	20.			0.0	F		GSa	95Ho03	*		
	9770.0	20.3			-2.8	o		GSa		07St12		
	9713.7	15.				3		GSa		10St14	*	
$^{261}\text{Bh}(\alpha)^{257}\text{Db}$	10562.1	25.	10500	50	-1.2	o			GSa		89Mu09	
	10507.3	75.			-0.1	o		Bka	06Fo02	*		
	10492.1	75.			0.2	8		Bka		08Ne08	*	
	10504.3	40.			-0.1	8		GSa		10He11	*	
* $^{261}\text{Rf}(\alpha)^{257}\text{No}$	Two events with $E_\alpha=8500(+70-30)$ keV										GAu	**
* $^{261}\text{Rf}(\alpha)^{257}\text{No}$	From direct production (fusion-evaporation)										11Ha13	**
* $^{261}\text{Rf}(\alpha)^{257}\text{No}$	Decay chain of $^{265}\text{Sg}$ , observation is independent of previous item										12Ha05	**
* $^{261}\text{Rf}^m(\alpha)^{257}\text{No}^p$	In addition 60% $E_\alpha=8380(30)$ keV										98Tu01	**
* $^{261}\text{Rf}^m(\alpha)^{257}\text{No}^p$	Single event, decay time 103.2 s										08Ga08	**
* $^{261}\text{Sg}(\alpha)^{257}\text{Rf}$	F : $\alpha$ 's to 157 level summed with conversion electron										FGK10a	**
* $^{261}\text{Sg}(\alpha)^{257}\text{Rf}$	$E_\alpha=9410(15)$ to 157(1) level										10St14	**
* $^{261}\text{Bh}(\alpha)^{257}\text{Db}$	$E_\alpha=10346(75)$ one event; error estimated by evaluator										06Fo02	**
* $^{261}\text{Bh}(\alpha)^{257}\text{Db}$	Highest $E_\alpha$ ; error estimated; others 10054, 10285, 10113, 10165, 9989										08Ne08	**
* $^{261}\text{Bh}(\alpha)^{257}\text{Db}$	Average of 2 highest 10331, 10355 as read from graph; error estimated										10He11	**
$^{262}\text{Db}(\alpha)^{258}\text{Lr}^p$	8794.5	20.	8806	11	0.6	7			Bka		71Gh01	
	8815.8	20.			-0.5	7			88Gr30			
	8804.7	20.			0.1	7		GSa		99Dr09		
	8875.8	20.			-1.4	o		RIa		09Mo12		
	8814.8	30.5			-0.3	7		RIa		14Ha04		
$^{262}\text{Sg}(\alpha)^{258}\text{Rf}$	9599.8	15.2				3		GSa		10Ac.A		
$^{262}\text{Bh}(\alpha)^{258}\text{Db}$	10216.2	25.	10319	15	4.1	F			GSa		89Mu09	*
	10300.5	25.4			0.7	o		GSa	97Ho14			
	10231.4	25.4			3.5	B		Bka		06Fo02		
	10239.2	30.			2.7	B		Bka		08Ne08	*	
$^{262}\text{Bh}^m(\alpha)^{258}\text{Db}$	10319.5	15.				9		GSa		09He20	*	
	10531.1	25.4	10530	50	0.1	o			GSa		89Mu09	
	10605.2	25.4			-1.4	o		GSa	97Ho14			
	10508.7	76.2			0.5	o		Bka		06Fo02	*	
	10544.3	76.2			-0.2	9		Bka		08Ne08		
10534.1	15.			0.0	9		GSa		09He20			
* $^{262}\text{Bh}(\alpha)^{258}\text{Db}$	F : not highest line, see reference										97Ho14	**
* $^{262}\text{Bh}(\alpha)^{258}\text{Db}$	$E_\alpha=10096, 10025, 10125$ keV										08Ne08	**
* $^{262}\text{Bh}(\alpha)^{258}\text{Db}$	$E_\alpha=10008(15)$ to 156.5 level										09He20	**
* $^{262}\text{Bh}^m(\alpha)^{258}\text{Db}$	Single event, error estimated by evaluator										GAu	**



**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{263}\text{Rf}(\alpha)^{259}\text{No}^p$	8022.2	40.6	8022	29	0.0	6					93Gr.C
	8022.2	40.6			0.0	6					99Ga.A
$^{263}\text{Db}(\alpha)^{259}\text{Lr}^p$	8484.3	27.				8			Bka		92Kr01
$^{263}\text{Sg}(\alpha)^{259}\text{Rf}$	9393.1	40.	9400	60	0.3	o			Bka		74Gh04
	9403.3	60.				5			Bka		06Gr24 *
$^{263}\text{Sg}(\alpha)^{259}\text{Rf}^q$	9200.2	40.	9200	60	-0.1	o			Bka		74Gh04
	9149.4	60.9			0.8	o			Bka		94Gr08
	9198.1	60.9				6			Bka		06Gr24 *
$^{263}\text{Sg}^m(\alpha)^{259}\text{Rf}^p$	9391.1	20.	9390	13	-0.1	8			GSa		98Ho13
	9382.9	50.8			0.1	o			Bka		03Gi05
	9393.1	35.			-0.1	8			RIa		04Mo40 *
	9388.0	20.			0.1	8			Bka		04Fo08
	9198.1	11.			17.5	F			GSa		10Ni14 *
$^{263}\text{Hs}(\alpha)^{259}\text{Sg}$	10733.5	60.				12			Bka		09Dr02
$^{263}\text{Hs}^m(\alpha)^{259}\text{Sg}$	11058.5	60.				12			Bka		09Dr02 *
* $^{263}\text{Sg}(\alpha)^{259}\text{Rf}$	Two events $E_\alpha=9290$ and $9230$ keV										06Gr24 **
* $^{263}\text{Sg}(\alpha)^{259}\text{Rf}^q$	Four events $E_\alpha=9010, 9100, 9060$ and $9060$ keV										06Gr24 **
* $^{263}\text{Sg}^m(\alpha)^{259}\text{Rf}^p$	Also lower $E_\alpha=9130, 9040, 9150$ keV										04Mo40 **
* $^{263}\text{Sg}^m(\alpha)^{259}\text{Rf}^p$	F : the $\alpha$ chain originating from $^{267}\text{Hs}$ is in conflict with other data										10Ni14 **
* $^{263}\text{Hs}^m(\alpha)^{259}\text{Sg}$	Assignment assumed by evaluator										GAu **
$^{264}\text{Bh}(\alpha)^{260}\text{Db}^p$	9767.3	20.	9760	18	-0.4	6			GSa		95Ho04 *
	9636.0	60.			2.1	U			Lza		04Ga29 *
	9737.8	35.5			0.6	6			RIa		04Mo26 *
$^{264}\text{Hs}(\alpha)^{260}\text{Sg}$	10870	210	10591	20	-1.3	o			GSa		87Mu15 *
	10590.8	20.				4			GSa		95Ho.B
	10966.4	80.			-4.7	B			RIa		11Sa41 *
* $^{264}\text{Bh}(\alpha)^{260}\text{Db}^p$	Three more events in reference $E_\alpha=9365, 9514$ and $9113$ keV										02Ho11 **
* $^{264}\text{Bh}(\alpha)^{260}\text{Db}^p$	Three more events $E_\alpha=9501, 9481, 9440$ keV										04Ga29 **
* $^{264}\text{Bh}(\alpha)^{260}\text{Db}^p$	Six events; also two $E_\alpha=9830$ keV										04Mo26 **
* $^{264}\text{Hs}(\alpha)^{260}\text{Sg}$	$Q_\alpha=11000(+100-300)$ derived from T(1/2), one event only										87Mu15 **
* $^{264}\text{Hs}(\alpha)^{260}\text{Sg}$	Also $E_\alpha=10610(40)$ keV										11Sa41 **
$^{265}\text{Sg}(\alpha)^{261}\text{Rf}^m$	8945.3	60.	8980	50	0.7	F			DbA		94La22 *
	8904.7	30.			1.5	F			GSa		96Ho13 *
	8975.7	30.			0.1	o			GSa		98Tu01
	9077.3	30.			-1.9	F			GSa		98Tu01 *
	9036.6	50.8			-1.1	o			GSa		03Tu05
	8985.9	50.			-0.1	6			GSa		08Du09
	8975.7	50.8			0.1	6			RIIm		12Ha05
$^{265}\text{Sg}^m(\alpha)^{261}\text{Rf}^p$	8823.5	50.	8810	40	-0.2	o			GSa		03Tu05
	8813.3	40.			0.0	o			GSa		06Dv01
	8823.5	50.			-0.2	6			GSa		08Dv02
	8843.8	40.			-0.7	o			RIa		08Mo09
	8823.5	50.			-0.2	o			RIa		12Ha05
	8792.9	81.2			0.3	6			RIa		13Su04
	9381.9	50.				11			Lza		04Ga29
$^{265}\text{Bh}(\alpha)^{261}\text{Db}^p$	10524.2	25.	10470	15	-2.1	o			GSa		87Mu15
	10468.3	20.			0.1	o			GSa		95Ho03
	10459.2	15.			0.7	o			GSa		99He11 *
	10470.3	15.2				4			GSa		09He20 *
$^{265}\text{Hs}^m(\alpha)^{261}\text{Sg}$	10712.0	20.	10700	15	-0.6	o			GSa		95Ho03
	10734.3	15.2			-2.3	o			GSa		99He11 *
	10699.8	15.				4			GSa		09He20
* $^{265}\text{Sg}(\alpha)^{261}\text{Rf}^m$	F : average but probably due to several groups, see reference										98Tu01 **
* $^{265}\text{Sg}(\alpha)^{261}\text{Rf}^m$	F : this event is not trusted, see reference from same group										02Ho11 **
* $^{265}\text{Sg}(\alpha)^{261}\text{Rf}^m$	F : most probably not from $^{265}\text{Sg}$										GAu **
* $^{265}\text{Hs}(\alpha)^{261}\text{Sg}$	Also $E_\alpha=10426(15)$ keV										99He11 **
* $^{265}\text{Hs}(\alpha)^{261}\text{Sg}$	Most intense line, also $E_\alpha=10282(15), 10428(15), 10573(15)$ keV										09He20 **
* $^{265}\text{Hs}^m(\alpha)^{261}\text{Sg}$	Also $E_\alpha=10726(15)$ keV										99He11 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference	
$^{266}\text{Sg}(\alpha)^{262}\text{Rf}$	8762.0	50.	8800#	100#	0.8	F			DbA	94La22	*	
	8904.1	40.			-2.6	F			GSa	98Tu01	*	
	8853.4	50.			-1.1	F			GSa	02Tu05	*	
$^{266}\text{Bh}(\alpha)^{262}\text{Db}^p$	9432	50	9380	30	-1.1	9			Bka	00Wi15	*	
	9245.3	81.2			1.6	9			Lza	06Qi03		
	9371.2	50.			0.1	9			Rla	09Mo12	*	
$^{266}\text{Hs}(\alpha)^{262}\text{Sg}$	10335.7	20.3	10346	16	0.5	4			GSa	01Ho06		
	10360	24			-0.6	4			GSa	11Ac.A		
$^{266}\text{Hs}^m(\alpha)^{262}\text{Sg}^p$	10592.6	20.				7			GSa	11Ac.A		
$^{266}\text{Mt}(\alpha)^{262}\text{Bh}$	10995.7	25.				10			GSa	97Ho14		
$^{266}\text{Mt}^m(\alpha)^{262}\text{Bh}^m$	11269.7	50.	11920	50	13.0	B			GSa	84Mu07	*	
	11168.1	30.			25.0	B			GSa	89Mu16		
	11918.6	50.				10			GSa	97Ho14	*	
$*^{266}\text{Sg}(\alpha)^{262}\text{Rf}$	Average of two groups										02Tu05	**
$*^{266}\text{Sg}(\alpha)^{262}\text{Rf}$	F : no $\alpha$ decay from $^{266}\text{Sg}$ , all re-assigned to $^{265}\text{Sg}$										08Dv02	**
$*^{266}\text{Bh}(\alpha)^{262}\text{Db}^p$	Also $E_\alpha=9770(40)$ , $9080(40)$ keV from $^{278}\text{Ed}$ decay chain										08Mo09	**
$*^{266}\text{Mt}^m(\alpha)^{262}\text{Bh}^m$	One $E_\alpha$ only; may be ground state										AHW	**
$*^{266}\text{Mt}^m(\alpha)^{262}\text{Bh}^m$	One $E_\alpha=11739(50)$ , one 11306; several smaller										AHW	**
$^{267}\text{Sg}(\alpha)^{263}\text{Rf}^p$	8325.0	50.				8			GSa	08Dv02		
	$^{267}\text{Bh}(\alpha)^{263}\text{Db}^p$	8964.5	30.5	8970	26	0.2	10			Bka	00Wi15	
8984.8		50.			-0.3	10			GSa	02Tu05		
$^{267}\text{Hs}(\alpha)^{263}\text{Sg}$	10015.3	60.	10038	13	0.4	6			DbA	95La20	*	
	10032.6	20.			0.2	6			GSa	98Ho13		
	10035.7	50.			0.0	o			Bka	03Gi05		
	10069.1	35.			-0.9	6			Rla	04Mo40	*	
	10034.6	20.			0.1	6			Bka	04Fo08	*	
$^{267}\text{Hs}(\alpha)^{263}\text{Sg}^m$	10145.2	50.			-2.2	U			GSa	10Ni14	*	
	9978.8	20.	9987	13	0.4	7			GSa	98Ho13	*	
	9979.8	35.			0.2	7			Rla	04Mo40	*	
$^{267}\text{Hs}^m(\alpha)^{263}\text{Sg}^p$	9997.0	20.			-0.5	7			Bka	04Fo08	*	
	9979.8	20.				9			Bka	04Fo08	*	
$^{267}\text{Ds}(\alpha)^{263}\text{Hs}$	11776.8	50.8				13			Bka	95Gh04	*	
$*^{267}\text{Hs}(\alpha)^{263}\text{Sg}$	Selecting two events at $E_\alpha=9860,9870$ ; and one $E_\alpha=9740(60)$ keV										95La20	**
$*^{267}\text{Hs}(\alpha)^{263}\text{Sg}$	Selecting four events at $E_\alpha=9970, 9890, 9900, 9910$ keV										04Mo40	**
$*^{267}\text{Hs}(\alpha)^{263}\text{Sg}$	Selecting two events at $E_\alpha=9880, 9888$ keV										04Fo08	**
$*^{267}\text{Hs}(\alpha)^{263}\text{Sg}$	Directly produced $^{267}\text{Hs}$ ; daughter and grand-daughter also conflicting										10Ni14	**
$*^{267}\text{Hs}(\alpha)^{263}\text{Sg}^m$	And one $E_\alpha=9749(20)$ at 13 ms										98Ho13	**
$*^{267}\text{Hs}(\alpha)^{263}\text{Sg}^m$	Selecting 7 events at $E_\alpha=9830, 9820, 9820, 9860, 9820, 9830, 9830$ keV										04Mo40	**
$*^{267}\text{Hs}(\alpha)^{263}\text{Sg}^m$	Selecting 2 events at $E_\alpha=9864, 9830$ keV										04Fo08	**
$*^{267}\text{Ds}(\alpha)^{263}\text{Hs}$	Maybe the upper isomer at about 250 keV excitation energy										AHW	**
$^{268}\text{Hs}(\alpha)^{264}\text{Sg}$	9622.9	16.				8			GSa	10Ni14		
$^{268}\text{Mt}(\alpha)^{264}\text{Bh}^p$	10395.5	20.	10438	19	2.1	o			GSa	95Ho04	*	
	10431.9	20.			0.3	8			GSa	02Ho11	*	
	10476.7	50.			-0.8	8			Rla	04Mo26	*	
	10268	20			8.5	B			Bka	04Fo08	*	
$*^{268}\text{Mt}(\alpha)^{264}\text{Bh}^p$	Two events $E_\alpha=10221$ coinc. $E(\gamma)=93$ and $10259$ ; event #3 $E_\alpha=10097$ keV could be decay of an isomer with lifetime=171 ms										95Ho04	**
$*^{268}\text{Mt}(\alpha)^{264}\text{Bh}^p$	Average of event 95Ho04 $E_\alpha=10259$ and present 10294 keV										02Ho11	**
$*^{268}\text{Mt}(\alpha)^{264}\text{Bh}^p$	Also $E_\alpha=10340$ keV										04Mo26	**
$*^{268}\text{Mt}(\alpha)^{264}\text{Bh}^p$	One event only										04Fo08	**

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{269}\text{Sg}(\alpha)^{265}\text{Rf}$	8699.6	100.	8650	50	-1.0	7					10E106
	8628.5	60.9			0.4	7			DbA		15U102
$^{269}\text{Hs}(\alpha)^{265}\text{Sg}^m$	9369.6	30.	9280	30	-3.0	B			GSa		96Ho13 *
	9349.3	30.			-2.3	o			GSa		02Ho11 *
	9278.2	50.8			0.0	o			GSa		03Tu05 *
	9207.1	30.5			2.4	o			GSa		06Dv01
	9268.0	50.8			0.2	7			GSa		08Dv02
	9349.2	71.1			-1.0	o			RIa		08Mo09
	9288.3	40.6			-0.2	7			RIa		13Su04
$^{269}\text{Ds}(\alpha)^{265}\text{Hs}^m$	11280.1	20.				5			GSa		95Ho03
$^{*269}\text{Hs}(\alpha)^{265}\text{Sg}^m$	Event number 2 only; first event rejected, see reference										02Ho11 **
$^{*269}\text{Hs}(\alpha)^{265}\text{Sg}^m$	Two events $E_\alpha=9230, 9180$ both following $300 \mu\text{s } ^{273}\text{Ds}$										02Ho11 **
$^{*269}\text{Hs}(\alpha)^{265}\text{Sg}^m$	Three events $E_\alpha=9180, 9100, 8880$ ; latter probably due to energy loss										03Tu05 **
$^{270}\text{Db}(\alpha)^{266}\text{Lr}$	8019.0	30.5	8260#	200#	4.8	D			GSa		14Kh04 *
$^{270}\text{Bh}(\alpha)^{266}\text{Db}$	9064.5	81.2				10			DbA		07Og02
$^{270}\text{Hs}(\alpha)^{266}\text{Sg}$	9324.3	52.8	9070	40	-4.8	F			GSa		03Tu05 *
	9024	52			1.5	o			GSa		06Dv01 *
	9013.8	50.			1.1	7			GSa		08Dv02 *
	9123.4	77.			-0.7	7			GSa		10Gr04 *
	9155.8	80.			-1.1	7			DbA		13Og03
$^{270}\text{Mt}(\alpha)^{266}\text{Bh}$	10181.1	70.				10			RIa		08Mo09
$^{270}\text{Ds}(\alpha)^{266}\text{Hs}$	11196.2	50.8	11117	28	-1.6	o			GSa		01Ho06
	11117.0	28.4				5			GSa		11Ac.A
$^{270}\text{Ds}^m(\alpha)^{266}\text{Hs}$	12333	50	12510	50	3.5	B			GSa		01Ho06
	12508.6	20.				5			GSa		11Ac.A
$^{270}\text{Ds}^m(\alpha)^{266}\text{Hs}^m$	11318	50	11410	50	1.7	o			GSa		01Ho06
	11405.2	52.				6			GSa		11Ac.A
$^{*270}\text{Db}(\alpha)^{266}\text{Lr}$	Trends from Mass Surface TMS suggest $^{270}\text{Db}$ 240 less bound										GAu **
$^{*270}\text{Hs}(\alpha)^{266}\text{Sg}$	F : re-assigned to $^{269}\text{Hs}$										06Dv01 **
$^{*270}\text{Hs}(\alpha)^{266}\text{Sg}$	4 events at 8850(1.62s), 8900(0.05s), 8920(0.5s), 8880(0.4s)										GAu **
$^{*270}\text{Hs}(\alpha)^{266}\text{Sg}$	2 events at 8760(0.27s), 8810(0.27s); $E_\alpha=8880(50)$ for both experiments										GAu **
$^{*270}\text{Hs}(\alpha)^{266}\text{Sg}$	Symmetrized from $E_\alpha=9020(+50-100)$ ; independent from previous item										GAu **
$^{271}\text{Sg}(\alpha)^{267}\text{Rf}^p$	8658	80	8670	50	0.2	o			DbA		04Og12
	8668.2	81.2				10			DbA		06Og05
$^{271}\text{Bh}(\alpha)^{267}\text{Db}$	9490.3	162.4	9420	50	-1.4	8			DbA		12St.A
	9409.1	71.0			0.3	8			GSa		13Ru11
$^{271}\text{Hs}(\alpha)^{267}\text{Sg}^p$	9439.6	50.7				10			GSa		08Dv02
$^{271}\text{Ds}(\alpha)^{267}\text{Hs}$	10869.8	20.	10870	18	0.0	7			GSa		98Ho13
	10870.8	35.			0.0	7			RIa		04Mo40 *
$^{271}\text{Ds}^m(\alpha)^{267}\text{Hs}$	10937.8	20.				7			Bka		04Fo08 *
	10803.8	50.	10938	20	2.7	F					12Zh04 *
$^{271}\text{Ds}^m(\alpha)^{267}\text{Hs}^m$	10899.2	20.	10899	13	0.0	8			GSa		98Ho13
	10880.8	50.			0.3	o			Bka		03Gi05
	10883.0	35.			0.4	8			RIa		04Mo40 *
	10903.3	20.			-0.2	8			Bka		04Fo08 *
$^{*271}\text{Ds}(\alpha)^{267}\text{Hs}$	Decay chain number 6 for the long-lived isomer, GAU interpretation										04Mo40 **
$^{*271}\text{Ds}^m(\alpha)^{267}\text{Hs}$	Decay chain number 6, GAU interpretation										04Fo08 **
$^{*271}\text{Ds}^m(\alpha)^{267}\text{Hs}$	F : $\alpha$ escaped ?										GAu **
$^{*271}\text{Ds}^m(\alpha)^{267}\text{Hs}^m$	GAU : average of decay chains number 2, 5, 10, 13 for short-lived isomer										04Mo40 **
$^{*271}\text{Ds}^m(\alpha)^{267}\text{Hs}^m$	Decay chains number 5 and 7, GAU interpretation										04Fo08 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{272}\text{Bh}(\alpha)^{268}\text{Db}$	9303.0	20.				8			Bka		15Ga24 *
$^{272}\text{Bh}(\alpha)^{268}\text{Db}^p$	9154.9	60.	9160	50	0.0	o			DbA		04Og03
	9144.7	60.9			0.2	o			DbA		12Og02
	9126.4	30.4			0.6	9			DbA		13Og01 *
	9187.3	30.4			-0.6	9			GSa		13Ru11 *
$^{272}\text{Rg}(\alpha)^{268}\text{Mt}$	10981.9	20.	11197	13	10.8	B			GSa		95Ho04 *
	11191.9	20.			0.3	9			GSa		02Ho11 *
	11184.7	35.			0.4	9			RIa		04Mo26 *
	11207.1	20.3			-0.5	9			Bka		04Fo08 *
$^{*272}\text{Bh}(\alpha)^{268}\text{Db}$	Average 9.16(2) 9.20(5) 9.18(5) 9.16(5)										15Ga24 **
$^{*272}\text{Bh}(\alpha)^{268}\text{Db}^p$	Average $E_\alpha$ of 20 events; error increased										13Og01 **
$^{*272}\text{Bh}(\alpha)^{268}\text{Db}^p$	Average $E_\alpha$ of 8 events; error increased										13Ru11 **
$^{*272}\text{Rg}(\alpha)^{268}\text{Mt}$	B : one event only; E(K) in coincidence may explain disagreement										GAu **
$^{*272}\text{Rg}(\alpha)^{268}\text{Mt}$	Two events $E_\alpha=11008$ and $11046$ keV										02Ho11 **
$^{*272}\text{Rg}(\alpha)^{268}\text{Mt}$	Also others up to $E_\alpha=11560$ keV										04Mo26 **
$^{*272}\text{Rg}(\alpha)^{268}\text{Mt}$	One event only										04Fo08 **
$^{273}\text{Hs}(\alpha)^{269}\text{Sg}$	9732.9	40.	9700	50	-0.6	8					10EI06
	9671.9	40.6			0.6	8			DbA		15Uu02
$^{273}\text{Ds}(\alpha)^{269}\text{Hs}$	9875.0	20.	11370	50	29.8	F			GSa		96Ho13 *
	11248.1	30.			2.4	F			GSa		96Ho13 *
	11519.1	60.			-3.0	F			DbA		96La12 *
	11366.9	20.3				8			GSa		02Ho11 *
	11311.0	70.			1.1	U			RIa		08Mo09
	11194.3	81.2			3.5	B			RIa		13Su04
$^{*273}\text{Ds}(\alpha)^{269}\text{Hs}$	F : this event is distrusted, see reference										02Ho11 **
$^{*273}\text{Ds}(\alpha)^{269}\text{Hs}$	F : event number 2, probably to excited state in $^{269}\text{Hs}$										GAu **
$^{*273}\text{Ds}(\alpha)^{269}\text{Hs}$	F : this event is distrusted, see reference; average 4 others $E_\alpha=11720$ keV										02Ho11 **
$^{*273}\text{Ds}(\alpha)^{269}\text{Hs}$	And one $E_\alpha=11080$ keV										02Ho11 **
$^{274}\text{Bh}(\alpha)^{270}\text{Db}$	8930.6	101.5	8950	50	0.4	o			DbA		11Og04 *
	8890.0	50.7			1.2	6			DbA		13Og04
	8971.2	30.5			-0.4	6			GSa		14Kh04
$^{274}\text{Mt}(\alpha)^{270}\text{Bh}^p$	9904.8	101.5				12			DbA		07Og02
$^{274}\text{Rg}(\alpha)^{270}\text{Mt}$	11477.9	70.				11			RIa		08Mo09 *
$^{*274}\text{Bh}(\alpha)^{270}\text{Db}$	All results from this work were first published in reference										10Og01 **
$^{*274}\text{Rg}(\alpha)^{270}\text{Mt}$	Also one $E_\alpha=11150(70)$ keV										08Mo09 **
$^{275}\text{Hs}(\alpha)^{271}\text{Sg}$	9437.5	71.0	9440	50	0.0	o			DbA		04Og12
	9437.5	60.9				11			DbA		06Og05
$^{275}\text{Mt}(\alpha)^{271}\text{Bh}$	10482.8	90.	10480	50	0.0	U			DbA		04Og03
	10503.0	60.			-0.4	U			DbA		12St.A
	10482.7	10.1				9			GSa		13Ru11
$^{276}\text{Mt}(\alpha)^{272}\text{Bh}$	10253	65	10100	9	-2.3	o			DbA		04Og03 *
	10212	84			-1.3	U			DbA		12Og02 *
	10160	16			-3.7	B			DbA		13Og01 *
	10102.3	10.1			-0.2	9			GSa		13Ru11 *
	10093.0	20.			0.4	9			Bka		15Ga24 *
$^{276}\text{Mt}^m(\alpha)^{272}\text{Bh}$	10354	84				9			DbA		12Og02 *
$^{*276}\text{Mt}(\alpha)^{272}\text{Bh}$	$E_\alpha=9710(60)$ to 434 and 363 keV levels										13Ru11 **
$^{*276}\text{Mt}(\alpha)^{272}\text{Bh}$	$E_\alpha=9670(80)$ to 434 and 363 keV levels, and one 9260(64) keV										13Ru11 **
$^{*276}\text{Mt}(\alpha)^{272}\text{Bh}$	Average for 18 events $E_\alpha=9622(16)$ to 434 and 363 keV levels										13Ru11 **
$^{*276}\text{Mt}(\alpha)^{272}\text{Bh}$	$E_\alpha=9530(10)$ to 434 keV level; also $E_\alpha=9600(10)$ to 362 keV level										13Ru11 **
$^{*276}\text{Mt}(\alpha)^{272}\text{Bh}$	Average of 9 events $E_\alpha=9570-9630=9590$ to 362 keV level										15Ga24 **
$^{*276}\text{Mt}(\alpha)^{272}\text{Bh}$	also one event $E_\alpha=9890(20)$ to 60 keV level										15Ga24 **
$^{*276}\text{Mt}^m(\alpha)^{272}\text{Bh}$	$E_\alpha=9810(80)$ to 434 and 363 keV levels										13Ru11 **

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{277}\text{Ds}(\alpha)^{273}\text{Hs}^p$	10725.2	40.	10720	50	-0.2	10					10El06
	10704.8	40.6			0.2	10			Db		15Ut02
$^{277}\text{Cn}(\alpha)^{273}\text{Ds}$	11622.2	30.				9			GSa		96Ho13 *
	11821.0	30.	11620	50	-4.0	F			GSa		96Ho13 *
	11486.2	40.6			2.7	C			RIa		04MoZU
	11486.2	40.			2.7	B			RIa		08Mo09 *
	11232.4	81.2			7.8	B			RIa		13Su04
* $^{277}\text{Cn}(\alpha)^{273}\text{Ds}$	And one $E_\alpha=11170(20)$ $Q_\alpha=11334.(20.)$ keV										
* $^{277}\text{Cn}(\alpha)^{273}\text{Ds}$	F : this event is distrusted, see reference										
* $^{277}\text{Cn}(\alpha)^{273}\text{Ds}$	And one $E_\alpha=11090(70)$ keV										
$^{278}\text{Mt}(\alpha)^{274}\text{Bh}$	9689.7	190.	9630	50	-1.2	U			Db		11Og04
	9527.3	71.			2.0	7			Db		12Og06
	9689.6	30.4			-1.2	7			Db		13Og04
	9588.2	30.			0.8	7			GSa		14Kh04
	10846.3	81.2				13			Db		07Og02
$^{278}\text{Rg}(\alpha)^{274}\text{Mt}$	11850.8	40.				12			RIa		08Mo09 *
	11992.8	61.	11850	50	-2.8	o			RIa		12Mo25 *
* $^{278}\text{Ed}(\alpha)^{274}\text{Rg}$	Also one $E_\alpha=11520(40)$ keV										
* $^{278}\text{Ed}(\alpha)^{274}\text{Rg}$	Post-deadline, disagrees with previous result from same group										
$^{279}\text{Ds}(\alpha)^{275}\text{Hs}^p$	9841.3	60.9	9840	50	0.0	o			Db		04Og12
	9841.3	60.9				13			Db		06Og05
$^{279}\text{Rg}(\alpha)^{275}\text{Mt}$	10521.1	162.3				10			Db		04Og03
$^{280}\text{Rg}(\alpha)^{276}\text{Mt}$	10128.6	60.	10146	7	0.3	o			Db		04Og03 *
	10128.6	60.			0.3	U			Db		12Og02 *
	10148.8	10.2			-0.3	10			GSa		13Ru11 *
	10088.0	15.2			3.8	B			Db		13Og01 *
	10148.8	10.2			-0.3	10			Bka		15Ga24 *
	10124.8	20.			1.0	10			Bka		15Ga24 *
	* $^{280}\text{Rg}(\alpha)^{276}\text{Mt}$	$E_\alpha=9750(60)$ to 237 keV level									
* $^{280}\text{Rg}(\alpha)^{276}\text{Mt}$	$E_\alpha=9750(60)$ to 237 keV level										
* $^{280}\text{Rg}(\alpha)^{276}\text{Mt}$	$E_\alpha=9770(10)$ to 237 keV level										
* $^{280}\text{Rg}(\alpha)^{276}\text{Mt}$	Average of 19 events $E_\alpha=9710(15)$ to 237 keV level										
* $^{280}\text{Rg}(\alpha)^{276}\text{Mt}$	$E_\alpha=9770(10)$ to 237 keV level										
* $^{280}\text{Rg}(\alpha)^{276}\text{Mt}$	Highest $E_\alpha=9980(20)$										
$^{281}\text{Ds}(\alpha)^{277}\text{Hs}^p$	8957.8	180.	8850	50	-2.1	F			Db		99Og10 *
	8825.9	100.			0.5	F			Db		04Mo15 *
	8856.3	30.4			-0.1	o			GSt		10Du06
	8853.3	25.			0.0	5			GSt		11Ga19
	8853.3	15.			0.0	5			GSa		12Ho12
	9449.8	15.				5			GSa		12Ho12 *
$^{281}\text{Ds}^m(\alpha)^{277}\text{Hs}^m$	9454.8	40.6	9900#	400#	9.0	D			Db		13Og04 *
$^{281}\text{Rg}(\alpha)^{277}\text{Mt}$	10459.2	40.	10450	50	-0.1	11					10El06
$^{281}\text{Cn}(\alpha)^{277}\text{Ds}$	10448.9	40.6			0.1	11			Db		15Ut02
* $^{281}\text{Ds}(\alpha)^{277}\text{Hs}^p$	F : wrong $\alpha$ chain, see $^{285}\text{Cn}$ and $^{289}\text{Fl}$										
* $^{281}\text{Ds}(\alpha)^{277}\text{Hs}^p$	F : non tracable information										
* $^{281}\text{Ds}^m(\alpha)^{277}\text{Hs}^m$	Assignment of $^{293}\text{Lv}^m$ - $^{289}\text{Fl}^m$ - $^{285}\text{Cn}^m$ - $^{281}\text{Ds}^m$ - $^{277}\text{Hs}^m$ chain is tentative										
* $^{281}\text{Rg}(\alpha)^{277}\text{Mt}$	Trends from Mass Surface TMS suggest $^{281}\text{Rg}$ 450 less bound										

**Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)**

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{282}\text{Rg}(\alpha)^{278}\text{Mt}^p$	9129.7	101.4	9170	50	0.8	o			Db		11Og04
	9139.8	50.7			0.6	9			Db		13Og04
	9180.4	30.4			-0.2	9			GS		14Kh04 *
$^{282}\text{Cn}(\alpha)^{278}\text{Ds}$	9667.4	100.	10170#	200#	10.1	F			Db		04Mo15 *
$^{282}\text{Ed}(\alpha)^{278}\text{Rg}$	10783.2	81.2				14			Db		07Og02
* $^{282}\text{Rg}(\alpha)^{278}\text{Mt}^p$	also observed $E_\alpha=8860(30)$ keV										14Kh04 **
* $^{282}\text{Cn}(\alpha)^{278}\text{Ds}$	F : non tracable information										GAU **
$^{283}\text{Cn}(\alpha)^{279}\text{Ds}^p$	9677	70	9660	50	-0.4	o			Db		04Og07
	9677.0	60.9			-0.4	o			Db		04Og12
	9677.0	60.9			-0.4	15			Db		06Og05
	9606.0	60.9			1.0	15					07Ei02
	9656.7	15.2			0.0	15			GS		07Ho18
	9788.6	100.			-2.7	U			Bk		09St21
$^{283}\text{Ed}(\alpha)^{279}\text{Rg}^p$	10265.4	90.	10380	50	2.2	U			Db		04Og03
	10376.9	10.1				12			GS		13Ru11
$^{284}\text{Cn}(\alpha)^{280}\text{Ds}$	9301.3	50.	9600#	200#	6.0	F			Db		01Og01 *
	9269.0	100.			3.3	F			Db		04Mo15 *
$^{284}\text{Ed}(\alpha)^{280}\text{Rg}$	10254.6	20.3	10280	50	0.5	11			GS		13Ru11 *
	10305.3	20.			-0.5	11			Bk		15Ga24 *
* $^{284}\text{Cn}(\alpha)^{280}\text{Ds}$	F : no $\alpha$ observed in later work by same group; re-assigned to $^{285}\text{Cn}$										04Og07 **
* $^{284}\text{Cn}(\alpha)^{280}\text{Ds}$	F : non tracable information										GAU **
* $^{284}\text{Ed}(\alpha)^{280}\text{Rg}$	Highest $E_\alpha=10110(10)$ , error increased										13Ru11 **
* $^{284}\text{Ed}(\alpha)^{280}\text{Rg}$	Highest $E_\alpha=10160(20)$										15Ga24 **
$^{285}\text{Cn}(\alpha)^{281}\text{Ds}$	8793.7	50.	9320	50	10.5	F			Db		99Og10 *
	8793.7	100.			10.5	F			Db		04Mo15 *
	9290.6	60.9			0.5	o			Db		04Og07
	9280.5	50.			0.7	6			Db		07Og01
	9341.3	30.4			-0.5	o			GSt		10Du06
	9341.3	30.4			-0.5	6			GSt		11Ga19
	9314.9	15.			0.0	6			GS		12Ho12
$^{285}\text{Cn}^m(\alpha)^{281}\text{Ds}^m$	9845.4	15.2				6			GS		12Ho12 *
$^{285}\text{Ed}(\alpha)^{281}\text{Rg}$	9878.9	81.1	10010	50	2.6	B			Db		11Og04 *
	10026.9	62.			-0.4	o			Db		12Og02 *
	9767.3	183.			4.8	B			Db		12Og06
	10031.0	162.			-0.4	o			Db		13Og01
	9990.4	41.			0.4	o			Db		13Og04
	10008.7	20.3				11			Db		16Fo16 *
	10558.4	50.7				12			Db		15Ut02
* $^{285}\text{Cn}(\alpha)^{281}\text{Ds}$	F : one event at 15.4 m, later work yields much shorter half-lives										GAU **
* $^{285}\text{Cn}(\alpha)^{281}\text{Ds}$	F : non tracable information										GAU **
* $^{285}\text{Cn}^m(\alpha)^{281}\text{Ds}^m$	Assignment of $^{293}\text{Lv}^m-^{289}\text{Fl}^m-^{285}\text{Cn}^m-^{281}\text{Ds}^m-^{277}\text{Hs}^m$ chain is tentative										12Ho12 **
* $^{285}\text{Ed}(\alpha)^{281}\text{Rg}$	And $E_\alpha=9480(110)$ keV										11Og04 **
* $^{285}\text{Ed}(\alpha)^{281}\text{Rg}$	Also $E_\alpha=9740(80)$ and $9480(110)$ keV										12Og02 **
* $^{285}\text{Ed}(\alpha)^{281}\text{Rg}$	reanalyzed data of 11Og04, 12Og02, 12Og06, 13Og01 and 13Og04										16Fo16 **
*	unweighted average of the 4 highest energy events as selected by										GAU **
*	the evaluator: s06=9857 s11=9902 s12=9845 s13=9867 in Table 2										GAU **
$^{286}\text{Ed}(\alpha)^{282}\text{Rg}$	9766.9	100.	9790	50	0.5	10			Db		11Og04
	9817.5	111.6			-0.6	10			Db		12Og06
	9432.1	304.3			7.1	B			GS		14Kh04

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{286}\text{Fl}(\alpha)^{282}\text{Cn}$	10142.1	100.	10370	30	2.3	F			DbA		04Mo15 *
	10172.5	314.4			0.6	o			DbA		04Og07
	10345	60			0.4	o			DbA		04Og12
	10334.7	60.9			0.6	11			DbA		06Og05
	10375.4	40.			-0.1	11			BkA		09St21
	10456.5	100.			-0.8	11			BkA		10El06
* $^{286}\text{Fl}(\alpha)^{282}\text{Cn}$	F : non traceable information										GAU **
$^{287}\text{Fl}(\alpha)^{283}\text{Cn}$	10435.7	20.3	10160	50	-5.5	F			DbA		99Og07 *
	10182.2	70.			-0.4	o			DbA		04Og07
	10161.9	60.8			0.0	o			DbA		04Og12
	10161.9	60.8				16			DbA		06Og05
$^{287}\text{Ef}(\alpha)^{283}\text{Ed}$	10740	90	10760	50	0.5	o			DbA		04Og03
	10740	60			0.5	13			DbA		12St.A
	10780.5	50.7			-0.3	13			GSa		13Ru11
* $^{287}\text{Fl}(\alpha)^{283}\text{Cn}$	F : 2 evts at 1.32 s, 14.4 s, later work yields T=1.7 s										GAU **
$^{288}\text{Fl}(\alpha)^{284}\text{Cn}$	9968.8	50.	10072	13	2.1	F			DbA		01Og01 *
	9958.8	100.			1.1	F			DbA		04Mo15 *
	10090.3	80.			-0.2	o			DbA		04Og07
	10090.3	70.			-0.3	o			DbA		04Og12
	10080.2	60.8			-0.1	11			DbA		07Og01
	10090.3	30.			-0.6	o			GSt		10Du06
	10090.3	30.			-0.6	11			GSt		11Ga19
	10067.0	15.			0.3	11			GSa		12Ho12
$^{288}\text{Ef}(\alpha)^{284}\text{Ed}$	10607.6	60.8	10750	50	2.9	B			DbA		04Og03
	10627.8	60.			2.5	B			DbA		12Og02
	10727.2	82.1			0.5	U			DbA		13Og01 *
	10698.8	10.1			1.1	B			GSa		13Ru11 *
	10754.6	20.3				12			BkA		15Ga24 *
* $^{288}\text{Fl}(\alpha)^{284}\text{Cn}$	F : T=1800(+2100-600) ms, later work yields shorter half-lives re-assigned to $^{289}\text{Fl}$										GAU **
* $^{288}\text{Fl}(\alpha)^{284}\text{Cn}$	F : non traceable information										GAU **
* $^{288}\text{Ef}(\alpha)^{284}\text{Ed}$	Highest $E_\alpha=10578(81)$ ; average 24 events would yield $E_\alpha=10480$ $Q_\alpha=10627.8$										13Og01 **
* $^{288}\text{Ef}(\alpha)^{284}\text{Ed}$	Average of 2 highest $E_\alpha=10560(10)$ $10540(10)$										13Ru11 **
* $^{288}\text{Ef}(\alpha)^{284}\text{Ed}$	Average of 2 highest $E_\alpha=10610(20)$ $10600(20)$										15Ga24 **
$^{289}\text{Fl}(\alpha)^{285}\text{Cn}$	9846.5	50.7	9970	50	2.4	F			DbA		99Og10 *
	9846.5	101.4			2.4	F			DbA		04Mo15 *
	9958.1	60.			0.2	o			DbA		04Og07
	9958.1	50.			0.2	7			DbA		07Og01
	10008.8	30.			-0.9	o			GSt		10Du06
	10008.8	30.			-0.9	7			GSt		11Ga19
	9956.0	15.2			0.2	7			GSa		12Ho12
	$^{289}\text{Fl}^m(\alpha)^{285}\text{Cn}^m$	10169.9	15.				7		GSa		12Ho12 *
$^{289}\text{Ef}(\alpha)^{285}\text{Ed}$	10455.0	90.	10510	50	1.1	o			DbA		11Og04
	10522.8	62.			-0.2	o			DbA		12Og02 *
	10404.2	71.			2.1	o			DbA		12Og06
	10546.2	81.			-0.7	o			DbA		13Og01
	10607.0	152.			-1.9	o			DbA		13Og04
	10510.7	20.3				12			DbA		16Fo16 *
* $^{289}\text{Fl}(\alpha)^{285}\text{Cn}$	F : one event at 30.4 s, later work yields much shorter half-lives										GAU **
* $^{289}\text{Fl}(\alpha)^{285}\text{Cn}$	F : non traceable information										GAU **
* $^{289}\text{Fl}^m(\alpha)^{285}\text{Cn}^m$	Assignment of $^{293}\text{Lv}^m_{-289}\text{Fl}^m_{-285}\text{Cn}^m_{-281}\text{Ds}^m_{-277}\text{Hs}^m$ chain is tentative										12Ho12 **
* $^{289}\text{Ef}(\alpha)^{285}\text{Ed}$	Also $E_\alpha=10310(90)$ keV										12Og02 **
* $^{289}\text{Ef}(\alpha)^{285}\text{Ed}$	reanalyzed data of 11Og04, 12Og02, 12Og06, 13Og01 and 13Og04										16Fo16 **
*	unweighted average of the 3 highest energy events selected by										GAU **
*	the evaluator: s06=10370 s12=10364 s14=10362 in Table 2										GAU **

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 030002-50)

Item	Input value		Adjusted value		$v_i$	Dg	Signf.	Main infl.	Lab	$F$	Reference
$^{290}\text{Ef}(\alpha)^{286}\text{Ed}$	10454.4	40.6							GSa		14Kh04
$^{290}\text{Lv}(\alpha)^{286}\text{Fl}$	10920.9	100.	11000	70	1.6	F			DbA		04Mo15 *
	11002.0	81.1			0.0	o			DbA		04Og07
	10991.8	81.1			0.1	o			DbA		06Og05
	10999.9	65.9				12			DbA		12Og06
$^{*290}\text{Lv}(\alpha)^{286}\text{Fl}$	F : non tracable information										GAu **
$^{291}\text{Lv}(\alpha)^{287}\text{Fl}$	10890	70	10890	50	0.0	o			DbA		04Og07
	10890	70				17			DbA		06Og05
$^{292}\text{Lv}(\alpha)^{288}\text{Fl}$	10707.0	50.	10774	15	1.3	F			DbA		01Og01 *
	10676.5	100.			1.0	F			DbA		04Mo15 *
	10808.3	71.0			-0.5	12			DbA		04Og12
	10772.8	15.2			0.1	12			GSa		12Ho12
$^{*292}\text{Lv}(\alpha)^{288}\text{Fl}$	F : daughter and grand-daughter re-assigned to $^{289}\text{Fl}$ and $^{285}\text{Cn}$										GAu **
$^{*292}\text{Lv}(\alpha)^{288}\text{Fl}$	F : non tracable information										GAu **
$^{293}\text{Lv}(\alpha)^{289}\text{Fl}$	10676	60	10680	50	0.1	o			DbA		04Og07
	10686.1	60.8			-0.1	8			DbA		07Og01
	10679.0	15.2			0.0	8			GSa		12Ho12
$^{293}\text{Lv}^m(\alpha)^{289}\text{Fl}^m$	10647.6	15.2				8			GSa		12Ho12 *
$^{293}\text{Eh}(\alpha)^{289}\text{Ef}$	11182.9	81.1	11290	50	2.2	o			DbA		11Og04 *
	10949.7	71.			6.9	B			DbA		12Og06 *
	11253.9	91.2			0.8	o			DbA		13Og04 *
	11293.4	20.3				13			DbA		16Fo16 *
$^{*293}\text{Lv}^m(\alpha)^{289}\text{Fl}^m$	Assignment of $^{293}\text{Lv}^m_{-289}\text{Fl}^m_{-285}\text{Cn}^m_{-281}\text{Ds}^m_{-277}\text{Hs}^m$ chain is tentative										12Ho12 **
$^{*293}\text{Eh}(\alpha)^{289}\text{Ef}$	reanalyzed data of 11Og04, 12Og02, 12Og06, 13Og01 and 13Og04										16Fo16 **
*	unweighted average of the 8 highest energy events selected by										GAu **
*	the evaluator: s02=11140 s03=11080 s07=11142 s08=11114 s12=11183										GAu **
*	s13=11203 s14=11059 s16=11190 in Table 2										GAu **
$^{294}\text{Eh}(\alpha)^{290}\text{Ef}$	11202.6	40.6					12		GSa		14Kh04
$^{294}\text{Ei}(\alpha)^{290}\text{Lv}$	11800.9	100.	11840	70	0.4	F			DbA		04Mo15 *
	11810.9	60.			0.5	o			DbA		04Og12
	11810.9	60.			0.5	o			DbA		06Og05
	11840.3	65.9				13			DbA		12Og06
$^{*294}\text{Ei}(\alpha)^{290}\text{Lv}$	F : non tracable information										GAu **
$^{295}\text{Ei}(\alpha)^{291}\text{Lv}$	11810.4	71.0	11700#	200#	-2.2	F			DbA		04Og05 *